

Homework 3

ASEN 5090 Introduction to GNSS

Justin Le

September 25, 2026

Problem 1: GPS Observations

(a.)

% Read rinex file

```
rinexData = rinexread('NIST00USA_R_20262310000_01D_30S_MO.rnx');
```

```
GPS_rinexData = rinexData.GPS;
```

```
PRN_number = 5;
```

```
PRN_index = GPS_rinexData.SatelliteID == PRN_number;
```

```
PRN05_GPS_rinexData = GPS_rinexData(PRN_index,:);
```

```
[NIST_GPS_week, NIST_GPS_TOW] = utc2gpstime(PRN05_GPS_rinexData.Time);
```

```
ephem_time_s = NIST_GPS_week.*604800 + NIST_GPS_TOW;
```

% Nan gap the discontinuous

```
gap_idx = [false; diff(ephem_time_s) > 1.5*median(diff(ephem_time_s))];
```

```
PRN05_GPS_rinexData.C1C(gap_idx) = NaN;
```

```
PRN05_GPS_rinexData.S1C(gap_idx) = NaN;
```

```
PRN05_GPS_rinexData.C1W(gap_idx) = NaN;
```

```
PRN05_GPS_rinexData.D1C(gap_idx) = NaN;
```

(b.)




Figure 1: C1C Pseudorange for PRN5

(c.)




Figure 2: S1C SNR for PRN5

(d.)




Figure 3: C1C and C1W Pseudorange for PRN5




Figure 4: C1C and C1W Residuals for PRN5

When examining the plots for C1C and C1W, we notice that they follow each other trends very well. Their reported pseudoranges deviate very minorly, when examining their residuals there are only differences of up to 0.7m between the two pseudoranges. This is reasonable as C1C and C1W are the C/A code and P code respectively. It is natural that they would be reporting similar numbers. The variations between them are a result from noise and multipathing, which would lead to the very minor variations that can be seen from Figure 4.

Problem 2: GPS Positions Based on Ephemeris Data

(a.)

```
% Use read clean to read in ephermides
clean_GPSbroadcast = read_clean_GPSbroadcast('brdc2310.26n', true);

% Read in sp3 ephem like HW2
sp3 = read_sp3('IGS00PSFIN_20262310000_01D_15M_ORB.SP3');
satPRN = extract_sp3_prn(sp3);

[health,satPos,satVel,satClkCorr] = eph2pvt2025(clean_GPSbroadcast, [satPRN(5).
    week_number satPRN(5).TOW_s], 5);
```

(b.)




Figure 5: Ephemeris vs Precise




Figure 6: Ephemeris vs Precise

Taking a look at these results, the precise positioning found from SP3 and the ephemeris broadcast in Figure 5, we see that they line up very well, by further using the data obtained by SP3 as truth, we find residuals of the ephemeris broadcast to be up to $\approx 1.5\text{m}$ from Figure 6. Compared to the results from HW2 on the almanac's residuals, which were up to 1 km, this is far more accurate. It shows that the almanac would be useful for checking visibility of satellites in the sky, however poor for positioning. The resulting residuals we observe here can begin to be used for positioning.

(c.)




Figure 7: Clock Bias

Problem 3: GPS Expected Range

(a.)




Figure 8: PRN5 Azimuth




Figure 9: PRN5 Elevation




Figure 10: PRN5 Azimuth




Figure 11: PRN5 Range

(b.)




Figure 12: PRN5 Range Initial vs Expected




Figure 13: PRN5 Range Initial vs Expected Residual

From here, we can find the largest difference from the expected range and the initial range we obtain from ephemeris to be about 78.9m. This is a rather large deviation between the ranges, showing how there is still correction needed from the broadcasted ephemeris received from a ground station/gps receiver.

Problem 4: Observed Minus Computed (Expected) Range




Figure 14: PRN5 C1C Pseudorange vs Expected Range




Figure 15: PRN5 C1C Pseudorange vs Expected Range Residuals

We are able to observe the differences between the reported pseudoranges and the expected range calculated previously in Figure 14 and Figure 15. They generally match up, however there are residuals of up to 85 km. The biggest differences between them are likely a result of the correction being done to find the expected range. The ephemeris broadcast, conversion to position, and subsequent correction we performed would be much more accurate than the pseudorange that is being reported. The pseudorange after all is affected by propagation effects in the atmosphere (tropospheric effects, ionospheric effects) which cause these errors observed between the two ranging measurements being reported.

TODO: provide justification.

Problem 5: Repeat for another PRN




Figure 16: PRN11 C1C Pseudorange vs Expected Range




Figure 17: PRN11 C1C Pseudorange vs Expected Range Residuals

Here PRN 11 is chosen, this is in a different GPS block than PRN 5 with notable different visibility patterns in the same time frame for NIST.