# Homework 4

ASEN 5335 Aerospace Environment

Justin Le
October 22, 2026

## Problem 1: Field Magnitudes

Using the WMM model from the NOAA calculator, we find the magnetic field in Boulder, CO to be,

| Model Used: | WMM-2025 | | | | | | |
|---|---|---|---|---|---|---|---|
| Latitude: | 40° 0' 59" N | | | | | | |
| Longitude: | 105° 17' 10" W | | | | | | |
| Elevation: | 5430.0 ft Mean Sea Level | | | | | | |
| **Date** | **Declination ( + E \| - W )** | **Inclination ( + D \| - U )** | **Horizontal Intensity** | **North Comp (+ N \| - S)** | **East Comp (+ E \| - W)** | **Vertical Comp (+ D \| - U)** | **Total Field** |
| 2026-10-18 | 7° 40' 8" | 66° 0' 55" | 20,760.4 nT | 20,574.7 nT | 2,770.4 nT | 46,662.2 nT | 51,072.1 nT |
| **Change/year** | -0° 4' 50"/yr | -0° 2' 59"/yr | -8.9 nT/yr | -5.0 nT/yr | -30.2 nT/yr | -129.4 nT/yr | -121.9 nT/yr |
| **Uncertainty** | 0° 22' | 0° 12' | 133 nT | 137 nT | 89 nT | 141 nT | 138 nT |

Figure 1: Boulder Magnetic Field

We repeat with the same model for McMurdo Station Anarctica to find,

| Model Used: | WMM-2025 | | | | | | |
|---|---|---|---|---|---|---|---|
| Latitude: | 77° 50' 46" S | | | | | | |
| Longitude: | 166° 40' 5" E | | | | | | |
| Elevation: | 33.0 ft Mean Sea Level | | | | | | |
| **Date** | **Declination ( + E \| - W )** | **Inclination ( + D \| - U )** | **Horizontal Intensity** | **North Comp (+ N \| - S)** | **East Comp (+ E \| - W)** | **Vertical Comp (+ D \| - U)** | **Total Field** |
| 2026-10-18 | 140° 5' 50" | -80° 5' 50" | 10,647.5 nT | -8,168.0 nT | 6,830.2 nT | -60,990.6 nT | 61,913.0 nT |
| **Change/year** | -0° 12' 10"/yr | 0° 2' 29"/yr | 33.8 nT/yr | -1.7 nT/yr | 50.6 nT/yr | 66.8 nT/yr | -60.0 nT/yr |
| **Uncertainty** | 0° 34' | 0° 12' | 133 nT | 137 nT | 89 nT | 141 nT | 138 nT |

Figure 2: McMurdo Magnetic Field

Based on the inclination observed at McMurdo, this tells us that the magnetic fields will pass through the station based on the inclination and the latitude of the station.

# Problem 2: Dipole Field

## (i).

Assuming a dipole field model with $L = 4$ $B_0 = 31,000$ nT, $R_E = 6371$ km, and at the equator ($\lambda = 0°$), we can use the following equations to calculate the magnetic field magnitude,

$$r = LR_E \cos^2 \lambda$$
$$|B| = B_0(\frac{R_E}{r})^3\sqrt{1 + 3\sin^2 \lambda}$$

$$\therefore r = 25,484,000$$
$$\boxed{\therefore |B| = 484.4 \text{ nT}}$$

## (ii).

Repeating the above equations only for $\lambda = 30°$,

$$\therefore r = 19,113,000$$
$$\boxed{\therefore |B| = 1518.9 \text{ nT}}$$

# Problem 3: Magnetopause

## (i).

To find the standoff distance of the Earth to the magnetosphere, we can use the equation,

$$\rho_{sw} v_{sw}^2 = \frac{(2B_{mag})^2}{2\mu_0} = \frac{4B_0^2}{2\mu_0 L^6}$$
$$B_{mag} = B_0 \left(\frac{R_E}{r}\right)^3 = \frac{B_0}{L^3}, \qquad B_0 \approx 31{,}000 \text{ nT}$$
$$L = \left(\frac{2B_0^2}{\mu_0 \rho_{sw} v_{sw}^2}\right)^{1/6}$$

Given $v_{sw} = 400$ km/s, and $\mu_0 = 4\pi \times 10^{-7}$, and $\rho_{sw} = n m_p = 5$ proton/cm$^3$ $*1.672 \times 10^{-27}$ kg $= 8.36 \times 10^{-21}$ kg/m$^3$.

$$\boxed{L = 10.2}$$

## (ii).

We can repeat the calculation above but instead using $v = 1000$ km/s, and $n = 10$ proton/cm$^3$, for a new $\rho_{sw} = 1.67 \times 10^{-20}$ kg/m$^3$.

$$\boxed{L = 6.7}$$

# Problem 4: The Ring Current

## (i).

The reason why there is a current in the first place is because of protons drifting west, while electrons drift east. This causes a net westward current which is the ring current. Additionally, drift speed scales with energy, this leads to the current being stronger where hotter particles (resulting in more energy) sitting at a higher altitude in the atmosphere at about 3-6 $R_E$.

## (ii).

Finding how long it takes for an 85 keV electron to drift all the around Earth at $L = 4$, we can use the equations,

$$\mathbf{v}_{d\nabla} = \frac{mv_\perp^2}{2q}\frac{\mathbf{B}\times\nabla B}{B^3}$$
$$E = \frac{1}{2}mv_\perp^2$$

If we assume the particle remains in the equator and all its energy is perpendicular ($v_\parallel = 0$). At the equator $|\mathbf{B}\times\nabla B| = B|\nabla B|$. The drift speed is then,

$$v_d = \frac{2E}{2q}\frac{B|\nabla B|}{B^3} = \frac{E}{qB}\frac{|\nabla B|}{B}$$

To find the gradient we use the dipole field at the equator ($\lambda = 0°$) from Problem 2 and take the derivative with respect to $r$,

$$B = B_0\left(\frac{R_E}{r}\right)^3$$
$$\frac{dB}{dr} = -3B_0\frac{R_E^3}{r^4} = -\frac{3B}{r}$$
$$\therefore |\nabla B| = \frac{3B}{r}$$

Substituting the gradient back into the drift speed,

$$v_d = \frac{E}{qB}\frac{3B}{rB} = \frac{3E}{qBr}, \qquad B = \frac{B_0}{L^3}, \quad r = LR_E$$

The time to drift all the way around the Earth is the circumference of the drift path divided by the drift speed,

$$T = \frac{2\pi r}{v_d} = \frac{2\pi qBr^2}{3E} = \frac{2\pi qB_0R_E^2}{3EL}$$

Given $L = 4$, $B_0 = 31,000$ nT, $R_E = 6371$ km, and $E = 85$ keV, we have $B = 484.4$ nT (from Problem 2) and $r = 25,484,000$ m.

$$v_d = \frac{3(85,000)}{(484.4 \times 10^{-9})(25,484,000)} = 20,660 \text{ m/s}$$

$$T = \frac{2\pi(25,484,000)}{20,660} = 7,750 \text{ s}$$

$$\boxed{\therefore T \approx 2.15 \text{ hours}}$$

# Problem 5: Drift due to Gravity

## (i.)

Under the influence of gravity, protons will drift east and electrons will drift west.

## (ii.)

We find the magnitude of the drift velocity due to gravity is with the equation,

$$\mathbf{v}_{dg} = \frac{m\mathbf{g} \times \mathbf{B}}{qB^2}$$

## (iii.)

To find how long it will take to drift around the Earth under gravity, we can use the drift to to gravity equation and assuming a particle drifting around the equator,

$$\mathbf{v}_{dg} = \frac{m\mathbf{g} \times \mathbf{B}}{qB^2} \quad , \quad v_{dg} = \frac{mg}{qB}$$

$$g = \frac{g_{Earth}}{L^2} = \frac{9.81}{4^2} = 0.613 \text{ m/s}^2$$

$$T = \frac{2\pi r}{v_{dg}} = \frac{2\pi qBr}{mg}$$

$$v_{dg} = \frac{(1.673 \times 10^{-27})(0.613)}{(1.602 \times 10^{-19})(484.4 \times 10^{-9})} = 1.32 \times 10^{-2} \text{ m/s}$$

$$T = \frac{2\pi(25,484,000)}{1.32 \times 10^{-2}} = 1.21 \times 10^{10} \text{ s}$$

$$\boxed{\therefore T \approx 384 \text{ years}}$$

Here we find that the drift to to gravity is far slower than the gradient drift from Problem 4, and that when calculating ring current this can be assumed to be a negligible effect.

# Problem 6: Geomagnetic Indices





Figure 3: Distribution Function

## (i.)

Using Figure 3, we can identify the most common Kp in this distribution as $Kp = 1$, as the most frequent occurrence in this distribution is at 10.

## (ii.)

Examining what values will be at level 9, we find the ratio of the Kp = 90 out of the total measurements.

$$\boxed{\frac{39}{464{,}616} = 0.0084\%}$$

## (iii.)

When repeating the above for July 2004, we find,





Figure 4: Distribution Function for July 2004

Here we find that the most common Kp is 11, so most common Kp is $Kp = 1.1$, and of there are no occurrences of Kp level 9,

# Problem 7: Geomagnetic Storm

## (i.)

We can recreate the Halloween Storm and obtain the following,




Figure 5: DST Index for the Halloween Storm

## (ii.)

Using this information, we can calculate the ring current magnitude with,

$$B = \frac{\mu_0 I}{2R}, \; I = \frac{2RB}{\mu_0}$$

$$\mu_0 = 4\pi \times 10^{-7} \text{ Tm/A}$$

$$B = 400 \times 10^{-9} \text{ T}$$

$$R = 5R_E = 5(6371) \times 10^3 \text{ m}$$

$$\boxed{\therefore I = 20 \times 10^6 \text{ A}}$$

# Problem 8: Key Parameters in the Solar Wind Driving a Magnetic Storm

## (i.)

The most critical parameter for describing solar wind is the interplanetary magnetic field. This is because for a geomagnetic storm to occur, the magnetic field field must point southward and also be sustained for an amount of time while also being stronger than usual, typically a result from coronal mass ejections. The interplanetary magnetic field essentially determines the conditions needed for a magnetic storm.

## (ii.)

The positive surge that results from the magnetopause current of the Chapman Ferraro current. This is because when a CME hits Earth, the higher solar wind pressure compresses the magnetosphere inward, which is the magnetic storm's initial phase. The current forms because the solar wind gyrating around the magnetopause field lines. This cancels out the Earth's filed outside of the magnetosphere and increases the field inside of it, which leads to a eastward current as opposed to the usual westward current, causing a northward field and shows the surge in positive DST.

# Problem 9: The Core

## (i.)

Aaron Eckhart's description of the Earth's core. His description is indeed accurate with a sold inner core and a swirling outer core. Physics 101, hot metal moving fast drives EM field, the hot metal swirling does indeed drive the EM field present on Earth, this can be seen in how its turbulent nature has previously led to the shifting and inverting. EM field does shield us from cosmic radiation, due to physics the EM field repels the cosmic radiation from Earth.

## (ii.)

Microwave radiation will literally cook our planet, solar wind is not microwaves, and most of the radiation that would impact us would be absorbed by the atmosphere. Everybody on Earth is dead in a year, this is just a ridiculous timeline, the weakened field would not led to extinction. Static discharges in the atmosphere will lead to superstorms, the EM field would have no impact on the weather effects seen in the lower parts of the atmosphere and the surface of Earth.

# Problem 10: Magnetospheric Multiscale

## (i.)

The primary objective of the MSS mission is to study the reconnection of magnetic field lines in Earth's magnetosphere. This is how the solar wind's magnetic field connects to Earth's and releases stored energy. It aims to look at small scale physics looking at how particles get accelerated and its turbulence.

## (ii.)

The reason why there were four spacecraft is because that allows for measuring data in 3D. They are arranged in a tetrahedron, where by taking 4 points of measurements not on the same plane, it allows measurements of 3 dimensions over time.

## (iii.)

All four spacecraft, carry the same instruments. The Hot Plasma Suite, which carry electron spectrometers (4 dual electron spectrometers, 4 dual ion spectrometers, and 1 HPCA), Energy Particle Detectors (2 FEEPS and 1 EIS), the FIELDS suite (4 spin-plane double probes, 2 axial double probes, 2 fluxgate magnetometers, 1 search coil magnetometer, and 1 EDI), which was discussed in a previous homework, 2 ASPOC, which is the active spacecraft potential control to prevent the charge potential from building on the spacecraft. There are multiple instruments on each spacecraft because the spacecrafts rotate around its axis as spin stabilization keeps the spacecraft attitude steady without reaction wheels, resulting in sensors sweeping across the sky each rotation.

## (iv.)

There are two phases in the mission, dayside and the magnetotail. They exist because there are two different conditions, in dayside the solar wind field lines will reconnect with Earth, at the magnetotail field lines stretched by solar storm reconnects. In dayside phase, the perigee would be $\approx 1.2R_E$, the apogee is $\approx 12R_E$. Meanwhile in the magnetotail phase the perigee is $\approx 1.2R_E$ but then the apogee is $\approx 25R_E$