

Problem 2. $\forall i((0 \leq i < N) \rightarrow (p_i \wedge q_i)) \rightarrow \forall i((0 \leq i \leq 2N) \rightarrow (x_0 \leq x_i))$ holds for every $N \in \mathbb{N}$

Proof: We prove this statement using induction on N .

Basis: For the basis of induction, we establish that $\forall i((0 \leq i < 0) \rightarrow (p_i \wedge q_i)) \rightarrow \forall i((0 \leq i \leq 0) \rightarrow (x_0 \leq x_i))$. The antecedent of this conditional is vacuously true, since $0 \leq i < 0$ is false for any $i \in \mathbb{N}$. Therefore, it will suffice to prove that the consequent, $\forall i((0 \leq i \leq 0) \rightarrow (x_0 \leq x_i))$, is true. Suppose that $0 \leq m \leq 0$. Then $m = 0$. Consequently, $x_0 \leq x_m$ holds trivially, because $x_0 \leq x_0$. Thus, the theorem holds when $N = 0$.

IH: For the induction hypothesis, suppose that $\forall i((0 \leq i < N) \rightarrow (p_i \wedge q_i)) \rightarrow \forall i((0 \leq i \leq 2N) \rightarrow (x_0 \leq x_i))$.

IS: Suppose that $\forall i((0 \leq i < N + 1) \rightarrow p_i \wedge q_i)$. Furthermore, suppose that $0 \leq m \leq 2(N + 1)$. Either $m < N + 1$ or $m \geq N + 1$. We consider each case in detail:

i) If $m < N + 1$, then, since $\forall i((0 \leq i < N + 1) \rightarrow p_i \wedge q_i)$ holds by assumption, we must have $\forall i((0 \leq i < N) \rightarrow p_i \wedge q_i)$ trivially. Since $\forall i((0 \leq i < N) \rightarrow p_i \wedge q_i)$, $\forall i((0 \leq i \leq 2N) \rightarrow (x_0 \leq x_i))$ holds by the *IH*. Since $0 \leq m < N + 1$, we must have $0 \leq m \leq 2N$. Therefore, because $0 \leq m \leq 2N$ and $\forall i((0 \leq i \leq 2N) \rightarrow (x_0 \leq x_i))$, we have $x_0 \leq x_m$.

ii) If $m \geq N + 1$, then it is easy to check that either $m = 2i + 1$ or $m = 2i + 2$ for some $i < N + 1$. Since $i < N + 1$, $p_i \wedge q_i$ by our initial supposition. If $m = 2i + 1$, then, since p_i holds, $x_{2i} \leq x_m$. If $m = 2i + 2$, then, since q_i holds, $x_{2i} \leq x_m$ as well. Thus, in either of our two cases, $x_{2i} \leq x_m$. Since $i < N + 1$, $2i \leq 2N$ trivially, which means that $x_0 \leq x_{2i}$ by the *IH*. Since $x_0 \leq x_{2i}$ and $x_{2i} \leq x_m$, we must have $x_0 \leq x_m$.

In both of the two cases, (i) and (ii), we see that $x_0 \leq x_m$, which means that $\forall i((0 \leq i \leq 2(N + 1)) \rightarrow (x_0 \leq x_i))$. \square

Problem 3. $\forall i((i \geq 0) \rightarrow (p_i \wedge q_i))$ is inductive.

Proof: We have already shown that $\forall i((i \geq 0) \rightarrow p_i)$ is inductive, so we must prove the same thing for the predicate q .

Suppose that in a certain state, we satisfy $\forall i((i \geq 0) \rightarrow (p_i \wedge q_i))$. Suppose that $i \geq 0$. Then, q_i holds by supposition, which means that $x_{2i} \leq x_{2i+2}$. Now, suppose that we transition to a new state. We want to show that in this new state, $x'_{2i} \leq x'_{2i+2}$. This transition was either a push or a pop. We consider each case:

i) If the transition was a push, $x'_{2i} = \min(x_{2i}, x_{2i-1})$ and $x'_{2i+2} = x'_{2(i+1)} = \min(x_{2i+2}, x_{2i+1})$. So, we want to show that $\min(x_{2i}, x_{2i-1}) \leq \min(x_{2i+2}, x_{2i+1})$. This gives us four more sub-cases:

a) First of all, $x_{2i} \leq x_{2i+2}$ by the fact that q_i holds. Therefore, if $\min(x_{2i}, x_{2i-1}) = x_{2i}$ and $\min(x_{2i+2}, x_{2i+1}) = x_{2i+2}$, then $\min(x_{2i}, x_{2i-1}) \leq \min(x_{2i+2}, x_{2i+1})$.

b) Now, if $\min(x_{2i}, x_{2i-1}) = x_{2i-1}$ and $\min(x_{2i+2}, x_{2i+1}) = x_{2i+2}$, then $x_{2i-1} \leq x_{2i}$ trivially. We have already seen from case (a) that $x_{2i} \leq x_{2i+2}$. Since $x_{2i} \leq x_{2i+2}$ and $x_{2i-1} \leq x_{2i}$, we must have $x_{2i-1} \leq x_{2i+2}$,

which means $\min(x_{2i}, x_{2i-1}) \leq \min(x_{2i+2}, x_{2i+1})$.

c) Since p is an inductive property, we must have $x_{2i} \leq x_{2i+1}$. Therefore, if $\min(x_{2i}, x_{2i-1}) = x_{2i}$ and $\min(x_{2i+2}, x_{2i+1}) = x_{2i+1}$, then $\min(x_{2i}, x_{2i-1}) \leq \min(x_{2i+2}, x_{2i+1})$.

d) Finally, $x_{2i-1} \leq x_{2i+1}$, since q holds by supposition. Therefore, if $\min(x_{2i}, x_{2i-1}) = x_{2i-1}$ and $\min(x_{2i+2}, x_{2i+1}) = x_{2i+1}$, then $\min(x_{2i}, x_{2i-1}) \leq \min(x_{2i+2}, x_{2i+1})$.

In all four cases, we showed that $x'_{2i} \leq x'_{2i+2}$, which means that q_i is an inductive property across all push operations.

ii) If the transition was a pop, then we want to show that $\min(x_{2i+1}, x_{2i+2}) \leq \min(x_{2i+3}, x_{2i+4})$. Again, we can split this up into four sub-cases:

a) First of all, $x_{2i+2} \leq x_{2i+3}$ since p_i holds by assumption, so if $\min(x_{2i+1}, x_{2i+2}) = x_{2i+2}$ and $\min(x_{2i+3}, x_{2i+4}) = x_{2i+3}$, then $\min(x_{2i+1}, x_{2i+2}) \leq \min(x_{2i+3}, x_{2i+4})$.

b) Also, $x_{2i+2} \leq x_{2i+4}$ since q_i holds by assumption, so if $\min(x_{2i+1}, x_{2i+2}) = x_{2i+2}$ and $\min(x_{2i+3}, x_{2i+4}) = x_{2i+4}$, then $\min(x_{2i+1}, x_{2i+2}) \leq \min(x_{2i+3}, x_{2i+4})$.

c) If $\min(x_{2i+1}, x_{2i+2}) = x_{2i+1}$ and $\min(x_{2i+3}, x_{2i+4}) = x_{2i+4}$, then $x_{2i+1} \leq x_{2i+2}$. We already showed in case (b) that $x_{2i+2} \leq x_{2i+4}$, so this means that $x_{2i+1} \leq x_{2i+2}$. Thus, $\min(x_{2i+1}, x_{2i+2}) \leq \min(x_{2i+3}, x_{2i+4})$.


d) Finally, if $\min(x_{2i+1}, x_{2i+2}) = x_{2i+1}$ and $\min(x_{2i+3}, x_{2i+4}) = x_{2i+3}$, then again we have $x_{2i+1} \leq x_{2i+2}$. In case (a) we showed $x_{2i+2} \leq x_{2i+3}$, so $x_{2i+1} \leq x_{2i+3}$. Therefore, $\min(x_{2i+1}, x_{2i+2}) \leq \min(x_{2i+3}, x_{2i+4})$ in this case as well.

Whether we push or pop, the property q is preserved. Since $p_i \wedge q_i$ holds at the initial state and $p_i \wedge q_i$ holds at every state we can transition into from a state satisfying $p_i \wedge q_i$, we conclude that the property is inductive. \square

Problem 4. First, we expand the formula $\varphi = pRFq$ into disjunctive normal form:

$$\begin{aligned}
 pRFq &\iff Fq \wedge (p \vee X(pRFq)) \\
 Fq \wedge (p \vee X(pRFq)) &\iff (q \vee X(Fq)) \wedge (p \vee X(pRFq)) \\
 (q \vee X(Fq)) \wedge (p \vee X(pRFq)) &\iff (p \wedge q) \vee (p \wedge X(Fq)) \vee (q \wedge X(pRFq)) \vee (X(Fq) \wedge X(pRFq)) \\
 &\iff (p \wedge q \wedge X(true)) \vee (p \wedge X(Fq)) \vee (q \wedge X(pRFq)) \vee (true \wedge X(Fq \wedge pRFq))
 \end{aligned}$$

Now, we can create the corresponding Buchi automata:



Problem 5.