

PREOPERATIVE DIAGNOSES,1. Metastatic carcinoma of the bladder.,2. Bowel obstruction.,POSTOPERATIVE

DIAGNOSES,1. Metastatic carcinoma of the bladder.,2.

Bowel obstruction.,PROCEDURE: , Port insertion through the right subclavian vein percutaneously under radiological

guidance.,PROCEDURE DETAIL: ,The patient was electively taken to the operating room after obtaining an informed

consent. A time-out process was followed. Antibiotics were

given. Then, the patient's right deltopectoral area was

prepped and draped in the usual fashion. Xylocaine 1% was

infiltrated. The right subclavian vein was percutaneously

cannulated without any difficulty. Then using the Seldinger technique, the catheter part of the port, which was a

single-lumen port, was passed through the introducer under x-ray guidance and placed in the junction of the superior vena

cava and the right atrium.,A pocket had been fashioned and a

single-lumen drum of the port was connected to the catheter,

which had been trimmed and affixed to the pectoralis fascia

with couple of sutures of Vicryl. Then, the fascia was closed

using subcuticular suture of Monocryl. The drum was

aspirated and irrigated with heparinized saline and then was

put in the pocket and the skin was closed. A dressing was

applied including the needle and the port with the catheter so

that the floor could use the catheter right away.,The patient

tolerated the procedure well and was sent to recovery room in

satisfactory condition. A chest x-ray was performed that

showed that there were no complications of procedure and

that the catheter was in right place.