

PROCEDURE:, Colonoscopy.,PREOPERATIVE DIAGNOSIS:
, Follow up adenomas.,POSTOPERATIVE DIAGNOSES:,1.
Two colon polyps, removed.,2. Small internal hemorrhoids.,3.
Otherwise normal examination of cecum.,MEDICATIONS: ,
Fentanyl 150 mcg and Versed 7 mg slow IV
push.,INDICATIONS: , This is a 60-year-old white female with
a history of adenomas. She does have irregular bowel
habits.,FINDINGS: , The patient was placed in the left lateral
decubitus position and the above medications were
administered. The colonoscope was advanced to the cecum
as identified by the ileocecal valve, appendiceal orifice, and
blind pouch. The colonoscope was slowly withdrawn and a
careful examination of the colonic mucosa was made,
including a retroflexed view of the rectum. There was a 4 mm
descending colon polyp, which was removed with jumbo
forceps, and sent for histology in bottle one. There was a 10
mm pale, flat polyp in the distal rectum, which was removed
with jumbo forceps, and sent for histology in bottle 2. There
were small internal hemorrhoids. The remainder of the
examination was normal to the cecum. The patient tolerated
the procedure well without complication.,IMPRESSION:,1.
Two colon polyps, removed.,2. Small internal hemorrhoids.,3.
Otherwise normal examination to cecum.,PLAN: , I will await
the results of the colon polyp histology. The patient was told
the importance of daily fiber.