

Contest Duration: 2025-10-25(Sat) 23:00 (<http://www.timeanddate.com/worldclock/fixedtime.html?iso=20251025T2100&p1=248>) - 2025-10-26(Sun) 00:40 (<http://www.timeanddate.com/worldclock/fixedtime.html?iso=20251025T2240&p1=248>) (local time) (100 minutes)

[Back to Home \(/home\)](/home)

[🏠 Top \(/contests/abc429\)](/contests/abc429)

[📋 Tasks \(/contests/abc429/tasks\)](/contests/abc429/tasks)

[❓ Clarifications \(/contests/abc429/clarifications\)](/contests/abc429/clarifications)

[📊 Results ▼](#)

[🏆 Standings \(/contests/abc429/standings\)](/contests/abc429/standings)

[🏆 Virtual Standings \(/contests/abc429/standings/virtual\)](/contests/abc429/standings/virtual)

[📖 Editorial \(/contests/abc429/editorial\)](/contests/abc429/editorial)

[💬 Discuss \(https://codeforces.com/blog/entry/147752\)](https://codeforces.com/blog/entry/147752)



C - Odd One Subsequence

[Editorial \(/contests/abc429/tasks/abc429_c/editorial\)](/contests/abc429/tasks/abc429_c/editorial)



Time Limit: 2 sec / Memory Limit: 1024 MiB

Score : 300 points

Problem Statement

You are given an integer sequence of length N , $A = (A_1, A_2, \dots, A_N)$.

Find the number of triples of integers (i, j, k) satisfying $1 \leq i < j < k \leq N$ that satisfy the following condition:

Exactly two distinct values are contained in A_i, A_j, A_k . That is, two of A_i, A_j, A_k are equal, and the remaining one is different.

Constraints

- $3 \leq N \leq 2 \times 10^5$
- $1 \leq A_i \leq N$
- All input values are integers.

Input

The input is given from Standard Input in the following format:

2026-01-02 (Fri)
05:32:06 +11:00

N $A_1 \ A_2 \ \dots \ A_N$

Output

Print the number of triples of integers that satisfy the condition.

Sample Input 1

[Copy](#)

```
5
3 2 5 2 2
```

[Copy](#)

Sample Output 1

[Copy](#)

```
6
```

[Copy](#)

For example, $(i, j, k) = (1, 2, 4)$ satisfies the condition because exactly two distinct values, 2 and 3, are contained in $A_1 = 3$, $A_2 = 2$, and $A_4 = 2$.

Including this, the six triples $(i, j, k) =$

$(1, 2, 4), (1, 2, 5), (1, 4, 5), (2, 3, 4), (2, 3, 5), (3, 4, 5)$ satisfy the condition.

Therefore, print 6.

Sample Input 2

[Copy](#)

```
3
1 1 1
```

[Copy](#)

Sample Output 2

[Copy](#)

```
0
```

[Copy](#)

There may be no triples that satisfy the condition.

[/#telegram](#))

[url=https%3A%2F%2Fatcoder.jp%2Fcontests%2Fabc429%2Ftasks%2Fabc429_c%3Flang%3Den&title=C%20-](https://atcoder.jp/contests/abc429/tasks/abc429_c3)
:)

[Terms of service \(/tos\)](#) [Privacy Policy \(/privacy\)](#) [Information Protection Policy \(/personal\)](#) [Company \(/company\)](#)
[FAQ \(/faq\)](#) [Contact \(/contact\)](#)

Copyright Since 2012 ©AtCoder Inc. (<http://atcoder.co.jp>) All rights reserved.