

Contest Duration: 2025-08-09(Sat) 22:00 (<http://www.timeanddate.com/worldclock/fixedtime.html?iso=20250809T2100&p1=248>) - 2025-08-09(Sat) 23:40 (<http://www.timeanddate.com/worldclock/fixedtime.html?iso=20250809T2240&p1=248>) (local time) (100 minutes)

[Back to Home \(/home\)](/home)

[🏠 Top \(/contests/abc418\)](/contests/abc418)

[📋 Tasks \(/contests/abc418/tasks\)](/contests/abc418/tasks)

[❓ Clarifications \(/contests/abc418/clarifications\)](/contests/abc418/clarifications)

[📊 Results ▼](#)

[🏆 Standings \(/contests/abc418/standings\)](/contests/abc418/standings)

[🏆 Virtual Standings \(/contests/abc418/standings/virtual\)](/contests/abc418/standings/virtual)

[📖 Editorial \(/contests/abc418/editorial\)](/contests/abc418/editorial)

[💬 Discuss \(https://codeforces.com/blog/entry/145383\)](https://codeforces.com/blog/entry/145383)



F - We're teapots

[Editorial \(/contests/abc418/tasks/abc418_f/editorial\)](/contests/abc418/tasks/abc418_f/editorial)



Time Limit: 3 sec / Memory Limit: 1024 MiB

Score : 550 points

Problem Statement

There are N teapots arranged in a row, numbered from 1 to N from left to right.

There is a sequence of integers (a_1, \dots, a_N) , initially with values $a_1 = \dots = a_N = -1$.

You will fill each teapot with either tea or coffee so that the following conditions are all satisfied:

- For any two adjacent teapots, at least one of them contains tea.
- For any integer i satisfying $1 \leq i \leq N$, if $a_i \neq -1$, then exactly a_i of teapots $1, \dots, i$ contain coffee.

You are given Q queries, which you should process in the given order.

The j -th query ($1 \leq j \leq Q$) is as follows:

- Change the value of a_{X_j} to Y_j . Then, print the number, modulo 998244353, of ways to fill the teapots satisfying the conditions.

Constraints

- $2 \leq N \leq 2 \times 10^5$
- $1 \leq Q \leq 2 \times 10^5$
- $1 \leq X_j \leq N$ ($1 \leq j \leq Q$)
- $-1 \leq Y_j \leq X_j$ ($1 \leq j \leq Q$)

2026-01-02 (Fri)
05:28:26 +11:00

- All input values are integers.

Input

The input is given from Standard Input in the following format:

```
N Q
X1 Y1
⋮
XQ YQ
```

Output

Print Q lines.

The j -th line ($1 \leq j \leq Q$) should contain the value to be printed for the j -th query.

Sample Input 1

[Copy](#)

```
5 6
1 1
4 2
1 0
4 -1
5 1
5 5
```

[Copy](#)

Sample Output 1

[Copy](#)

```
5
3
1
8
4
0
```

[Copy](#)

- After the operation in the first query, $a = (1, -1, -1, -1, -1)$. The ways to fill the teapots satisfying the conditions are the following five ways:
 - Put coffee in teapot 1, and tea in the others.
 - Put coffee in teapots 1, 3, and tea in the others.
 - Put coffee in teapots 1, 3, 5, and tea in the others.
 - Put coffee in teapots 1, 4, and tea in the others.
 - Put coffee in teapots 1, 5, and tea in the others.

2026-01-02 (Fri)
05:28:26 +11:00

- After the operation in the second query, $a = (1, -1, -1, 2, -1)$. The ways to fill the teapots satisfying the conditions are the following three ways:
 - Put coffee in teapots 1, 3, and tea in the others.
 - Put coffee in teapots 1, 3, 5, and tea in the others.
 - Put coffee in teapots 1, 4, and tea in the others.
- After the operation in the third query, $a = (0, -1, -1, 2, -1)$. The ways to fill the teapots satisfying the conditions are the following one way:
 - Put coffee in teapots 2, 4, and tea in the others.
- After the operation in the fourth query, $a = (0, -1, -1, -1, -1)$. The ways to fill the teapots satisfying the conditions are the following eight ways:
 - Put coffee in none of the teapots and tea in all of them.
 - Put coffee in teapot 2, and tea in the others.
 - Put coffee in teapots 2, 4, and tea in the others.
 - Put coffee in teapots 2, 5, and tea in the others.
 - Put coffee in teapot 3, and tea in the others.
 - Put coffee in teapots 3, 5, and tea in the others.
 - Put coffee in teapot 4, and tea in the others.
 - Put coffee in teapot 5, and tea in the others.
- After the operation in the fifth query, $a = (0, -1, -1, -1, 1)$. The ways to fill the teapots satisfying the conditions are the following four ways:
 - Put coffee in teapot 2, and tea in the others.
 - Put coffee in teapot 3, and tea in the others.
 - Put coffee in teapot 4, and tea in the others.
 - Put coffee in teapot 5, and tea in the others.
- After the operation in the sixth query, $a = (0, -1, -1, -1, 5)$. The number of ways to fill the teapots satisfying the conditions is zero.

/#telegram)

#url=https%3A%2F%2Fatcoder.jp%2Fcontests%2Fabc418%2Ftasks%2Fabc418_f%3Flang%3Den&title=F%20-

[Rule \(/contests/abc418/rules\)](/contests/abc418/rules) [Glossary \(/contests/abc418/glossary\)](/contests/abc418/glossary)

[Terms of service \(/tos\)](/tos) [Privacy Policy \(/privacy\)](/privacy) [Information Protection Policy \(/personal\)](/personal) [Company \(/company\)](/company)
[FAQ \(/faq\)](/faq) [Contact \(/contact\)](/contact)

Copyright Since 2012 ©AtCoder Inc. (<http://atcoder.co.jp>) All rights reserved.

2026-01-02 (Fri)
 05:28:26 +11:00