

Contest Duration: 2025-07-05(Sat) 22:00 (<http://www.timeanddate.com/worldclock/fixedtime.html?iso=20250705T2100&p1=248>) - 2025-07-05(Sat) 23:40 (<http://www.timeanddate.com/worldclock/fixedtime.html?iso=20250705T2240&p1=248>) (local time) (100 minutes)

[Back to Home \(/home\)](#)

[🏠 Top \(/contests/abc413\)](#)

[📋 Tasks \(/contests/abc413/tasks\)](#)

[❓ Clarifications \(/contests/abc413/clarifications\)](#)

[📊 Results ▼](#)

[🏆 Standings \(/contests/abc413/standings\)](#)

[🏆 Virtual Standings \(/contests/abc413/standings/virtual\)](#)

[📖 Editorial \(/contests/abc413/editorial\)](#)

[💬 Discuss \(https://codeforces.com/blog/entry/144450\)](https://codeforces.com/blog/entry/144450)



D - Make Geometric Sequence

[Editorial \(/contests/abc413/tasks/abc413_d/editorial\)](#)



Time Limit: 2 sec / Memory Limit: 1024 MiB

Score : 425 points

Problem Statement

You are given an integer sequence $A = (A_1, A_2, \dots, A_N)$ of length N . It is guaranteed that for any i ($1 \leq i \leq N$), A_i is not 0.

Determine whether there exists a permutation $B = (B_1, B_2, \dots, B_N)$ of A such that B forms a geometric sequence.

A sequence $S = (S_1, S_2, \dots, S_N)$ is a geometric sequence if there exists a real number r such that $S_{i+1} = rS_i$ for all integers $1 \leq i < N$.

Solve T test cases per input file.

Constraints

- $1 \leq T \leq 10^5$
- $2 \leq N \leq 2 \times 10^5$
- $-10^9 \leq A_i \leq 10^9$ ($1 \leq i \leq N$)
- $A_i \neq 0$ ($1 \leq i \leq N$)
- The sum of N over all test cases in a single input file is at most 2×10^5 .
- All input values are integers.

2026-01-02 (Fri)
05:26:33 +11:00

Input

The input is given from standard input in the following format:

```
T
testcase1
testcase2
⋮
testcaseT
```

where testcase_i is the i -th test case ($1 \leq i \leq T$), and each test case is given in the following format:

```
N
A1 A2 ... AN
```

Output

Output T lines. The i -th line ($1 \leq i \leq T$) should contain Yes if A can be rearranged to form a geometric sequence in the i -th test case, and No otherwise.

Sample Input 1

[Copy](#)

```
3
5
1 8 2 4 16
5
-16 24 54 81 -36
7
90000 8100 -27000 729 -300000 -2430 1000000
```

[Copy](#)

Sample Output 1

[Copy](#)

```
Yes
No
Yes
```

[Copy](#)

In the first test case, the rearrangement $(16, 8, 4, 2, 1)$ of A forms a geometric sequence with common ratio $r = \frac{1}{2}$. Thus, print Yes on the first line.

In the second test case, no rearrangement of A satisfies the condition. Thus, print No on the second line.

2026-01-02 (Fri)
05:26:33 +11:00

#telegram)

url=https%3A%2F%2Fatcoder.jp%2Fcontests%2Fabc413%2Ftasks%2Fabc413_d%3Flang%3Den&title=D%20-
ence)

[Rule \(/contests/abc413/rules\)](/contests/abc413/rules) [Glossary \(/contests/abc413/glossary\)](/contests/abc413/glossary)

[Terms of service \(/tos\)](/tos) [Privacy Policy \(/privacy\)](/privacy) [Information Protection Policy \(/personal\)](/personal) [Company \(/company\)](/company)
[FAQ \(/faq\)](/faq) [Contact \(/contact\)](/contact)

Copyright Since 2012 ©AtCoder Inc. (<http://atcoder.co.jp>) All rights reserved.