

Contest Duration: 2025-11-29(Sat) 23:00 (<http://www.timeanddate.com/worldclock/fixedtime.html?iso=20251129T2100&p1=248>) - 2025-11-30(Sun) 00:40 (<http://www.timeanddate.com/worldclock/fixedtime.html?iso=20251129T2240&p1=248>) (local time) (100 minutes)

[Back to Home \(/home\)](#)

 [Top \(/contests/abc434\)](#)

 [Tasks \(/contests/abc434/tasks\)](#)

 [Clarifications \(/contests/abc434/clarifications\)](#)  [Results ▾](#)

 [Standings \(/contests/abc434/standings\)](#)

 [Virtual Standings \(/contests/abc434/standings/virtual\)](#)  [Editorial \(/contests/abc434/editorial\)](#)

 [Discuss \(<https://codeforces.com/blog/entry/148758>\)](#) 

E - Distribute Bunnies

[Editorial \(/contests/abc434/tasks/abc434_e/editorial\)](#)  / 

Time Limit: 2 sec / Memory Limit: 1024 MiB

Score : 500 points

Problem Statement

There are N rabbits numbered 1 to N on a number line. Rabbit i is at coordinate X_i .

Multiple rabbits may be at the same coordinate.

Each rabbit has a parameter called **jumping power**, and rabbit i has jumping power R_i .

Now, all rabbits will jump exactly once. When a rabbit at coordinate x with jumping power r jumps, it moves to coordinate $x + r$ or coordinate $x - r$.

If you can freely choose which coordinate each rabbit jumps to, find the maximum possible number of distinct coordinates where rabbits are present after all rabbits have jumped.

Constraints

- $1 \leq N \leq 2 \times 10^5$
- $-10^9 \leq X_i \leq 10^9$
- $1 \leq R_i \leq 10^9$
- All input values are integers.

Input

The input is given from Standard Input in the following format:

2026-01-02 (Fri)
05:34:00 +11:00

```
N  
X1 R1  
X2 R2  
:  
XN RN
```

Output

Output the maximum possible number of distinct coordinates where rabbits are present after jumping.

Sample Input 1

[Copy](#)

```
3  
4 1  
2 3  
4 5
```

[Copy](#)

Sample Output 1

[Copy](#)

```
3
```

[Copy](#)

If each rabbit moves as shown below, it is possible to achieve 3 as the number of distinct coordinates where rabbits are present after jumping, and this is the maximum possible.

- Rabbit 1 moves to $4 - 1 = 3$.
- Rabbit 2 moves to $2 + 3 = 5$.
- Rabbit 3 moves to $4 - 5 = -1$.

Sample Input 2

[Copy](#)

```
6  
2 1  
3 2  
6 1  
5 2  
4 3  
4 1
```

[Copy](#)

Sample Output 2

[Copy](#)

2026-01-02 (Fri)

05:34:00 +11:00

[Copy](#)

If each rabbit moves as shown below, it is possible to achieve 4 as the number of distinct coordinates where rabbits are present after jumping, and this is the maximum possible.

- Rabbit 1 moves to $2 - 1 = 1$.
- Rabbit 2 moves to $3 + 2 = 5$.
- Rabbit 3 moves to $6 + 1 = 7$.
- Rabbit 4 moves to $5 + 2 = 7$.
- Rabbit 5 moves to $4 - 3 = 1$.
- Rabbit 6 moves to $4 - 1 = 3$.

Sample Input 3

[Copy](#)

```
10
1000000000 1000000000
1000000000 1
-1000000000 1000000000
-1000000000 1
0 1
2 1
1 2
4 1
3 2
4 3
```

[Copy](#)

Sample Output 3

[Copy](#)

```
9
```

[Copy](#)

'#telegram)

?url=https%3A%2F%2Fatcoder.jp%2Fcontests%2Fabc434%2Ftasks%2Fabc434_e%3Flang%3Den&title=E%20-

[Rule \(/contests/abc434/rules\)](#) [Glossary \(/contests/abc434/glossary\)](#)

[Terms of service \(/tos\)](#) [Privacy Policy \(/privacy\)](#) [Information Protection Policy \(/personal\)](#) [Company \(/company\)](#)
[FAQ \(/faq\)](#) [Contact \(/contact\)](#)

Copyright Since 2012 ©AtCoder Inc. (<http://atcoder.co.jp>) All rights reserved.