Statistical Sciences

DoSS Summer Bootcamp Probability
Module 9

Ichiro Hashimoto

University of Toronto

July 24, 2025

Outline

- Counterexamples

Counterexamples

Recall: A random variable $X \in L^p$ if $\|X\|_{L^p} = (E|X|^p)^{1/p} < \infty$.
$X_n \to X$ in L^p if $\lim_{n\to\infty} \|X_n - X\|_{L^p} = 0$

Monotonicity of L^p Convergence

If $q > p > 0$, L^q convergence implies L^p convergence

Counterexample to the Converse:

Strategy: Find X_n s.t. $X_n \to 0$ in L^p but $\mathbb{E}|X_n|^q \geq 1$

$$X_n = \begin{cases} n^\alpha & \text{w.p } n^{-1} \\ 0 & \text{w.p } 1-n^{-1} \end{cases}$$

Find α later

$E|X_n|^p = n^{\alpha p - 1}$, $\mathbb{E}|X_n|^q = n^{\alpha q - 1}$

Let α s.t. $\alpha p - 1 < 0 \leq \alpha q - 1$

$\iff \underline{q^{-1} \leq \alpha < p^{-1}}$

For any α satisfying this, $X_n \to 0$ in L^p while $\mathbb{E}|X_n|^q \geq 1$

Counterexamples

Recall: X_n converges to X in probability if for any $\epsilon > 0$ $\lim_{n\to\infty} P(|X_n - X| > \epsilon) = 0$.

L^p convergence implies Convergence in Probability

If $X_n \to X$ in L^p, then $X_n \to X$ in probability.

Counterexample to the Converse:

Strategy Find X_n s.t. $\mathbb{E}|X_n|^p \to \infty$ and $X_n \xrightarrow{p} 0$

$$X_n = \begin{cases} n^{1+\frac{1}{p}} & \text{w.p } \frac{1}{n} \\ 0 & \text{w.p } 1-\frac{1}{n} \end{cases}$$

Then for $\forall\, \varepsilon \in (0,1)$, $\mathbb{P}(|X_n| > \varepsilon) = \mathbb{P}(X_n = n^{1+\frac{1}{p}}) = \frac{1}{n} \to 0$.

Thus, $X_n \xrightarrow{p} 0$ holds.

However, $E|X_n|^p = n^{p+1} \cdot \frac{1}{n} = n^p \to \infty.$

Counterexamples

Recall: X_n converges to X in probability if for any $\epsilon > 0$ $\lim_{n\to\infty} P(|X_n - X| > \epsilon) = 0$.

a.s. Convergence implies Convergence in Probability

If $X_n \to X$ almost surely, then $X_n \to X$ in probability.

Counterexample to the Converse:

$\Omega = [0,1)$, $\mathbb{P}$ uniform on $[0,1)$

$$X_{n,m}(\omega) = \begin{cases} 1 & \text{if } \omega \in [\frac{m}{n}, \frac{m+1}{n}) \\ 0 & \text{else.} \end{cases}$$

$0 \leq m \leq n-1$

For $\varepsilon \in (0,1)$, $\mathbb{P}(|X_{n,m}| > \varepsilon) = \mathbb{P}(X_{n,m} = 1) = \frac{1}{n} \to 0$

For each n by taking $m = 0 \to n-1$



For fixed $\omega \in [0,1)$,

$X_{n,m}(\omega)$ takes both 0 and 1

while m runs from 0 to $n-1$ for each n.

Thus, $X_{n,m}(\omega)$ takes both 0 and 1 infinitely many times,

which means $X_{n,m}(\omega)$ does not converge to 0 everywhere.

This example is also counterexample to the following claim:

Claim: If $X_n \to X$ in L^p for any p,

then $X_n \to X$ a.s.

This is since $\mathbb{E} X_{n,m}^p = 1^p \cdot \frac{1}{n} = \frac{1}{n} \to 0$.

while $X_{n,m} \not\to 0$ a.s.

Counterexamples

Recall: X_n converges to X in distribution if for any continuity point x of $P(X \leq x)$, $\lim_{n\to\infty} P(X_n \leq x) = P(X \leq x)$ holds.

Convergence in Probability implies Convergence in Distribution

If $X_n \to X$ in probability, then $X_n \to X$ in distribution.

Counterexample to the Converse:

Counterexamples

Recall: X_n converges to X in distribution if for any continuity point x of $P(X \leq x)$, $\lim_{n\to\infty} P(X_n \leq x) = P(X \leq x)$ holds.

Convergence in Probability implies Convergence in Distribution

If $X_n \to X$ in probability, then $X_n \to X$ in distribution.

Special case when the Converse holds:

Counterexamples

Monotone Convergence Theorem

If $X_n \geq 0$ and $X_n \nearrow X$, then $EX_n \nearrow EX$

Counterexample when X_n is not lower bounded:

Counterexamples

Dominated Convergence Theorem

If $X_n \to X$ a.s. and $|X_n| \leq Y$ a.s. for all n and Y is integrable, then $EX_n \to EX$

Counterexample when X_n is not dominated by an integrable random variable: