

Written Homework 2**Exercise 1**

[Temperature model] When Dr. Watson picks up the loot, he realizes that the loot is an ice cube of temperature -20° , in which some curious object is encapsulated. In order to obtain what is inside the ice cube, Holmes and Watson put the ice cube in a room of constant temperature $T_R = 30^\circ$. It is known in this situation, the rate of change of the temperature of the ice cube T_C satisfies

$$\frac{dT_C}{dt} = -k(T_C - T_R),$$

where k is a constant.

After one minute, the temperature of the ice cube rises to -15° . How long will it take for the temperature of the ice cube to become 0° so that it melts?

Hint: we need to figure out the constant k in order to solve this problem.

Exercise 2

Find the general solution of

1. $xy' + y = x^3$
2. $y' + ay = b$
3. $\frac{dr}{d\theta} = (r + e^\theta) \tan \theta$
4. $\frac{dy}{dx} - \frac{2xy}{x^2 + 1} = 1$

Exercise 3

Find a particular solution of

1. $xy' + 3y = \frac{\sin x}{x^2}$, $x \neq 0$, $y(\frac{\pi}{2}) = 1$
2. $y' - y = e^x$, $y(0) = 1$

The differential equation of the form

$$\frac{dy}{dx} + P(x)y = Q(x)y^n \tag{1}$$

is called the **Bernoulli Equation**. If $n = 0$, then this is the linear first order differential equation; otherwise, we can not solve this ODE by the method of integrating factor. However, if we multiply

$$(1 - n)y^{-n}$$

on both sides of (1), we get

$$(1 - n)y^{-n} \frac{dy}{dx} + (1 - n)P(x)y^{1-n} = (1 - n)Q(x).$$

Notice that

$$\frac{d(y^{1-n})}{dx} = (1 - n)y^{-n} \frac{dy}{dx},$$

Equation (1) becomes

$$\frac{d(y^{1-n})}{dx} + (1 - n)P(x)y^{1-n} = (1 - n)Q(x). \quad (2)$$

Now let $u = y^{1-n}$, then (2) becomes

$$\frac{du}{dx} + (1 - n)P(x)u = (1 - n)Q(x) \quad (3)$$

which can be solved by the integrating factor method.

Use the above method to solve Exercise 4 and 5:

Exercise 4

Solve

1. $xy' + y = y^2 \log x$
2. $y' + xy = \frac{x}{y^3}, \quad y \neq 0$

Exercise 5

Find a particular solution of $y' + \frac{1}{x}y = \frac{y^2}{x}, \quad y(0) = 1$.

Exercise 6

Solve

$$\frac{dx}{dy} + 2yx = e^{-y^2}.$$

Exercise 7

Solve for a particular solution

$$(x - \sin y)dy + (\tan y)dx = 0, \quad y(1) = \frac{\pi}{6}$$

Hint: Look at the last exercise.