

Random Walks in Number Theory: The Magic and Mystery of Numbertheoretic Fourier Series

Yu Fu, Jia Yu, Junxian Li (Project Leader), A.J. Hildebrand (Faculty Mentor)



Illinois Geometry Lab Open House, December 11, 2014

Weyl Sums

Weyl sums are sums of the form


$$S(N) = \sum_{n=1}^N e^{2\pi i f(n)},$$

where $f(n)$ is some real-valued smooth function. **Examples:**

- **Gauss sums:** $f(n) = n^2/p$, p prime.
- **Generalized Gauss sums:** $f(n) = an^b$, a, b constants.

Properties: Such curves of these sums show fractal-like behavior, and swirls called "curlicues". The graphs arise in physics (e.g. thermodynamics).

Weyl Sums Gallery



References

- Berry, M. V. ; Goldberg, J. *Renormalisation of curlicues*, Nonlinearity 1 (1988): 1-26
- Moore, Ross. R. ; van der Poorten, A. J. *On the thermodynamics of curves and other curlicues*. Proceedings of conference on geometry and physics (1989): 89-0031
- Dekking, M. ; Mendès-France, M. *Uniform distribution modulo one: a geometrical viewpoint*. J. für Reine Angew. Math. 329 (1981): 143-153

Numbertheoretic Fourier Series

Definition

Numbertheoretic Fourier series are sums of the form

$$g(t) = \sum_{n=1}^{\infty} \frac{f(n)}{n} e^{2\pi i nt},$$

as $0 \leq t \leq 1$, where $f(n)$ is some natural numbertheoretic function. In particular we consider the following functions:


$$\mu(n)n^b, \sin(an^b), e^{2\pi i an^b}, \{\sqrt{n}\}.$$

We compare their behavior with that of random coefficients $f(n)$.

Observations

- Fourier series with Moebius coefficients $\mu(n)$, or more generally $\mu(n)n^b$, behave like series with random coefficients. Their plots show chaotic behavior.
- Fourier series with coefficients $f(n) = e^{2\pi i an^b}$ show interesting spiral-type features.
- Close-up images of these graphs reveal intriguing fractal-type features.
- Fourier series with random complex coefficients $f(n)$ with $|f(n)| = 1$ show chaotic behavior.

Numbertheoretic Fourier Series Gallery



Motivation and Background


- If $f(n)$ is randomly chosen in $\{-1, 1\}$ or uniformly distributed in $[-1, 1]$, then

$$\sum_{n=1}^{\infty} \frac{f(n)}{n} e^{2\pi i nt}$$

converges for all t with probability 1.

- If $f(n) = \mu(n)$, then $\sum_{n=1}^{\infty} \frac{\mu(n)}{n} e^{2\pi i nt}$ converges for all t (Bateman-Chowla (1963)).
- The geometric behavior of the latter series was investigated by Froberg (1966) and Bohman-Froberg (1995).

Fourier Series Close-ups Gallery



References

- Bateman, P. T. ; Chowla , S. *Some special trigonometrical series related to the distribution of prime numbers*. J. London Math. Soc. 38 (1963): 372-374
- Bohman, J. ; Froberg, C. -E. *Heuristic investigation of chaotic mapping producing fractal objects*. BIT Numerical Mathematics, (1995): 609-615
- Froberg, Carl-Erik. *Numerical studies of the Moebius power series*. Nordisk Tidskr. Informations-Behandling 6 (1966): 191-211