


“Skin α -Synuclein Aggregation Seeding Activity as a Novel Biomarker for Parkinson Disease”

Presented by: Jackie Lindstrom

Parkinson's Disease (PD) Background

- Second most common neurodegenerative disease¹
- Characterized by symptoms such as⁶
 - Shaking or tremors at rest
 - Bradykinesia (slowness of movement)
 - Muscle stiffness
 - Cognitive changes such as dementia, depression, and/or anxiety
- No existing cure
- Existing treatments target symptoms, not the source of disease




Diagnosis

Current diagnostic techniques are lacking. By the time of diagnosis a large amount of dopaminergic neurons have already died. Therefore, development of early diagnostic techniques are critical for early distribution of treatment.

α -synuclein

α -synuclein: a small, insoluble protein, composed of 140 amino acids and encoded by the SNCA gene²

- Normal role: involved with vesicle tracking, vesicle docking, and neurotransmitter release, however still uncertain
- PD brains have abnormal α -synuclein behavior... which is why PD is classified as a synucleinopathy
 - Lewy Bodies: clumps of α -synuclein and other proteins that aggregate in regions of the brain
 - Oligomerization of α -synuclein is hypothesized to be toxic
- Autopsies show neuronal death (specifically in dopaminergic neurons) that could be **linked** to the aggregation of **α -synuclein**
 - Neuronal death in basal ganglia region of brain
- Immunohistochemistry and immunofluorescence has detected α -synuclein in peripheral tissues.... however inaccurate for diagnosis



****Found that misfolded a-synuclein exhibits “prion-like aggregation seeding ability”, meaning that it will trigger other proteins to also misfold, and the clumps of abnormal proteins will proliferate. This is a possible explanation for how neuronal death spreads through the brain****


Article Goal:

Assess α -synuclein presence in peripheral skin tissue as a contender for a synucleinopathic disease biomarker. This could be beneficial because....

- Could show confirmation of PD *before* the disease progresses far and symptoms show
- Would be a minimally invasive technique, as peripheral tissue is easier to access than the brain
- Would provide a way to monitor changes induced by therapies that target α -synuclein (which currently exist!)


Experimental Information

- Skin samples from 130 autopsies
 - Cadavers had a variety of neurodegenerative diseases: PD, Lewy body dementia, multiple system atrophy, and tauopathic diseases, along with controls
 - Diagnosis was confirmed by neuropathological tissue exams
 - Skin samples from 41 biopsies
 - Diagnosed PD patients and controls only
 - Obtained from leg or posterior cervical region of neck
1. First confirmed presence of a-syn by IHC and IF
 2. Used real-time quaking-induced conversion (RT-QuIC)
 - The presence of an abnormal a-syn protein prion protein will form aggregates known as amyloids that fluoresce
 3. Used protein misfolding cyclic amplification (PMCA assays)
 - A technique conceptually similar to PCR that involves leaving prion with normal protein, so that the normal protein is converted. Ultrasound then breaks up protein so to repeat



RT-QuIC & PMCA Assay:

- Increased levels of fluorescence for synucleopathies compared to control



“To our knowledge, this study has demonstrated for the first time that skin a-synuclein has aggregation seeding activity that was significantly higher in individuals with PD and other synucleopathies than in those with tauopathies and NNCs”

Limitations

- Compared to PD and “normal” cases, not enough non-PD synucleopathies examined
- More skin biopsies would be helpful
- (!)Remember, diagnosis of PD from clinical symptoms is not precise, so the separation of PD s non-PD may not be reliable
 - Follow up with autopsies

Sources

- (1) Li, S.; Le, W. Milestones of Parkinson's Disease Research: 200 Years of History and Beyond. *Neuroscience Bulletin* **2017**, *33* (5), 598–602. <https://doi.org/10.1007/s12264-017-0178-2>.
- (2) Jankovic, J.; Tan, E. K. Parkinson's Disease: Etiopathogenesis and Treatment. *Journal of Neurology, Neurosurgery & Psychiatry* **2020**, *91* (8), 795–808. <https://doi.org/10.1136/jnnp-2019-322338>.
- (3) Galvan, A.; Devergnas, A.; Wichmann, T. Alterations in Neuronal Activity in Basal Ganglia-Thalamocortical Circuits in the Parkinsonian State. *Frontiers in Neuroanatomy* **2015**, *9*. <https://doi.org/10.3389/fnana.2015.00005>.
- (4) Simon, D. K.; Tanner, C. M.; Brundin, P. Parkinson Disease Epidemiology, Pathology, Genetics, and Pathophysiology. *Clinics in Geriatric Medicine* **2020**, *36* (1), 1–12. <https://doi.org/10.1016/j.cger.2019.08.002>.
- (5) Ingelsson, M. Alpha-Synuclein Oligomers—Neurotoxic Molecules in Parkinson's Disease and Other Lewy Body Disorders. *Frontiers in Neuroscience* **2016**, *10*. <https://doi.org/10.3389/fnins.2016.00408>.
- (6) NHS Choices. Diagnosis - Parkinson's disease <https://www.nhs.uk/conditions/parkinsons-disease/diagnosis/>.
- (7) Gątarek, P.; Pawełczyk, M.; Jastrzębski, K.; Głabiński, A.; Kałużna-Czaplińska, J. Analytical Methods Used in the Study of Parkinson's Disease. *TrAC Trends in Analytical Chemistry* **2019**, *118*, 292–302. <https://doi.org/10.1016/j.trac.2019.05.047>.
- (8) Tambasco, N.; Romoli, M.; Calabresi, P. Levodopa in Parkinson's Disease: Current Status and Future Developments. *Current Neuropharmacology* **2018**, *16* (8), 1239–1252. <https://doi.org/10.2174/1570159x15666170510143821>.
- (9) Gravit, L. The Promise and Potential of Stem Cells in Parkinson's Disease. *Nature* **2021**, *597* (7878), S8–S10. <https://doi.org/10.1038/d41586-021-02622-3>.
- (10) Axelsen, T. M.; Woldbye, D. P. D. Gene Therapy for Parkinson's Disease, an Update. *Journal of Parkinson's Disease* **2018**, *8* (2), 195–215. <https://doi.org/10.3233/jpd-181331>.
- (11) Wang, Z.; Becker, K.; Donadio, V.; Siedlak, S.; Yuan, J.; Rezaee, M.; Incensi, A.; Kuzkina, A.; Orrú, C. D.; Tatsuoka, C.; Liguori, R.; Gunzler, S. A.; Caughey, B.; Jimenez-Capdeville, M. E.; Zhu, X.; Doppler, K.; Cui, L.; Chen, S. G.; Ma, J.; Zou, W.-Q. Skin α -Synuclein Aggregation Seeding Activity as a Novel Biomarker for Parkinson Disease. *JAMA Neurology* **2021**, *78* (1), 30. <https://doi.org/10.1001/jamaneurol.2020.3311>.