ATOM User Guide

Version 6.2.0, November 14, 2026

José Luís Martins
INESC MN, Lisboa, Portugal

jlmartins@inesc-mn.pt

jose.l.martins@tecnico.ulisboa.pt

Section 4 adapted from SIESTA documentation by Alberto García

Contents

1 Introduction **2**

1.1 What the code does . . . 2

1.2 Resources for ab-initio DFT and pseudopotentials . . . 2

2 Installation **3**

2.1 Step 1: Downloading and extracting the archive . . . 3

2.2 Step 2: Generating the documentation . . . 3

2.3 Step 3: Compiling the code . . . 4

3 Running the code **5**

3.1 Validation . . . 5

3.2 Running pseudopotential construction with all tests and plots . . . 5

3.2.1 First command line argument . . . 6

3.2.2 Other command line arguments . . . 7

3.3 Tested pseudopotentials . . . 7

3.4 Running step by step . . . 9

3.4.1 The all-electron calculation . . . 9

3.4.2 Constructing the pseudopotential . . . 12

3.4.3 Visualizing the pseudopotential . . . 15

3.4.4 Applying the Kleinman-Bylander transformation . . . 15

3.4.5 Visualizing the KB operators . . . 18

3.4.6 Logarithmic derivative plots . . . 18

3.4.7 Testing the pseudopotentials . . . 19

4 The atom.dat input file **20**

4.1 All-electron calculations . . . 20

4.2 Pseudopotential generation . . . 21

4.3 Core corrections . . . 22

4.4 Format of the file . . . 22

5 Computational details **25**

5.1 Radial grid . . . 26

5.2 Solving radial equations . . . 26

5.3 Atomic orbitals . . . 27

5.4 Fourier transforms . . . 28

1 Introduction

This is a modernized version of an old code that generates and tests pseudopotentials to be used in Density Functional Theory (DFT) calculations of electronic structure of molecules, solids, etc...

The original code was written by Sverre Froyen at the University of California Berkeley. Phys. Rev. B 26, 1738 (1982), https://doi.org/10.1103/PhysRevB.26.1738. A 1985 version of the code can be found in `Doc/Historic/atomsve.f`, it can still be compiled and run! That code had major enhancements by Norman J. Troullier at the University of Minnesota, Minneapolis. Phys. Rev. B 43, 1993 (1991), https://doi.org/10.1103/PhysRevB.43.1993 Since then the code has been maintained and improved by José Luís Martins. Peter Schuster and Manuel Maria Alemany wrote new plotting packages. GGA were introduced by Carlos Balbas and José Soler in a collaboration with the SIESTA code.

The present version is a "translation" from old FORTRAN (Fortran 70) to modern Fortran (90 $\rightarrow$ present), with detailed documentation and added enhancements by José Luís Martins. Another modernized version of the original Sverre Froyen code can be found at `https://siesta-project.org/siesta/Documentation/Pseudopotentials`, is maintained by Alberto García, who added some features and made substantial structural changes to the April 1990 (5.0) Minnesota version. Both versions can use the same input file as the original 1982 code.

1.1 What the code does

The program's basic capabilities are:

- All-electron DFT atomic calculations for arbitrary electronic configurations and a few Local Density Approximation (LDA) and Generalized Gradient Approximations (GGA) functionals.
- Generation of ab-initio pseudopotentials, and their transformation to the Kleinman-Bylander form with atomic basis functions.
- Atomic calculations in which the effect of the core is represented by a previously generated pseudopotential. These are useful to make sure that the pseudopotential correctly reproduces the all-electron results for the valence complex.

1.2 Resources for ab-initio DFT and pseudopotentials

We will not review here the background of density functional pseudopotentials. But, in this case more than ever, there is a lot to be gained from reading the original literature, for both the basics, and what is implemented in the code. Here are a few basic references:

- The book by Richard Martin "Electronic Structure: Basic Theory and Practical Methods" (Cambridge University Press) reviews the field and has a chapter on pseudopotentials.
- Some seminal papers on ab-initio pseudopotentials, (but certainly not an exhaustive or historic list of the development of the key concepts.):

  Hamann, Schluter, Chiang, Phys. Rev. Lett. 43, 1494 (1979)
  Kerker, J. Phys. C 13, L189-94 (1980)
  Bachelet, Hamann, Schluter, Phys. Rev. B 26, 4199 (1982)

- Troullier-Martins elaboration of Kerker method:

  Troullier, Martins, Phys. Rev. B 43, 1993 (1991)

- Partial core corrections:

  Louie, Froyen, Cohen, Phys. Rev. B 26, 1738 (1982)

- Reviews of plane-wave pseudopotential ab-initio calculations:

  W. E. Pickett, "Pseudopotential Methods in Condensed Matter Applications", Computer Physics Reports 9, 115 (1989)

  M. C. Payne, M. P. Teter, D. C. Allan, T. A. Arias and J. D. Joannopoulos, "Iterative minimization techniques for ab initio total-energy calculations: molecular dynamics and conjugate gradients", Rev. Mod. Phys. 64, 1045, (1992)

2 Installation

2.1 Step 1: Downloading and extracting the archive

The code is available from José Luís Martins GitHub:

`https://github.com/jlm785/pseudopotential`

If you downloaded the `pseudopotential-6.x.y.tgz` file (where `x` and `y` are major and minor version numbers) just extract it,

```
$ tar xzf pseudopotential-6.x.y.tgz
```

and you will have the code in a `pseudopotential-6.x.y` directory If you cloned the git you already have the relevant code in the main directory. As always releases are stable, git clones are just snapshots. Do not clone if you notice active modifications.

2.2 Step 2: Generating the documentation

The documentation is in the `atom/Doc` directory, where `atom/` stands for `pseudopotential-6.x.y`. It includes this file, `user_guide.tex` in `atom/Doc/UserGuide`, with a description of the key input file `atom.dat` and key output and intermediate files. To get the pdf just run

```
atom/Doc/UserGuide$ pdflatex user_guide.tex
```

and repeat to have the correct references,

```
atom/Doc/UserGuide$ pdflatex user_guide.tex
```

to get a `user_guide.pdf` file.

A navigable description of the whole code for developers can be generated from the information in the folder `Doc/doxy`. For that detailed description you need doxygen (`https://www.doxygen.nl`) and graphviz (`https://graphviz.org/`). Both are available on most distributions. To check that your computer has doxygen and graphviz installed run the commands

```
atom/Doc$ dpkg -s doxygen
```

```
atom/Doc$ dpkg -s graphviz
```

on a Debian based distribution (Ubuntu et al), in other distributions use equivalent tools (yum, rpm, dnf, zypper,...) or the relevant GUI.

If they are not available just install them

```
atom/Doc$ sudo apt install doxygen
```

```
atom/Doc$ sudo apt install graphviz
```

again for Debian based distros, use equivalent tools (yum, rpm, dnf, zypper,...) or the relevant GUI. Finally run doxyfile in the doxy directory

```
atom/Doc/doxy$ doxygen Doxyfile
```

You will get two folders `atom/Doc/html` and `atom/Doc/latex` If you know what you are doing you can edit `atom/Doc/doxy/Doxyfile` to change the defaults. Finally it is useful to create a link to the file `atom/Doc/html/index.html`

```
atom/Doc$ ln -s html/index.html documentation.html
```

Opening that link in your browser will allow you to see the documentation for every file in the present code, with the relevant call tree and a short description of all the arguments of the subroutines.

2.3 Step 3: Compiling the code

The code is regularly tested with several Fortran compilers, `ifx` (from Intel oneAPI), `gfortran` (from gnu). Once in a while it also tested with other compilers, such as `pgfortran`. As the performance is not an issue for these "simple" one-dimensional (radial Schrödinger) calculations just use the compiler that is more familiar for you.

The good news is that the code is self-contained, it just needs the compiler,

```
atom$ cd Src
```

```
atom/Src$ gfortran atom_all.f90 lib_*/*.f90 lib_blas_pack/*.f -o atom_all.exe
```

will work out of the box. But you may want to change the compilation flags, for example to optimize for your computer. That is easily achieved with `make`. All the folders have an appropriate `Makefile` you just need edit the preamble file `atom/Src/make.inc`. In principle you have to make just two choices. The first is to identify which CPU you are using, as it narrows the choice of possible compilers. This is done by commenting/uncommenting the lines below `# compiler for the job`. The second is the compiler you want, and is the key decision. Again comment/uncomment the lines below `# Suggestions for compilers`. For any other modifications of the file it is assumed you know what you are doing... The CPW2000, `elk` and WANNIER90 use a `make.inc` file so you can find plenty of examples in those codes for specific combinations of computers and compilers.

To compile just go to `atom/Src` and type make

```
atom/Src$ make
```

In a few seconds you should have the main library `libatom_`*compiler*`.a` and a few executables `atom_`*compiler*`.exe`, etc... where *compiler* is the name of the chosen compiler. As you probably are not interested in comparing compilers you can create links, for example

```
atom$ ln -s Src/atom_all_gfortran.exe atom_all.exe
```

```
atom$ ln -s Src/atom_gfortran.exe atom.exe
```

etc...

The code relies on `gnuplot` (`http://www.gnuplot.info`) to produce graphics interactively. Just install it on your machine (apt, yum, rpm, dnf, zypper,...) if you want to see them as the code runs. If `gnuplot` is not installed, there is no problem, you will have in your folder the data required to make plots with `gnuplot` or with `matplotlib` on another machine.

3 Running the code

I assume you have a working directory separate from the source. In the following I will call it `atom/WORK`, and assume you created the links in `atom/`.

This is a code with a long history. Keeping it back-compatible (the original code still runs) made it easier to check that modifications did not introduce bugs, at the cost of some compromises, mainly in the number and constraints on the format of files. The old code and its additions communicated through files. The new version of the code still uses those file formats of several decades ago. The good news about the present version, is that, except for their presence in the working directory, you do not have to worry about them in most cases.

You can run the whole pseudopotential generation and testing procedure at once, or run step by step. Each step is one of the rectangular boxes in Fig. 1, the whole generation and testing procedure runs all the steps in a single shot. As the code uses intermediary files you can go back and repeat each of the steps you want, for example to modify the default values of parameters or generate new graphics.

WARNING: the defaults of the code are "educated guesses" for the majority of atoms. Any pseudopotential should be thoroughly tested before being used. For a few elements the code has *tested pseudopotentials*, compared with all-electron calculations, and it always tells you if you are using one of them or an educated guess (Section 3.3).

3.1 Validation

The compiled code should be tested. Go to the `Validation` folder and run the validation script (edit the path to the executable in `run.sh` if you did not compile with `ifx`),

```
atom$ cd Validation
```

```
atom/Validation$ bash run.sh
```

go for a coffee, and when you return you should have the pseudopotentials for all atoms with the default choices. **WARNING:** the defaults of the code are "educated guesses" for the majority of atoms, in particular transition metals, Lanthanides, and Actinides. Any pseudopotential should be thoroughly tested before being used! For the elements with tested pseudopotentials (Section 3.3) the validation uses the first (default) one.

3.2 Running pseudopotential construction with all tests and plots

Assuming you have gnuplot installed and you copied the executables to the code root, you can run the whole pseudopotential generation and testing with a simple command:

```
atom/WORK$ ../atom_all.exe
```

the code will check if there is a `atom.dat` file in the working directory. This is the original input file of the old code and its strictly formatted structure is described in Section 4. If the file is present it will use it, otherwise the program will ask for the chemical symbol of the element

you want, and, if the element has tested pseudopotentials (Section 3.3), which one you want, write a `atom.dat` file with default values. In both cases it will then proceed with the reference all-electron calculation, the pseudopotential generation, finding atomic orbitals basis set and non-local projectors, generate plots and perform tests on the pseudopotential.

After running the code you will find many files on your working directory. These are described in more detail in the next subsections. Briefly, you get the "input" file `atom.dat` and the detailed output file `atom.out`. The most important files are those that are read by other codes, listed in Table 1, where *Xy* is the chemical symbol.

Table 1: Pseudopotential files written for other codes.

file	code
Xy`_LDA_TM.UPF`, *Xy*`_GGA_TM.UPF`	QUANTUM ESPRESSO
Xy`.psp8`	ABINIT
Xy`.psml`	SIESTA and other codes that use libpsml
Xy`.psf`	SIESTA
Xy`_POTKB_F.DAT`	CPW2000
Xy`_POTRE.DAT`	PARSEC

Notice that *Xy*`.psf` and *Xy*`_POTRE.DAT` are very similar. The first part of the internal file `psd.pot` is similar to those, however it includes at the end the information about the partial core kinetic energy density. Finally you get many graphics related files, `*.pdf` and `*.gp`, and a few `*.dat` internal data files. For details see the description of running the code step by step below.

3.2.1 First command line argument

You can pass arguments in the command line, for example

```
atom/WORK$ ../atom_all.exe -h
```

will print a short help text and stop.

You can also run in a full interactive mode

```
atom/WORK$ ../atom_all.exe -x
```

where you are asked to override the default values, and therefore can be cumbersome. Overriding the defaults is easier by running individually each main component of the code. The interactive mode is described below in the section for the step by step run. In the interactive mode a "wrong answer" will cause the program to ignore it and proceed with the default value, eventually giving a second chance to the user to enter an answer in the accepted range.

The most useful first argument is the chemical symbol,

```
atom/WORK$ ../atom_all.exe Si
```

where you can swap "Si" for the chemical symbol of any element that interests you from Hydrogen to Oganesson. All the work will be done with default values without prompting the user. If the element has tested pseudopotentials the default is the first of them (for carbon the `HARD` one), otherwise the defaults are educated guesses and the program says so. Read the **Warning** above about default values.

3.2.2 Other command line arguments

After the chemical symbol you can give other arguments, in any order, as the program identifies them by their content (lower or upper case, at most four characters each, except the functional):

- `xcLDA/xcGGA` chooses the exchange and correlation functional, the LDA with the Ceperley-Alder (Perdew-Zunger) parametrization or the PBE GGA. The default is the LDA. Other functionals (the Perdew-Wang 92 LDA) are offered in the interactive mode (Section 3.3). `LDA`, `GGA` or `PBE` without the `xc` stop the program with a hint;
- `yes/no` forces using or not using the partial core correction for exchange and correlation;
- an argument with a `Z` as second character (`SZ/SZP/DZ/DZP`) chooses the basis set, see below;
- any other argument is the label of a tested pseudopotential, for example `HARD` or `SOFT` for carbon, or `NONE` for the educated guess of the old tables (Section 3.3).

For example `../atom_all.exe C soft`, `../atom_all.exe C HARD xcLDA`, `../atom_all.exe Si xcGGA` or `../atom_all.exe C DZ none no`. If the label is not one of the tested pseudopotentials of that element the program lists the available ones and stops. If the element has no tested pseudopotentials the label is ignored, with a message.

Default values for the reference atomic configuration and pseudopotential radii can be overridden by careful editing the `atom.dat` file. Although the choice of using the partial core-correction for exchange and correlation can be altered in that way, it is easier to do it with the `yes/no` argument. Remember that changing any of those parameters of a tested pseudopotential (the partial core correction included) gives a pseudopotential that was not tested.

The code generates "SIESTA-like" confined atomic orbitals. By default two basis sets are built and written to the *Xy*`_POTKB_F.DAT` file (and to `plot_kb.dat`): set 1 is double-zeta with polarization (DZP) and set 2 is single-zeta (SZ). The second zeta is the relaxed energy derivative ϕ of the first one. The atomic orbitals have angular momenta $l = 0, \ldots, l_a$, where l_a is the larger of the highest occupied l and $l_{\max} - 1$, $l_{\max}$ being the highest angular momentum of the pseudopotential, and the polarization orbitals have $l_a + 1$ (not above f). There is a single polarization orbital (in SZP and DZP), the perturbative polarization of the first l_a orbital ψ as in SIESTA. The basis set argument, which takes the values `SZ/SZP/DZ/DZP` (lower case is accepted), replaces the two default sets by a single set of that kind; any other value with a `Z` as second character stops the program. See section 5.3 for the details on how these orbitals are constructed.

3.3 Tested pseudopotentials

For most elements the default parameters of the pseudopotential (reference configuration, core radii, partial core correction and local component of the Kleinman-Bylander form) come from internal tables that are educated guesses. For a few elements there are *tested pseudopotentials*, sets of parameters that were compared with all-electron calculations of molecules and solids. An element may have one or several of them, each identified by a label of up to four characters. For example carbon has a `HARD` pseudopotential, with a $2s^2 2p^2 3d^0$ reference configuration, core radii of 1.10 bohr, partial core correction and the d potential as local component, and a `SOFT` one, with $2s^2 2p^2$, core radii of 1.50 bohr, no partial core correction and the p potential as local component. The list for an element, with a short description of the tests, is shown when you are asked to choose (see below); all of them are in the file `Src/lib_atom/atom_p_tbl_tested.f90`.

When the code is run without prompting the user, for example `../atom_all.exe C`, the first tested pseudopotential of the element is used. Another one is chosen by giving its label as an argument (Section 3.2.2), for example `../atom_all.exe C SOFT`, and the label `NONE` gives the educated guess of the old tables. When the user is prompted for the chemical symbol (`atom_all.exe -x`, or `atom_all.exe` or `atom.exe` without an `atom.dat` file) the program also asks which pseudopotential to use and which exchange and correlation functional,

```
Tested pseudopotentials for C:

   1   HARD   diamond LAPW, C2H2 all-electr.
   2   SOFT   diamond
   3   NONE   old tables, core radii: hard tested (triple bond)

Enter your choice (default 1)

Exchange and correlation functional:

   1   LDA   ca   Ceperley-Alder, Perdew-Zunger parametrization
   2   LDA   pw   Perdew-Wang 92
   3   GGA   pb   Perdew-Burke-Ernzerhof (PBE)

Enter your choice (default 1)
```

and writes `atom.dat` with the parameters of the choices. Elements without tested pseudopotentials skip the first question.

The choice is not written in `atom.dat`, as you can edit that file and the pseudopotential would no longer be the tested one. Instead, when the pseudopotential is constructed (`psd_gen.exe` or `atom_all.exe`) the parameters actually used are compared with those of the tested pseudopotentials of the element: the valence configuration (principal and angular quantum numbers and occupations, added over spin or j), the core radii (within 0.005 bohr), the partial core correction with its parameters `cfac` and `rcfac`, and the parameter `cfactau` of the partial core kinetic energy density, which is independent of the partial core correction (Section 4.3). The exchange and correlation functional and the relativistic or spin-polarized character of the calculation are not compared, so you can generate the tested pseudopotential with another functional, for example `../atom_all.exe C HARD xcGGA` (check with which functional it was tested). The program reports the result,

```
The parameters are those of the tested pseudopotential C HARD
tests: diamond LAPW, C2H2 all-electr.
```

or

```
The parameters differ from the tested C pseudopotentials:  HARD  SOFT
This pseudopotential was not tested
core radii: hard tested (triple bond), test local and other choices
```

or, for elements without tested pseudopotentials,

```
There are no tested pseudopotentials for P, this one was not tested
core radii: hard guess, test local and other choices
```

Any change of a tested pseudopotential, a core radius in `atom.dat` or the partial core correction forced with `yes/no`, gives a pseudopotential that was not tested.

When the pseudopotential was not tested, the last line gives the confidence on the core radii of the old table, if those are the radii used (otherwise it says that the core radii differ from the old table, and that all the choices should be tested). That confidence refers only to the core radii: `hard guess` means they were never used, `hard used` that they were used in calculations, for example `hard used (oxides)` for sodium, and `hard tested` or `fully tested` that they were compared with all-electron calculations (`hard` because the radii of the old table are on the small side). The choice of the local component and the other parameters were not tested, hence the warning.

The result is written as `var=HARD` (or `var=NONE` if the pseudopotential was not tested) in the title of `psd.pot`, after the configuration and core radii, and from there it goes to the titles of *Xy*`_POTKB_F.DAT` and `pseudokb.dat` and to the *Xy*`.psml` file. `kb_conv.exe` reads it from `psd.pot` and uses the local component of the tested pseudopotential, or the one of the old table (not tested) if the pseudopotential was not tested. Files `psd.pot` written by older versions of the code have no label and are treated as not tested.

To add a tested pseudopotential you only have to edit `atom_p_tbl_tested.f90`, writing all its parameters (not only the ones that differ from the old tables). The first label of an element is its default. `NONE`, `LIST`, `YES`, `NO`, `LDA`, `GGA`, `PBE`, anything with a `Z` as second character and anything starting with `XC` cannot be used as labels.

3.4 Running step by step

While `atom_all` runs everything at once, the same results can be obtained by a few smaller programs that communicate through files, as described below. Running them step by step makes it easy to repeat one of the steps, for example with other core radii or another local component, without repeating the others. Figure 1 shows the programs and the files each of them reads and writes.

3.4.1 The all-electron calculation

The starting point of a pseudopotential construction is an all-electron calculation. This is done by running the "atom" code:

```
atom/WORK$ ../atom.exe
```

the program searches for a file `atom.dat` in the `WORK` directory with the information on the atomic configuration and options of the calculation such as the exchange and correlation model. This is the key input file and is described in Section 4.

If the file `atom.dat` is not present, the code asks the user the chemical symbol of the element to be calculated, and, if there are tested pseudopotentials for that element, which of them to use (Section 3.3), and the exchange and correlation functional, writes an `atom.dat` file with the default values and runs the calculation. The user may edit the `atom.dat` file to change the default choices, taking into account that the file is read with fixed formats (back-compatibility), so the columns matter (Section 4.4).

If the validation step of Section 3.1 was run, the user can find `atom.dat` for all atoms in the folders `Validation/res_`*Xy* where *Xy* is the chemical symbol. Modifying this file is the starting point for the construction of a good pseudopotential.




Figure 1: Programs (boxes) and the files through which they communicate when the code is run step by step. `atom`, `psd_gen`, `atom_test` and `kb_test` read `atom.dat`; the dashed boxes at the bottom list the files written for other codes (Table 1). `atom_all` runs the whole chain at once.

At the end of the `atom.exe` calculation a file `datafile.dat` is written with the information of the atomic configuration and self-consistent potential to be used by the pseudopotential generation code. In the interactive mode the output goes to the default output, in contrast to the non-interactive all at once mode where the output goes to the `atom.out` file. The output is largely self-explanatory. All energies are in Rydberg unless explicitly stated, and all lengths in Bohr radii. At the top of the output (example below) we have the date the code was run; the exchange-correlation functional, "ca" meaning the LDA based on the Ceperley-Alder homogeneous electron gas results; the nuclear and electronic charges and the numbers of core and valence orbitals; and the atomic configuration with quantum numbers and occupations (`occ`)

```
 atom-LDAca04-Nov-26           Si      file generated by atm_write 04-Nov-26
**********************

 Si pseudopotential generation
 -----------------------------

 correlation = ca   nonspin-polarized

 nuclear charge             = 14.000000
 number of core orbitals    =  3
 number of valence orbitals =  3
 electronic charge          = 14.000000
 ionic charge               =  0.000000


 input data for orbitals
```

```
 i    n    l    s     j     occ

 1    1    0   0.0   0.0    2.0000
 2    2    0   0.0   0.0    2.0000
 3    2    1   0.0   0.0    6.0000
 4    3    0   0.0   0.0    2.0000
 5    3    1   0.0   0.0    2.0000
 6    3    2   0.0   0.0    0.0000
```

Next we have the description of the radial mesh used in the calculation,

```
 radial grid parameters

 r(1) = .0 , r(2) =  0.887E-06 , ... , r(2686) = 119.814
 a =  6.000  b = 200.000
```

indicating that we have 2686 points on the grid, the first being the origin, the second $r(2)$ at 0.887×10^{-6} bohr, and the last at almost 120 bohr. See Section 5.1 for how the radial grid is constructed from the values of a and b.

For each orbital we have a list

- `a extr`, the values of $u(r) = r\psi(r)$ at its extrema (in relativistic calculations u is the major component, and the minor component is listed as `b extr`);
- `r extr`, the radii of those extrema;
- `r zero`, the radii where $\psi = 0$ (nodes);
- `r 90/99 %`, the radii that contain 90% and 99% of the charge of the orbital.

For the $3p$ orbital of Si one obtains

```
 n = 3  l = 1  s = 0.0
        a extr          0.264  -0.620
        r extr          0.316   2.146
        r zero          0.718
        r 90/99 %       4.463   6.726
```

showing extrema at 0.316 and 2.146 bohr, and a node at 0.718 bohr. It also shows that 90% of the electron density in that orbital is within 4.463 bohr and 99% within 6.726 bohr.

Next we have the number of iterations of the self-consistent screening potential and, for each orbital, the eigenvalue, the kinetic energy and the potential energy of the interaction with the nucleus (in Rydberg; the eigenvalue also includes the Hartree and exchange-correlation energies, so it is not the sum of the other two),

```
 Total number of iterations needed for electron screening potential is 16



 Si output data for orbitals
```

```
 ----------------------------

 nl    s      occ         eigenvalue    kinetic energy      pot energy

 1s   0.0    2.0000    -130.36911374     183.01377674    -378.73491523
 2s   0.0    2.0000     -10.14892755      25.89954327     -71.62102263
 2p   0.0    6.0000      -7.02876330      24.42537956     -68.74331327
 3s   0.0    2.0000      -0.79662747       3.23745219     -17.68692632
 3p   0.0    2.0000      -0.30705182       2.06135800     -13.62572590
 3d   0.0    0.0000       0.00231324       0.00232980      -0.38199473
```

and finally a control of the virial theorem. For non-relativistic calculations the virial sum should be zero. For relativistic calculations this (non-relativistic) theorem does not hold. In this non-relativistic example the virial sum is -0.00018 Ry (about 2.5 meV), a measure of the numerical accuracy of the calculation.

```
virial theorem(nonrelativistic)
 --------------

 kinetic energy  *  2      =     1149.95307571
 potential energy          =    -1151.36048794
 virial correction         =        1.40722738
 ---------------------------------------------
 virial sum                =       -0.00018485
```

3.4.2 Constructing the pseudopotential

The key operation is the construction of the pseudopotential. To generate the Troullier-Martins pseudopotential run

```
atom/WORK$ ../psd_gen.exe
```

The program reads the all-electron results from `datafile.dat`, presents a short summary of the results of the all-electron calculation and the current values for core radii, and asks the user if they should be modified.

```
   Found a file with all-electron results for Si


   Calculation was done with Ceperley-Alder LDA


   Based on a non-spin-polarized and non-relativistic calculation


   Calculation was run on 04-Nov-26 with code version 6.1.0



   The current value of core radii are:
```

```
   l =  0  rc =        1.800
   l =  1  rc =        1.800
   l =  2  rc =        1.800

   Do you want to modify them? (y/n)
```

If you answer yes, the program lists for each l the current core radius (from `atom.dat`), the value from an internal table of default values, and the radii of the outermost node and of the outermost maximum of the all-electron valence wave-function. It then asks for the maximum angular momentum of the pseudopotential channels, indicating the accepted range, and for the core radius of each channel, asking for confirmation of values that look suspicious. If you give a value out of the accepted range, the program will proceed with the current value.

The next question is about the partial core correction for exchange and correlation: the user is presented with the current situation and is asked if they want to change it. Next the user is asked if they want to change the radius where the partial core sets in. That radius $r_{\rm pc}$ is the outermost point where the core electron density is $c_{\rm fac}$ times the valence electron density, $\rho_{\rm core}(r_{\rm pc}) = c_{\rm fac}\,\rho_{\rm valence}(r_{\rm pc})$ (with a correction for ionized reference configurations), and the default value is $c_{\rm fac} = 1$ (`cfac` in the question). If a radius is given explicitly in `atom.dat`, the question is about that radius (`rcfac`) instead. For more details see Section 4.3.

The code will next ask if the user wants to change $c_{{\rm fac}\tau}$ (`cfactau`), which plays the same role for the partial core kinetic energy density: inside the outermost radius where $\rho_{\rm core} = c_{{\rm fac}\tau}\,\rho_{\rm valence}$ the core kinetic energy density is replaced by a smooth function.

Once the questions are answered, the code will print the parameters that describe the Troullier-Martins pseudopotential, and the core kinetic energy before and after truncation,

```
Si pseudopotential generation using the Improved Troullier and Martins method
 ------------------------------------------------------------

 nl    s    eigenvalue      rc          cdrc       delta       gamma       alpha_j

 1s   0.0   -0.796627    1.796507    0.369768   -1.387513    0.708147   -0.100295   -0.135680
 2p   0.0   -0.307052    1.796507    0.206765   -0.943628   -0.236975   -0.008022    0.021175
 3d   0.0    0.002313    1.796507    0.000000   -8.732549   -0.883722   -0.086774    0.195276
 core kinetic energy =    564.38

 kinetic energy density truncated
 inside r =  1.204
 ac =  6.954 bc = -4.133 cc =  0.491

 truncated core kinetic energy =     63.31
```

it will then list the characteristics of the pseudo-wave-functions, as for the all-electron orbitals (the pseudo-wave-functions have no nodes, so they are labelled 1s, 2p, 3d),

```
n = 1  l = 0  s = 0.0
        a extr         0.737
        r extr         1.778
        r zero
```

```
       r 90/99 %      3.306   4.835

 n = 2  l = 1  s = 0.0
       a extr         0.620
       r extr         2.146
       r zero
       r 90/99 %      4.463   6.726

 n = 3  l = 2  s = 0.0
       a extr         0.150
       r extr        80.446
       r zero
       r 90/99 %     99.081 111.157
```

and the core radii (bohr), eigenvalues (Rydberg), and the pseudopotentials at the origin and at their minimum (most negative) value, with the radius of that minimum.

```
 test of eigenvalues

 rcut =  s   1.80  p   1.80  d   1.80
 eval =  -0.79661  -0.30704   0.00231



 l    vps(0)    vpsmin      at r

 s     0.065    -5.501     1.392
 p    -6.064    -6.064     0.000
 d   -15.757   -15.757     0.000
```

Next it will show the eigenvalue, kinetic energy and bare potential energy of the pseudo-wavefunctions,

```
Si output data for orbitals
 ---------------------------

 nl    s      occ         eigenvalue    kinetic energy      pot energy

 1s   0.0    2.0000      -0.79661328       0.51034020      -3.73919585
 2p   0.0    2.0000      -0.30703675       0.76934634      -3.25652725
 3d   0.0    0.0000       0.00231324       0.00232980      -0.10914135
```

the eigenvalues should agree with the all-electron ones (here to about 10^{-5} Ry). Finally it will show the energy contributions for the pseudo-atom.

```
 total energies
 --------------

 sum of eigenvalues        =       -2.20730006
 kinetic energy from ek    =        2.55937308
```

```
el-ion interaction energy =      -13.99144619
el-el  interaction energy =        5.86809482
vxc    correction         =       -2.51140668
virial correction         =        0.17637063
exchange + corr energy    =       -1.92764767
kinetic energy from ev    =        2.55936317
potential energy          =      -10.05099904
-------------------------------------------
total energy              =       -7.49163587
```

At the end of the calculation the user has five new files: an unformatted `pseudo.dat` file used later in tests, a `plot.dat` file used for later plotting, a formatted `psd.pot` file used to pass the information for the Kleinman-Bylander transformation, and the files *Xy*`_POTRE.DAT` for PARSEC and *Xy*`.psf` for SIESTA. The file `psd.pot` is therefore the key output file of the program. Before the construction the program also tells you if the parameters are those of a tested pseudopotential, and that information is written in the title of `psd.pot` (Section 3.3).

3.4.3 Visualizing the pseudopotential

If `gnuplot` is installed the user can compare pseudo and all-electron wave-functions, observe the pseudopotential and respective Fourier transforms by running

```
atom/WORK$ ../plot_show.exe
```

The program will read `plot.dat` (or another file chosen by the user) and allow the user to change the range of the axes of the plots, and save a `pdf` version of the plots. If the user doesn't clean the `*.gp` files at the last question, the `WORK` directory will have several `command_`*name*`.gp` and *name*`.gp` files. The user can see the plots again with gnuplot

```
atom/WORK$ gnuplot command_name.gp
```

and can edit the command file to change the plot. The user can also use the data in *name*`.gp` files to generate plots with other software, including xmgrace and matplotlib. The script `Tools/atom_plot_mpl.py` draws the same figures with matplotlib from the `fig_`*name*`.dat` files. If gnuplot is not installed, the program just quits.

3.4.4 Applying the Kleinman-Bylander transformation

In plane-wave codes the pseudopotential is used in its Kleinman-Bylander form (Kleinman and Bylander, Phys. Rev. Lett. 48, 1425 (1982)). That transformation plus the generation of an efficient atomic basis set can be achieved by

```
atom/WORK$ ../kb_conv.exe
```

the program will read `psd.pot` and print the orbital energies, the default cutoff radii of the atomic basis functions (`r_siesta` is the radius of the energy-shift rule of SIESTA, `r_c` the radius used, see Section 5.3), the default local pseudopotential and the default Fourier grid.

```
  Orbital energies

    n     l            e

    1     0        -0.796617
```

```
  2     1         -0.307040
  3     2          0.002313

Basis cutoff radii (bohr)

 l    r_siesta      r_c
 0     4.9746     7.2132
 1     6.1764     7.1028
 2                7.1028   perturbative polarization of l =  1

The current parameters for the Kleinman-Bylander
non-local pseudopotential transformation are:

The maximum of all potentials will be used as local

The Fourier grid will range from q = 0.0000 to   40.0000  with steps of     0.0100
Enough for a cutoff of    200.000 Ha.
```

If the title of `psd.pot` identifies a tested pseudopotential (Section 3.3) the default local component is the one of that pseudopotential, otherwise it comes from an internal table, and for most elements it is the smoothed maximum of all the pseudopotentials, as in the example. The code will ask the user if they accept the defaults and in the negative case allows the choice of new parameters.

The first question, and a key decision, is the choice of the local component of the pseudopotential. A bad choice can introduce "ghost states" and the code will check if they appear, with the method of Gonze, Käckell and Scheffler, Phys. Rev. B 41, 12264 (1990). If a ghost state is found you are strongly warned in the output. A first rule of thumb is that the "best" local pseudopotential would be the one with highest angular momentum $l_{\max}$, as it would reduce the number of projectors, and in most cases an eventual $l_{\max}+1$ pseudopotential is more similar to the $l_{\max}$ than to $l<l_{\max}$ pseudopotentials. A second rule of thumb is that the smoothed maximum of all the pseudopotentials (the default of the internal table) makes all the projectors attractive and therefore dramatically reduces the possibility of having ghost states: the local potential is then the least attractive one, and its bound states lie above the reference energies.

The next question is about the Fourier grid (Section 5.4). Unless you have a good reason, the default grid is good enough.

The third question is about the confined atomic orbital basis. Read Section 5.3 for details of those basis sets before answering the questions.

The code then prints basic information about the calculation,

```
Constructing KB operator and basis functions

 Kleinman-Bylander pseudopotential conversion
 and Bessel/Fourier transform program. Version  6.1.0

 Run on  07-Nov-26

 Size of arrays:   lmax =     2    mxdnr =     2686

 The semi-local pseudopotential for Si was created on 05-Nov-26
```

```
with atomic program version atom 6.1.0
Pseudopotential type:   Improved Troullier - Martins potential

Pseudo parameters:
3s(  2.00)  rc= 1.803p(  2.00)  rc= 1.803d(  0.00)  rc= 1.80


The Fourier grid will range from q = 0.0000 to   40.0000  with steps of     0.0100

Total valence electrons =       4.000000

Orbital energies

  n     l             e

  1     0         -0.796617
  2     1         -0.307040
  3     2          0.002313
```

The code then prints a **suggestion** for the energy cutoff in the plane-wave code,

```
Suggested range of kinetic energy cutoff
for plane-wave calculations:

lower  cutoff (bands)             12  Ha
higher cutoff (pressure)          20  Ha
```

and then it will show the ratio $\int u_l^2\, \delta V_l\, dr / \int u_l^2\, |\delta V_l|\, dr$, where $u_l = r\psi_l$ is the pseudo-wave-function and $\delta V_l = V_l - V_{\rm local}$ the difference between the semi-local and local pseudopotentials.

```
ratio for l=   0          is      -1.0000
ratio for l=   1          is      -1.0000
ratio for l=   2          is      -1.0000
```

a negative value indicates an attractive projector. In this case all values are -1 due to the choice of the smoothed maximum of the pseudopotentials. A rule of thumb is that a small absolute value may indicate trouble with ghost states. The real test for ghost states is shown next,

```
Ghost State Test (X. Gonze et al)

l     j    0-node-eigen    1-node-eigen       True-eigen   inorm

0            -0.743217       -0.024325       -0.796617    -1
1            -0.244156        0.001389       -0.307040    -1
2             0.002313        0.005753        0.002313    -1
```

`0-node-eigen` and `1-node-eigen` are the two lowest eigenvalues of the local potential for that l, `True-eigen` is the reference eigenvalue, and `inorm` is the sign of the numerator of the ratio above. A ghost state is reported if `inorm` is -1 and the reference eigenvalue lies above the 0-node eigenvalue, or if `inorm` is 1 and it lies above the 1-node eigenvalue (Gonze, Käckell and Scheffler). In that case a warning will be displayed.

The final part of the output gives the information about the confined atomic orbitals,

```
Atomic basis functions

psi: confined orbital, E is its eigenvalue (Ry)
phi: relaxed energy derivative of the previous psi, E = <phi|H|phi> (Ry)
pol: perturbative polarization of the first function of l-1, E = <pol|H|pol> (Ry)
     w(r) = exp(-(r_c-r_w)/(r-r_w)),  r_w =  0.000 r_c

set  l  zeta  kind  nodes     r_c          E         err
  1  0    1   psi     0     7.2132    -0.795849
  1  0    2   phi     1     7.2132     0.259734   1.29E-06
  1  1    1   psi     0     7.1058    -0.297367
  1  1    2   phi     1     7.1058     0.492662   1.09E-06
  1  2    1   pol     0     7.1058     0.314735
  2  0    1   psi     0     7.2132    -0.795849
  2  1    1   psi     0     7.1058    -0.297367
```

where the first set (DZP) has 5 radial functions (13 orbitals counting the $2l + 1$ values of m) and the second set (SZ) has 2 radial functions (4 orbitals). The column `err` is an estimate of the relative error of ϕ, and `r_c` is rounded to a point of the radial grid.

At the end of the KB transformation the program writes an unformatted `pseudokb.dat` used later in tests, a `plot_kb.dat` file used for later plotting (which also contains the atomic basis functions), and the files for other codes: Xy`_POTKB_F.DAT` for CPW2000, Xy`_LDA_TM.UPF` (or `_GGA_`) for QUANTUM ESPRESSO, Xy`.psp8` for ABINIT and Xy`.psml` for SIESTA, where Xy is the chemical symbol of the relevant element.

3.4.5 Visualizing the KB operators

If `gnuplot` is installed the user can compare the pseudo wave-functions with the atomic basis functions, observe the KB operators and respective Fourier transforms, and have an estimate of the rate of convergence of the total energy with plane-wave basis size by running

```
atom/WORK$ ../plot_kb_show.exe
```

The program will read the `plot_kb.dat` file or another file specified by the user. See Section 3.4.3 for dealing with the files generated by the program. The basis functions can also be plotted, one figure for each basis set and angular momentum, with `python3 Tools/plot_basis.py` (matplotlib), which reads `plot_kb.dat`.

3.4.6 Logarithmic derivative plots

If `gnuplot` is installed the user can plot, as functions of the energy, the logarithmic derivatives of the radial wave-functions at a given radius with

```
atom/WORK$ ../plot_ln_show.exe
```

The program will read the `pseudo.dat`, `pseudokb.dat` and `datafile.dat` files (or the files specified by the user). See Section 3.4.3 for dealing with the files generated by the program.

This is a test based on scattering theory. Two potentials scatter in the same way at a radius r_0,

for an energy E and angular momentum l, if the logarithmic derivatives

$$\frac{\partial}{\partial r} \log \psi_l(r; E)\Big|_{r_0} \tag{1}$$

of the solutions $\psi_l(r; E)$ of the radial Schrödinger equation that are regular at the origin are the same. For r_0 outside the core radii, the logarithmic derivatives of the all-electron and pseudo potentials agree at the reference eigenvalues, because the wave-functions are equal there, and norm conservation (the never cited Topp-Hopfield theorem) makes their first energy derivatives also agree, so they are very close near them. The plots show how far in energy that agreement extends, for the semi-local and the Kleinman-Bylander forms. A ghost state shows up as an extra pole of the KB curve.

3.4.7 Testing the pseudopotentials

Pseudopotential testing means using the pseudopotential in an environment that is different from the one used to generate it, and for which we have reliable data without the pseudopotential approximation. The simplest test is the atom in other configurations. For example after generating a pseudopotential for silicon with the s^2p^2 ground state configuration, one compares all-electron and pseudopotential calculations for the ionized configurations s^1p^2 and s^2p^1, and for the excited configurations s^1p^3, $s^1p^2d^1$ and $s^2p^1d^1$.

By editing the `atom.dat` file and running

```
atom/WORK$ ../atom.exe
```

the user can either see the all-electron results for a different configuration (`ae` keyword on the first line and deleting the second line, back-compatibility as you should have guessed) or see the result of a different configuration on the pseudopotential described in `pseudo.dat` (`pt` keyword on the first line, keeping the second line). Running

```
atom/WORK$ ../kb_test.exe
```

the user achieves the same type of test for the KB pseudopotential described by `pseudokb.dat`. If you are worried about ghost states, this is the most direct test: a ghost state near or below the occupied levels changes the eigenvalues and energies of the test configurations. A ghost far above them may go unnoticed here, but it still appears as an extra pole of the KB curve in the logarithmic derivative plots (Section 3.4.6). Note that `atom.exe` always rewrites `datafile.dat`, also in a pseudopotential test (`pt`); keep a copy of the all-electron one if you want to run `psd_gen.exe` or `plot_ln_show.exe` again.

A series of tests of ionization and promotion energies can be done with the original `atom.dat` by running

```
atom/WORK$ ../atom_test.exe
```

At the end the program will print a comparison of the ionization and promotion energies calculated with the all-electron potential, the semi-local pseudopotential and the pseudopotential in KB form, for silicon

```
Comparison of ionization energies (eV)
nl      all-electron   pseudo     KB-proj
3s       15.189       15.174       15.174
3p        7.840        7.835        7.836
Comparison of excitation energies (eV)
nl      nl     all-electron   pseudo      KB-proj      charge
```

```
   3s      3p        6.750        6.746        6.746      -1.00  1.00
   3s      3d       13.080       13.063       13.063      -1.00  1.00
   3p      3d        5.861        5.857        5.857      -1.00  1.00
```

Discrepancies of a few hundredths of an eV are to be expected; anything larger may indicate a bad pseudopotential.

4 The atom.dat input file

The description of that file, Section 4, was mostly written by Alberto García for the SIESTA documentation (version 3.2.2, January 2006) and adapted to describe the present code.

4.1 All-electron calculations

Assume we want to find the orbital eigenvalues, total energy, and/or charge density of Si in its ground state for some functional. You already have the compiled executables from SIESTA or from the ATOM code that you can find in the `Src/` folder. You should now go to a "working" directory and try the following: Our input file is named `atom.dat` and contains the lines (see Sect. 4.4 for more details):

```
  ae     Si   ground state all-electron
n=Si c=ca
     14.0       0.0       0.0     120.0       6.0      80.0
   3    2
   3    0     2.00      0.00
   3    1     2.00      0.00

#2345678901234567890123456789012345678901234567890      Ruler
```

We can run the calculation by typing *pathtoexecutables*`/atom.exe > atom.out` or *pathtoexecutables*`/atom_`*compilername*`.exe > atom.out` Here we are redirecting the output to `atom.out`. As described in Section 2.3, the present release of the code adds the compiler name by default, for example `atom_gfortran.exe` for code compiled with `gfortran`.

It is interesting to peruse the `atom.out` file. In particular, it lists the orbital eigenvalues (in Rydbergs, as almost every other energy in the program):

```
 nl    s      occ         eigenvalue    kinetic energy      pot energy

 1s   0.0    2.0000    -130.36911227     183.01377610    -378.73491457
 2s   0.0    2.0000     -10.14892691      25.89954256     -71.62102165
 2p   0.0    6.0000      -7.02876265      24.42537869     -68.74331196
 3s   0.0    2.0000      -0.79662738       3.23745216     -17.68692614
 3p   0.0    2.0000      -0.30705175       2.06135786     -13.62572529
```

(For a relativistic or spin-polarized calculation, there would be "up" and "down" flags in the `s` column above.)

4.2 Pseudopotential generation

To generate a pseudopotential for Si, using the Troullier-Martins scheme repeat the calculation with relativistic effects (although they are quite small) and use the LDA (Ceperley-Alder flavor). The input file contains the lines (see Sect. 4.4 for more details):

```
  pg     Si     Silicon
       tm2
n=Si c=car
     14.0       0.0       0.0     120.0       6.0      80.0
   3    4
   3    0     2.00      0.00
   3    1     2.00      0.00
   3    2     0.00      0.00
   4    3     0.00      0.00
   1.90     1.90     1.90     1.90

#2345678901234567890123456789012345678901234567890123456789012345678901234567890      Ruler
```

Note the two extra lines with respect to an all-electron calculation. The pseudopotential core radii for all channels are 1.90 bohrs. Even though they are nominally empty in the ground state, we include the $3d$ and $4f$ states in order to generate the corresponding pseudopotentials.

It is certainly cumbersome to type a fixed format input file. The good news is that running `/atom_`*compilername*`.exe` in a folder without the `atom.dat` and entering the chemical symbol of the element will generate such a file with default values, which can be modified for subsequent runs. Notice that the default values may not be the best, unless they are those of a tested pseudopotential (Section 3.3), so that file is just a template to be edited!

We can run the calculation by typing again *pathtoexecutables*`/atom.exe > atom.out` or *pathtoexecutables*`/atom_`*compilername*`.exe > atom.out` The `atom.out` file now has the spin-orbit effect and we see in the effect on eigenvalues that is small. The empty scattering orbitals may have a small positive eigenvalue.

nl	s	occ	eigenvalue	kinetic energy	pot energy
1s	0.5	2.0000	-130.51541377	0.00000000	-380.32924747
2s	0.5	2.0000	-10.18537055	0.00000000	-71.99156489
2p	-0.5	2.0000	-7.05518951	0.00000000	-69.08320077
2p	0.5	4.0000	-7.00775181	0.00000000	-68.71538886
3s	0.5	2.0000	-0.79937163	0.00000000	-17.74263422
3p	-0.5	0.6667	-0.30807126	0.00000000	-13.66178946
3p	0.5	1.3333	-0.30567131	0.00000000	-13.60785806
3d	-0.5	0.0000	0.00231324	0.00000000	-0.38199583
3d	0.5	0.0000	0.00231324	0.00000000	-0.38199582
4f	-0.5	0.0000	0.00340155	0.00000000	-0.35430233
4f	0.5	0.0000	0.00340155	0.00000000	-0.35430233

Running the pseudopotential generation by typing *pathtoexecutables*`/psd_gen.exe > psd.out` or *pathtoexecutables*`/psd_gen_`*compilername*`.exe > psd.out` will give you (just accept the defaults) a new default output where you can find the pseudopotential eigenvalues.

```
 nl     s      occ         eigenvalue    kinetic energy      pot energy

 1s    0.5    2.0000      -0.79936317       0.50555149      -3.74109919
 2p   -0.5    0.6667      -0.30805322       0.77243757      -3.26354801
 2p    0.5    1.3333      -0.30566087       0.76702417      -3.25195654
 3d   -0.5    0.0000       0.00231324       0.00233363      -0.10914167
 3d    0.5    0.0000       0.00231324       0.00233363      -0.10914167
 4f   -0.5    0.0000       0.00340158       0.00342993      -0.10122710
 4f    0.5    0.0000       0.00340158       0.00342993      -0.10122710
```

which are almost identical to the all-electron values.

4.3 Core corrections

The program can generate pseudopotentials with the non-linear exchange-correlation correction proposed in S. G. Louie, S. Froyen, and M. L. Cohen, Phys. Rev. B 26, 1738 (1982).

The pseudocore charge density equals the core charge density outside a given radius r_{pc} and has the smooth form

$$4\pi\rho_{pc}(r) = \exp\left(a + br^2 + cr^4\right)$$

inside that radius, with the parameters a, b and c calculated by the program so that the density and its first and second derivatives are continuous at r_{pc}. (The older form $Ar\sin(br)$ for $r^2\rho_{pc}$, with only the first derivative continuous, gave problems in the generation of GGA pseudopotentials and is no longer used.) The same smooth form replaces the core kinetic energy density inside its own radius.

To use core corrections in the pseudopotential generation the jobcode in the first line should be `pe` instead of `pg`.

The radius r_{pc} should be given in the sixth slot in the last input line (see above). If it is negative or zero (or blank), the radius is then computed using the fifth number in that line (`rcore_flag`, see the example input file above) and the following criterion: at r_{pc} the core charge density equals `rcore_flag`*(valence charge density). It is *highly recommended* to set an explicit value for the pseudocore radius r_{pc}, rather than letting the program provide a default.

If `rcore_flag` is input as negative, the full core charge is used. If `rcore_flag` is input as zero, it is set equal to one, which will be thus the default if `pe` is given but no numbers are given for these two variables.

The seventh number in that line plays the role of `rcore_flag` for the partial core kinetic energy density. If it is zero or blank it is set equal to one. If it is negative the full core kinetic energy density is used, which is only useful for tests with very large cutoffs.

The output file contains the radius used and the parameters a, b and c (`ac`, `bc`, `cc`) used for the matching.

4.4 Format of the file

For historical reasons, the input file is in a rigid column format. Fortunately, most of the column fields line up, so the possibility of errors is reduced. We will begin by describing in detail a very simple input file for an **all-electron calculation** for the ground state of Si.

The file for Si could be:

```
  ae     Si   ground state all-electron
 n=Si c=ca
      0.0       0.0       0.0       0.0       0.0       0.0
   3    2
   3    0     2.00      0.00
   3    1     2.00      0.00

#2345678901234567890123456789012345678901234567890      Ruler
```

- The first line specifies:
  - The calculation code (`ae` here stands for "all-electron").
  - A title for the job (here `Si ground state all-electron`).

  (format 3x,a2,a50)

- Second line:
  - Chemical symbol of the nucleus (here `Si`, obviously)
  - Exchange-correlation type. Here, `ca` stands for Ceperley-Alder. The "best" LDA choice should be `ca`. The Perdew-Wang 92 parametrization of the LDA is `pw`. It is also possible to use a gradient-corrected functional: `pb` indicates use of the PBE scheme by Perdew, Burke, and Ernzerhof (PRL 77, 3865 (1996)). The exchange-correlation energy and potential are computed using a modified Soler-Balbás package.
  - The character `r` next to `ca` is a flag to perform the calculation relativistically, that is, solving the Dirac equation instead of the Schrodinger equation. The full range of options is:
    - `s` : Spin-polarized calculation, non-relativistic.
    - `r`: Relativistic calculation, obviously polarized.
    - (blank) : Non-polarized (spin ignored), non-relativistic calculation.

  (format 3x,a2,3x,a2,a1,2x)

- Third line. Its use is somewhat esoteric and for most calculations it may contain just zeros (six numbers) as the code will use the default values.

  The first number is the nuclear charge, Z. You can invent elements with fractional charge. If it is zero the program will get it from the chemical symbol of the previous line. The second and third number are there for historical reasons. Their values are never used. The fourth to sixth numbers specify the integration mesh. Again if they are zero the code uses default values. The fourth number is the maximum range of the radius values $r_{\max}$. The sixth number B controls the increment between consecutive grid points. $r(n+1) \sim (1+1/B)r(n)$, in other words it takes about B points to increase the radius in the mesh by about e. The fifth number A controls the first non-zero point in the mesh. $r(1) = 0$, $r(2) \simeq exp(-A)/(Z * B)$.

  The full specification of the quasi-logarithmic mesh is $a = exp(-A)/Z$, $b = 1/B$, $r(i) = a * (exp(b * (i-1)) - 1)$, $r(i) < r_{\max}$.

The rest of the file is devoted to the specification of the electronic configuration:

- Fourth line:
  Number of core and valence orbitals. For example, for Si, we have $1s$, $2s$, and $2p$ in the core (a total of 3 orbitals), and $3s$ and $3p$ in the valence complex (2 orbitals).

  (format 2i5)

- Fifth, sixth... lines: (there is one line for each valence orbital)

  - `n` (principal quantum number)
  - `l` (angular momentum quantum number)
  - Occupation of the orbital in electrons.

  (format 2i5,3f10.3)

  (There are two f input descriptors to allow the input of "up" and "down" occupations in spin-polarized calculations (see example below)). There is a third float point number that can fix the eigenvalue of the orbital. It has not been used in years, Forget it exists.

  The valence orbitals can be given in any order. The same orbital cannot be given twice, nor can a valence orbital be one of the core orbitals (the program stops in those cases). Two valence orbitals with the same angular momentum and different principal quantum numbers are allowed.

The different treatment of core and valence orbitals in the input for an all-electron calculation is purely cosmetic. The program "knows" how to fill the internal orbitals in the right order, so it is only necessary to give their number. That is handy for heavy atoms... Overzealous users might want to check the output to make sure that the core orbitals are indeed correctly treated.

For compatibility with the old code, the file should end with an empty line. The new code doesn't care about it.

For a **pseudopotential test calculation**, the format is the same, except that the job code is `pt` instead of `ae` and that the second line with the "flavor" of a generation run (for example `tm2`, see below) must be present.

For a **pseudopotential generation run**, in addition to the electronic configuration chosen for the generation of the pseudopotentials (which is input in the same manner as above), one has to specify the "flavor" (generation scheme) and the set of core radii r_c for the construction of the pseudowavefunction. Here is an example for Si using the Troullier-Martins scheme:

```
  pg     Si     Pseudopotential
       tm2     2.00
n=Si c=ca
      0.0       0.0       0.0       0.0       0.0       0.0
   3    3
   3    0     2.00      0.00
   3    1     0.50      0.00
   3    2     0.50      0.00
     1.12      1.35      1.17       0.0       0.0       0.0

#2345678901234567890123456789012345678901234567890123456789012345678901234567890   Ruler
```

Apart from the `pg` (pseudopotential generation) job code in the first line, there are two extra lines:

- Second line:

  Flavor and radius at which to compute logarithmic derivatives for test purposes.

  The flavor can be one of : tm2 Improved Troullier-Martins

  (format 8x, a3, f9.3)

- The last line (before the blank line) specifies, as in SIESTA and the old versions of the code:

  – The values of the r_c in atomic units (bohrs) for the s, p, d, and f orbitals. They are given by angular momentum, whatever the order of the valence orbitals in the previous lines (it is a good practice to input the valence orbitals in the order of increasing angular momentum, so that there is no possible confusion).
  (format 4f10.5)

  – Two extra fields (2f10.5) which are relevant only if partial core corrections are used (see Sect 4.3). If they are zero or blank default values are used.

  – In the most recent versions of the code an extra field (f10.5) is used for the definition of the matching radius of the partial core kinetic energy density.

  The fields are read in fixed columns: the core radii in columns 1–40, `rcore_flag` in columns 41–50, r_{pc} in 51–60 and the kinetic energy density flag in 61–70. Any of the last three can be left blank. Versions of the code from 2021 to 2026 wrote these three numbers in columns 51–80 (with columns 41–50 blank); such files should be edited to move them ten columns to the left.

In the `tm2` example above, only s ,p, and d r_c's are given. Here is an example for Silicon in which we are only interested in the s and p channels for our pseudopotential:

```
  pg     Si     s and p only
       tm2     2.00
n=Si c=ca
      0.0       0.0       0.0       0.0       0.0       0.0
   3    2
   3    0     2.00      0.00
   3    1     2.00      0.00
     1.80      1.80      0.00       0.0       0.0       0.0

#2345678901234567890123456789012345678901234567890123456789012345678901234567890   Ruler
```

The new version of the code has an extension for the case a g orbital (or higher angular momentum) is present in the valence. In that case the line has one r_c field for each angular momentum up to the largest one in the valence, for example five fields (format 5f10.5) for a g orbital, and the core correction fields follow immediately after them (columns 51–80 in that example). Pseudopotentials are generated up to g ($l = 4$).

This completes the discussion of the more common features of the input file.

5 Computational details

Internally the code uses Rydberg atomic units (energies in Ry, lengths in bohr). The files for CPW2000 (Xy`_POTKB_F.DAT`) are written in Hartree.

5.1 Radial grid

The radial grid points are given by

$$r(n) = \frac{e^{-A}}{Z}\Big(e^{(n-1)/B} - 1\Big), \qquad n = 1, \ldots, n_{\max}, \tag{2}$$

where A and B are the constants given in `atom.dat` (defaults $A = 6$, $B = 200$, printed as `a` and `b` in the output), Z is the atomic number, and $n_{\max}$ is the last point with $r(n_{\max}) \le r_{\max}$, the maximum extension of the radial grid, which can also be set in `atom.dat` (default $r_{\max} = 120$ bohr). Near the origin the grid is almost linear,

$$r(n) \simeq (n-1)\frac{e^{-A}}{BZ}, \tag{3}$$

with a spacing inversely proportional to Z, as required by the $1s$ orbital. For large radii it is logarithmic,

$$\frac{r(n)}{r(n-1)} \simeq e^{1/B} \simeq 1 + \frac{1}{B}, \tag{4}$$

so the number of points is

$$n_{\max} \simeq 1 + B\ln\big(1 + r_{\max} Z e^{A}\big), \tag{5}$$

about 2700 for Si with the default values, and doubling $r_{\max}$ adds only $B\ln 2 \simeq 0.7B$ points. The accuracy is mostly controlled by B. The radial equations are integrated in the variable n, with $dr/dn = (r + e^{-A}/Z)/B$.

5.2 Solving radial equations

The non-relativistic (Schrödinger) and relativistic (Dirac, two components) radial equations are solved on the radial grid.

- At the start of the self-consistent atomic calculation no good guess of the eigenvalues is available, and the Schrödinger equation (also in relativistic calculations) is discretized by finite differences in n. The lowest eigenvalues and eigenvectors of the resulting tridiagonal matrix are found with the LAPACK routines `dstebz` (bisection) and `dstein` (inverse iteration). This requires no initial guess, and gives the starting eigenvalues and charge density.

- In the self-consistent iterations each bound state is found by integrating outwards from the origin up to the classical turning point, and inwards from a "practical infinity" (where $r\sqrt{V - \epsilon}$ is large) down to the same point. The two solutions are matched in value, and the eigenvalue is corrected from the discontinuity of the derivative (first-order perturbation theory), with the number of nodes used to bracket the eigenvalue.

- The integration uses a predictor-corrector method in the variable n: the predictor is the 5th order Adams-Bashforth and the corrector the 5th order Adams-Moulton formula. They need 5 starting points, which are obtained from the expansion of the wave-functions near the origin, $u(r) \simeq r^{l+1}(1 + a_1 r + a_2 r^2)$ for the Schrödinger equation, and $r^s(1 + a_1 r + a_2 r^2)$, $r^s(b_0 + b_1 r + b_2 r^2)$ for the major and minor components of the Dirac equation, with $s = \sqrt{\kappa^2 - (\alpha Z)^2}$. (The original code, from which this one derives, used $\alpha Z/4$ in that expansion, the factor of 2 between Hartree and Rydberg units having been applied in the wrong direction. It was found by comparing with the analytical eigenvalues of $-Z/r$.)

- The confined orbitals of the atomic basis (section 5.3) vanish at the cutoff radius r_c. They are found by integrating outwards up to r_c and adjusting the energy (secant method and bisection, with the number of nodes) until the wave-function vanishes at r_c.

- The perturbative polarization orbitals are solutions of a linear inhomogeneous equation. It is solved directly with the Numerov method in the variable n for $u(r)/\sqrt{dr/dn}$, integrating outwards from the origin and inwards from r_c, and matching at $0.9\,r_c$.

It has been checked that with increased radial grid quality the eigenvalues of $-Z/r$ converge to the analytical values, for both the Schrödinger and the Dirac equations.

5.3 Atomic orbitals

The code produces atomic orbitals that are used by the CPW2000 code for very good starting guesses of the plane-wave iterative diagonalization, for calculations in the atomic-orbital basis, and for the modified tight-binding (MTB) interpolation. They are written in real space to `plot_kb.dat`, with step 0.05 bohr (each one followed by a line with its basis set, zeta, kind, nodes, cutoff radius and energy), and, as Fourier transforms (section 5.4), to *Xy*`_POTKB_F.DAT`.

The atomic orbitals have angular momenta $l = 0, \ldots, l_a$, where l_a is the larger of the highest occupied l and $l_{\max} - 1$, $l_{\max}$ being the highest angular momentum of the pseudopotential, and the polarization orbital has $l_a + 1$ (not above f). For example, $l_a = 0$ for H, Li, Be and Na (polarization p), $l_a = 1$ for Mg, K, Ca and the p-block (polarization d), $l_a = 2$ for the transition metals and La, Ac (polarization f), and $l_a = 3$ for Ce–Yb, Th–U and Lu (no polarization orbital). A single-zeta (SZ) set has one orbital ψ for each l, a double-zeta (DZ) set adds a second orbital ϕ, and the "P" sets add one polarization orbital. The orbitals are similar to the ones used in the SIESTA code, and are of three types:

- The first orbital ψ for each l is a confined orbital, as the first SIESTA orbital, with a cutoff radius r_c beyond which it is zero. The radial equation is first solved at an energy $\epsilon_l + \delta E$, a shift δE above the eigenvalue ϵ_l of the valence state. The node of that solution is the "SIESTA radius" $r_S(l)$. In this code $\delta E = 0.02$ Ry (parameter `ESHIFT` in `atom_kb_basis_cutoff.f90`). The default of SIESTA was also 0.02 Ry in version 4.1, and is 0.01 Ry (0.136 eV) in version 5.4.2 (input `PAO.EnergyShift`). Where SIESTA uses $r_c = r_S(l)$, this code uses

$$r_c = \min\bigl(\max(F_l\, r_S(l), R_{\min}), R_{\max}\bigr), \tag{6}$$

  with $F_0 = 1.45$, $F_l = 1.15$ for $l > 0$, $R_{\min} = 6$ bohr and $R_{\max} = 9$ bohr. These constants minimize the difference between the atomic-orbital and plane-wave total energies of the elemental solids calculated with CPW2000, and are parameters in `atom_kb_basis_rule.f90`. The orbital ψ is the solution of the radial equation, with $\psi(r_c) = 0$, and with the soft confining potential of J. Junquera, O. Paz, D. Sánchez-Portal and E. Artacho, Phys. Rev. B 64, 235111 (2001),

$$W(r) = \begin{cases} A_W \exp\Bigl(-\frac{(1-f)\,r_c}{r - f r_c}\Bigr)\frac{1}{r_c - r}, & \text{if } f r_c < r < r_c, \\ 0, & \text{otherwise,} \end{cases} \tag{7}$$

  with $A_W = 40$ Ry and $f = 0.9$, the same values as the SIESTA defaults (soft confinement is the default of SIESTA 5.4.2, but not of version 4.1).

- The second orbital ϕ for angular momentum l differs from the SIESTA recipe ("split-norm"). It is similar to the energy derivative of the first orbital, $\partial\psi/\partial\epsilon$, in the spirit of the linearized methods (O. K. Andersen, Phys. Rev. B 12, 3060 (1975)), but vanishes at r_c, which $\partial\psi/\partial\epsilon$ cannot do for a confined orbital. It is the solution of

$$(H-\epsilon)\phi(r) = \psi(r) - \frac{w(r)\psi(r)}{\langle\psi \mid w \mid \psi\rangle}, \qquad \langle\psi \mid \phi\rangle = 0, \qquad \phi(r_c) = 0, \tag{8}$$

  where H includes the confining potential, ϵ is the eigenvalue of ψ, and $w(r) = \exp(-r_c/r)$ for $r < r_c$. It is calculated as $\phi = (\partial\psi_\lambda/\partial\lambda)/\langle\psi \mid w \mid \psi\rangle$, where ψ_λ is the confined orbital of $H + \lambda w$, by finite differences in λ with Richardson extrapolation. It has one node and the same cutoff radius as ψ.

- The polarization orbital is the perturbative polarization of the first orbital ψ of l_a, as in SIESTA (E. Artacho, D. Sánchez-Portal, P. Ordejón, A. García and J. M. Soler, Phys. Stat. Sol. (b) 215, 809 (1999)). It is the solution of

$$(H_{l_a+1} - \epsilon)\, u(r) = -r^p\, u_\psi(r), \qquad u(r_c) = 0, \tag{9}$$

  with the cutoff radius, confinement and eigenvalue ϵ of ψ ($u = rR$, and H_{l_a+1} is the radial hamiltonian for angular momentum $l_a + 1$). The SIESTA papers write the source with $p = 1$ (the response to an electric field), but the SIESTA code (version 5.4.2, routine `polarization` in `atom.F`) integrates the source in the grid variable without the factor $(dr/dn)^{3/2}$, which on its logarithmic grid is effectively $p = -1/2$. That gives a more compact orbital and lower total energies: for Ge (DZP, 40 Ha cutoff) the atomic-orbital energy is 102.7, 79.0 and 74.1 meV/atom above the plane-wave energy for $p = 1$, 0 and $-1/2$. The default is $p = -1/2$ (parameter `POLPOW` in `atom_kb_polarization.f90`), which reproduces the SIESTA polarization orbitals. For the elemental solids a single perturbative polarization orbital costs on average 0.7 to 2.6 mHa/atom, depending on the group of elements, compared with a confined $l_a + 1$ orbital plus its ϕ, with three (p), five (d) or seven (f) fewer orbitals per atom.

A bespoke basis set (number of orbitals and cutoff radius for each l) can be defined in the interactive mode of `kb_conv`. A cutoff radius of 0 asks for the orbital ϕ of the previous orbital, and -1 asks for the perturbative polarization orbital.

5.4 Fourier transforms

The *Xy*`_POTKB_F.DAT` file contains the local potential, the Kleinman-Bylander projectors, the core and valence charge densities, the core kinetic energy density and the atomic orbitals as functions of $q = k\,\delta q$, $k = 0, \ldots, 4000$, with $\delta q = 0.01$ bohr^{-1} ($q_{\max} = 40$ bohr^{-1}, enough for wave-function cutoffs up to 200 Ha). For an atomic orbital $R(r)$ of angular momentum l the file contains

$$\tilde{R}(q) = \sqrt{\frac{2l+1}{4\pi}}\sqrt{\frac{2}{\pi}}\int_0^{r_c} r^2 R(r)\, j_l(qr)\, dr, \tag{10}$$

where j_l is the spherical Bessel function. The integrals are calculated with the 8-point closed Newton-Cotes formula in the variable n, where the grid spacing is smaller than $0.1/q_{\max}$, and beyond that radius on a uniform grid with spacing $0.1/q_{\max}$, with the function interpolated by cubic splines.