Gravity from the GeoVac spectral action: continuum results and the discrete-substrate program

GeoVac Project

May 28, 2026

Abstract

We organize the GeoVac project's gravity arc (sprints G1 through G8, plus the discrete-substrate opening G4-3, all completed 2026-05-28) as a standalone document. On the Fock-projected unit three-sphere S^3 the Camporesi–Higuchi spinor spectrum $|\lambda_n| = n + 3/2$ with multiplicity $2(n+1)(n+2)$ produces a spectral zeta identity $\zeta_{\text{unit}}(-k) = 0$ for every $k \geq 0$, so the Chamseddine–Connes spectral action on S^3_R is two-term up to exponentially small corrections, and its two-term form has a formal extremum at $u_{\text{crit}} = R\Lambda = 1/\sqrt{6}$ (G1; the exact action is monotonic in R). Chamseddine and Connes [15] already showed that the expansion terminates after two terms at every order, up to rapidly decaying corrections ("all higher order terms vanish" on $S^3 \times S^1$); the present paper's contribution is an alternative algebraic proof of the vanishing, through the Bernoulli identity $B_{2k+1}(3/2) = (2k+1)/4^k$. The two-term exactness propagates to $S^3 \times S^1_\beta$ by heat-kernel factorization, recovering Einstein–Hilbert plus a cosmological constant with no higher-curvature corrections at any order (G2), and survives the $\beta \to \infty$ decompactification to a zero-temperature de Sitter vacuum on $S^3 \times \mathbb{R}_\tau$ with closed-form minimum action density $s_{\min} = -\Lambda/(6\sqrt{6})$ (G5). The two-term exactness is spinor-bundle-specific: scalar and transverse-traceless tensor heat traces inherit the standard Connes–Chamseddine infinite Seeley–DeWitt expansion (G3, closed-form proof $a_k^{\Delta} = 2\pi^2/k!$). On the Euclidean Schwarzschild cigar the S^2 Dirac heat trace at the spatial horizon section reproduces standard CC continuum behavior with $a_1 = -4\pi/3$ (G4-1), and the Sommerfeld–Cheeger conical-defect contribution combined with the replica method gives $S_{\text{BH}} = A\Lambda^2/(12\pi)$ (G4-2). The kinetic graviton sector of the $(1,1)$ irrep of $SO(4) = SU(2) \times SU(2)$ admits physical modes with positive kinetic eigenvalue after gauge-orbit classification via $V = i[X, D_0]$, satisfying the necessary condition for graviton dynamics (G6-Diag-Full). The subsequent Fierz–Pauli analysis (G6-FP) decomposes the $(1,1)$ subspace under the diagonal $SU(2)$ into $J{=}0 \oplus J{=}1 \oplus J{=}2$ and proves a J-blindness theorem: the spectral action $S^{(2)}$ has identical eigenvalue spectra across all three J-sectors, by Schur's lemma from the $SO(4)$ invariance of the trace functional. The Fierz–Pauli decomposition (TT vs. trace vs. longitudinal) therefore requires the metric identification $\delta D \leftrightarrow \delta g_{\mu\nu}$, not just the spectral action. The Newton constant $G_{\text{eff}} = 3\pi/\Lambda^2$ and the cosmological constant $\Lambda_{cc} = 6\Lambda^2$ emerge from the G2 coefficients (G7), traced over the rank-4 Dirac spinor bundle; G_{eff} then equals the entropy-route G_N of G4-2, with the standard cosmological-constant-scale gap inherited from Connes–Chamseddine. For an arbitrary cutoff function f with Mellin moments $\phi(s) = \int_0^\infty f(x) x^{s-1} dx$, the gravity predictions generalize to $G_{\text{eff}} = 3\pi/(\phi(1)\Lambda^2)$, $\Lambda_{cc} = 6\phi(2)/\phi(1) \cdot \Lambda^2$, and $R_{\text{crit}}\Lambda = \sqrt{\phi(1)/(6\phi(2))}$; *no* simple combination is cutoff-independent, classifying the cutoff function as external Class 1 calibration data (G8). Finally, we report on the extended discrete-substrate program. The substrate $\mathcal{G}_{\text{cigar}} = \mathbb{Z}_+(a)|_{N_\rho} \times \mathbb{Z}/N_\phi \times \text{Fock}(S^2, l_{\max})$ is operational at sprint scale through five closed sub-sprints (G4-3a / cleanup / b / c / d, plus the G4-3d-UV extension): Hermitian polar Laplacian built, variable warp realized, and the leading Weyl term $K(t) \sim A/(4\pi t)$ reproduced at the scale of the boundary correction (the 1% agreement once quoted at $t \approx 0.1$, $N_\phi = 192$ compared against the bulk term alone and omitted the $\sim 5.6\%$ Dirichlet perimeter term). The wedge-lattice proper conical-defect test (G4-3c-proper) has the correct sign at every apex angle α but shows that on the *scalar* side the literal Sommerfeld–Cheeger $(1/12)(1/\alpha - \alpha)$ extraction sits below the bulk-subtraction discretization floor at sprint

scale. On the *spinor* side, however — the natural object for the Dirac spectral triple — the situation reverses dramatically: the G4-4 sub-sprint sequence (14 sub-sprints across G4-4a/b/c/d/e/f) closes the warped Dirac program at sprint scale, identifying the spinor conical-defect tip coefficient $\Delta_K^{\mathrm{Dirac,tip}}(\alpha) = -(1/12)(1/\alpha - \alpha)$ (opposite sign from scalar SC, identical magnitude), with the discrete value converging to it as $a \to 0$ (the 2.1×10^{-5} agreement at $\alpha = 2/5$, $t = 2.0$ is the point where a radial sweep crosses the target) and extracting the replica-method derivative $d\Delta_K^{\mathrm{Dirac}}/d\alpha|_{\alpha=1} = +1/6$ at 96.69% recovery. The half-integer angular momentum + anti-periodic spinor BC cleanly extracts the conical-defect tip term where the integer scalar BC does not (BC-sector diagnostic: spinor 99.4% vs scalar 66.3% on the same substrate). The load-bearing structural quantities for the extended S_{BH} derivation (G4-5 discrete replica method + G4-6 full S_{BH} calculation) are quantitatively validated; the remaining work is integration over t with cutoff regularization, a bounded remaining program end-to-end.

Contents

1 Introduction **3**

2 The GeoVac substrate **5**

3 Spectral action on S^3_R: two-term exactness (G1) **6**

4 Thermal product on $S^3 \times S^1_\beta$ (G2) **7**

5 Bundle-specific two-term exactness (G3) **7**

6 Bekenstein–Hawking entropy from the conical defect (G4-1, G4-2) **8**
6.1 S^2 Dirac structural analysis (G4-1) . . . 8
6.2 Conical-defect / replica method (G4-2) . . . 9

7 Decompactified de Sitter vacuum on $S^3 \times \mathbb{R}_\tau$ (G5) **10**

8 Graviton diagnostic on the $(1,1)$ irrep (G6-Diag-Full) **10**
8.1 Setup . . . 10
8.2 Verification . . . 10

9 Fierz–Pauli decomposition within $(1,1)$ (G6-FP) **11**
9.1 Dimensions . . . 11
9.2 J-blindness theorem . . . 11
9.3 Implications for graviton physics . . . 12

10 Newton constant and cosmological constant from G2 (G7) **12**

11 Cutoff-function dependence (G8) **13**

12 Discrete-substrate program (G4-3) **14**
12.1 Discrete substrate . . . 14
12.2 Constant-warp factorization . . . 15
12.3 Sub-sprint sequence (updated) . . . 15
12.4 Honest scope at G4-3 closure . . . 16
12.5 Connection to G4-2 continuum derivation . . . 16
12.6 Connection to G8 cutoff dependence . . . 16

13 G4-4: sprint-scale closure of the warped Dirac program **17**
13.1 Architectural inputs from G4-3 . . . 17
13.2 G4-4a: constant-warp Dirac . . . 17
13.3 G4-4b: variable-warp Dirac . . . 17
13.4 G4-4c: conical-defect spinor and the headline tip coefficient . . . 18
13.5 G4-4d, G4-4e, G4-4f: Seeley-DeWitt, BC sectors, replica derivative . . . 19
13.6 Implications for the extended S_{BH} program . . . 20
13.7 G4-5: discrete replica method for S_{BH} . . . 20
13.7.1 G4-5a / G4-5a-refined: tip-only replica integration . . . 20
13.7.2 G4-5b: bulk Weyl extraction is structurally 4D, not 2D . . . 21
13.7.3 G4-5c: joint warp + conical-defect Dirac — bit-exact F6 + $r_h^2\Lambda^2/3$ at UV 21
13.7.4 G4-5d: cutoff dependence and sector-wise Mellin moment map . . . 22
13.7.5 G4-5 headline closure and implications for G4-6 . . . 23
13.7.6 Post-G4-5 follow-on push (v3.19.0): the $\alpha > 1$ Möbius form (a finite-a artifact) and the Mellin map . . . 24
13.7.7 Post-v3.19.0 follow-on thread (v3.20.0): closed-form per-t UV target and F14 closure . . . 28

14 R2: replica-weight-harmless convergence of the cigar-tip entropy (L6) **29**

15 Discussion: structural-skeleton-scope reading **33**

16 Open questions **35**

1 Introduction

The GeoVac project [3, 5, 6] is a spectral graph-theory program in which the Fock-projection of the unit three-sphere S^3 replaces the continuous Hilbert space of one-electron quantum mechanics with a discrete substrate $\mathcal{G}_{S^3}$ whose vertices carry the standard (n, l, m, s) quantum numbers. On these labels the continuum S^3 Laplacian acts exactly as the diagonal $SO(4)$ Casimir, with spectrum n^2-1 and multiplicity n^2, which Fock's energy shell maps to the hydrogenic spectrum; the discrete graph Laplacian $D-A$ on this substrate is a different operator, with spectrum in $[0, 8]$, that does not converge to it [1, 3, 4] (corrected 2026-09-25 from "the discrete graph Laplacian ... has spectrum n^2-1 [retracted 2026-09-25: graph-laplacian-converges-to-s3] exactly"). The framework's spectral triple $(\mathcal{A}_{GV}, \mathcal{H}_{GV}, D_{GV})$ has been shown in [6, 9] to satisfy the three J-relations at finite cutoff ($J^2 = -I$, $JD = +DJ$, $J\mathcal{O}J^{-1} = \mathcal{O}$; the chirality-doubled model truncation is even, sign triple $(-, +, +)$, with no KO label, while the true Camporesi–Higuchi window that converges is odd like the continuum; the order-zero/one conditions fail on both; corrected 2026-09-28 from "KO-dim 3") and to converge in the van Suijlekom state-space Gromov–Hausdorff distance to the round-S^3 Camporesi–Higuchi spectral triple as $n_{\max} \to \infty$.

This paper organizes a sprint sequence (G1 through G8 plus the opening of G4-3, all completed on 2026-05-28) that asks how this discrete substrate behaves under the Chamseddine–Connes spectral action [13, 16] when extended to higher-dimensional product geometries relevant to gravity. The motivating question was: *can the GeoVac substrate support a sensible gravity reading at the spectral action level?* The short answer is yes, with the standard caveats of the Connes–Chamseddine program (cutoff function as external input, cosmological-constant-scale gap inherited from standard CC). The detailed answer organizes naturally into a continuum-level arc (sprints G1 through G8) followed by the opening of a discrete-substrate arc (G4-3).

The continuum arc produces the following structural results.

1. **Two-term exactness identity** (G1): $\zeta_{\mathrm{unit}}(-k) = 0$ for every $k \geq 0$ on the Camporesi–Higuchi spinor spectrum. The Chamseddine–Connes spectral action on S^3_R becomes $S(R, \Lambda) =$

$\phi(3/2)(\Lambda R)^3 - \frac{1}{4}\phi(1/2)(\Lambda R) + O(e^{-\Lambda^2 R^2})$, with no higher-curvature power corrections at any order (all-order vanishing first shown by Chamseddine–Connes; the Bernoulli route is an alternative proof). The two-term form has a formal extremum at $R_{\text{crit}}\Lambda = 1/\sqrt{6}$; the exact action is monotonic in R. (SYMBOLIC PROOF; `tests/test_paper51_zeta_unit_neg_k.py`.)

2. **Thermal product factorization** (G2): the spectral action on $S^3 \times S^1_\beta$ factorizes by the heat-kernel identity $K_{S^3\times S^1}(t) = K_{S^3}(t) \cdot K_{S^1}(t)$ and reproduces Einstein–Hilbert plus a cosmological constant with the GeoVac substrate's two-term exactness preserved. (INTERNAL THEOREM.)

3. **Bundle-specific exactness** (G3): the two-term exactness is specific to the Camporesi–Higuchi spinor bundle on S^3. The scalar and transverse-traceless tensor heat traces on S^3 have the standard Connes–Chamseddine infinite Seeley–DeWitt expansion. We derive the closed form $a_k^\Delta = 2\pi^2/k!$ for the scalar Laplacian on unit S^3. (INTERNAL THEOREM — classical θ_3/Poisson asymptotics; the reduction and Taylor/normalization algebra are symbolic, the closed form matched numerically at 50 dps; `tests/test_paper51_scalar_ak.py`.)

4. S^2 **Dirac and Bekenstein–Hawking** (G4-1, G4-2): The S^2 Dirac with integer spectrum $|\lambda_n| = n+1$, multiplicity $4(n+1)$, has a full Connes–Chamseddine Seeley–DeWitt expansion (not two-term). Combined with the Sommerfeld–Cheeger conical-defect contribution $(1/12)(1/\alpha - \alpha)$ at the cigar tip, the replica method gives $S_{\text{BH}} = A\Lambda^2/(12\pi)$, with Newton constant $G_N = 3\pi/\Lambda^2$ (Planck-scale; equal to the action-route G_{eff} of G7 once the G2 trace is taken over the rank-4 Dirac bundle, as Wald's theorem requires in the two-term-exact Einstein-gravity sector — see §6). (INTERNAL THEOREM, continuum-level.)

5. **Decompactified de Sitter vacuum** (G5): the $\beta \to \infty$ decompactification of the G2 product gives a zero-temperature de Sitter vacuum on $S^3 \times \mathbb{R}_\tau$ with closed-form minimum action density $s_{\min} = -\Lambda/(6\sqrt{6})$. (SYMBOLIC PROOF; `tests/test_paper51_dS_vacuum.py`.)

6. **Graviton diagnostic** (G6-Diag-Full): within the $(1,1)$ irrep of $SO(4) = SU(2) \times SU(2)$, physical graviton modes (after gauge-orbit classification via $V = i[X, D_0]$) have positive kinetic eigenvalue at every tested $n_{\max} \in \{1,2,3\}$, satisfying the necessary condition for graviton dynamics on the discrete substrate. (MEASURED, sprint-scale.)

7. **Fierz–Pauli decomposition and J-blindness** (G6-FP): the $(1,1)$ subspace decomposes under the diagonal $SU(2) \subset SU(2)_L \times SU(2)_R$ as $J{=}0 \oplus J{=}1 \oplus J{=}2$ (trace + antisymmetric + graviton). The spectral-action second variation $S^{(2)}$ is J-blind: identical eigenvalues across all three sectors, with cross-sector multiplicities scaling as 1:3:5 (Schur's lemma). The Fierz–Pauli decomposition (TT vs. trace vs. longitudinal) requires the metric identification, not just the spectral action. This structurally closes the graviton program at the spectral-action level. (INTERNAL THEOREM; `tests/test_paper51_j_blindness.py`.)

8. **Newton constant and cosmological constant** (G7): $G_{\text{eff}} = 3\pi/\Lambda^2$ and $\Lambda_{cc} = 6\Lambda^2$ for Gaussian cutoff. The cosmological-constant-scale gap of $\sim 10^{120}$ between predicted $\Lambda_{cc}/M^2_{\text{Planck}} = 6$ and observed $\sim 10^{-120}$ is the standard Connes–Chamseddine issue, inherited not GeoVac-specific. (INTERNAL THEOREM.)

9. **Cutoff dependence** (G8): for arbitrary cutoff f, $G_{\text{eff}} = 3\pi/(\phi(1)\Lambda^2)$, $\Lambda_{cc} = 6\phi(2)/\phi(1) \cdot \Lambda^2$, $R_{\text{crit}}\Lambda = \sqrt{\phi(1)/(6\phi(2))}$. No simple combination is cutoff-independent. The cutoff function is external Class 1 calibration data alongside Yukawas, $Z_2/\delta m$ multi-loop QED counterterms, and the Born rule. (INTERNAL THEOREM; `tests/test_paper51_cutoff_mellin.py`.)

The discrete-substrate arc opens with G4-3:

9. **Discrete warped substrate** (G4-3): the near-horizon cigar geometry $D^2 \times S^2_{r_h}$ has discrete substrate $\mathcal{G}_{\text{cigar}} = \mathbb{Z}_+(a)|_{N_\rho} \times \mathbb{Z}/N_\phi \times \text{Fock}(S^2, l_{\max})$. The constant-warp factorization $K_{\text{cigar}}(t) = K_{D^2}(t) \cdot K_{S^2_{r_h}}(t)$ holds at the operator level. The conical-defect parameter $\alpha = N_\phi/N_0$ identifies the discrete analog of the off-shell tip deformation used in the replica method. Sub-sprints G4-3a through G4-6 are named as a substantial program commitment, well beyond sprint scale. (MEASURED, sprint-scale; `tests/test_paper51_L6_full.py` F1–F7 panel.)

The paper is organized as follows. Section 2 recalls the GeoVac substrate background needed for the gravity arc. Sections 3 through 11 document the continuum sprints in order, with Section 9 presenting the Fierz–Pauli decomposition and the J-blindness theorem. Section 12 opens the discrete-substrate program. Section 15 discusses the structural-skeleton-scope reading of the framework's gravity predictions, and section 16 lists open questions.

Companion documents. The detailed numerical machinery underlying each sprint is documented in Paper 28 §4.7–§4.17 [5]; this document organizes that material as a self-contained narrative for the gravity / spectral action / NCG audience.

2 The GeoVac substrate

The GeoVac substrate on unit S^3 is the Fock-projected discrete graph with vertex set indexed by (n, l, m) for $n \geq 1$, $0 \leq l < n$, $|m| \leq l$, and edges given by the standard ladder operators of the S^3 hyperspherical harmonics [3, 4]. The integer spectrum $\lambda_n = n^2 - 1$ is the continuum S^3 Laplacian's, carried exactly on the vertex labels by the diagonal $SO(4)$ Casimir; the graph Laplacian's own spectrum lies in $[0, 8]$ and does not converge to it (Paper 7). The hydrogen spectrum follows from $n^2 - 1$ through Fock's energy shell; the matched constant $\kappa = -1/16$ (its coincidence with a Fock-projection conformal weight is an Observation, not a fit) fixes only the graph's extremal eigenvalue.

The Camporesi–Higuchi spinor bundle on S^3 has Dirac operator D_{CH} with spectrum

$$|\lambda_n^{CH}| = n + 3/2, \qquad g_n = 2(n+1)(n+2), \tag{1}$$

where g_n is the multiplicity per n-shell. This is the spinor analog of the bosonic Laplacian spectrum $\lambda_n = -(n^2 - 1)$ on S^3.

The truncated spectral triple $(\mathcal{A}_{GV}, \mathcal{H}_{GV}, D_{GV})$ at finite cutoff $n_{\max}$ has Hilbert space $\mathcal{H}_{GV} = \bigoplus_{n \leq n_{\max}} \mathcal{H}_n$ with dimension $\sum_{n \leq n_{\max}} g_n = \frac{2}{3} n_{\max}(n_{\max}+1)(n_{\max}+2)$. At $n_{\max} = 3$ this gives 40 modes, matching Paper 2's structural cutoff $\Delta^{-1} = g_3^{\text{Dirac}} = 40$ [2].

Spectral action. The Chamseddine–Connes spectral action on a generic spectral triple is

$$S[D] = \text{Tr}\, f(D^2/\Lambda^2) = \sum_n g_n\, f(\lambda_n^2/\Lambda^2), \tag{2}$$

where f is a smooth positive cutoff function and Λ is a mass scale. For the GeoVac substrate on S^3_R (radius R), $\lambda_n = (n + 3/2)/R$, so $\lambda_n^2/\Lambda^2 = (n+3/2)^2/(R\Lambda)^2$, and the action depends only on the dimensionless combination $u = R\Lambda$.

Cutoff function. We will work primarily with the Gaussian cutoff $f(x) = e^{-x}$ until Section 11, where we generalize to arbitrary f. For the Gaussian cutoff, $\phi(s) := \int_0^\infty f(x) x^{s-1} dx = \Gamma(s)$, so $\phi(k) = (k-1)!$ for integer k and $\phi(1/2) = \sqrt{\pi}$, $\phi(3/2) = \sqrt{\pi}/2$.

3 Spectral action on S^3_R: two-term exactness (G1)

Theorem 3.1 (Spectral zeta identity at non-positive integers — SYMBOLIC PROOF; `tests/test_paper51_ze`
For the Camporesi–Higuchi spectrum on unit S^3, the spectral zeta function

$$\zeta_{\text{unit}}(s) := \sum_{n\geq 0} g_n (n+3/2)^{-2s} \tag{3}$$

satisfies $\zeta_{\text{unit}}(-k) = 0$ for every integer $k \geq 0$.

Proof. $\zeta_{\text{unit}}(s) = 2\sum_{n\geq 0}(n+1)(n+2)(n+3/2)^{-2s}$, which equals $2\sum_{n\geq 0}[(n+3/2)^2 - 1/4](n+3/2)^{-2s}$. This separates as $\zeta_{\text{unit}}(s) = 2\zeta_H(2s-2, 3/2) - \frac{1}{2}\zeta_H(2s, 3/2)$, where ζ_H is the Hurwitz zeta. By standard Hurwitz functional equations, $\zeta_H(-m, a) = -B_{m+1}(a)/(m+1)$ for non-negative integer m. At $s = -k$:

$$\zeta_{\text{unit}}(-k) = -\frac{2B_{2k+3}(3/2)}{2k+3} + \frac{B_{2k+1}(3/2)}{2(2k+1)}.$$

From the shift formula $B_n(x+1) = B_n(x) + nx^{n-1}$ and the identity $B_{2k+1}(1/2) = 0$ (standard for all $k \geq 0$), one obtains $B_{2k+1}(3/2) = (2k+1)/4^k$. Substituting: the first term gives $-2(2k+3)/[4^{k+1}(2k+3)] = -1/(2\cdot 4^k)$ and the second gives $(2k+1)/[4^k \cdot 2(2k+1)] = 1/(2\cdot 4^k)$. Their sum is zero at every $k \geq 0$. □ □

The identity in Theorem 3.1 forces the Mellin-transform expansion of $\mathrm{Tr}\, f(D^2/\Lambda^2)$ on S^3_R to truncate at low order.

Corollary 3.2 (Two-term exactness of the spectral action — SYMBOLIC PROOF; `tests/test_paper51_zeta_`
For Gaussian cutoff on S^3_R, the spectral action equals

$$S(R, \Lambda) = \phi(3/2)(\Lambda R)^3 - \frac{1}{4}\phi(1/2)(\Lambda R) + O(e^{-\Lambda^2 R^2}). \tag{4}$$

There are no higher-curvature power corrections at any order.

The corresponding extremum equation is $\frac{d}{du}[\phi(3/2)u^3 - \frac{1}{4}\phi(1/2)u] = 0$, giving $3\phi(3/2)u^2 = \frac{1}{4}\phi(1/2)$, or $u^2_{\text{crit}} = \phi(1/2)/(12\phi(3/2))$. For Gaussian cutoff this gives $u^2_{\text{crit}} = (\sqrt{\pi})/(12\cdot\sqrt{\pi}/2) = 1/6$, so $R_{\text{crit}}\Lambda = 1/\sqrt{6}$.

Remark 3.3 (Relation to Chamseddine–Connes 2008)**.** Chamseddine and Connes [15] showed that the spectral action on S^3 and $S^3 \times S^1$ is given by two terms up to rapidly decaying corrections, so that all higher Seeley–DeWitt terms vanish (checking a_4, a_6 against the heat-kernel expansion as "remarkable cancellations"). The Bernoulli argument of Theorem 3.1 is an alternative algebraic proof of the same all-order vanishing: the identity $B_{2k+1}(3/2) = (2k+1)/4^k$ (from $B_{2k+1}(1/2) = 0$ plus the shift formula) forces exact pairwise cancellation in the Hurwitz decomposition at every non-positive integer. Under the standard CC volume normalization the two-term-exact Dirac SD coefficients $a_0^{D^2} = 4\pi^2$, $a_1^{D^2} = -2\pi^2$ on S^3 are mixed-Tate periods over $\mathbb{Q}$ [14]; the scalar series $a_k^{\Delta} = 2\pi^2/k!$ likewise. The full inheritance corollary (GeoVac S^3 M2 sector lives in the strictly smaller pure-Tate sub-ring $\bigoplus_k \pi^{2k}\cdot\mathbb{Q}$) is recorded as Paper 32 [6] § VIII Corollary; the sprint memo `debug/sprint_mixed_tate_test_memo.md` (June 2026) contains the detailed verification.

Remark 3.4 (Structural uniqueness of two-term exactness)**.** The identity $\zeta_{\text{unit}}(-k) = 0$ is *universal* across all odd-dimensional spheres with the CH Dirac: verified symbolically for S^3, S^5, and S^7 through $k = 7$ (`debug/g4_structural_s5_comparison.py`). The mechanism on S^3 is transparent: $g_n = 2[(n+3/2)^2 - 1/4]$ decomposes into two Hurwitz zetas, and $B_{2k+1}(3/2) = (2k+1)/4^k$ (from $B_{2k+1}(1/2) = 0$ plus the shift formula at $3/2 = 1/2 + 1$) produces exact

pairwise cancellation at every order. On S^5 and S^7 the degeneracy polynomials (degree 4 and 6 respectively, all-even in $m = n + d/2$) produce analogous cancellations via the Bernoulli multi-step formula $B_{2m+1}(p + 1/2) = (2m+1)\sum_{l=0}^{p-1}(l + 1/2)^{2m}$.

The *physical* uniqueness of S^3 is not that it alone terminates, but that it terminates at the **minimum** number of terms. On S^{2m+1}, the spectral action has exactly $(m+1)$ power-law terms: cosmological constant (Λ^{2m+1}) + Einstein–Hilbert (Λ^{2m-1}) + Gauss–Bonnet-type (Λ^{2m-3}) + …, with no further corrections at any order. For S^3 $(m = 1)$: exactly **two** terms — *pure Einstein gravity*, no R^2 or higher-curvature corrections. For S^5 $(m = 2)$: three terms including R^2. For S^7 $(m = 3)$: four terms including R^2 and R^3. Only S^3 produces a spectral action that is purely Einstein–Hilbert + cosmological, making it the unique natural geometry for gravity without higher-curvature corrections.

Remark 3.5 (Honest scope of the extremum)**.** The IR-regime extremum at $u_{\text{crit}} = 1/\sqrt{6}$ is a property of the truncated two-term asymptotic expansion. The exact mode sum $\text{Tr}\, f(D^2/\Lambda^2)$ is monotonic in R; the extremum is a feature of the asymptotic, interpreted in standard CC fashion as a classical action selection.

4 Thermal product on $S^3 \times S^1_\beta$ (G2)

The Euclidean thermal product geometry $S^3 \times S^1_\beta$ has Dirac operator $D = D_{S^3} \otimes \sigma_1 + I \otimes \sigma_2\,(-i\partial_\tau)$ with $\tau \in [0, \beta)$, acting on the rank-4 bundle $\mathbb{C}^2 \otimes \mathbb{C}^2$ (two-component S^3 spinors tensored with $\mathbb{C}^2$; odd-dimensional S^3 has no chirality operator, so a rank-2 product is not a Dirac operator). Then $D^2 = D^2_{S^3} \otimes I - \partial^2_\tau$ and the heat kernel factorizes with a factor 2:

$$K_{S^3 \times S^1}(t) = 2\, K_{S^3}(t) \cdot K_{S^1}(t) = 2\, K_{S^3}(t) \cdot \frac{\beta}{\sqrt{4\pi t}}. \tag{5}$$

Corollary 4.1 (4D spectral action with exact two-term structure — INTERNAL THEOREM)**.** *The Chamseddine–Connes spectral action on $S^3_R \times S^1_\beta$ for Gaussian cutoff equals*

$$S(R, \beta, \Lambda) = \frac{\beta R^3}{2}\Lambda^4 - \frac{\beta R}{4}\Lambda^2 + O(\textit{exp small}). \tag{6}$$

The expression has the exact Einstein–Hilbert plus cosmological constant form, with no higher-curvature corrections at any order.

The classical de Sitter extremum (varying R at fixed Λ and β) gives $\frac{d}{dR}[\beta R^3/2 \cdot \Lambda^4 - \beta R/4 \cdot \Lambda^2] = 0$, yielding $R_{\text{crit}}\Lambda = 1/\sqrt{6}$ exactly as in G1. The matching against Einstein–Hilbert plus cosmological constant identifies the two terms with the GeoVac-determined values of Newton's constant and the cosmological constant (G7, Section 10).

5 Bundle-specific two-term exactness (G3)

The two-term exactness in Sections 3 and 4 is *specific to the Camporesi–Higuchi spinor bundle* on S^3. To prove this we exhibit closed forms for the heat traces on three sectors: the spinor sector (Dirac), the scalar Laplacian, and the transverse-traceless (TT) tensor sector.

Dirac sector (spinor bundle). From Paper 28 Theorem on two-term exactness:

$$K_{D^2}(t) = \frac{\sqrt{\pi}}{2}t^{-3/2} - \frac{\sqrt{\pi}}{4}t^{-1/2} + O(e^{-\pi^2/t}), \tag{7}$$

where the exponentially-small correction comes from Jacobi ϑ_2 modular transformation on the half-integer-shifted Camporesi–Higuchi spectrum.

Scalar sector. The bosonic Laplacian on unit S^3 has spectrum $\lambda_n^\Delta = n(n+2)$ with multiplicity $(n+1)^2$.

Theorem 5.1 (Closed-form Seeley–DeWitt coefficients for scalar Laplacian on unit S^3 — INTERNAL THEOREM (classical θ_3/Poisson asymptotics; algebra symbolic, closed form matched at 50 dps); `tests/test_paper51_scalar_ak.py`). *The heat trace satisfies*

$$K_\Delta(t) = \frac{\sqrt{\pi}}{4}\frac{e^t}{t^{3/2}} + O(e^{-\pi^2/t}) = \sum_{k=0}^{\infty} a_k^\Delta\, t^{k-3/2}\cdot(4\pi)^{-3/2}, \tag{8}$$

with $a_k^\Delta = 2\pi^2/k!$ *for all* $k \geq 0$.

The proof uses the Jacobi ϑ_3 modular transformation on the integer-spectrum bosonic sum, exploiting the structural difference between half-integer (Dirac) and integer (Laplacian) spectra.

TT-tensor sector. The Lichnerowicz Laplacian on transverse-traceless symmetric 2-tensors on unit S^3 has spectrum $(n+1)(n+3)-2$ with multiplicity polynomial in n. The heat trace has the form

$$K_{TT}(t) = e^{3t}\left[\frac{\sqrt{\pi}}{2}t^{-3/2} - 4\sqrt{\pi}t^{-1/2} + 4\right] + 6e^{2t} + O(e^{-\pi^2/t}), \tag{9}$$

which is pure integer ϑ_3 content: writing the spectrum as m^2-3 ($m \geq 3$) with multiplicity $2(m^2-4)$, the full-lattice sum is a ϑ_3 series, and the $+4$ and $6e^{2t}$ terms are the excluded $m = 1, 2$ modes.

Remark 5.2 (Structural distinction)**.** The two-term exactness theorem holds only for the Dirac sector (half-integer spectrum, ϑ_2 modular content). The scalar Laplacian (integer spectrum, ϑ_3 content) gives a complete infinite Seeley–DeWitt series $a_k^\Delta = 2\pi^2/k!$. The TT-tensor sector mixes both. Hence GeoVac's clean Einstein–Hilbert plus cosmological constant from G1/G2 lives in the spinor sector sourced by the Fock projection; the graviton sector inherits standard CC continuum expansion with infinite higher-curvature corrections.

6 Bekenstein–Hawking entropy from the conical defect (G4-1, G4-2)

The Euclidean Schwarzschild cigar geometry is

$$ds^2 = d\rho^2 + \rho^2 d\phi^2 + r(\rho)^2 d\Omega_2^2, \tag{10}$$

where (ρ, ϕ) parameterize the disk-with-tip and the ϕ-period is 2π for smooth tip (and $2\pi\alpha$ for $\alpha \neq 1$ off-shell). At the horizon $\rho = 0$, the spatial sphere has radius $r(0) = r_h = 2M$.

6.1 S^2 Dirac structural analysis (G4-1)

The S^2 Dirac operator has integer spectrum $|\lambda_n^{S^2}| = n+1$ with multiplicity $4(n+1)$ for $n \geq 0$. The corresponding squared-Dirac spectrum is $\lambda_n^2 = (n+1)^2$ with the same multiplicity, summing to the heat trace

$$K_{S^2}(t) = \sum_{n\geq 0} 4(n+1)e^{-(n+1)^2 t}. \tag{11}$$

This is the standard CC infinite Seeley–DeWitt expansion (*not* two-term exact): the integer Dirac spectrum on S^2 uses the ϑ_3 branch, not the half-integer ϑ_2 branch specific to the S^3 Camporesi–Higuchi spectrum. The leading behavior is $K_{S^2}(t) \sim 2/t$ (Weyl law), with $a_1^{S^2\ \text{Dirac}} = -4\pi/3$ as the first curvature correction (verified numerically).

Remark 6.1 (Pattern crystallizes)**.** The two-term exactness on the GeoVac substrate is *not* a generic property of S^3. It is specific to the half-integer Camporesi–Higuchi spinor spectrum coupled to the Jacobi ϑ_2 modular structure. Integer-spectrum bundles (scalar on S^3, integer Dirac on S^2, TT-tensor mixture) all show the standard infinite Seeley–DeWitt series.

6.2 Conical-defect / replica method (G4-2)

The Sommerfeld–Cheeger formula [20, 17] for the heat trace of the 2D Laplacian on a disk with conical tip of apex angle $2\pi\alpha$ is

$$K_{\text{cone}}(t) = K_{\text{bulk}}(t) + \frac{1}{12}\left(\frac{1}{\alpha} - \alpha\right) + O(t). \tag{12}$$

The leading $(1/12)(1/\alpha - \alpha)$ is the topological conical-defect contribution at the tip, vanishing at $\alpha = 1$ (smooth tip).

Combined with the S^2 Dirac heat trace from G4-1, the tip contribution to the cigar action at Gaussian cutoff is

$$I_E^{\text{conical}}(\alpha, \Lambda, r_h) = \frac{r_h^2 \Lambda^2}{6}\left(\frac{1}{\alpha} - \alpha\right) + O(1/\Lambda^2). \tag{13}$$

Applying the replica method $S_{\text{BH}} = -\frac{dI_E^{\text{conical}}}{d\alpha}|_{\alpha=1}$:

$$S_{\text{BH}} = -\left.\frac{dI_E^{\text{conical}}}{d\alpha}\right|_{\alpha=1} = \frac{r_h^2\Lambda^2}{3} = \frac{A\Lambda^2}{12\pi}, \tag{14}$$

with horizon area $A = 4\pi r_h^2$.

Matching $S_{\text{BH}} = A/(4G_N)$**:**

$$G_N = \frac{3\pi}{\Lambda^2}, \tag{15}$$

Planck-scale Newton constant. For the Gaussian cutoff this equals the $G_{\text{eff}} = 3\pi/\Lambda^2$ of the Einstein–Hilbert route (G7, §10) once the G2 trace is taken over the rank-4 Dirac bundle.

Remark 6.2 (The former factor of 2 was the G2 spinor rank)**.** An earlier printing traced G2 over a rank-2 bundle, giving $G_{\text{eff}} = 6\pi/\Lambda^2$, twice the entropy-route value; the rank-4 Dirac trace removes the factor. A first reading had attributed it to a scalar-vs-Dirac conical-defect coefficient. That reading was *incorrect*: the 2D Dirac and scalar cones share the conical-coefficient *magnitude* $(1/12)\,|1/\alpha - \alpha|$ (the discrete spinor extraction carries the opposite sign in the anti-periodic convention; §13), confirmed three independent ways — by central charge ($c = 1$ for both a free 2D Dirac fermion and a free real scalar), by Fursaev–Miele [24] ("spin 1/2 resembles the scalar case"), and by the discrete-substrate spinor extraction of G4-4 ($-1/12$, approached as $a \to 0$, the anti-periodic half-integer spectrum being structurally essential). Wald's theorem [26] required the two routes to agree: the Dirac-sector spectral action is two-term exact (G1–G2: Einstein–Hilbert plus cosmological constant, no higher curvature), so the geometry is pure Einstein gravity, where the G in the action and the G in $S_{\text{BH}} = A/(4G)$ are *identically equal*. The mismatch was the 4D Dirac spinor trace $a_0 = 4 = 2 \otimes 2$, which G4-2's factorized $D^2 \times S^2$ setup carried and the earlier G2 trace did not; with it the two routes agree for the Gaussian cutoff. The load-bearing structural finding remains the emergence of $A\Lambda^2$ from the conical tip via the replica method.

7 Decompactified de Sitter vacuum on $S^3 \times \mathbb{R}_\tau$ (G5)

The $\beta \to \infty$ decompactification of G2 sends $S^1_\beta \to \mathbb{R}_\tau$. The heat-kernel density satisfies $K_{\mathbb{R}}(t) = 1/(2\sqrt{\pi t})$, matching K_{S^1}/β in the limit. The spectral action density becomes

$$s(R, \Lambda) = \frac{R^3}{2}\Lambda^4 - \frac{R}{4}\Lambda^2, \tag{16}$$

with extremum at $R_{\text{crit}}\Lambda = 1/\sqrt{6}$ (same as G1, G2).

Theorem 7.1 (Closed-form de Sitter vacuum action density — SYMBOLIC PROOF; `tests/test_paper51_dS_`
The minimum action density at the de Sitter extremum is

$$s_{\min} = -\frac{\Lambda}{6\sqrt{6}} \quad (at\ R_{\text{crit}} = 1/(\sqrt{6}\Lambda)). \tag{17}$$

The decompactified vacuum is the zero-temperature de Sitter vacuum on $S^3 \times \mathbb{R}_\tau$, with the Stefan–Boltzmann T^4 thermal correction sitting in the exponentially-small (IR) part of the heat trace, structurally distinct from the UV asymptotic.

8 Graviton diagnostic on the $(1, 1)$ irrep (G6-Diag-Full)

The natural place to look for gravitons on the GeoVac substrate is within the $(j_L, j_R) = (1, 1)$ irrep of $SO(4) = SU(2)_L \times SU(2)_R$. This is a 9-dimensional irrep at each n-shell, containing the symmetric traceless rank-2 tensor representation that hosts spin-2 graviton modes.

8.1 Setup

For variations $\delta D = X$ around the unperturbed Camporesi–Higuchi Dirac D_0, the kinetic eigenvalue is $A_\lambda = a_\lambda(4\lambda^2/\Lambda^4 - 2/\Lambda^2)$ in Gaussian-cutoff spectral action. The cross-sector modes are gauge-orbit tangents

$$V = i[X, D_0], \tag{18}$$

while the within-sector modes are physical.

8.2 Verification

At $n_{\max} \in \{1, 2, 3\}$, the physical (within-sector) $(1, 1)$ modes have positive kinetic eigenvalue:

- $|\lambda| = 5/2$ ($n_{\max} = 1$): $A_\lambda = +0.127$ (at $\Lambda^2 = 6$)
- $|\lambda| = 7/2$ ($n_{\max} = 2$): $A_\lambda = +0.133$
- $|\lambda| = 9/2$ ($n_{\max} = 3$): $A_\lambda = +0.066$

No sign changes or instabilities are observed; the Gaussian-regulator suppression behavior is consistent with standard CC continuum expectations. The cross-sector (gauge) blocks have mixed signs (gauge-orbit curvature, not physical). A 5-point finite-difference stencil verifies the analytical eigenvalues to relative difference 10^{-9}.

Proposition 8.1 (Necessary condition for graviton dynamics — MEASURED, sprint-scale ($n_{\max} \leq 3$)). *The GeoVac substrate admits physical* $(1, 1)$ *modes with positive kinetic eigenvalue at every tested* $n_{\max} \in \{1, 2, 3\}$*, after gauge-orbit classification via* $V = i[X, D_0]$*. The necessary condition for graviton dynamics is satisfied.*

Honest scope. This is a necessary condition, not sufficient. The Fierz–Pauli decomposition within the $(1,1)$ irrep is addressed in Section 9.

9 Fierz–Pauli decomposition within $(1,1)$ (G6-FP)

The $(1,1)$ irrep of $SO(4) = SU(2)_L \times SU(2)_R$ decomposes under the diagonal $SU(2) \subset SU(2)_L \times SU(2)_R$ as

$$(1,1) = J{=}0 \ \oplus \ J{=}1 \ \oplus \ J{=}2. \tag{19}$$

Physically, $J{=}0$ (1 dim) is the trace/conformal mode, $J{=}1$ (3 dim) is the antisymmetric tensor (not a metric perturbation), and $J{=}2$ (5 dim) is the symmetric traceless tensor containing the graviton.

9.1 Dimensions

$n_{\max}$	$\dim H$	$J{=}0$	$J{=}1$	$J{=}2$	Total
1	16	6	14	26	46
2	40	16	40	72	128
3	80	26	66	118	210
4	140	36	92	164	292

Convention note: $\dim H$ here sums the CH spinor degeneracies with the shell index starting at $n = 0$ ($\dim H = \frac{2}{3}(n_{\max}{+}1)(n_{\max}{+}2)(n_{\max}{+}3)$), while §II counts shells from $n = 1$ — so §II's "$n_{\max} = 3$ gives 40 modes" corresponds to this table's $n_{\max} = 2$ row. Same space, shell labels offset by one.

Within-sector modes (where $\|[D_0, V]\|^2 = 0$) scale as 1:1:3 across $J = 0, 1, 2$ rather than the naïve 1:3:5, because Hermitianisation $(V + V^T)/2$ eliminates 4 of 9 modes per within-sector block (each block contributes $1 + 1 + 3 = 5$ modes instead of $1 + 3 + 5 = 9$).

9.2 J-blindness theorem

Theorem 9.1 (J-blindness of $S^{(2)}$ — INTERNAL THEOREM; `tests/test_paper51_j_blindness.py`). *The Gaussian spectral-action second variation $S^{(2)}$ restricted to the $(1,1)$ subspace of* $\mathrm{Herm}(H_{n_{\max}})$ *has identical eigenvalue spectra across $J = 0, 1, 2$. For cross-sector modes, eigenvalue multiplicities scale as* $\dim(J) = 1{:}3{:}5$.

Proof. The spectral action $S = \mathrm{Tr}(f(D^2/\Lambda^2))$ is invariant under unitary conjugation $D \mapsto UDU^*$, hence under the $SO(4)$ symmetry of D_0 and in particular under the diagonal $SU(2)$. The second variation $S^{(2)}$ is therefore equivariant under diagonal $SU(2)$. By Schur's lemma, $S^{(2)}$ is proportional to the identity on each irreducible J-component within a given sector pair (a, b), with proportionality constant

$$w(\lambda_a, \lambda_b) = \begin{cases} e^{-\lambda_a^2/\Lambda^2}(4\lambda_a^2/\Lambda^4 - 2/\Lambda^2) & \lambda_a = \lambda_b, \\ \dfrac{(\lambda_a + \lambda_b)^2}{\Lambda^4} \dfrac{\Lambda^2(a_a - a_b)}{\lambda_b^2 - \lambda_a^2} - \dfrac{a_a + a_b}{\Lambda^2} & \lambda_a \neq \lambda_b, \end{cases} \tag{20}$$

where $a_k = e^{-\lambda_k^2/\Lambda^2}$. This weight depends only on the eigenvalues λ_a, λ_b of D_0, not on the angular quantum numbers or on J. □

Numerical verification: the analytical formula (20) matches the five-point central-difference stencil to relative error 1.24×10^{-6} at $n_{\max} = 2$ (archived sprint measurement; the live regression `tests/test_paper51_j_blindness.py` checks the stencil match at the 10^{-4} level at $n_{\max} = 1$ — stencil noise sets that floor — and the eigenvalue-set/multiplicity identities of the theorem at $n_{\max} = 1, 2$).

Eigenvalue table at $n_{\max} = 3$, $\Lambda^2 = 6$. At $n_{\max} = 3$, the $J = 2$ sector has dim = 118 and the $S^{(2)}$ eigenvalue clusters are:

$S^{(2)}$ value	J=0 mult	J=1 mult	J=2 mult	Ratio
+0.1334	2	2	6	1:1:3
+0.1274	2	2	6	1:1:3
+0.1214	4	12	20	1:3:5
+0.0961	4	12	20	1:3:5
+0.0656	2	2	6	1:1:3
−0.0253	4	12	20	1:3:5
−0.0743	4	12	20	1:3:5
−0.1594	4	12	20	1:3:5

Each eigenvalue appears in *every* J-sector at the same value. Within-sector modes ($\|[D_0, V]\|^2 = 0$) show the 1:1:3 multiplicity pattern; cross-sector modes show the full 1:3:5.

9.3 Implications for graviton physics

The J-blindness theorem draws a sharp scope boundary:

- **Accessible from $S^{(2)}$:** existence of positive kinetic modes (Prop. 8.1), bit-exact integer kinetic spectrum $(2k)^2$, correct Seeley–DeWitt a_1 sign structure, 2% approach to the Lichnerowicz ratio 13/6.
- **Not accessible from $S^{(2)}$:** distinction between TT (2 DOF), longitudinal (3 DOF), and trace (1 DOF) within $(1, 1)$; the Fierz–Pauli mass structure; the identification of which modes are physical gravitons.
- **Source of the missing structure:** the metric identification $\delta D \leftrightarrow \delta g_{\mu\nu}$ (inner fluctuation / Connes–Chamseddine dictionary), which breaks $SO(4) \to SO(3)$ and distinguishes symmetric from antisymmetric perturbations. Whether this identification arises on the discrete substrate is not shown here; Paper 38's state-space Gromov–Hausdorff convergence concerns the commutative factor and does not supply it.

The theorem unifies three earlier diagnostics: GD-5's irrep-blindness (special case for within-sector modes), GD-1's $9 \to 2$ reduction as a continuum phenomenon (consistent with J-blindness: the spectral action alone does not reduce degrees of freedom within $(1, 1)$), and GD-4's helicity question (irrelevant at the $S^{(2)}$ level since J-blindness makes all helicities equivalent).

Structural closing of G6 at the spectral-action level. The J-blindness theorem shows that $S^{(2)}$ has extracted all information available from the $(1, 1)$ subspace via the spectral action alone. Further graviton physics (TT isolation, propagator, linearised Einstein equations) requires the metric identification, which is a propinquity-level construction.

10 Newton constant and cosmological constant from G2 (G7)

Matching the G2 spectral action coefficients in Equation (6) to Einstein–Hilbert plus a cosmological constant,

$$S_{\text{EH}} = -\frac{1}{16\pi G_{\text{eff}}} \int R_{\text{scalar}} \sqrt{g} + \frac{\Lambda_{cc}}{8\pi G_{\text{eff}}} \int \sqrt{g}, \tag{21}$$

with the volume forms on $S^3_R \times S^1_\beta$ giving Vol $= 2\pi^2 R^3 \beta$ and $\int R_{\text{scalar}} \sqrt{g} = 12\pi^2 R\beta$, the coefficient matching gives

$$\Lambda^2 \text{ term}: \quad G_{\text{eff}} = 3\pi/\Lambda^2, \tag{22}$$

$$\Lambda^4 \text{ term}: \quad \Lambda_{cc} = 6\Lambda^2. \tag{23}$$

Proposition 10.1 (Newton constant and cosmological constant — INTERNAL THEOREM). *The GeoVac spectral action on $S^3_R \times S^1_\beta$ with Gaussian cutoff produces Newton constant $G_{\text{eff}} = 3\pi/\Lambda^2$ (Planck-scale) and cosmological constant $\Lambda_{cc} = 6\Lambda^2$.*

The cosmological-constant-scale gap. Setting $\Lambda = M_{\text{Planck}}$ gives $\Lambda_{cc}/M^2_{\text{Planck}} = 6$, in contrast to the observed $\sim 10^{-120}$. This 10^{120} gap is the standard Connes–Chamseddine issue [16] and is inherited by GeoVac, not produced by GeoVac. The framework does not *add* a cosmological-constant problem; it inherits the standard one.

Background extremality consistency check. The Gaussian integral $\int_0^\infty (4u^2-2)e^{-u^2}\,du = 4\sqrt{\pi}/4 - 2\sqrt{\pi}/2 = 0$ confirms G6-Diag-Full's physical $(1,1)$ eigenvalue formula vanishes against the Gaussian-weighted background trace, consistent with background-extremality of the spectral action.

11 Cutoff-function dependence (G8)

Sprints G1 through G7 assume the Gaussian cutoff $f(x) = e^{-x}$. The question we ask in G8 is: how do the gravity predictions change for arbitrary cutoff f, and which combinations (if any) are cutoff-independent?

Mellin moments. For arbitrary cutoff f, define

$$\phi(s) := \int_0^\infty f(x)\, x^{s-1}\, dx \qquad \text{(Mellin transform of } f\text{)}. \tag{24}$$

The relevant moments for spectral-action matching are $\phi(1)$ (Einstein–Hilbert scale) and $\phi(2)$ (cosmological-constant scale).

General formulas.

Theorem 11.1 (Cutoff-dependent gravity predictions — INTERNAL THEOREM; `tests/test_paper51_cuto`
For arbitrary smooth positive cutoff f with finite Mellin moments $\phi(1), \phi(2)$, the GeoVac spectral action on $S^3_R \times S^1_\beta$ produces

$$G_{\text{eff}}(f,\Lambda) = \frac{3\pi}{\phi(1)\,\Lambda^2}, \tag{25}$$

$$\Lambda_{cc}(f,\Lambda) = \frac{6\,\phi(2)}{\phi(1)}\,\Lambda^2, \tag{26}$$

$$R_{\text{crit}}\,\Lambda = \sqrt{\frac{\phi(1)}{6\,\phi(2)}}. \tag{27}$$

Table 1: Gravity predictions for three cutoff functions.

Cutoff $f(x)$	$\phi(1)$	$\phi(2)$	$G_{\text{eff}}\Lambda^2$	Λ_{cc}/Λ^2	$R_{\text{crit}}\Lambda$
Gaussian e^{-x}	1	1	3π	6	$1/\sqrt{6}$
Polynomial e^{-x^2}	$\sqrt{\pi}/2$	$1/2$	$6\sqrt{\pi}$	$6/\sqrt{\pi}$	$\sqrt{\sqrt{\pi}/6} \approx 0.544$
Sharp $\Theta(1-x)$	1	$1/2$	3π	3	$1/\sqrt{3}$

Three specific cutoffs.

No cutoff-independent gravity prediction. Direct algebra on Theorem 11.1 shows that every combination of G_{eff} and Λ_{cc} depends on $\phi(1)$ and/or $\phi(2)/\phi(1)$:

- $G_{\text{eff}} \cdot \Lambda^2 = 3\pi/\phi(1)$ (depends on $\phi(1)$)
- $\Lambda_{cc}/\Lambda^2 = 6\,\phi(2)/\phi(1)$ (depends on ratio)
- $G_{\text{eff}} \cdot \Lambda_{cc} = 18\pi\,\phi(2)/\phi(1)^2$ (depends on both)
- $R_{\text{crit}}\Lambda = \sqrt{\phi(1)/(6\,\phi(2))}$ (depends on ratio)

Class 1 calibration data. The cutoff function f is *external Class 1 calibration data*, alongside the standard-model Yukawa couplings, the Z_2 and δm multi-loop QED renormalization counterterms, and the Born rule. The framework determines:

- the structural form of $S[D]$ (spectrum, Seeley–DeWitt coefficients);
- the proportionality of G_{eff} and Λ_{cc} to specific ϕ-moments;
- the functional form of $G_{\text{eff}}(f), \Lambda_{cc}(f), R_{\text{crit}}(f)$.

The framework does *not* determine:

- the numerical value of G_{eff} or Λ_{cc} in absolute units;
- which cutoff function to use.

This is consistent with standard CC [13, 16]: the cutoff function is external input; the framework determines the structural shape of $S[D]$.

12 Discrete-substrate program (G4-3)

Sections 3 through 11 produce the continuum gravity arc on S^3 and product geometries. G4-3 opens the *discrete-substrate* arc: rebuild the gravity-arc results on GeoVac's discrete substrate rather than at the continuum level. Adjacent discrete-substrate approaches to black-hole entropy (causal set theory [30], LQG [28], CDT [29]) provide complementary comparison points; the GeoVac discrete-substrate program targets the spectral-action / Camporesi–Higuchi Dirac route on a warped-product geometry built from the Fock projection on S^2.

12.1 Discrete substrate

For the cigar's near-horizon limit with constant warp $r(\rho) = r_h$:

$$\mathcal{G}_{\text{cigar}}(a, N_\rho, N_\phi, l_{\max}, r_h) = \mathbb{Z}_+(a)|_{N_\rho} \times \mathbb{Z}/N_\phi \times \text{Fock}(S^2, l_{\max}), \tag{28}$$

where:

- $\mathbb{Z}_+(a)|_{N_\rho}$ is the N_ρ-truncated nonnegative integer lattice with spacing a, $\rho_k = ka$;
- $\mathbb{Z}/N_\phi$ is the discrete ϕ-circle with N_ϕ sites;
- $\text{Fock}(S^2, l_{\max})$ is the standard S^2 Fock projection with $l \le l_{\max}$ angular cutoff.

The conical-defect parameter is $\alpha = N_\phi/N_0$ where N_0 is the reference smooth-tip count.

12.2 Constant-warp factorization

At $r(\rho) = r_h$ constant, the Laplacian factorizes:

$$\Delta_{\text{cigar}} = \Delta_{D^2}(\rho, \phi) + \frac{1}{r_h^2}\,\Delta_{S^2}. \tag{29}$$

The two factors act on independent sectors, so by construction:

$$K_{\text{cigar}}(t) = K_{D^2}(t) \cdot K_{S^2_{r_h}}(t). \tag{30}$$

This factorization holds at the operator level on the discrete substrate.

12.3 Sub-sprint sequence (updated)

G4-3 opens an extended discrete-substrate sub-sprint sequence. Status as of this version of the paper:

- **G4-3a**: constant-warp substrate + scoping (sprint-scale, DONE)
- **G4-3a-cleanup**: Hermitian discrete polar Laplacian via $u = \sqrt{\rho}\, f$ substitution and symmetric finite differences. Bessel-zero continuum convergence verified; Hermiticity bit-exact (DONE)
- **G4-3b**: variable warp $r(\rho) = r_h\sqrt{1 + (\rho/r_h)^2}$ for asymptotic Schwarzschild. Asymmetric tridiagonal radial operator preserves self-adjointness numerically; heat-trace ratios in physical range (DONE)
- **G4-3c** (naive N_ϕ sweep): tested whether the original sweep at fixed period 2π extracts the Sommerfeld–Cheeger $(1/12)(1/\alpha - \alpha)$ coefficient. DONE, PARTIAL-POSITIVE: framework operational but the sweep varies angular resolution, not apex angle. Proper wedge follow-on indicated.
- **G4-3c-proper**: wedge-lattice with apex angle $2\pi\alpha$ and fixed angular resolution $h_\phi = 2\pi/N_0$, $N_\phi(\alpha) = \alpha N_0$. Sign of the topological residual correct at every α; magnitude $\sim 1/4$ of the SC target with strong $\alpha < 1$ vs $\alpha > 1$ asymmetry attributable to UV-truncation mismatch between branches. DONE, NEGATIVE-WITH-STRUCTURAL-DIAGNOSIS; sharpens G4-5 (discrete replica method) as the proper SC-extraction target.
- **G4-3d**: continuum-limit heat-kernel asymptotics verification via a-sweep at fixed $N_\phi = 24$. Weyl law $K(t) = A/(4\pi t) + \cdots$ verified in the IR regime ($t \in \{0.5, 1.0\}$) at the 5–7% level; UV regime saturated by angular truncation (DONE, POSITIVE-IR).
- **G4-3d-UV**: same a-sweep substrate refined in $N_\phi \in \{24, 48, 96, 144, 192\}$. The bulk Weyl term is matched within 1% at $t = 0.1$, $N_\phi = 192$ and within 7% across $t \in [0.01, 0.1]$, a comparison that omits the Dirichlet perimeter term (about 5.6% at $R = 10$, $t = 0.1$). The G4-3d Sec. 5 prediction that the small-t saturation is angular truncation (NOT Hermiticity) is confirmed quantitatively at the 3× gain factor on the $N_\phi = 96$ vs $N_\phi = 24$ panel (DONE, POSITIVE-UV-VERIFIED). A high-m overshoot mechanism is identified at fine N_ϕ (ratio $4/\pi^2$ between discrete and continuum azimuthal Laplacian at the truncation edge), structural to the uniform-ϕ FD angular discretization.
- **G4-4**: warped Dirac spectrum on the discrete substrate (a major sub-program beyond sprint scale). Architectural scoping memo (G4-4 first-move) closed; sub-sprint sequence G4-4a/b/c/d/e/f named with load-bearing falsifiers F1 (constant-warp factorization), F2 (chirality grading), F3 (continuum recovery at small t).

- **G4-5**: discrete replica method (beyond sprint scale). *Now identified as the proper target for literal Sommerfeld–Cheeger extraction* per the G4-3c-proper structural diagnosis.
- **G4-6**: full discrete-substrate S_{BH} derivation from the G4-4 + G4-5 fusion (beyond sprint scale).

The end-to-end discrete-substrate derivation of S_{BH} from G4-4 onwards remains a substantial multi-stage commitment after the G4-3a→G4-3d-UV closure.

12.4 Honest scope at G4-3 closure

Five of seven sub-sprints in the G4-3 sequence (G4-3a / cleanup / b / c / d, plus the G4-3d-UV extension) are closed at sprint scale. The substrate is intact and operational: constant-warp factorization at the operator level holds by construction, variable-warp tip-regular substrate built, conical-defect parameterization defined via $N_\phi(\alpha)$ wedge lattice, leading Weyl term reproduced at the level of the omitted boundary correction. The remaining sub-sprints G4-4, G4-5, G4-6 form the *extended commitment* to derive S_{BH} from the discrete substrate.

What is NOT extracted at sprint scale:

- Literal Sommerfeld–Cheeger $(1/12)(1/\alpha - \alpha)$ coefficient extraction (off by factor 2–4 at proper wedge lattice with $N_0 = 120$; the bulk subtraction inherits UV-asymmetry between $\alpha < 1$ and $\alpha > 1$ branches that is structurally larger than the SC tip term at this discretization).
- Discrete spinor Dirac on the cigar substrate (G4-4 extended target).
- Discrete replica method (G4-5 extended target).

The G4-3c-proper finding is the sharpening result: the literal Sommerfeld–Cheeger extraction is structurally below the bulk subtraction's discretization floor at sprint scale, even with the proper apex-angle wedge lattice. G4-5 (discrete replica method) is named as the correct path.

12.5 Connection to G4-2 continuum derivation

G4-2 (Section 6) derived $S_{\mathrm{BH}} = A\Lambda^2/(12\pi)$ at the continuum level. G4-3 onwards rebuilds this on the discrete substrate, with three target identifications:

1. discrete conical contribution $\to (1/12)(1/\alpha - \alpha)$ in the continuum limit;
2. discrete Dirac on warped substrate $\to$ G4-1's S^2 Dirac at constant warp;
3. discrete replica method gives $S_{\mathrm{BH}} \to A\Lambda^2/(12\pi)$.

12.6 Connection to G8 cutoff dependence

The Λ^2 prefactor in S_{BH} was naively expected to inherit G8's cutoff dependence as $\propto \phi(2)/\phi(1)$ for arbitrary cutoff; the G4-5d measurement REJECTED that prediction (65% deviation across three cutoff families) — under the proper-time estimator the 2D tip follows the log-regulated $\phi(0)$, while in the spectral action the 4D cigar's leading tip term carries $\phi(1)$, per the sector-wise map 2D tip $\leftrightarrow$ $\phi(0)$ / EH $\leftrightarrow$ $\phi(1)$ / Λ_{cc} $\leftrightarrow$ $\phi(2)$ (Theorem 13.1). In the spectral action the Λ^2 prefactor of S_{BH} carries $\phi(1)$, not $\phi(2)$; $\phi(0)$ belongs to the proper-time estimator used on the substrate.

13 G4-4: sprint-scale closure of the warped Dirac program

The G4-3 sub-sprint sequence (§12) closes the discrete substrate. G4-4 is the next sub-sprint of the extended S_{BH} program: building the *Dirac operator* on the discrete substrate and extracting the load-bearing structural quantities for the replica method.

This section reports a substantial collapse of the scoping estimate: the G4-4 program was sized as a major sub-program for sub-sprints G4-4a/b/c/d/e/f, but the load-bearing structural quantities for S_{BH} are now *quantitatively validated at sprint scale* across 14 sub-sprints completed in a single focused session.

13.1 Architectural inputs from G4-3

G4-4 builds on the G4-3a-cleanup Hermitian polar Laplacian and the G4-3b variable warp profile $r(\rho) = r_h\sqrt{1+(\rho/r_h)^2}$. The continuum 4D Dirac on the warped cigar is

$$D_{\mathrm{cigar}} = D_{D^2}\otimes I_{S^2} + \gamma^5_{D^2}\otimes\frac{D_{S^2}}{r(\rho)} + \frac{r'(\rho)}{r(\rho)}\,\gamma^\rho\otimes I_{S^2}, \tag{31}$$

with squared form (per Camporesi 1996)

$$D^2_{\mathrm{cigar}} = D^2_{D^2}\otimes I + I\otimes\frac{D^2_{S^2}}{r(\rho)^2} + \left(\frac{r'(\rho)}{r(\rho)}\right)^2 + \text{cross terms with } \gamma^\rho. \tag{32}$$

13.2 G4-4a: constant-warp Dirac

At $r(\rho) = r_h$ constant, the warped product reduces to $D^2_{\mathrm{cigar}} = D^2_{D^2}\otimes I + I\otimes D^2_{S^2}/r_h^2$ exactly. The heat trace factorizes:

$$K^{\mathrm{Dirac}}_{\mathrm{cigar}}(t) = K^{\mathrm{Dirac}}_{D^2}(t)\cdot K^{\mathrm{Dirac}}_{S^2}(t). \tag{33}$$

Three load-bearing falsifiers F1 (factorization), F2 (chirality grading), F3 (continuum recovery at small t) closed at sprint scale:

- F1 bit-exact at machine precision (rel err $\sim 10^{-16}$) across three panel sizes (Hilbert dim up to 115,200).
- F2 verified at the Cl(2,0) gamma algebra (residuals 0.00) AND at the operator level on every Fourier block of the explicit linear Dirac (canonical chirality-graded $\sqrt{L}$ construction).
- F3 spinor Weyl ratio at $t = 0.1$, $N_\phi = 192$: $R_{\mathrm{Dirac}}(0.1) = 0.996$ against the bulk term; the continuum value of the same ratio, perimeter term included, is 0.944, so the substrate sits about 5.5% above it; rank-2 ratio $K^{\mathrm{Dirac}}/K^{\mathrm{scalar}} = 1.9998$ bit-essentially-exact.

(MEASURED, sprint-scale; F1–F7 are exercised as a live panel in `tests/test_paper51_L6_full.py`, and the F3 value is pinned — as $a_0^{\mathrm{Dirac}} = 2R_{\mathrm{Dirac}}(0.1) = 1.992$ — in `tests/test_paper51_g44_headlines.py`.)

The lowest-mode validation $|\lambda^{\mathrm{Dirac}}_{\mathrm{min}}|$ converges to π/R (the half-integer angular momentum Bessel zero $j_{1/2,1} = \pi$) at 0.5% at the finest UV panel, operationally verifying the anti-periodic spinor BC at the eigenvalue level.

13.3 G4-4b: variable-warp Dirac

Three substantive findings across four G4-4b sub-sprints.

F6 Riemannian-limit (LOAD-BEARING). At constant warp, `VariableWarpDirac` reduces bit-exact to G4-4a's `WarpedDiracConstant` (residual $\sim 10^{-16}$).

F7 factorization-loss structural form. At smooth-tip warp, the factorization $K_{\text{cigar}} = K_{D^2} \cdot K_{S^2}$ is broken by $\sim (r'/r)^2$ corrections. Empirical sweep at fixed substrate, $r_h \in \{50, 100, 200\}$:

$$\frac{\Delta_{\text{fact}}(t)}{K_{\text{const}}(t)} \approx 0.110 \cdot t \cdot \left(\frac{R}{r_h}\right)^4 + O\left(\left(\frac{R}{r_h}\right)^6\right). \tag{34}$$

The perturbative power-law slope is **3.994 ± 0.001**, matching the Taylor prediction $\alpha = 4$ at 0.1% precision. The mass-deviation integral driver $|\langle 1/r^2 - 1/r_h^2 \rangle_\rho|$ gives CV = 0.000 at leading order. The spin-connection-squared driver $\langle (r'/r)^2 \rangle$ is equivalent at leading order — empirically validating the structural prediction that factorization-loss IS the spin-connection content.

Cone-Dirac saturation. The asymptotic-free recovery limit $r_h \to 0$ does NOT saturate at the naive "all-modes-massless" value $4(l_{\max}+1)(l_{\max}+2) \cdot K_{\text{disk}}$, because the lattice spacing a fixes a finite S^2 mass $(n+1)^2/a^2$ at the apex regardless of r_h. The CORRECT saturation is the *cone-Dirac heat trace* (singular-tip warp $r(\rho) = \rho$):

$$K_{\text{var}}^{\text{Dirac}}(t; r_h \to 0) = K_{\text{cone}}^{\text{Dirac}}(t), \tag{35}$$

to 6+ digits at $r_h = 10^{-3}$ (by construction: the lattice excludes $\rho = 0$, so $\sqrt{\rho^2 + r_h^2} \to \rho$). This is a substantive structural finding: **the discrete-substrate gravity has two scales** — the substrate UV a and the warp scale r_h — and the asymptotic-free condition is local (holds at large ρ, fails at the apex where $r(\rho_1) \approx a$). (MEASURED, sprint-scale; `tests/test_paper51_g44_headlines.py` recomputes the independent cone-Dirac reference $r(\rho) = \rho$ at the G4-4b-c substrate $N_\rho = 20$, $a = 0.3$, $N_\phi = 12$, $l_{\max} = 3$ and pins the match at 6×10^{-8} relative — 7 digits, inside the 6-digit claim — with the $O(r_h^2)$ convergence and the plateau ratio $K_{\text{cone}}/K_{\text{disk}}(t{=}0.5) = 50.848$.)

Level 1.5 spin connection. Adding the scalar $(r'/r)^2$ correction to D^2 (the leading scalar piece of the γ^ρ-mixing cross term) gives a 0.4–3.9% modification to K_{Lvl1} across the variable-warp regime. F6 extension at constant warp bit-exact. A hidden bug in `warp_derivative_over_warp()` (a hardcoded analytical smooth-tip formula instead of FD on the actual profile) was caught by the F6 bit-exact reduction — a load-bearing demonstration that bit-exact reductions function as correctness infrastructure.

13.4 G4-4c: conical-defect spinor and the headline tip coefficient

The class `DiscreteWedgeDirac` (apex angle $2\pi\alpha$, anti-periodic spinor BC) is the spinor analog of T1's G4-3c-proper scalar wedge. At $\alpha = 1$, the wedge-Dirac reduces bit-exact to G4-4a's disk-Dirac.

The headline structural identification. The spinor conical-defect tip coefficient is identified, with the discrete value converging to it as the radial mesh refines:

$$\boxed{\Delta_K^{\text{Dirac,tip}}(\alpha) = -\frac{1}{12}\left(\frac{1}{\alpha} - \alpha\right)} \tag{36}$$

Best extraction (G4-4c week 3): $\alpha = 2/5$, $t = 2.0$, recovery = 1.000021 (relative error 2.1×10^{-5}). Nine of eleven tested α values reach $> 99.5\%$ recovery somewhere in the t-sweep. (MEASURED, sprint-scale; the best-cell recovery is recomputed at the production substrate and pinned in `tests/test_paper51_g44_headlines.py`.) This is the spinor conical-defect heat-kernel coefficient: *identical in magnitude* to the scalar Sommerfeld [20]/Dowker (1977) [18]/Cheeger–Solodukhin [17, 21] tip coefficient and *opposite in sign* under the anti-periodic (spinor) convention of the discrete extraction; the published spinor-on-cone treatment is Fursaev–Miele [24] ("spin 1/2 resembles the scalar case"), and the shared magnitude is the content of the G4-2 remark.

The scalar/spinor disparity. G4-3c-proper (T1) extracted only $\sim 28\%$ of the SCALAR SC slope at the same sprint-scale substrate. Why does the spinor case work where the scalar case fails? The G4-4e BC sectors diagnostic provides the answer:

Sector	Mean SC recovery	Notes
Spinor (anti-periodic, $m \in \mathbb{Z} + \frac{1}{2}$)	**99.4%**	(§13.4)
Scalar (periodic, $m \in \mathbb{Z}$)	66.3%	(T1, this paper §12)
Ratio	1.50×	—

The scalar has a zero-mode at $m = 0$ (no centrifugal barrier in the $u = \sqrt{\rho}\, f$ representation; effective centrifugal $-1/4$ attractive) whose contribution is structurally asymmetric in $\alpha \leftrightarrow 1/\alpha$ at finite discretization. The spinor (anti-periodic) lacks any zero-mode; $m_{\text{eff}} = 1/2$ at its ground mode has zero effective centrifugal in the u-rep. **The anti-periodic spinor BC with half-integer angular momentum is structurally essential for clean conical-defect SC extraction on a discrete substrate.** (MEASURED, sprint-scale; the spinor row — per-α recoveries and the 99.4% panel mean at $t = 1.0$ — is recomputed and pinned in `tests/test_paper51_g44_headlines.py`.)

$\alpha > 1$ structural asymmetry. A clean negative landed: at $\alpha > 1$ (excess-angle / saddle-cone), the recovery plateaus at **bit-identical 67.88% across** $N_0 \in \{120, 240, 480\}$ — azimuthal refinement does NOT help. Radial refinement does: the B4 $a \to 0$ diagnostic (§13.7.6) shows the recovery climbing toward 1 as the radial mesh refines, so the gap is a finite-a artifact and the continuum slope at $\alpha > 1$ is the plain Sommerfeld–Cheeger continuation.

13.5 G4-4d, G4-4e, G4-4f: Seeley-DeWitt, BC sectors, replica derivative

Three brief sprints validating cross-cutting structural quantities.

G4-4d Seeley-DeWitt extraction. At the sweet spot $t = 0.1$, $N_\phi = 192$:

$$a_0^{\text{Dirac}} = 1.992 \quad \text{(bulk term: 2.0; continuum with perimeter term: 1.888).} \tag{37}$$

The comparison with 2.0 omits the Dirichlet perimeter term; against the continuum value of the same estimator (1.888 at $R = 10$, $t = 0.1$) the substrate is 5.5% high, and its $a \to 0$ sweep at $N_\phi = 192$ $(2.024, 1.992, 1.979, 1.973)$ has not reached it, so 1.992 is a crossing of the bulk value, not a 99.6% recovery. (MEASURED; recomputed at the paper's own substrate $N_\rho = 200$, $a = 0.05$, $N_\phi = 192$, $t = 0.1$ and pinned in `tests/test_paper51_g44_headlines.py`.) Subleading a_1 (boundary) is small at the sweet spot; a clean identification requires multi-window fitting or continuum extrapolation, deferred. The naive polynomial fit at small t is biased by the UV overshoot identified in G4-3d-UV (the $4/\pi^2$ factor between discrete and continuum azimuthal Laplacian at the truncation edge).

G4-4f replica derivative. The replica-method load-bearing quantity:

$$\left.\frac{d\Delta_K^{\text{Dirac}}}{d\alpha}\right|_{\alpha=1} = -\frac{1}{12}\left(-\frac{1}{\alpha^2} - 1\right)\Bigg|_{\alpha=1} = +\frac{1}{6}. \tag{38}$$

Central finite difference on the discrete substrate ($\alpha_\pm = 1 \pm k/N_0$, $k \in \{1, 2, 4, 6, 12, 24\}$, $N_0 = 120$) gives, at sweet-spot $t = 5.0$:

$$\left[\frac{dK_{\text{wedge}}^{\text{Dirac}}}{d\alpha} - K_{\text{disk}}^{\text{Dirac}}\right]_{\alpha=1}^{\text{discrete}} = 0.1612 \quad \text{(continuum: 0.1667, recovery } \mathbf{96.69\%}\text{).} \tag{39}$$

(MEASURED, sprint-scale.)

13.6 Implications for the extended S_{BH} program

The replica method derives the BH entropy as

$$S_{\mathrm{BH}} = -\frac{dI_E}{d\alpha}\bigg|_{\alpha=1}, \quad I_E = \int_0^\infty \frac{dt}{t}\, K(\alpha, t)\, f(t\Lambda^2). \tag{40}$$

The load-bearing quantities for this calculation are:

1. The conical-defect tip coefficient $\Delta_K^{\mathrm{Dirac,tip}}(\alpha)$ (Eq. 36);
2. The replica derivative at $\alpha = 1$ (Eq. 38);
3. The cutoff regularization function $f(t\Lambda^2)$ (Class 1 calibration, §11).

Items 1 and 2 are approached on the discrete substrate: the tip coefficient converges as the radial mesh refines, and the replica derivative is extracted at 96.69%. Item 3 is the standard Connes–Chamseddine cutoff issue, structural to the framework and external to the discrete substrate.

Sprint-scale closure implication. Before this session, the extended S_{BH} program (G4-5 discrete replica method + G4-6 full S_{BH} derivation, a substantial end-to-end commitment per G4-4 scoping) rested on an architectural sketch. **The remaining program work is integration over t with proper cutoff regularization, not foundational uncertainty**; the load-bearing quantities (tip coefficient $-1/12$ and replica derivative $+1/6$) are approached at sprint scale, the first converging as $a \to 0$ and the second extracted at 96.69%.

13.7 G4-5: discrete replica method for S_{BH}

The G4-4c tip coefficient (Eq. 36) and G4-4f replica derivative (Eq. 38) are the load-bearing inputs for the replica-method derivation of S_{BH} on the discrete substrate. G4-5 integrates these over t against a Connes–Chamseddine cutoff function $f(t\Lambda^2)$ and produces S_{BH} directly. The sub-sprint sequence G4-5a/b/c/d landed in parallel after the G4-4 sprint-scale closure of this session, yielding two substantive structural reframings of the Connes–Chamseddine cutoff-function dependence of S_{BH}.

13.7.1 G4-5a / G4-5a-refined: tip-only replica integration

The tip-only contribution

$$S_{\mathrm{tip}}(\Lambda) = +\frac{1}{2}\int_{t_{\mathrm{UV}}}^{t_{\mathrm{IR}}} \frac{dt}{t}\, f(t\Lambda^2)\, \Delta'(t), \quad \Delta'(t) = \frac{dK^{\mathrm{Dirac}}_{\mathrm{wedge}}}{d\alpha}\bigg|_{\alpha=1} - K^{\mathrm{Dirac}}_{\mathrm{disk}}(t). \tag{41}$$

G4-5a first move (sprint-scale t-grid $\{0.1, \ldots, 20\}$, Gaussian cutoff $f(x) = e^{-x}$): S_{tip} finite, positive, Λ-decreasing across $\Lambda \in \{0.5, 1, 1.5, 2\}$. Sanity checks all pass. Discrete/continuum ratio 0.13–0.37, structurally gapped by the missing UV t-window.

G4-5a-refined extended the t-grid to $\{0.0025, 0.005, \ldots, 10\}$ (12 points; UV cutoff $t_{\mathrm{UV}} = a^2$). Recovery ratio improves **uniformly** at every Λ:

Λ	G4-5a (rough)	G4-5a-refined (rough)	G4-5a-refined (exact Mellin)
0.5	0.37	**0.585**	**0.635**
1.0	0.26	0.517	0.572
1.5	0.18	0.464	0.522
2.0	0.13	0.424	0.484

Mean recovery doubled. Exact Gaussian Mellin ($M_0(\Lambda) = E_1(a^2\Lambda^2) - E_1(R^2\Lambda^2)$) pushes three of four Λ values past 0.5; $\Lambda = 2$ misses by 3%. The residual gap is identified as the T2 G4-3d-UV azimuthal-truncation overshoot: per-t tip recovery is 1.3% at $t = 0.0025$, 16.5% at $t = 0.005$, 36.2% at $t = 0.01$, ..., 76.3% at $t = 0.1$ (the G4-5a starting point). The $4/\pi^2$ discrete/continuum azimuthal-Laplacian ratio at the truncation edge is the structural origin — closure requires either a spectral azimuthal discretization (DST/Fourier) or N_ϕ refinement beyond 192.

13.7.2 G4-5b: bulk Weyl extraction is structurally 4D, not 2D

A substantive structural reframing landed: the bulk Λ^4 (cosmological constant) term in the Connes–Chamseddine spectral action is *not* present in the disk-Dirac $K^{\text{Dirac}}_{D^2}(t)$ at leading order. The 2D Seeley–DeWitt expansion of the disk-Dirac is

$$K^{\text{Dirac}}_{D^2}(t) \sim \frac{2A_{D^2}}{4\pi t} + a_1 t^{-1/2} + a_2 + O(t), \tag{42}$$

giving a leading $1/t$ Mellin moment, not $1/t^2$. The bulk Λ^4 piece arises only in the *4D embedding* $D^2 \times S^2$ (G4-5c). Empirical fit of $I_{\text{CC}}(\Lambda) = c_4\Lambda^4 + c_2\Lambda^2 + c_0 + c_{-2}\Lambda^{-2}$ on $K^{\text{Dirac}}_{D^2}$ gives:

- $c_4 = +0.74$ (vs naive 4D prediction $A_{D^2}a_0/(8\pi^2) \approx 2.53$): not present in 2D as a structural piece, accounting for the 0.29 ratio;
- $c_2 = -141$ to -195 (sign correct, OoM correct vs $-A_{D^2}/(2\pi) \approx -50$; factor 3–4 magnitude off);
- $c_0 = +10354$ to $+10577$ (matching $A_{D^2}/(2\pi a^2) \approx 20000$ at ratio 0.53 — the UV-cutoff Weyl term).

This sharpens §13.7.5: the bulk Weyl Λ^4 emerges only when the disk-Dirac is tensored with $S^2_{r_h}$. **The 4D embedding is structurally required for the cosmological-constant prediction**, not an optional extension. G4-5c verifies this empirically (Sec. 13.7.3) by extracting the joint S_{BH} at the cigar geometry.

13.7.3 G4-5c: joint warp + conical-defect Dirac — bit-exact F6 + $r_h^2\Lambda^2/3$ at UV

The class `JointWarpConicalDirac` (driver level) combines the wedge azimuthal structure of G4-4c with the variable-warp radial structure of G4-4b. Per (n, k_ϕ) block: $H = L_{\text{disk}}(m_{\text{eff}}) + \text{diag}((n+1)^2/r(\rho)^2)$ with $m^2_{\text{eff}} = (2/h_\phi)^2 \sin^2(\pi(k+1/2)/N_\phi)$ and $h_\phi = 2\pi\alpha/N_\phi$. The wedge enters through Fourier-mode m_{eff}; the warp enters through the radial mass diagonal — they act on different tensor factors, so the combination is mechanical.

F6 extension LOAD-BEARING (bit-exact both directions):

- F6-A: at $\alpha = 1$, joint reduces to G4-4b's `VariableWarpDirac.smooth_tip` with rel_err $= 0$ exactly (structural identity, not just machine precision);
- F6-B: at constant warp $r(\rho) = r_h$, joint reduces to `DiscreteWedgeDirac` $\times$ `S2DiracSpectrum` product with rel_err $\leq 1.2 \times 10^{-13}$ across $\alpha \in \{0.5, 1, 1.5\}$ and $t \in \{0.05, 0.1, 0.5, 1\}$.

S_{BH} extraction: the discrete replica derivative of the joint heat trace integrated against Gaussian cutoff gives, at $r_h = 2$:

Λ	$S_{\rm BH}^{\rm discrete}$	continuum $r_h^2\Lambda^2/3$	ratio
0.5	+9.69	+0.333	29.08
1.0	+7.75	+1.333	5.81
2.0	+4.55	+5.333	**0.85**

The Λ-monotone descent of the ratio ($29.1 \to 5.81 \to 0.85$) is the substrate-UV/IR signature: at $\Lambda \cdot R \lesssim 1$, the Gaussian $e^{-t\Lambda^2}$ does not fully suppress large-t (IR) contributions and the integral over-counts. **At $\Lambda = 2$, the discrete extraction lands within the factor-2 band of the continuum $r_h^2\Lambda^2/3$ prediction, recovery** $= 0.85$. This is consistent with G4-2's continuum derivation (Sec. 6) at one Λ cell; the recovery is Λ-dependent ($29.1, 5.81, 0.85$) and not converged. The $\alpha > 1$ branch (a finite-a artifact, B4) does not affect the $\alpha = 1$ extraction.

13.7.4 G4-5d: cutoff dependence and sector-wise Mellin moment map

The original G8 prediction (Sec. 11) was $S_{\rm BH}(f) \propto \phi(2)$ for any cutoff f. G4-5d tests this on three cutoff classes (Gaussian e^{-x}, sharp $\Theta(1-x)$, polynomial e^{-x^2}). Discrete result at $\Lambda = 1$:

Cutoff	$S_{\rm tip}^{\rm disc}(\Lambda = 1)$	predicted ratio vs Gauss	observed
Gaussian, $f(x) = e^{-x}$, $\phi(2) = 1$	0.1303	1 (reference)	1
Sharp, $f = \Theta(1-x)$, $\phi(2) = 1/2$	0.1907	1/2	**1.464**
Polynomial, $f = e^{-x^2}$, $\phi(2) = 1/2$	0.1425	1/2	**1.094**

The naive G8 prediction $S_{\rm sharp}/S_{\rm Gauss} = 0.5$ is **rejected at 65%** deviation. Empirical ordering $S_{\rm sharp} > S_{\rm poly} > S_{\rm Gauss}$ does NOT match $\phi(2)$ ordering (which would be sharp = polynomial = 0.5· Gaussian).

The structural reframing. The topological tip contribution to $S_{\rm BH}$ is dimensionless and R-independent, so it inherits no polynomial x^k weight from the heat-kernel expansion. Under the proper-time estimator used on the substrate the relevant moment is therefore $\phi(0)$ (logarithmically regulated), not $\phi(2)$; in the spectral action the 4D tip carries $\phi(1)$ (Theorem 13.1). The empirical ordering $S_{\rm sharp} > S_{\rm poly} > S_{\rm Gauss}$ matches the qualitative $\phi(0)$ ordering exactly: sharp gives log-divergent $\phi_{\rm sharp}(0) \to \infty$ regulated by the UV cutoff at $t = 1/\Lambda^2$; polynomial gives intermediate decay; Gaussian gives the smallest $\phi_{\rm Gauss}(0) = \gamma_{\rm E}$ (Euler–Mascheroni-class log-divergence).

This promotes G8 from a single Mellin-scaling law to a **sector-wise moment map**:

Sector	Wilson coefficient	Mellin moment
Bulk R^0 (cosmological constant)	$\Lambda_{\rm cc}$	$\phi(2)$
Bulk R^1 (Einstein–Hilbert)	$G_{\rm eff}^{-1}$	$\phi(1)$
2D tip, proper-time estimator	(tip coeff 1/12)	$\boldsymbol{\phi(0)}$

This is a refinement of G8 for the proper-time estimator used on the substrate; in the spectral action itself the 4D tip carries $\phi(1)$ (Theorem 13.1). G7's $G_{\rm eff} = 3\pi/(\phi(1)\Lambda^2)$ is the $\phi(1)$-moment result; G7's $\Lambda_{\rm cc} = 6\phi(2)/\phi(1)\Lambda^2$ is the $\phi(2)$-moment result; G4-5d adds the $\phi(0)$-moment result for $S_{\rm BH}$. **The cutoff-function calibration data (Class 1 per §11) controls all three sectors simultaneously, but through different Mellin moments.**

Theorem 13.1 (Sector-wise Mellin moment map — INTERNAL THEOREM; `tests/test_paper51_cutoff_me`
Let f be a cutoff function with regulated Mellin moments $\phi(s) = \int_0^\infty f(x)\, x^{s-1}\, dx$, and let the Chamseddine–Connes spectral action on the Euclidean Schwarzschild cigar be decomposed into bulk and topological-tip sectors via the replica method. Then each sector's cutoff dependence is controlled by a single Mellin index k:

(i) *The cosmological-constant sector (proportional to $a_0 \sim t^{-d/2}$) scales with $\phi(d/2) = \phi(2)$ in $d = 4$.*

(ii) *The Einstein–Hilbert sector (proportional to $a_2 \sim t^{-(d-2)/2}$) scales with $\phi((d-2)/2) = \phi(1)$ in $d = 4$.*

(iii) *The topological tip sector (the Sommerfeld–Cheeger contribution). On the bare 2D cone the tip term is t-independent, so in the spectral action it contributes $\frac{1}{6}f(0)$, independent of the Mellin moments. On the 4D cigar it multiplies the Dirac transverse kernel $K_{S^2}(t) = 2r_h^2/t - 1/3 + O(t)$, so its leading contribution to S_{BH} is $(r_h^2\Lambda^2/3)\,\phi(1)$, the Einstein–Hilbert moment, and the subleading piece is $-f(0)/18$.*

Proof. Write the cutoff as a Laplace transform, $f(x) = \int_0^\infty e^{-sx} g(s)\,ds$, so that $S = \mathrm{Tr}\, f(D^2/\Lambda^2) = \int_0^\infty g(s)\, K(s/\Lambda^2)\,ds$ with $K(t) = \mathrm{Tr}\, e^{-tD^2}$. A heat-trace term $c\,t^{-\beta}$ with $\beta > 0$ contributes $c\,\Lambda^{2\beta}\int_0^\infty g(s)s^{-\beta}ds = c\,\Lambda^{2\beta}\phi(\beta)/\Gamma(\beta)$, and a t^0 term contributes $c\,f(0)$. The Seeley–DeWitt expansion $K(t) \sim \sum_n a_n\, t^{(n-d)/2}$ then gives (i) and (ii) for the $n = 0$ and $n = 2$ bulk coefficients.

For the replica tip sector: the Sommerfeld–Cheeger contribution to the heat trace on a cone of apex angle $2\pi\alpha$ is a *topological* correction $\Delta K^{\mathrm{SC}}(\alpha, t) = c(\alpha) \cdot K_{S^{d-2}}(t)$, where $c(\alpha) = (1/12)(1/\alpha - \alpha)$ for spinors and $K_{S^{d-2}}(t)$ is the transverse-sphere heat kernel; its replica derivative is the constant $1/6$. On the bare 2D cone the tip term is t^0 and contributes $f(0)/6$. On the 4D cigar, $\frac{1}{6}(2r_h^2/t - 1/3)$ contributes $\frac{1}{3}r_h^2\Lambda^2\phi(1) - f(0)/18$; the leading term is the $A\Lambda^2/(12\pi)$ of G4-2 at the Gaussian cutoff. □ □

Remark 13.2 (The substrate's cutoff ordering)**.** The G4-5d measurement weighted the discrete substrate's tip with the proper-time functional $\int (dt/t)\, f(t\Lambda^2)\, \mathrm{tip}(t)$, which for a constant tip is the log-regulated $\phi(0)_{\mathrm{reg}}$ and orders the cutoffs sharp > polynomial > Gaussian (sub-5% across 9 channels, archived sprint measurement; the qualitative ordering is pinned in `tests/test_paper51_cutoff_mellin.py`). That is a property of this estimator, not of the spectral action, in which the continuum 2D tip carries $f(0)$ and the 4D leading tip carries $\phi(1)$.

13.7.5 G4-5 headline closure and implications for G4-6

Combining G4-5a-refined + G4-5b + G4-5c + G4-5d:

Operationally extracted on the discrete substrate. $S_{\mathrm{BH}}^{\mathrm{discrete}}(r_h = 2, \Lambda = 2) = 4.55$ (G4-5c), with recovery 0.85 vs continuum $r_h^2\Lambda^2/3 = 5.33$. F6 load-bearing reductions bit-exact in both reduction directions.

Structurally identified.

1. The tip-only contribution to S_{BH} extracts at the discrete substrate level via the replica method (G4-5a-refined, recovery 0.42–0.59).
2. The bulk Λ^4 cosmological term is structurally *absent* in 2D and emerges only in the 4D cigar embedding (G4-5b).
3. On the 4D cigar the leading tip contribution to S_{BH} carries $\phi(1)$, like Einstein–Hilbert; the cosmological constant is at $\phi(2)$; the bare 2D tip carries $f(0)$ (Theorem 13.1).

Implications for G4-6 (full S_{BH} closure). The remaining program work for G4-6 is:

- Subleading $O(a^2/r_h^2)$ UV corrections via multi-substrate continuum extrapolation;
- Subleading $O(r_h^2/R^2)$ IR corrections via boundary regularization;

- (Resolved since by B4: the $\alpha > 1$ asymmetry is a finite-a artifact.)
- Refinement of the azimuthal discretization (DST/Fourier) to close the T2 high-m overshoot in the tip-recovery diagnostic.

The end-to-end G4-6 commitment shrank further once the $\alpha > 1$ asymmetry was traced to a finite-a artifact (B4, Sec. 13.7.6 below). The end-to-end S_{BH} program is now substantially smaller than the original G4-4 scoping estimate because the structural quantities have crystallized into a single sector-wise moment map.

The structural-skeleton-scope reading at G4-5 closure. The framework predicts the structural form of S_{BH}:

$$S_{\mathrm{BH}}^{\mathrm{structural}} = \frac{r_h^2 \Lambda^2}{3} \cdot \phi(1)[f] \tag{43}$$

at leading order, with a subleading $-f(0)/18$. The cutoff moment is *not* $\phi(2)$, which is what G4-5d rejected. The framework predicts the $r_h^2\Lambda^2/3$ *structural form*; it does NOT predict M_{tip} (cutoff function calibration data, Class 1). Consistent with the structural-skeleton-scope reading elsewhere in the framework [7, 8].

13.7.6 Post-G4-5 follow-on push (v3.19.0): the $\alpha > 1$ Möbius form (a finite-a artifact) and the Mellin map

A second 5-agent parallel dispatch following the v3.18.0 G4-5 closure landed five non-overlapping tracks targeting open issues plus the extended G4-6 scoping. The new content is one fixed-a closed form (the $\alpha > 1$ Möbius form, later shown to be a finite-a artifact) and one empirical cutoff ordering (the sector-wise Mellin moment map).

The $\alpha > 1$ Möbius modification. The G4-4c week 2 result (N_0-independent 67.88% recovery at $\alpha > 1$, Sec. 13.4) is fitted, at fixed radial spacing a, by the closed form

$$\mathrm{slope}^{\mathrm{Dirac,ex}}(\alpha) \;=\; -\frac{1}{12} \cdot \frac{\alpha}{2\alpha - 1} \qquad (\alpha > 1), \tag{44}$$

equivalently

$$\Delta_K^{\mathrm{Dirac,ex}}(\alpha) \;=\; \frac{\alpha^2 - 1}{24(\alpha - 1/2)}. \tag{45}$$

The modification factor $F(\alpha) = \alpha/(2\alpha - 1)$ is a Möbius transformation with three structural features:

1. $F(1) = 1$ (smooth-disk limit; matches the continuum Sommerfeld–Cheeger formula at $\alpha = 1$);
2. $F(\alpha) \to 1/2$ as $\alpha \to \infty$ (asymptotic spinor double-cover monodromy signature);
3. $F'(1)$ finite (smooth at the deficit/excess transition).

Literature check. A literature grounding sprint (task #26 of the post-v3.19.0 follow-on queue, 2026-05-29) found that the Möbius factor $F(\alpha) = \alpha/(2\alpha - 1)$ has no analog in the standard published spinor-on-cone literature (Cheeger [17], Dowker [27], Fursaev–Miele [24]), where the spinor heat-kernel coefficient at conical defect resembles the antisymmetric scalar Cheeger formula $-(1/12)(1/\alpha - \alpha)$ with no Möbius modification at $\alpha > 1$. The discrete substrate computes a result that is either (i) a yet-unidentified continuum excess-angle correction missed in the standard derivations, or (ii) a discrete-substrate-specific UV artifact; the latter is

partially ruled out by the G4-4c week 2 N_0-independence at $N_0 \in \{120, 240, 480\}$ but not eliminated. Mechanism identification is a named follow-on. Note: the v3.19.0 first-pass attribution to a "Fursaev–Solodukhin spinor double-cover correction" (via an arXiv ID that turned out to be a different paper) was retracted during the post-v3.19.0 audit; the Fursaev–Solodukhin 1995 paper [23] cited elsewhere in this paper is correctly attributed (Riemannian geometry of conical defects, BH entropy) but does not contain the Möbius modification.

Mechanism RESOLVED (B4, 2026-05-29): finite-a artifact, continuum is Sommerfeld–Cheeger. The open question above is settled in favour of option (ii). The L6 bookkeeping sprint re-measured the $\alpha > 1$ recovery with the clean machinery – spectral azimuthal eigenvalues $m_{\text{eff}} = (k + \frac{1}{2})/\alpha$ plus a *radial continuum extrapolation* $a \to 0$ at several t – the one axis the prior robustness checks (t-shape, azimuthal discretisation, R; all at fixed $a \approx 0.05$) never varied. At $\alpha = 2$, $t = 1$, the recovery climbs monotonically away from the Möbius value $\alpha/(2\alpha - 1) = 0.667$ toward the Sommerfeld–Cheeger value 1 as $N_\rho = 200 \to 3200$ $(0.675, 0.769, 0.837, 0.885, 0.919)$, with $(1 - \text{recovery}) \sim \sqrt{a}$ (error ratio $1.4085, 1.4143, 1.4164$ over successive halvings, approaching $\sqrt{2} = 1.4142$). The $\sqrt{a}$ rate is the signature of the genuine cone singularity at $\alpha \neq 1$ (a true deficit/excess-angle apex), contrasting the smooth-apex $O(a)$ at $\alpha = 1$. Thus recovery $= 1 - C\sqrt{a} \to 1$: the continuum excess-angle slope is the standard Sommerfeld–Cheeger continuation $-\frac{1}{12}$, and the Möbius factor $\alpha/(2\alpha - 1)$ is the *fixed-a appearance* of this quantity at the substrate scale $a \approx 0.05$ – a discrete-substrate artifact, not a continuum correction. This explains why no published spinor-on-cone result, and no mechanism sprint (Fursaev–Solodukhin attribution, Routes A/B/C, harmonic conjugate), ever found a Möbius mechanism: there was none to find. The N_0 $(= N_\phi)$ independence checked earlier probes the azimuthal count, not the radial a, so it could not have caught the artifact. The $\alpha > 1$ thread is retired as a finite-a artifact. (Memo: `debug/l6_b4_moebius_continuum_diagnostic_memo.md`.) The harmonic- conjugate algebraic structure recorded below is therefore a property of the fixed-a form, not of a continuum object.

Algebraic structure: harmonic conjugate. A structural sharpening identified post-v3.20.0 (Task 11 of the post-v3.20.0 continuation thread, 2026-05-29; memo `debug/sprint_moebius_mode_counting_di` the Möbius factor $F(\alpha) = \alpha/(2\alpha - 1)$ is the unique Möbius transformation satisfying

$$\frac{1}{\alpha} + \frac{1}{F(\alpha)} = 2, \tag{46}$$

i.e., F is the harmonic conjugate of α with respect to 1 (harmonic mean of α and F equals exactly 1, the smooth-disk value). Three structural consequences: $F(1) = 1$; $F(F(\alpha)) = \alpha$ (involution); $F(\alpha) = \alpha$ only at $\alpha = 1$. This is algebraically equivalent to the spinor-double-cover sketch $\alpha_{\text{eff}} = 2\alpha - 1$ via $F = \alpha/\alpha_{\text{eff}}$, but gives an additional constraint — the harmonic mean of base and effective angles equals the smooth-disk angle. It is an algebraic property of the fitted fixed-a form; since that form is a finite-a artifact (B4), there is no continuum mechanism behind it.

Substrate-level soft-IR identity (a $t \approx 1$ coincidence). A subsequent production-code mode-decomposition diagnostic (Task 25 of the post-v3.20.0 continuation threads, 2026-05-29; memo `debug/sprint_moebius_mode_decomposition_memo.md`) identifies the substrate-level content of the harmonic-conjugate constraint. On the spectral azimuthal substrate (Sec. 13.7.7; classes `DiscreteDiskDiracSpectral` / `DiscreteWedgeDiracSpectral` added in `geovac/gravity/warped_dira` the discrete heat trace at apex angle $2\pi\alpha$ distributes its mass with soft-IR fraction $X(\alpha) :=$ soft_IR_frac(α) satisfying the structural identity

$$F(\alpha) = \frac{1}{2 \cdot (1 - X(\alpha))}, \tag{47}$$

which, combined with Eq. (46), is equivalent to

$$X(\alpha) \;=\; \frac{1}{2\alpha} \qquad \text{(asymptotic; } \alpha > 1\text{)}. \tag{48}$$

The relation is verified at sub-percent precision for $\alpha \geq 2$; at $\alpha = 3$ the match is **0.03%** (essentially exact). The mechanism interpretation: the lowest spinor eigenvalue on the wedge at apex $2\pi\alpha$ is $1/(2\alpha)$ (anti-periodic BC), the lowest-mode heat-trace contribution at intermediate t varies slowly with α, and the total heat trace scales linearly with α (mode count αN_0), so the soft-IR mass fraction scales as $1/(2\alpha)$ asymptotically, reproducing the Möbius form. The identity holds only near $t \approx 1$ (GD-2 audit below) and the form itself disappears as $a \to 0$ (B4), so it is a property of the fixed-a substrate, not a mechanism.

t-robustness audit (Sprint GD-2, 2026-05-29) — form robust, coefficient and soft-IR identity $t \approx 1$-tuned. Both the coefficient match and the soft-IR identity Eq. (48) were extracted at a single heat-trace time $t = 1.0$ (a deliberately chosen "sweet spot"). A robustness audit sweeping $t \in \{0.25, 0.5, 1, 2, 4\}$ separates the robust content from the sweet-spot artifact:

- *Robust.* At every t and $\alpha \in \{2, 3, 5\}$ the measured slope is far closer to the Möbius form $-(1/12)\alpha/(2\alpha - 1)$ than to the continuum Sommerfeld–Cheeger form $-(1/12)(1/\alpha - \alpha)$ (e.g. at $\alpha = 2$: measured ≈ -0.05, Möbius $= -0.056$, SC $= +0.125$, wrong sign). The $\alpha > 1$ slope is unambiguously Möbius-*shaped*, not SC-shaped — robustly in t.
- $t \approx 1$-*tuned.* The *exact* coefficient match (slope/Möbius) crosses unity near $t \approx 1$ but drifts to ~ 0.78 at $t = 0.25$ and ~ 1.13 at $t = 4$ (spread ~ 0.34); the soft-IR ratio $(1 - X)F$ likewise hits $1/2$ only at $t \approx 1$ and drifts 0.31–0.59 across the same window. The "sub-2% / 0.03%" precision was therefore a $t \approx 1$ crossing, not a t-robust coefficient, and the soft-IR identity Eq. (48) is a $t \approx 1$ coincidence rather than a structural mechanism.

The continuum tip is t-independent (1/6; Sec. 13.7.7), so the t-dependence of the substrate's tip extraction is a finite-a effect, and clean continuum coefficients appear only at a sweet-spot t. Net: the Möbius *functional form* is robust at fixed substrate spacing a; the exact coefficient and the soft-IR "mechanism" are sweet-spot artifacts, and the B4 $a \to 0$ diagnostic (above) subsequently resolved the form itself as a finite-a substrate artifact — the continuum $\alpha > 1$ slope is the plain Sommerfeld–Cheeger continuation, leaving no continuum Möbius mechanism to seek. A companion Reading-B test (Sprint GD-3) further finds the form *discretization*-robust: the FD-azimuthal and exact-spectral wedge substrates give Möbius-shaped slopes agreeing to ~ 4 significant figures at every tested (α, t) — so at fixed radial spacing the form does not depend on the azimuthal discretization; it is set by the discrete radial structure, which B4 shows is where it comes from, consistent with the literature's absence of a continuum Möbius (Reading B). A further probe (Sprint GD-6) rules out a subtraction-convention origin: sweeping the radial extent $R = N_\rho a \in \{5, 7.5, 10, 15\}$ leaves the slope *bit-identical* (slope/Möbius $= 1.012, 0.977, 0.977$ at $\alpha = 2, 3, 5$ for every R), so the $\Delta_K = K_{\text{wedge}} - \alpha K_{\text{disk}}$ subtraction cleanly isolates an R-independent tip (bulk R^2 and perimeter R cancel) — at fixed a the Möbius is a tip effect, not an edge/convention artifact. The form is thus robust in t, azimuthal discretization and R at fixed a, but not under radial refinement (B4); the exact-coefficient match to $\alpha/(2\alpha - 1)$ is $t \approx 1$-sharp. Drivers `debug/sprint_gd2_moebius_t_robustness.py`, `debug/sprint_gd2_moebius_slope_t.py`, `debug/sprint_gd3_reading_b_substrate.py`, `debug/sprint_gd6_` memos `debug/sprint_gd2_moebius_t_robustness_memo.md`, `debug/sprint_gd3_reading_b_memo.md`, `debug/sprint_gd6_moebius_convention_memo.md`.

Literature evidence for substrate-universal reading. A multi-source literature check (Task b' of the post-v3.20.0 thread 8, 2026-05-29; memo `debug/sprint_moebius_route_a_fursaev_miele_pdf_`

of the standard published spin-1/2 conical heat kernel literature returns *no Möbius modification* at excess angle. Specifically: the Fursaev–Miele 1996 abstract [24] states verbatim that the spin-1/2 result "resembles the scalar case," mirroring the standard antisymmetric Cheeger formula $-(1/12)(1/\alpha - \alpha)$ without modification. The Solodukhin 2011 Living Review [25] and the Beccaria–Tseytlin 2017 conically-deformed-sphere work [19] likewise do not surface the Möbius factor in their treatments. The unanimous literature signal supports the substrate-universal reading: the Möbius pattern is a structural feature of the discrete substrate (tied to the lowest spinor eigenvalue $1/(2\alpha)$ structure) rather than a continuum theorem missing from the published derivations.

Substrate-discretization-invariance verification. The v3.19.0 measurement convention was audited (Option 2 of Track β'' of post-v3.20.0 thread 9; memo `debug/sprint_multi_thread_day_2026_05_2` and the substrate-level identification is verified across substrate discretizations. Specifically, the v3.19.0 "slope" is defined as the *ratio* $\text{slope}^{v3.19} := \Delta_K(\alpha)/(1/\alpha - \alpha)$ (*not* the derivative $d\Delta_K/d\alpha$), measured at fixed $(R = 10, a = 0.05, N_\rho = 200, N_0 = 120, t = 1.0)$. Using this convention on the spectral substrate (introduced in Sec. 13.7.7; classes `DiscreteDiskDiracSpectral` and `DiscreteWedgeDiracSpectral`), the fresh measurement reproduces the v3.19.0 reported values bit-exactly to four decimal places:

α	v3.19.0 reported	spectral substrate (fresh)	rel. diff
1.5	-0.06466	-0.06466	0.0%
2.0	-0.05622	-0.05622	0.0%
3.0	-0.04883	-0.04883	0.0%

This reproduction to four decimal places confirms the measurement convention and its independence of the azimuthal discretization at fixed a; it says nothing about the $a \to 0$ limit, in which the form disappears (B4). A complementary direct derivative diagnostic ($d\Delta_K/d\alpha$ at $\alpha = 2$, on both FD and spectral substrates at $N_\rho \in \{100, 200, 400, 600\}$) gives a N_ρ-stable value $+0.052$, structurally consistent with the positive Δ_K value at excess angle but *not* the same observable as the v3.19.0 ratio. The previously-flagged "sign discrepancy" (Track α'' of thread 9) was resolved as a definition-versus-derivative observable difference, not a substrate property mismatch.

Empirical match. At $N_0 = 480$, $t = 1.0$:

α	measured slope	predicted slope	rel. err
1.5	-0.0647	-0.0625	-3.3%
2.0	-0.0562	-0.0556	-1.2%
3.0	-0.0488	-0.0500	$+2.4\%$

Average 2.3% relative error. The reciprocal-pair antisymmetry $\Delta_K(\alpha) + \Delta_K(1/\alpha) \neq 0$ is predicted to within $\sim 10\%$ across three reciprocal pairs, confirming the same mechanism explains both the absolute slope deficit and the broken reciprocal antisymmetry.

Implication for the G4-2 conical-replica derivation at $\alpha = 1$. *Unchanged.* The $\partial_\alpha I_E^{\text{conical}}|_{\alpha=1}$ derivative in the G4-2 replica derivation is taken from the deficit side ($\alpha \leq 1$), where the standard $-(1/12)(1/\alpha - \alpha)$ formula holds. The Möbius form is a finite-a artifact (B4) and carries no continuum correction at $\alpha \neq 1$.

Sector-wise Mellin moment map empirical confirmation. The G4-5d structural reframing (Sec. 13.7.4) was retested under the corrected $\phi(0)$ Mellin-moment prediction for the topological tip sector. Across $k \in \{0, 1, 2\}$, the empirical ordering $S_{\text{sharp}} > S_{\text{poly}} > S_{\text{Gauss}}$ matches the

qualitative $\phi(0)$ prediction at every Λ tested; the polynomial-vs-Gaussian channel (the cleanest quadrature comparison) matches the $\phi(0)$ prediction within 6% at every Λ. The mean deviation drops from 49% under the naive $\phi(2)$ prediction to 18% under the $\phi(0)$ prediction. The sector-wise moment map

Sector	Mellin moment
2D tip, proper-time estimator	$\boldsymbol{\phi(0)}$
Bulk Λ^1 (Einstein–Hilbert)	$\phi(1)$
Bulk Λ^0 (cosmological constant)	$\phi(2)$

is the substantive empirical confirmation. A 20-point quadrature follow-on would close the residual sharp-cutoff channel mismatch and upgrade the PARTIAL result to POSITIVE.

FD/spectral azimuthal discretizations at the lattice scale. Replacing the finite-difference azimuthal Laplacian (edge eigenvalue ratio $4/\pi^2 \approx 0.405$) with a spectral DST/Fourier discretization (exact m_{eff}) shifts the per-t tip recovery at $t = a^2 = 0.0025$ from 1.31% (FD undershoot, G4-5a-refined) to 813.4% (spectral overshoot) of the continuum value $+1/6$. That value is the per-t target at every t: the continuum cone tip term is t-independent (the area and perimeter terms are linear in α and cancel in the replica derivative). At $t = a^2$ both schemes are inside the lattice scale and fail in opposite directions; the discrepancy is a UV discretization failure, not a missing target.

IR cutoff cure clean negative. Three cutoff variants (Gaussian, polynomial, sharp) tested across six $\Lambda \in \{0.5, 1, 1.5, 2, 3, 5\}$. None achieves the factor-2 gate at all Λ. The structural reason: the substrate's tip(t) profile peaks at $t \approx 0.05$–0.1 (a substrate-fixed UV scale tied to lattice spacing $a = 0.05$ and $r_h = 2$), independent of Λ. The continuum prediction $r_h^2\Lambda^2/3$ has clean Λ^2 scaling that the discrete extraction cannot inherit at fixed substrate. The proper cure is substrate UV refinement ($a \to a/2$, a compute scaling to $N_\rho \to 400$ beyond sprint scale), not a cutoff-function adjustment. A side correction: extending the t-grid from G4-5c's 8 points to 13 points shifts the $\Lambda = 2$ recovery from 0.85 to **1.96**, identifying $\Lambda \in [2, 3]$ as the valid extraction window at the current substrate.

Net effect on the G4-6 estimate. Three of the four G4-6 target structural closures have evidence in hand, each re-read since: the $\alpha > 1$ Möbius form is a finite-a artifact (B4, below), not an analytic closure; the per-t tip recovery at the lattice scale fails in both directions against the t-independent target 1/6; and the 2D-substrate cutoff ordering follows $\phi(0)$ under the proper-time estimator, which is the moment of the bare 2D tip, not of the 4D S_{BH} (Theorem 13.1).

13.7.7 Post-v3.19.0 follow-on thread (v3.20.0): closed-form per-t UV target and F14 closure

A single-thread follow-on of five tasks closed the v3.19.0 open items. Three findings are substantive enough to record in this paper.

F14 closure — sector-wise Mellin moment map at sub-5%. The v3.19.0 PARTIAL closure at 47.9% max deviation on the sharp-cutoff channel improved under two refinements: a 20-point log-spaced t-grid with three sharp-edge anchors at $t = 1/\Lambda^2$, and explicit analytical edge insertion at the sharp-cutoff discontinuity. Maximum deviation dropped to 4.82%, mean 2.67% across nine channels (3 cutoffs × 3 Λ). The sector-wise Mellin moment map $\{$2D tip (proper-time estimator) $\leftrightarrow \phi(0)$, EH $\leftrightarrow \phi(1)$, $\Lambda_{cc} \leftrightarrow \phi(2)\}$ is now empirically closed at strict sub-5% across the panel; F14 graduates from PARTIAL to POSITIVE.

Per-t UV target (withdrawn). An earlier printing posited a continuum target $\mathrm{tip}^{\mathrm{cont}}(t) = 1/(24\pi t) + \mathrm{const}$ on the 2D cone, built by combining the dimensionless Sommerfeld–Cheeger derivative with a $1/(4\pi t)$ planar prefactor. That combination is dimensionally inconsistent: the Sommerfeld–Cheeger term is the t^0 coefficient of the 2D heat trace, and the exact continuum tip (Bessel zeros on the disk, $\alpha \in \{1/2, 2/3, 3/2, 2\}$, $t/R^2 \in \{0.01, 0.02, 0.05\}$) is 0.166667 at every t. The continuum per-t target is therefore 1/6, the "5.305" target and the "all schemes undershoot" reading are withdrawn, and the spectral scheme's 813% at $t = a^2$ is an overshoot of 1/6.

Möbius $\alpha > 1$ extrapolation lock at $\alpha = 10$. Eq. (44) extrapolated to $\alpha \in \{4, 5, 10\}$ (not in the original three-point fit set) gives mean relative error 1.71%, with $\alpha = 10$ landing at $\mathbf{-0.032}$% rel. err. – essentially exact at the asymptote $F(\alpha) \to 1/2$. The decreasing relative error with α (3.3% at $\alpha = 1.5 \to 0.032$% at $\alpha = 10$) confirms the Möbius asymptote is structurally locked at the substrate scale $a \approx 0.05$, not a three-point coincidence – but *at fixed a*, like every other robustness check. The B4 continuum extrapolation above (the "Mechanism RESOLVED" paragraph) shows the radial $a \to 0$ limit drives the recovery off this Möbius lock toward the Sommerfeld–Cheeger value 1, so the lock is a feature of the finite-a form, not of the continuum. *Mechanism no longer open: the form is a finite-a artifact, the continuum is plain Sommerfeld–Cheeger.*

14 R2: replica-weight-harmless convergence of the cigar-tip entropy (L6)

The discrete-substrate program (§12–§13) reduced the gravity entropy question to a single irreducible theorem (charter `debug/gravity_campaign_R2_theorem_charter_memo.md`): does the discrete cigar-tip spectral functional converge so that $S_{\mathrm{tip}} \to A/4$ with a rate? The near-horizon cigar factorizes at constant warp into $D_\alpha^2 \otimes S^2(r_h)$, and (G4-3 phase-1b) the conical defect α lives entirely in the disk while the horizon sphere S^2 is a passive area multiplier $A = 4\pi r_h^2$. The entropy is the replica derivative of the *heat-trace spectral functional* (not of a propinquity metric):

$$S_{\mathrm{tip}} = \phi(0)[\mathrm{tip}(t)]\,, \qquad \mathrm{tip}(t) = \left.\frac{dK}{d\alpha}\right|_{\alpha=1} - K_{\mathrm{disk}}(t), \qquad K(\alpha, t) = \mathrm{Tr}\, e^{-tD_\alpha^2}. \tag{49}$$

On the discrete substrate (SPECTRAL azimuthal, spinor momentum $m_{\mathrm{eff}} = (k + \frac{1}{2})/\alpha$), $K(\alpha, t) = 4\sum_{j\geq 0} h((j + \frac{1}{2})/\alpha, t)$ with $h(m, t)$ the radial heat trace, and $\frac{dK}{d\alpha}|_1 = -4\sum_j (j + \frac{1}{2})h'(j + \frac{1}{2})$. The per-mode *replica contribution* $C_j = -4(j + \frac{1}{2})h'(j + \frac{1}{2}) > 0$ carries the $(j + \frac{1}{2})$ *replica weight*, heavier than the bare heat trace.

Write $n = (N_\phi, N_\rho, a)$ for the discrete truncation (N_ϕ azimuthal modes, N_ρ radial sites, spacing a, $R = N_\rho a$ fixed), and $g_j^{(n)}(\alpha) := 4\, h_n\big((j + \frac{1}{2})/\alpha,\, t\big)$, so that $K_n(\alpha, t) = \sum_{j\geq 0} g_j^{(n)}(\alpha)$ and the per-mode replica contribution is $C_j(\alpha) = -g_j^{(n)\prime}(\alpha)$. The continuum objects g_j, K are the $n \to \infty$ limits.

Theorem 14.1 (L6: replica-weight-harmless convergence — INTERNAL THEOREM; `tests/test_paper51_L6`) *Fix $t > 0$ and $\delta \in (0, 1)$. As $n \to \infty$ the maps $\alpha \mapsto K_n(\alpha, t)$ converge to $\alpha \mapsto K(\alpha, t)$ in $C^1([1 - \delta, 1 + \delta])$. In particular*

$$\lim_{n\to\infty} \left.\frac{dK_n}{d\alpha}\right|_{\alpha=1} = \left.\frac{dK}{d\alpha}\right|_{\alpha=1} = \left.\frac{d}{d\alpha}\right|_{\alpha=1} \lim_{n\to\infty} K_n, \tag{50}$$

and the convergence rate of the replica derivative equals the rate of the undifferentiated heat trace: the replica weight $(j + \frac{1}{2})$ does not degrade it. Consequently the cigar-tip entropy $S_{\mathrm{tip}} = \phi(0)[\,\mathrm{tip}(t)\,]$ inherits the convergence with the same rate.

The charter architecture has three layers, each with a lemma below. The theorem's proof uses Lemmas 14.3 (transported: the norm-resolvent heat-trace rate) and 14.4 (the new content). Lemma 14.2, argued by transport and **[OPEN]** at its disk-with-cone factor (the propinquity backbone, see the Grade paragraph), concerns geometric faithfulness and is not used.

Lemma 14.2 (Layer 1: propinquity backbone; **[OPEN]**). *For each fixed α near 1, the truncated constant-warp metric spectral triples $\mathcal{T}_{n,\alpha}$ converge to the continuum tip triple $\mathcal{T}_\alpha$ in the van Suijlekom state-space Gromov–Hausdorff distance, $d_{\mathrm{GH}}(\mathcal{T}_{n,\alpha}, \mathcal{T}_\alpha) \le C_3\,\gamma_n \to 0$, with C_3 truncation-independent (the full-disk metric statement is deferred; see the open questions).*

Source. At constant warp $r(\rho) = r_h$ the cigar squared-Dirac is a pure tensor product $D_\alpha^2 \otimes I + I \otimes D_{S^2}^2/r_h^2$ (warp cross-term vanishes, §12), so the triple factorises as disk-with-cone $\otimes$ horizon S^2. The azimuthal S^1 is the Connes–van Suijlekom Toeplitz operator system ([10], Prop. 4.2, prop= 2); the radial interval is the bounded-interval operator system with the apex centrifugally screened (Lemma 14.3); the S^2 factor is Berezin–Toeplitz (Hawkins, as in Paper 38 L4); the tensor assembly is PURE_TENSOR (Paper 45). This transport is **[ARGUED]** and does not close: the disk-with-cone factor is genuinely new (the $1/\rho^2$ term couples radial and azimuthal parts, and it is not a group), and no reconstruction map for it is established (see the Grade paragraph); the prop= 2 property of the assembled disk operator system is confirmed by direct computation below. The backbone is *undifferentiated*: it certifies geometric faithfulness at each α, not the replica derivative.

Lemma 14.3 (Layer 2: pointwise heat-trace rate). *For each fixed (α, t), $K_n(\alpha, t) \to K(\alpha, t)$ as $n \to \infty$, with error bounded by the sum of (i) a radial discretisation term $O(a^p)$ from the finite-difference polar Laplacian and (ii) super-polynomially small azimuthal and radial truncation terms. The radial operator $L(m) = -\partial_\rho^2 + (m^2 - \frac{1}{4})/\rho^2$ on $(0, R]$ has, for every $m = (j+\frac{1}{2})/\alpha > 0$, eigenfunctions $\sim \rho^{|m|}$ vanishing at the apex, so $\rho = 0$ is a regular endpoint requiring no boundary condition and the only boundary datum is the IR Dirichlet condition at $\rho = R$.*

Source. The radial factor is the bounded-interval norm-resolvent convergence of Paper 47, specialised to the centrifugally screened operator $L(m)$. The apex regularity is the no-zero-mode fact that makes the spinor conical-defect extraction clean (§13, G4-4; anti-periodic BC + half-integer $m \Rightarrow m \neq 0$). The exponent is $p = 1$ on the production grid $\rho_i = ia$ (a Dirichlet boundary-placement effect, pinned below) and $p = 2$ on the boundary-corrected grid.

Lemma 14.4 (Layer 3: the replica weight is harmless — new). *There is a constant $c(\delta) = t/((1+\delta)^2 R^2) > 0$ and, for each (t, δ), a summable Gaussian-dominating sequence*

$$G_j \;=\; C(t,\delta)\,(j+\tfrac{1}{2})^2\, e^{-c(\delta)(j+\frac{1}{2})^2}, \qquad \sum_{j\ge 0} G_j < \infty, \tag{51}$$

such that $\big|g_j^{(n)}(\alpha)\big| \le G_j$ and $\big|g_j^{(n)\prime}(\alpha)\big| \le G_j$ for all $\alpha \in [1-\delta, 1+\delta]$ and all n large. Hence $\sum_j g_j^{(n)}$ and $\sum_j g_j^{(n)\prime}$ converge uniformly on $[1-\delta, 1+\delta]$.

Proof. The centrifugal floor obeys $\lambda_0(m) \ge m^2/R^2$ (continuum Bessel-zero bound $j_{m,1} \ge m$; on the lattice the floor is at least as steep, $\lambda_0(m) \ge c_a\, m^2/R^2$ with $c_a \to 1^+$ as $a \to 0$, confirmed numerically below at $c_a \approx 1.16$). The excited gaps $\lambda_k(m) - \lambda_0(m)$ grow like $k^2\pi^2/R^2$ uniformly in m (the high modes see the box, not the centrifugal barrier), so the relative partition function $Z(t) = \sum_k e^{-t(\lambda_k - \lambda_0)}$ is bounded uniformly in m. Therefore

$$h(m,t) \le Z(t)\, e^{-tm^2/R^2}, \qquad |h'(m,t)| \le t\, C'\,(m/R^2)\, Z(t)\, e^{-tm^2/R^2}, \tag{52}$$

the second from $|\partial_m \lambda_k| = 2m\langle \rho^{-2}\rangle_k \le C' m/R^2$ (Hellmann–Feynman; the screened modes concentrate near $\rho \sim R$ so $\langle \rho^{-2}\rangle = O(R^{-2})$). For $\alpha \in [1-\delta, 1+\delta]$ and $m_j = (j+\frac{1}{2})/\alpha$ one has

$m_j \geq (j+\frac{1}{2})/(1+\delta)$, hence $e^{-tm_j^2/R^2} \leq e^{-c(\delta)(j+\frac{1}{2})^2}$, while $|g_j^{(n)\prime}| = 4|h'(m_j)|(j+\frac{1}{2})/\alpha^2$ carries a polynomial factor $\leq (j+\frac{1}{2})^2/(1-\delta)^2$. Both $|g_j^{(n)}|$ and $|g_j^{(n)\prime}|$ are thus bounded by (51), which is summable because a Gaussian beats any polynomial. The Weierstrass M-test gives uniform convergence of both series. □

Proof of Theorem 14.1. By Lemma 14.3, $K_n(\alpha,t) \to K(\alpha,t)$ pointwise. By Lemma 14.4 the series $\sum_j g_j^{(n)\prime}(\alpha)$ converges uniformly on $[1-\delta, 1+\delta]$ to $\sum_j g_j'(\alpha)$. The standard theorem on differentiation of series (pointwise convergence of $\sum g_j$ plus uniform convergence of $\sum g_j'$) gives $K(\cdot,t) \in C^1([1-\delta,1+\delta])$ with $K'(\alpha,t) = \sum_j g_j'(\alpha)$, and $K_n \to K$ in C^1. Evaluating at $\alpha = 1$ exchanges $\lim_n$ with $d/d\alpha$. The rate is the maximum of the Lemma 14.3 radial rate $O(a^p)$ and the super-polynomial truncation tail of (51); the latter is negligible, so the joint rate is the undifferentiated heat-trace rate $O(a^p)$ — the replica weight does not appear. The entropy $S_{\text{tip}} = \phi(0)[\text{tip}(t)]$, a continuous functional of $\text{tip}(t) = K'(\cdot,t)|_1 - K_{\text{disk}}$, inherits the convergence. □

The dominating exponent $c(\delta) \to t/R^2$ as $\delta \to 0$ is exactly the envelope slope measured below; the proof's analytical prediction and the numerics agree.

Numerical verification. (`debug/l6_replica_weight_harmless.py`; lean tridiagonal solver cross-checked bit-exact against the production `DiscreteDiskDiracSpectral`, $R = 10$, $t = 1$.) *(i) Gaussian envelope.* $C_j \sim \exp(\text{slope} \cdot (j+\frac{1}{2})^2)$, measured slope -0.01129 vs predicted $-t/R^2 = -0.01000$; the replica weight is crushed, C_j/C_0 falling from 4.7 at j=4 to 1.6×10^{-20} at j=64, and the azimuthal partial sum reaches the total to machine precision by $K_{\text{az}} \approx 48$ (tail/total $= 4 \times 10^{-13}$). *(ii) Uniformity.* The envelope slope across $\alpha \in [0.9, 1.1]$ lies in $[-0.01134, -0.01123]$ – essentially constant; the C^1 requirement holds. Direct α-uniformity test (`debug/g4_5a_alpha_uniformity.py`): the ratio $(K_{\text{fine}} - K_{\text{coarse}})/K_{\text{coarse}}$ varies by CV= 0.013 across $\alpha \in \{0.90, 0.95, 1.00, 1.05, 1.10\}$ at $(a, a') = (0.05, 0.025)$, confirming the discretisation error is α-independent to within 1.3%. *(iii) Rate.* Azimuthal convergence is super-polynomial (residual 0 by K_{az}=64); the lattice rate is $O(a^{1.1})$ at production scale (FD polar Laplacian at the screened apex). Extended UV-refinement panel (`debug/g4_5a_tip_uv_convergence.py`, $R = 10$, $t = 10$, $a \in \{0.10, 0.05, 0.025, 0.0125\}$, $N_\rho \in \{100, 200, 400, 800\}$): $B(a) = \{0.1585, 0.1628, 0.1650, 0.1662, 0.1$ errors $\{-4.9\%, -2.3\%, -1.0\%, -0.3\%, +0.04\%\}$. The error *changes sign* between $a = 0.0125$ (-0.30%) and $a = 0.00625$ ($+0.045\%$), confirming convergence to 1/6 from both sides. Richardson at order 1.7 from the pre-crossover pair $(0.025, 0.0125)$ gives $B_\infty = 0.166665$ (-0.001%, five significant figures). The convergence order accelerates from $p \approx 1.1$ at coarse resolution through $p \approx 1.7$ at intermediate scales, consistent with the structural error-cancellation in the replica difference (the leading $O(a)$ correction is α-independent and drops out of tip $= dK/d\alpha - K_{\text{disk}}$); the sign flip at $a = 0.00625$ confirms the sub-leading $O(a^2)$ correction has opposite sign. The joint rate is gated by the lattice / Paper-47 norm-resolvent rate, *not* degraded by the replica weight. *(iv) Derivative consistency.* At a single truncation the per-mode analytic $dK_n/d\alpha = 41.501652$ and a finite difference of K_n, 41.501681, agree to 2.9×10^{-5}. This checks the derivative implementation at fixed n; it does not test the interchange of $\lim_n$ with $d/d\alpha$, which rests on the proof of Theorem 14.1 (the assembly also needs the per-mode derivative convergence to be locally uniform in the mode index, which Lemma 14.3, pointwise in α, does not by itself supply).

Backbone (Layer 1, prop=2). The disk truncated operator system $O_n = P_n C(\text{disk}) P_n$, built from compressed disk monomials $z^a \bar{z}^b$ ($z = \rho e^{i\phi}$), has propagation number 2 at all tested truncations $(K, J) \in \{(2,2), (2,3), (3,2), (3,3)\}$: O^2 spans the full $M_N(\mathbb{C})$ envelope (`debug/l6_disk_prop2.py`). The disk backbone inherits the Connes–van Suijlekom 2-step-generating property of the Toeplitz S^1 ([10], Prop. 4.2) by direct computation. This confirms propagation number 2 only; the backbone itself is **[OPEN]**.

Structural error cancellation. The per-eigenvalue convergence rate of the discrete radial Laplacian is $O(a^1)$ (verified: `debug/g4_5a_layer2_resolvent.py`, $p = 0.99$ from consecutive-ratio analysis). But the *tip* converges at $O(a^{1.7})$ – faster by nearly a full power of a. The mechanism: the leading $O(a)$ error is α-independent (it comes from the apex boundary placement at $\rho = a$, which is geometrically α-neutral), so it cancels in the replica difference $\text{tip} = dK/d\alpha - K_{\text{disk}}$. The entropy is a "self-cleaning" observable on the discrete substrate – it automatically subtracts the dominant lattice artifact. This structural bonus is not assumed in the proof (which controls the full C^1 norm), but it explains the numerical observation that the entropy converges faster than the raw spectral data.

M1 / area signature. The passive S^2 Dirac heat trace satisfies $K_{S^2}(t)\, t \to 2r_h^2 = A/(2\pi) = \text{Vol}(S^2)\, r_h^2/(2\pi)$, so the horizon area $A = \text{Vol}(S^2)\, r_h^2$ enters S_{tip} through the sphere factor. The master Mellin engine M1 signature $4/\pi = \text{Vol}(S^2)/\pi^2$ – the same constant as the Fejér-kernel constant of Paper 38's GH bound (a kernel constant, not the convergence rate; Paper 38/40) and the Stefan–Boltzmann Matsubara prefactor – is therefore worn by the cigar entropy *by construction*, through the horizon-sphere area. (The wedge Hilbert–Schmidt orthogonality $1/\pi^2$ prefactor of Paper 42 §11.7 is the square of a posited normalisation $\beta = 2\pi/\kappa_g$ and is not shown to be this signature.) This is a by-construction signature, not an independent coincidence; it ties the cigar entropy to the Bernoulli ladder / master Mellin engine.

Grade. The new L6 content is Lemma 14.4 (the replica weight is harmless: the explicit Gaussian-dominating sequence (51) controls the mode series and its α-derivatives uniformly), which is proved here, together with the Theorem 14.1 assembly (uniform C^1 convergence $\Rightarrow$ $\lim_n$ commutes with $d/d\alpha$; joint rate = heat-trace rate). The two other lemmas: Lemma 14.2 (propinquity backbone) is **[OPEN]** — prop= 2 is confirmed by direct computation, but no reconstruction map for the disk-with-cone factor is established (see the next paragraph and Paper 53 [11], which proves convergence only on the flat plane and leaves the disk and the cone open); Lemma 14.3 (norm-resolvent rate) cites Paper 47. Every load-bearing claim is additionally verified numerically, and the proof's analytical dominating exponent $c(\delta) \to t/R^2$ matches the measured envelope slope. L6 is a Layer-2 convergence theorem (rates, not bit-exact identities), presented as such.

The exponent p of Lemma 14.3 and the continuum target are pinned below. *(B1, rate.)* Comparison of the discrete lowest eigenvalue $\lambda_0(m; a)$ against the exact continuum value $(j_{m,1}/R)^2$ (Bessel zero) gives order $q = 1.00$ at every $m \in \{\frac{1}{2}, \frac{3}{2}, \frac{5}{2}\}$, and the tip itself converges at the same order $p = 0.97$ – so $p = 1$ on the production grid $\rho_i = ia$. The mechanism is *Dirichlet boundary placement*, not the centrifugal singularity: the $\rho_i = ia$ grid sets the IR wall at $(N_\rho + 1)a = R + a$, displaced $O(a)$ from R. This is confirmed two ways: (i) the free case $m = \frac{1}{2}$ (zero centrifugal potential) still converges at $O(a)$, ruling out the $1/\rho^2$ term; (ii) the corrected grid $a = R/(N_\rho + 1)$ (wall exactly at R) restores the standard central-difference $O(a^2)$ (error ratio 4, three orders of magnitude smaller). The $O(a)$ rate is therefore a fixable substrate detail; $L(m)$ inherits the plain-interval Paper-47 norm-resolvent convergence, at $O(a)$ on the production grid and $O(a^2)$ on the boundary-corrected grid. *(B2, target.)* With the correct $O(a)$ order, linear-in-a extrapolation of the tip gives 0.166656 (2-point) and 0.166582 (least squares), versus $1/6 = 0.166667$ – agreement to -0.006% and -0.05%. The continuum replica derivative $dK/d\alpha|_1$ is the G4-4c Sommerfeld–Cheeger spinor-cone value $1/6$; the earlier -1.4% residual was an artifact of an order-2 Richardson on an $O(a)$ sequence. Both sub-pieces are thus closed, with the boundary-placement diagnosis correcting an earlier "apex-edge" guess. The $D_\alpha^2 \otimes S^2$ backbone (Lemma 14.2) remains **[OPEN]**: no reconstruction map for the disk-with-cone factor is known; it does not affect the Theorem 14.1 convergence statement. A de-risking attempt located the difficulty: the genuinely-new factor is the 2D disk-with-cone itself (a manifold-with-boundary, not a pure radial⊗azimuthal tensor, since the $1/\rho^2$ term couples them, and not a

group, so the central-Fejér reconstruction does not transport); the unital Markov-normalized Cesàro–Berezin map $B_n(f) = P_n M_g P_n$, $g = (\sum_a w_a \langle \psi_a, f\rangle \psi_a)/(\sum_a w_a \langle \psi_a, 1\rangle \psi_a)$ with Cesàro weight $w_a = (1-\lambda_a/\Lambda)_+^s$, is exactly unital, but it is *not* positivity-preserving: a non-negative ring observable reconstructs with negative values at Cesàro order $s = 2$ and $s = 4$ (Paper 53 [11], **[MEASURED]**). An earlier version of this paragraph claimed interior positivity at $s \geq 2$ (checked only on one observable), an interior rate $\Lambda^{-1.30}$, and a boundary obstruction at $\rho = R$; the rate was withdrawn earlier, and the boundary obstruction was a floating-point artifact of evaluating the 0/0 Markov ratio exactly at $\rho = R$ (off that point the error decays). All three are withdrawn (2026-09-29), and with them the earlier statement that Lemma 14.2 was upgraded from cited to proved. The lemma is **[OPEN]**: Paper 53 proves state-space convergence only on the flat plane (by a positive squared-window reconstruction and translation invariance), and the cone $\alpha \neq 1$ has no translation group. Theorem 14.1 does not depend on it.

15 Discussion: structural-skeleton-scope reading

The gravity arc produces two distinct classes of result:

Structural findings (G1, G2, G3, G4-1, G4-2, G5, G6-Diag-Full): cutoff-independent properties of the framework (closed forms, theorems, structural mechanisms). These are framework predictions in the strong sense:

- $\zeta_{\text{unit}}(-k) = 0$ identity and two-term exactness in the spinor sector;
- heat-kernel factorization on $S^3 \times S^1_\beta$ producing exact Einstein–Hilbert plus cosmological constant;
- bundle-specific exactness (spinor only, not scalar or tensor);
- Bekenstein–Hawking entropy form $S = A/(4G)$ from conical-defect / replica;
- de Sitter vacuum on the decompactified product;
- physical $(1, 1)$ modes with positive kinetic eigenvalue.

Numerical gravity predictions (G7): specific values of $G_{\text{eff}} = 3\pi/\Lambda^2$, $\Lambda_{cc} = 6\Lambda^2$. These depend on the Gaussian cutoff choice, and G8 shows they generalize cleanly to arbitrary cutoff with ϕ-moment-dependent prefactors.

The structural-skeleton-scope reading. GeoVac determines the *structural skeleton* of gravity at the spectral action level. The framework does not autonomously generate Yukawa-like calibration data; the cutoff function and its ϕ-moments are external input. This is the same structural-skeleton-scope pattern observed across the framework [7, 8], sharpened here for the gravity sector.

Projection-by-projection sharpening of the structural-skeleton scope. A systematic walkthrough of Paper 34's twenty-eight projections (see `debug/sprint_projection_specific_calibration_sc` 2026-05-29) finds that only *five* projections carry parametric Class-D calibration data (spectral action cutoff function §III.6; gauge couplings §III.10; rest masses §III.14; fine structure constant §III.16; Born rule §III.28), and of those, only *two* show known internal structure in their calibration values (the Koide cone for charged leptons in §III.14, at sub-arcsecond precision; and the Paper 2 K-formula coincidence for α in §III.16, K to 4.8×10^{-7} and the cubic's root to 8.8×10^{-8}). The remaining 23 projections carry no parametric calibration to test. This 2-of-28 concentration is a sharp form of the structural-skeleton-scope statement: calibration data

is genuinely external for the overwhelming majority of projections; the two known exceptions are framework observations that have already been investigated and remain open as numerical observations without derivation. The gravity sector's cutoff function falls into the "no internal structure" category alongside the gauge couplings.

Why this matters. The Connes–Chamseddine spectral action program [13, 16] has long pursued "gravity from noncommutative geometry" but typically requires the cutoff function to be specified externally. GeoVac inherits this situation: structural shape predicted, calibration data external. The framework's contribution is not to circumvent this calibration issue but to organize the gravity-sector predictions into a clear separation of *what GeoVac determines* versus *what is external Class 1 calibration data*.

Comparison to alternative gravity readings. The discrete-substrate approach (G4-3 onwards) differs from continuum quantum-gravity programs (loop quantum gravity, causal dynamical triangulation, AdS/CFT, asymptotic safety) in two main ways:

1. The substrate is not a quantization of a classical gravitational degree of freedom; it is the Fock-projection of the hydrogen spectrum, with gravity emerging from the spectral action on product geometries.

2. The framework's calibration data (cutoff function, Yukawas, etc.) is treated as external Class 1 input, not derived from the substrate.

The discrete-substrate arc opening (G4-3) is the natural extended follow-on: if the continuum arc shows what the GeoVac substrate can say about gravity at the spectral action level, the discrete-substrate arc shows what it says *intrinsically*, before passing to the continuum limit.

Empirical verification at G4-3 closure. The G4-3d-UV sweep (Section 12.3) demonstrates that the discrete substrate reproduces the leading Weyl term at $t \approx 0.1$ when N_ϕ is refined into the UV regime, at the level of the Dirichlet perimeter correction that the comparison omits. The slight overshoot at fine N_ϕ and small t is identified as a structural artifact of the azimuthal FD discretization (ratio $4/\pi^2 \approx 0.405$ between discrete and continuum angular Laplacian at the truncation edge), not a Hermiticity or implementation issue. The framework's continuum-limit prediction is robust at the discrete-substrate level when the discretization choices are properly characterized.

Empirical verification at G4-4 closure. The G4-4 program (Section 13) extends this empirical verification from the Weyl law to the load-bearing quantities for S_{BH}. The headline: the spinor conical-defect tip coefficient $-\frac{1}{12}(1/\alpha - \alpha)$ is approached by the discrete substrate as $a \to 0$ (§13.4); the replica derivative $+1/6$ is extracted at **96.69%** (§13.5). Both results land at sprint scale on the same UV-refined panels that verified the Weyl law — there is one substrate, and the bridges to continuum gravity are layered on top.

The BC sector diagnostic. A structurally substantive observation from this session: the half-integer angular momentum structure (anti-periodic spinor BC, $m \in \mathbb{Z} + 1/2$) is *essential* for clean conical-defect SC extraction on a discrete substrate. The integer scalar BC has a zero-mode whose contribution breaks the $\alpha \leftrightarrow 1/\alpha$ symmetry at finite discretization, biasing the scalar SC extraction by $\sim 34\%$ in the same regime where the spinor extracts cleanly. **This is consistent with continuum NCG expectations** (the Dirac operator with its half-integer spinor bundle is the natural object on a spectral triple, not the scalar Laplacian), here seen on the discrete substrate.

16 Open questions

Q1 (closed at sprint scale, major program remaining). The discrete-substrate program is now closed at G4-3 + G4-4 sub-sprint scale (19 sub-sprints). The remaining commitment is G4-5 (discrete replica method) and G4-6 (full S_{BH} derivation), a substantial multi-stage program end-to-end.

Update from G4-4 sprint-scale closure: the original Q1 statement ("the literal Sommerfeld–Cheeger extraction is a G4-5 target, not sprint-scale") was correct for the *scalar* wedge-lattice diagnostic (G4-3c-proper, T1), where the discretization floor exceeds the SC tip term. However, on the *spinor* wedge (G4-4c), the SC tip coefficient $-(1/12)(1/\alpha-\alpha)$ is approached as the radial mesh refines on the same substrate. The half-integer angular momentum + anti-periodic BC of the spinor sector cleanly extracts the conical-defect tip term where the integer scalar sector does not. This refines Q1: G4-5/G4-6 builds on a quantitatively validated foundation, not on a sketch.

Open beyond sprint scale: (i) resolved: the $\alpha > 1$ recovery deficit is a finite-a artifact (B4); (ii) the joint variable-warp + conical-defect Dirac (cigar with conical singularity, not addressed in any of G4-4a/b/c separately); (iii) the integration over t with cutoff regularization (G4-5 + G4-6, the standard Connes–Chamseddine integration).

Q2 (resolved). The factor of 2 once found between G7's G_{eff} and G4-2's $G_N = 3\pi/\Lambda^2$ was the G2 spinor rank: the earlier G2 trace ran over a rank-2 bundle, and the rank-4 Dirac trace gives $G_{\mathrm{eff}} = 3\pi/\Lambda^2$, as Wald's theorem requires in the two-term-exact Einstein sector (see the G4-2 remark). It never came from a scalar-vs-Dirac cone coefficient.

Q3. The cosmological-constant-scale gap of $\sim 10^{120}$ is inherited from standard CC. Whether the discrete-substrate program (G4-3 onwards) can illuminate this gap is open. The naive expectation is that the discrete substrate inherits the same gap; surprises would be substantial.

Q3 sharpened (post-G4-5 sector-wise Mellin moment map). The sector-wise Mellin moment map {2D tip (proper-time estimator) $\leftrightarrow \phi(0)$, EH $\leftrightarrow \phi(1)$, $\Lambda_{cc} \leftrightarrow \phi(2)$} allows a precise framework-internal statement of the CC problem. From §10, the dimensionless cosmological constant in Planck units is $\Lambda_{cc} \cdot G_{\mathrm{eff}} = 18\pi\,\phi(2)/\phi(1)^2$. For Gaussian ($\phi(1) = \phi(2) = 1$), this is $18\pi \approx 57$; for polynomial or sharp cutoffs, $O(1)$. Observation requires $\Lambda_{cc} \cdot G_{\mathrm{eff}} \sim 10^{-122}$. *The precise framework-internal statement of the cosmological-constant problem is therefore the requirement*

$$\frac{\phi(2)}{\phi(1)^2} \;\approx\; 10^{-124} \quad \text{with} \quad \phi(0),\, \phi(1) \;=\; O(1).$$

No natural cutoff function in the standard family satisfies this; the identification of a cutoff that does would constitute a calibration-data selection principle distinct from the Higgs–Yukawa selection problem addressed by Sprint H1 [6]. A diagnostic scoping pass (`debug/sprint_cc_scoping_memo.md`, 2026-05-29) confirmed that none of the framework's structural mechanisms (master Mellin engine constraint, two-term exactness, discrete substrate, thermal cancellation, or inner-factor calibration) provides a ϕ-moment selection rule of the required magnitude.

Q4. The G6-Full sprint (2026-05-31) advanced beyond the necessary-condition diagnostic via the CG-algebraic approach (SO(4)-projected quadratic form on the $(1,1)$ subspace). Three structural findings: (i) the discrete Laplacian $\|[D,V]\|^2/\|V\|^2$ on the $(1,1)$ subspace has *bit-exact integer* eigenvalues $(2k)^2$ ($k = 1,2,3,4$ at $n_{\max} = 3$) with ratios 1:4:9:16; (ii) the Seeley–DeWitt a_1 coefficient (Einstein–Hilbert content) is positive for physical modes and negative for gauge modes; (iii) the Λ-sweep at $\Lambda^2 = 4$ gives first positive-eigenvalue ratio 2.209, within 2% of the Lichnerowicz ratio $13/6 = 2.167$. The Fierz–Pauli decomposition (Section 9) further establishes

that $S^{(2)}$ is J-blind within $(1,1)$ (Theorem 9.1): the spectral action has extracted all the graviton information it can, and the TT vs. trace distinction requires the metric identification.

Q5. The choice of cutoff function (Class 1 calibration data per G8) is structurally external. Whether GeoVac admits an intrinsic mechanism for selecting a preferred cutoff is open. The working hypothesis is that no such mechanism exists within the substrate alone; calibration is genuinely external, as elsewhere in the framework.

Q6 (closed). Does Bisognano–Wichmann wedge entanglement entropy reproduce $S_{\mathrm{BH}} \propto A$? **No.** The BW thermal-state entanglement entropy on the hemispheric wedge scales as $S \sim 2\log n_{\max}$ (best fit $S = 1.963 \log n_{\max} + 0.540$, $R^2 = 0.9999$; area-law fit $R^2 = 0.83$, strongly rejected), verified at $n_{\max} \in \{2, \ldots, 12\}$. The mechanism: the Boltzmann weights $e^{-\mathrm{two_m}_j}$ are exponentially suppressed for large $\mathrm{two_m}_j$; at every $n_{\max}$, 89–96% of the probability sits on the ground shell with degeneracy $n_{\max}(n_{\max}+1) \sim n_{\max}^2$, giving $S \approx \log(n_{\max}^2) = 2\log n_{\max}$. The Bekenstein–Hawking $S_{\mathrm{BH}} = A/(4G)$ of §6 therefore comes from the spectral-action replica mechanism, *not* from BW entanglement entropy. The two routes to S_{BH} in quantum gravity (entanglement vs. spectral action) are structurally distinguishable on the GeoVac spectral triple, and the framework selects the spectral-action route. Source: `debug/bh_phase0_entanglement_entropy_memo.md`.

Acknowledgments

The gravity arc consisting of sprints G1 through G8 plus the discrete-substrate opening G4-3 was completed on 2026-05-28 in a single session. The PI directed the sprint sequence; sprint memos in the GeoVac repository `debug/` directory document the detailed computational basis. This paper organizes the sprint sequence as a self-contained narrative for the gravity / spectral action / NCG audience. The detailed numerical machinery underlying each sprint is documented in Paper 28 §4.7–§4.17 [5].

References

[1] GeoVac Project. Paper 0 – Counting Angular Momentum Labels by Isotropic Packing. `papers/group3_foundations/Paper_0_Geometric_Packing.tex`.

[2] GeoVac Project. Paper 2 – The Fine Structure Constant as a Spectral Coincidence on the Hopf Fibration. `papers/group5_qed_gauge/paper_2_alpha.tex`.

[3] GeoVac Project. Paper 7 – The Dimensionless Vacuum: Recovering the Schrödinger Equation on a Scale-Invariant Label Lattice. `papers/group3_foundations/Paper_7_Dimensionless_Vacuum.tex`.

[4] GeoVac Project. Paper 24 – The Bargmann–Segal Lattice: π-Free Discretization of the Harmonic Oscillator on S^5. `papers/group3_foundations/paper_24_bargmann_segal.tex`.

[5] GeoVac Project. Paper 28 – QED on S^3: Spectral Geometry of Perturbative Transcendentals. `papers/group5_qed_gauge/paper_28_qed_s3.tex`.

[6] GeoVac Project. Paper 32 – The GeoVac Spectral Triple: A Synthesis Paper Locking the Framework into the Marcolli–van Suijlekom Gauge-Network Lineage. `papers/group1_operator_algebras/paper_32_spectral_triple.tex`.

[7] GeoVac Project. Paper 34 – GeoVac as a Two-Layer Framework: Projection Taxonomy and the Empirical Matches Catalog. `papers/group6_precision_observations/paper_34_projection_taxonomy.tex`.

[8] GeoVac Project. Paper 35 – Time as Projection: The Klein-Gordon Spectrum on $S^3 \times \mathbb{R}$ and the Algebraic / Observation Split. `papers/group6_precision_observations/paper_35_time_as_projection.tex`.

[9] GeoVac Project. Paper 38 – State-space Gromov–Hausdorff convergence of truncated Camporesi–Higuchi spectral triples on S^3. `papers/group1_operator_algebras/paper_38_su2_propinquity_convergence.tex`.

[10] GeoVac Project. Paper 44 – Operator-system extension of the BBB Krein spectral triple at finite cutoff: propagation number and envelope dependence. `papers/group1_operator_algebras/paper_44_lorentzian_operator_system.tex`.

[11] GeoVac Project. Paper 53 – Spectral truncation of the flat plane: state-space convergence with a squared window, and the disk-with-cone left open. `papers/group1_operator_algebras/paper_53_disk_propinquity.tex`.

[12] R. Camporesi and A. Higuchi, "On the eigenfunctions of the Dirac operator on spheres and real hyperbolic spaces," *J. Geom. Phys.* **20**, 1–18 (1996).

[13] A. H. Chamseddine and A. Connes, "The spectral action principle," *Comm. Math. Phys.* **186**, 731–750 (1997).

[14] F. Fathizadeh and M. Marcolli, "Periods and motives in the spectral action of Robertson–Walker spacetimes," *Comm. Math. Phys.* **356**, 641–671 (2017), arXiv:1611.01815.

[15] A. H. Chamseddine and A. Connes, "The uncanny precision of the spectral action," *Comm. Math. Phys.* **293**, 867–897 (2010). arXiv:0812.0165.

[16] A. H. Chamseddine and A. Connes, "Noncommutative geometry as a framework for unification of all fundamental interactions including gravity," *Fortsch. Phys.* **58**, 553–600 (2010).

[17] J. Cheeger, "Spectral geometry of singular Riemannian spaces," *J. Differential Geom.* **18**, 575–657 (1983).

[18] J. S. Dowker, "Quantum field theory on a cone," *J. Phys. A* **10**, 115–124 (1977).

[19] M. Beccaria and A. A. Tseytlin, "C_T for conformal higher spin fields from partition function on conically deformed sphere," *J. High Energy Phys.* **09**, 123 (2017).

[20] A. Sommerfeld, "Über verzweigte Potentiale im Raum," *Proc. London Math. Soc.* **28**, 395–429 (1896).

[21] S. N. Solodukhin, "The conical singularity and quantum corrections to entropy of black hole," *Phys. Rev. D* **51**, 609–617 (1995).

[22] V. P. Frolov and D. V. Fursaev, "Plenty of nothing: black hole entropy in induced gravity," *Phys. Rev. D* **56**, 2212–2225 (1997).

[23] D. V. Fursaev and S. N. Solodukhin, "On the description of the Riemannian geometry in the presence of conical defects," *Phys. Rev. D* **52**, 2133–2143 (1995), arXiv:hep-th/9501127.

[24] D. V. Fursaev and G. Miele, "Cones, Spins and Heat Kernels," *Nucl. Phys. B* **484**, 697–723 (1997), arXiv:hep-th/9605153.

[25] S. N. Solodukhin, "Entanglement Entropy of Black Holes," *Living Rev. Relativity* **14**, 8 (2011), arXiv:1104.3712.

[26] R. M. Wald, "Black hole entropy is the Noether charge," *Phys. Rev. D* **48**, R3427–R3431 (1993), arXiv:gr-qc/9307038.

[27] J. S. Dowker, "Heat kernels on curved cones," *Class. Quant. Grav.* **11**, L137–L140 (1994), arXiv:hep-th/9406002.

[28] C. Rovelli, *Quantum Gravity* (Cambridge University Press, Cambridge, 2004).

[29] J. Ambjørn, J. Jurkiewicz, and R. Loll, "Reconstructing the universe," *Phys. Rev. D* **72**, 064014 (2005).

[30] V. Homšak and S. Veroni, "Boltzmannian state counting for black hole entropy in causal set theory," *Phys. Rev. D* **110**, 026015 (2024), arXiv:2404.11670.

[31] M. Marcolli and W. D. van Suijlekom, "Gauge networks in noncommutative geometry," *J. Geom. Phys.* **75**, 71–91 (2014).

[32] GeoVac Project. Sprint G4-3 – Discrete warped-product substrate for cigar. `debug/g4_3_warped_substrate_memo.md`, May 2026.