

Krein-pointed proper quantum metric spaces and the bridge to Mondino–Sämman Lorentzian pre-length spaces, with application to the GeoVac hemispheric wedge

J. Loutey*

May 2026 (preprint)

Abstract

We construct a categorical bridge between Latrémolière’s pinned proper quantum metric space (PPQMS) framework [25] and Mondino–Sämman’s covered Lorentzian pre-length space (LPLS) framework [30] (descoped; see the Status note: the bridge’s domain category is instantiated on the Paper 45 substrate, whose Lipschitz seminorm is degenerate, so the metric-level theorems are open pending repair and the bridge survives as categorical design). The bridge is a contravariant functor

$$\mathbf{W} : \mathbf{KreinMetaMet}_{\text{pp}} \longrightarrow \mathbf{LorPLG}_{\text{cov}}$$

sending a Krein-pointed proper QMS $(\mathcal{A}^K, L^K, \mathcal{M}^L, \omega_W^L)$ to a covered LPLS $(\hat{\mathcal{M}}^L, \ell^L, \hat{\omega}_W^L, \hat{\mathcal{U}}^L)$ via Gelfand spectrum of the M-diagonal Abelian topography, Wick-rotated modular flow as time separation, the Bisognano–Wichmann vacuum as basepoint, and admissible-scaling truncated covers. The structural mismatch between Latrémolière’s 4-point relaxed forward metametric and Mondino–Sämman’s 3-point reverse triangle (the F2 mismatch) is resolved via the Connes–Rovelli thermal-time hypothesis [10]: the Krein metametric measures *thermal time* of the wedge KMS state, the synthetic time separation measures *geometric proper time*, and the Wick rotation between them effects the analytic continuation that maps the additive (forward-triangle) thermal-time structure to the super-additive (reverse-triangle) geometric-time structure.

Main theorem. The Wick-rotation functor $\mathbf{W}$ is well-defined and the Krein–Mondino–Sämman Bridge Theorem (Theorem 5.4) holds at theorem-grade rigor for the *structural* legs — (B1) structural correspondence on 15 rows and (B3) pre-compactness inheritance via Paper 44 [48] propagation-number bound — while the metric-level legs ride Paper 45’s degenerate Krein seminorm: (B2) the reverse triangle inequality holds on-orbit via modular-flow equality and is *structurally vacuous off-orbit* (the M-diagonal topography forces orbit-pair contradiction), a structural/categorical property rather than a metric measurement; and (B4) convergence transport via Plancherel-nested Berezin compatibility with the Mondino–Sämman Def 3.6 axioms is **descoped** — conditional pending substrate repair, not an established metric convergence (see the Status note).

GeoVac wedge application. The bridge applies cleanly to the Paper 43 [47] hemispheric wedge at every finite cutoff. Seven newly accessible theorems follow (Theorems 6.2–6.8 + Corollary 6.9), headlined by **T6** (descoped; see the Status note): a *proposed G2* metric-level closure for the GeoVac wedge, *conditional on the upstream metric repair* — it inherits Paper 45’s Krein-side Λ^L , which is a degenerate seminorm (an evaluation of a closed-form rate formula, not a distance), so T6 is not yet a metric-level closure. The synthetic rate panel

*Independent researcher. jloutey@gmail.com. This work was developed in an AI-augmented agentic workflow with Anthropic’s Claude. Implementation, internal memos, and bibliographic collation were performed collaboratively; all theorems, design choices, and editorial decisions are the author’s responsibility.

transports the bit-exact values $\{2.0746, 1.6101, 1.3223\}$ from that formula (rate-formula values, not metric measurements). What survives unconditionally is the categorical bridge design (below); the quantitative pLGH-convergence reading is the repair target.

Honest scope. The bridge is K^+ -weak-form only. The strong-form bridge extending **W** to the Paper 46 [50] Appendix B enlarged substrate with chirality-flipping generators (Q1') is deferred as a substantial NCG-mathematics follow-on, beyond sprint scale. The general G2 metric-level closure on non-wedge carriers remains the named open frontier; we close only the wedge-specific case via the bridge image on the synthetic side. The construction preserves **W**'s functorial (not isometric) character throughout: the Krein metametric $\mathfrak{D}^K$ and the synthetic time separation ℓ^L measure different physical quantities (thermal vs. geometric time) of the same wedge KMS state.

Place in the math.OA standalone series. Tenth in the GeoVac math.OA arc (siblings: Papers 29 [41], 32 [42], 38 [43], 39 [44], 40 [45], 42 [46], 43 [47], 44 [48], 45 [49], 46 [50], 47 [51]). The result composes Paper 45/46 verbatim on the Krein-algebraic side with the Mondino–Sämman synthetic-Lorentzian Gromov–Hausdorff framework on the synthetic side, mediated by the Connes–Rovelli thermal-time identification.

Status and scope (June 2026)

Three issues are established upstream of this paper: (i) the K^+ -restricted compressed-Dirac Lipschitz seminorm underlying Λ^L vanishes identically on the truncated operator system — Krein-self-adjointness of the anti-Hermitian spatial part $iD_{GV} \otimes I$ forces $\{J_L, D_{GV} \otimes I\} = 0$, so the K^+ compression annihilates the spatial Dirac, and the momentum-diagonal temporal multipliers commute with ∂_t (verified bit-exact at $(n_{\max}, N_t) \in \{(2, 3), (3, 5)\}$); (ii) the “Latrémolière Thm 5.5 / Def. 3.4 / Def. 3.5” references cited for the tunneling-pair bound do not exist in the cited source — the device actually used is van Suijlekom’s state-space Gromov–Hausdorff framework; (iii) the constant $2/(n_{\max} + 1)$ is the maximum of the Plancherel mass distribution, not a cb-norm of any map used in the proofs (the smoothing map is unital completely positive, with cb-norm 1).

Consequently the bridge functor’s domain category **KreinMetaMet**_{pp}, as instantiated on the Paper 45 substrate, is *degenerate as a metric object*: the Lipschitz seminorms feeding the Krein metametric have kernels far exceeding the scalars, so the instantiated objects are not quantum (proper) metric spaces in the cited frameworks. T3 (Theorem 6.4) transports formula values, not metric data; T6 (Theorem 6.8, the G2 metric-level closure for the GeoVac wedge) rests on the same degenerate quantity and is *descope*d. What survives is the categorical design — the Wick-rotation functor **W** and the Connes–Rovelli thermal-time reading resolving the F2 forward-vs-reverse triangle mismatch — as a proposal conditional on repair of the underlying substrate. The Lorentzian quantum-metric convergence question is an open named research target.

Contents

1	Introduction	4
1.1	Motivation: the F2 forward-vs-reverse triangle mismatch	4
1.2	Main result: the Wick-rotation functor and the Krein–Mondino–Sämman Bridge .	5
1.3	GeoVac wedge application: seven newly accessible theorems	6
1.4	Honest scope	7
1.5	Roadmap of the paper	7
1.6	Related work	8
2	Setup	8
2.1	Latrémolière’s pinned proper QMS framework	8

2.2	Krein operator-system substrate from Papers 44/45/46	9
2.3	Mondino–Sämman pre-length space framework	10
2.4	Bisognano–Wichmann vacuum and the four-witness Wick-rotation theorem	10
2.5	Connes–Rovelli thermal-time hypothesis	11
2.6	Notation and conventions	11
3	Krein-pointed proper quantum metric spaces	11
3.1	The Krein-Leibniz Lipschitz seminorm	12
3.2	The Krein-separable QLCMS and Krein-pinned QLCMS	12
3.3	The Krein L-Lipschitz ω_W^L -pinned exhaustive sequence and the approximate-unit theorem	13
3.4	Krein topography and Krein-pointed proper QMS	13
3.5	Krein M-tunnels with extent element	14
4	Krein-pointed propinquity hypertopology	15
4.1	Extent of a Krein M-tunnel	15
4.2	The Krein M-local quantum metametric	15
4.3	Coincidence	16
4.4	The Krein quantum metametric and hypertopology	16
4.5	Compact-case agreement at $N_t = 1$	16
4.6	Numerical panel	17
5	The Mondino–Sämman bridge: Wick-rotation functor	17
5.1	The F2 mismatch and three candidate resolutions	17
5.1.1	R1: Negate time	17
5.1.2	R2: Functorial composite of Wick-rotation + modular-flow	17
5.1.3	R3: Connes–Rovelli thermal time as unifying framework	17
5.2	Definition of the Wick-rotation functor	18
5.3	The Krein–Mondino–Sämman Bridge Theorem	18
5.4	Structural correspondence (B1)	19
5.5	Reverse triangle inequality (B2): Decomposition O on the M-diagonal topography	19
5.6	Pre-compactness inheritance (B3)	20
5.7	Convergence transport (B4) via Plancherel-nested Berezin	21
5.8	Riemannian-limit recovery cross-check	21
5.9	Honest scope of the bridge theorem	21
6	Application to the GeoVac hemispheric wedge	22
6.1	The GeoVac wedge as a Krein PPQMS	22
6.2	The seven newly accessible theorems	22
6.2.1	T1: The wedge as Mondino–Sämman covered LPLS	22
6.2.2	T2: Synthetic compactness of the wedge family	22
6.2.3	T3: Synthetic bit-exact panel inheritance	23
6.2.4	T4: Synthetic four-witness readings	23
6.2.5	T5: Synthetic Pythagorean orthogonality with $1/\pi^2$ M1 signature	24
6.2.6	T6: G2 metric-level closure for the GeoVac wedge	24
6.2.7	T6a: Synthetic three-carrier identification	25
6.3	Honest scope of the wedge application	25

7	Synthetic-Lorentzian compactness theorems	25
7.1	Synthetic compactness of the GeoVac wedge family	25
7.2	The first quantitative pLGH panel from operator-algebraic input	25
7.3	G2 metric-equivalent closure for the GeoVac wedge (proposed/descoped)	25
7.4	Synthetic three-carrier identification	26
7.5	Why this is genuinely new content for Mondino–Sämman pLGH theory	26
7.6	Related synthetic-Lorentzian convergence frameworks	26
8	Open questions	26
8.1	Q1’: Strong-form bridge on the enlarged substrate	26
8.2	G2 metric-level closure on the general (non-wedge) carrier	27
8.3	G3 cross-manifold $\mathcal{T}_{S^3} \otimes \mathcal{T}_{\text{Hardy}}(S^5)$	28
8.4	G4 inner-factor calibration data	28
8.5	Q2’: Non-commutative Mondino–Sämman pre-length spaces	28

1 Introduction

1.1 Motivation: the F2 forward-vs-reverse triangle mismatch

Two recent advances in the convergence theory of generalized Lorentzian geometric structures, both published in December 2025, provide complementary but *a priori* incompatible frameworks:

- **Latréolière’s pinned proper quantum metric space framework** [25] (arXiv:2512.03573, December 3, 2025) extends the Latréolière propinquity [22, 24] from quantum *compact* metric spaces to a quantum Gromov–Hausdorff *hypertopology* on the class of pointed proper quantum metric spaces. The resulting metametric $\mathbb{D}^K$ satisfies a 4-point *relaxed forward* triangle inequality with codomain $[0, \infty]$. The framework is operator-algebraic: objects are C^* -algebras with Lipschitz seminorms, topographies (Abelian C^* -subalgebras), and pin states; morphisms are topographic quantum M-isometries.
- **Mondino–Sämman’s covered Lorentzian pre-length space framework** [30] (arXiv:2504.10380 v4, December 9, 2025) extends the synthetic Lorentzian geometry of Kunzinger–Sämman pre-length spaces [19] to a pointed Lorentzian Gromov–Hausdorff (pLGH) convergence theory on covered LPLS. The synthetic time separation ℓ satisfies the 3-point *reverse* triangle inequality (causal-curve super-additivity of proper time) with codomain $\{-\infty\} \cup [0, \infty]$. The framework is synthetic-geometric: objects are sets with timelike-causal pre-orders, cover slabs, and basepoint events.

The two frameworks share many structural elements: pinning, exhaustive covers, slab bounds, ε -nets, pre-compactness, Gromov–Hausdorff convergence. They are aimed at different aspects of the same physics: generalized-geometric approximation of Lorentzian objects. However, the *algebraic shapes* of their central inequalities are incompatible:

	Latréolière 2512.03573	Mondino–Sämman 2504.10380
codomain	$[0, \infty]$	$\{-\infty\} \cup [0, \infty]$
triangle direction	forward	reverse
triangle structure	4-point relaxed	3-point
sign convention	state-distance (≥ 0)	proper time (≥ 0 when causal)

We call this the **F2 mismatch**. Naive attempts at a bridge fail: negating the time variable (R1) puts the codomain in the wrong half-line and the 4-point structure does not collapse to 3-point under sign reversal. The substantive question is whether a bridge can be *categorical*: a

functor sending Krein-algebraic objects (where relevant) to synthetic-Lorentzian objects (where relevant), with the F2 mismatch interpreted as a structural signature of two distinct projection mechanisms applied to the same physical object, rather than as a defect to be reconciled.

1.2 Main result: the Wick-rotation functor and the Krein–Mondino–Sämman Bridge

What this paper contributes and what it inherits. From Mondino–Sämman [30] we inherit the covered Lorentzian pre-length space framework, the synthetic time separation ℓ , the 3-point reverse triangle inequality, the pLGH-convergence axioms (Def. 3.6), and the synthetic pre-compactness theorem (Thm. 6.2). From Latrémolière [25] we inherit the pinned proper quantum metric space framework, the 4-point relaxed forward metametric $\mathfrak{D}^K$, and the hyper-topology of admissible truncated covers. From Connes–Rovelli [10] we inherit the thermal-time hypothesis identifying modular flow with a physical time parameter on a KMS state. *What this paper adds* is (i) the categorical bridge $\mathbf{W} : \mathbf{KreinMetaMet}_{\text{pp}} \rightarrow \mathbf{LorPLG}_{\text{cov}}$ between those constructions; (ii) the F2 resolution via the thermal-vs-geometric-time distinction (the forward/reverse mismatch is a structural signature of two projection mechanisms applied to one KMS state, not a defect of either framework); (iii) the Bridge Theorem’s *structural* legs (B1, B3) at theorem-grade rigor on the K^+ -weak-form substrate, with the metric-level legs (B2 structurally vacuous off-orbit, B4 descoped) riding Paper 45’s degenerate Krein seminorm; and (iv) the seven newly accessible theorems on the GeoVac wedge image (T1–T6a), headlined by T6’s wedge-specific metric-level *route toward* the published-open G2 problem of Paper 45 [49] § 1.4 (descoped — conditional on the descoped Krein seminorm; the synthetic-side transport is a proposed route, not a closure). The strong-form bridge on the Paper 46 [50] Appendix B enlarged substrate (Q1’) is explicitly deferred (Section 8.1); we do *not* claim that or the general non-wedge G2 metric-level closure.

We construct (Definition 5.3, Theorem 5.4) the Wick-rotation functor

$$\mathbf{W} : \mathbf{KreinMetaMet}_{\text{pp}} \longrightarrow \mathbf{LorPLG}_{\text{cov}},$$

contravariant by Gelfand duality, sending a Krein-pointed proper QMS $(\mathcal{A}^K, L^K, \mathcal{M}^L, \omega_W^L)$ to the covered LPLS

$$\mathbf{W}(\mathcal{A}^K, L^K, \mathcal{M}^L, \omega_W^L) = (\hat{\mathcal{M}}^L, \ell^L, \hat{\omega}_W^L, \hat{\mathcal{U}}^L),$$

where $\hat{\mathcal{M}}^L$ is the Gelfand spectrum of the M-diagonal Abelian topography, $\ell^L = \kappa_g \cdot \tau_{\text{mod}}^{\omega_W^L}$ is κ_g times the modular-flow time on the wedge boost orbits ($-\infty$ off-orbit), $\hat{\omega}_W^L$ is the Bisognano–Wichmann vacuum as character of the topography, and $\hat{\mathcal{U}}^L$ is the truncation cover.

The Krein–Mondino–Sämman Bridge Theorem (Theorem 5.4) states that the *structural* bridge properties (B1, B3) hold at theorem-grade rigor, while the *metric-level convergence* leg (B4) is **descoped** — it rests on the degenerate Krein metric of Paper 45 (the Status note), so it is a conditional transport pending substrate repair, not an established metric convergence:

- (B1) *Structural correspondence* — 15 rows of structural alignment (set, topology, time separation, basepoint, cover, scales, slabs, ε -nets, convergence, pre-compactness, etc.).
- (B2) *Reverse triangle inequality* on the bridge image — on-orbit via additivity of Tomita–Takesaki modular flow at integer K_α^W spectrum [46]; off-orbit *structurally vacuous* at the K^+ -weak-form bridge because the M-diagonal topography forces orbit-pair contradiction (Decomposition O Case (iii) is empty).
- (B3) *Pre-compactness inheritance* — Mondino–Sämman Thm 6.2 applies on the bridge image via the Paper 44 propagation-number = 2 cardinality bound.

- (B4) *Convergence transport (descoped)* — Krein-side convergence under the Latrémolière hypertopology *would* induce Mondino–Sämman pLGH convergence, with Plancherel-nested Berezin maps matching the MS Def 3.6 axioms one-to-one; but the premise $\mathfrak{D}^K \rightarrow 0$ is built from the degenerate Krein metric of Paper 45, so this transport is conditional pending substrate repair, not an established metric convergence.

The F2 mismatch is resolved via the Connes–Rovelli thermal-time hypothesis [10]: the Krein metametric $\mathfrak{D}^K$ measures *thermal time* of the wedge KMS state, the synthetic time separation ℓ^L measures *geometric proper time*, and the Wick rotation between them is the analytic continuation mapping additive thermal-time structure to super-additive geometric-time structure. Concretely, on the Bisognano–Wichmann wedge, $t_{\text{geometric}} = \kappa_g \cdot t_{\text{thermal}}$ by the four-witness Wick-rotation theorem of Paper 42 [46].

1.3 GeoVac wedge application: seven newly accessible theorems

The bridge is illustrated on the canonical Paper 43 [47] hemispheric wedge $W_L = P_W^{\text{spat}} \otimes P_{t \geq 0}$ on the compact-temporal Lorentzian Krein space $\mathcal{K}_{n_{\text{max}}, N_t, T}$, with Bisognano–Wichmann vacuum $\omega_W^L = e^{-K_\alpha^W} / Z$ and modular Hamiltonian $K_\alpha^W = J_{\text{polar}}^W$ of integer spectrum two $_m_j$. All five Krein PPQMS axioms verify by direct construction. Seven newly accessible theorems follow (Section 6):

- **T1** (wedge as MS-covered LPLS) — every finite cutoff produces a well-defined Mondino–Sämman covered Lorentzian pre-length space with uniform timelike diameter 2π across all cover levels;
- **T2** (synthetic compactness of wedge family) — pLGH pre-compactness for sequences of GeoVac wedge cutoffs;
- **T3** (synthetic bit-exact panel) — the synthetic pLGH-rate panel at $(n_{\text{max}}, N_t) \in \{(2, 3), (3, 5), (4, 7)\}$ inherits bit-identically the Paper 45 Krein-side Λ^L values $\{2.0746, 1.6101, 1.3223\}$ (conditional on the bridge; these are the descoped Paper 45 *rate-formula* evaluations, not metric-convergence measurements — the Lorentzian quantum-metric convergence is descoped);
- **T4** (synthetic four-witness readings) — the Paper 42 [46] four-witness Wick-rotation theorem’s six physical witnesses reduce to two synthetic-side equivalence classes distinguished by the rapidity-rate normalization κ_g ;
- **T5** (synthetic Pythagorean orthogonality with M1 signature) — the operator-algebraic Pythagorean orthogonality $\langle H_{\text{local}}, D_W^L \rangle_{HS} = 0$ of Paper 43 [47] § 10.2 transports to a synthetic-side foliation orthogonality, with the closed-form $1/\pi^2$ residual prefactor identified as the master Mellin engine M1 Hopf-base-measure signature (Paper 32 [42] § VIII);
- **T6** (proposed **G2 metric-equivalent route for the GeoVac wedge**, conditional/descoped) — the deepest design content: the de-compactification limit $T \rightarrow \infty$ *would* close at the pLGH-equivalent (metric-level-equivalent) level on the synthetic side for the GeoVac wedge, *conditional on the bridge* — which inherits the descoped Paper 45 degeneracy. It is therefore a *proposed* workaround for the published-open non-compact propinquity problem (Paper 45 [49] § 1.4 G2) at the GeoVac-wedge-specific level, not a closure;
- **T6a** (synthetic three-carrier identification) — Paper 47 [51] § 6 three-carrier identification (periodic / Dirichlet / non-compact) extends from operator-algebraic to synthetic-Lorentzian level.

1.4 Honest scope

The bridge **W** is constructed at the **K⁺-weak-form** level only (the substrate of Paper 45 [49]). The strong-form extension of **W** to the Paper 46 [50] Appendix B enlarged substrate (chirality-flipping generators M^{flip} with $\{J, M^{\text{flip}}\} = 0$) is **not** constructed here; it is the named open question **Q1'** (Section 8.1). On the enlarged substrate the off-orbit super-additivity ceases to be vacuous and becomes a load-bearing operator-algebraic statement, turning Q1' into a substantive NCG-mathematics target, beyond sprint scale.

The functor **W** is **functorial, not isometric**: $\mathbb{D}^K$ on the Krein-algebraic side and ℓ^L on the synthetic side measure different physical quantities of the same wedge KMS state. The forward-vs-reverse triangle direction is not a "mismatch to be reconciled" but a structural signature of two distinct projection mechanisms applied to the same physical object. This is the F2 resolution.

For the GeoVac wedge, the substantive **T6 route toward G2** is wedge-specific: it proposes (conditionally; descoped, as it inherits the degenerate Krein seminorm) a de-compactification limit at the propinquity-equivalent (pLGH) level on the synthetic side for the wedge specifically. The general non-compact Latrémolière propinquity problem (Paper 45 [49] § 1.4 G2 on arbitrary non-wedge carriers) remains the named NCG-mathematics frontier and is **not** closed by this paper. The Paper 47 [51] norm-resolvent closure at the spectral level remains the operator-algebraic gold standard for general carriers; T6 specializes the metric-level question to the wedge via the bridge.

1.5 Roadmap of the paper

Section 2 recalls the necessary background: Latrémolière's pinned proper QMS framework (§ 2.1), the Krein operator-system substrate of Papers 44/45/46 (§ 2.2), the Mondino–Sämman pre-length space framework (§ 2.3), the Bisognano–Wichmann vacuum and the four-witness Wick-rotation theorem of Paper 42 (§ 2.4), and the Connes–Rovelli thermal-time hypothesis (§ 2.5).

Section 3 establishes the Krein-pointed proper quantum metric space: a 4-tuple $\mathbb{X}^K = (\mathcal{A}^K, L^K, \mathcal{M}^L, \omega_W^L)$ in which the nine Latrémolière axioms transport at the *categorical/structural* level. (**Descoped.**) On the Paper 45 substrate the Krein Lipschitz seminorm L^K is degenerate (its kernel exceeds the scalars; Status note (i)), so $\mathbb{X}^K$ is a categorical-design construction, *not* an established proper quantum metric space in the cited sense; the metric-level content of § 3 is descoped accordingly.

Section 4 constructs the Krein hypertopology on the class of Krein-pointed proper QMS, with explicit M-tunnel extent, M-local metametric, 4-point relaxed triangle inequality, coincidence theorem, and inframetric structure. The compact-case agreement at $N_t = 1$ reduces the operator-system substrate and $\text{SU}(2)$ rate factor bit-exactly to the Paper 38 [43] $\text{SU}(2)$ state-space Gromov–Hausdorff construction (the joint rate carries a $\text{U}(1)$ $T/4$ offset; see Theorem 4.9).

Section 5 constructs the Wick-rotation functor and states the Bridge Theorem. The F2 mismatch is resolved via R3 (Connes–Rovelli thermal time) as the unifying framework. The candidate resolutions R1 (negate time) and R2 (sub-functors) are tested and found insufficient.

Section 6 applies the bridge to the GeoVac hemispheric wedge, with the seven newly accessible theorems.

Section 7 states the synthetic-Lorentzian compactness theorems, with T6 (G2 metric-level closure for the GeoVac wedge) as the headline.

Section 8 catalogues the open questions: Q1' strong-form bridge with enlarged substrate, G2 on the general non-wedge carrier, G3 cross-manifold, G4 inner-factor calibration.

1.6 Related work

The Latrémolière lineage [12, 13, 21–25] provides the operator-algebraic side of the bridge. The pre-2025 papers handle the compact case; the 2512.03573 hypertopology paper [25] is the first to extend to proper (locally compact, non-unital) quantum metric spaces. Our Krein lift inherits and extends this framework.

The Mondino–Sämman lineage [19, 29, 30] provides the synthetic-Lorentzian side. The 2504.10380 v4 paper [30] introduces the pointed extension with covered LPLS and pLGH convergence; earlier papers by Kunzinger–Sämman established the underlying Lorentzian pre-length-space framework. The Mondino–Ryborz–Sämman stability paper [29] (May 2026) is the most recent flagship in this lineage. A concurrent extension by Braun–Sämman [5] develops a Gromov reconstruction theorem and measured Gromov–Hausdorff convergence in Lorentzian geometry; the bridge constructed here targets the unmeasured pointed framework of [30], but the measured extension is a natural future bridge target.

Adjacent synthetic-Lorentzian convergence frameworks include: Sormani–Vega null distance [36], Sakovich–Sormani timed Gromov–Hausdorff [34], Minguzzi–Suhr bounded Lorentzian metric spaces [28], Müller’s Cauchy slabs [31], Allen–Burtscher metric GH [1], Che–Perales–Sormani timed Hausdorff [8]. These frameworks operate in distinct categories from the present bridge (Mondino–Sämman’s ε -net-of-causal-diamonds approach versus null-distance metric-GH or timed Hausdorff variants); they are cited for completeness but are not in direct competition with the bridge construction.

The Connes–Rovelli thermal-time hypothesis [10] supplies the unifying interpretation. Active research on Connes-spectral-triples plus Tomita–Takesaki modular structure as a NCG framework for general relativity (e.g., Bertozzini–Conti–Lewkeeratiyutkul [2]) is adjacent to R3 but does not target the specific Krein-pointed PPQMS $\rightarrow$ Mondino–Sämman pLGH bridge constructed here.

Recent NCG-side work on twisted spectral triples and pseudo-Riemannian spectral geometry includes Bizi–Brouder–Besnard [4], van den Dungen [39], Franco–Eckstein [14], Strohmaier [37], and the recent Nieuviarts twisted-spectral-triple paper [32] (December 2025 v2). The Nieuviarts framework is restricted to even KO-dimension and does not apply directly to our KO-dim 3 substrate; we cite it as an alternative-to-Wick-rotation route in the Lorentzian-extension program. Ketterer [17] (May 2026) and Kubota [18] (May 2026) are recent synthetic-side internal refinements; the former treats convergence of Lorentzian spaces and curvature bounds for generalized cones, the latter establishes a Lorentzian coarea inequality. We cite both as adjacent works in the synthetic-Lorentzian convergence literature.

The Hekkelman–McDonald papers [16] on Connes spectral triples and noncommutative integration provide non-compact propinquity machinery that does not extend to a metric-level closure on the $S^3 \times \mathbb{R}_t$ carrier; our T6 closure for the wedge sits on the synthetic side via the bridge rather than via direct non-compact propinquity.

To the author’s knowledge, no published paper constructs a categorical bridge between operator-algebraic Lorentzian propinquity and synthetic Lorentzian Gromov–Hausdorff frameworks. The bridge constructed here is novel.

2 Setup

2.1 Latrémolière’s pinned proper QMS framework

We recall the central definitions from Latrémolière [25].

Definition 2.1 (Latrémolière 2512.03573 Def 1.18: Leibniz seminorm). Let $\mathcal{A}$ be a C^* -algebra. A *Leibniz Lipschitz seminorm* L on $\mathcal{A}$ is a (typically unbounded) seminorm on a dense $*$ -subalgebra $\text{dom}(L) \subseteq \mathcal{A}$ satisfying $L(a^*) = L(a)$ and the Leibniz inequality $L(ab) \leq \|a\|L(b) + L(a)\|b\|$.

Definition 2.2 (Latrémolière 2512.03573 Def 1.22: separable QLCMS). A *separable quantum locally compact metric space* (QLCMS) is a pair $(\mathcal{A}, L)$ of a separable C*-algebra $\mathcal{A}$ and a Leibniz seminorm L satisfying:

(FM) Fortet–Mourier distance metrizes the weak-* topology on $\mathcal{S}(\mathcal{A})$;

(CB) the unit ball $\{a \in \text{dom}(L) : L(a) \leq 1\}$ is closed;

(B) there exists a strictly positive $h \in \mathcal{A}_{\text{sa}}$ and a state $\mu \in \mathcal{S}(\mathcal{A})$ with $\mu(h) = 1$ such that

$$\{hah : a = b + t\mathbf{1}, L(b) \leq 1, \mu(b) = -t\}$$

is bounded.

Definition 2.3 (Latrémolière 2512.03573 Def 1.26: pinned QLCMS). A *pinned QLCMS* is a triple $(\mathcal{A}, L, \mu)$ extending Definition 2.2 with $\mu \in \mathcal{S}(\mathcal{A})$ a pin state, such that $\{\phi \in \mathcal{S}(\mathcal{A}) : \text{mk}_L(\phi, \mu) < \infty\}$ is weak-* dense in $\mathcal{S}(\mathcal{A})$, where mk_L is the Monge–Kantorovich distance.

Lemma 2.4 (Latrémolière 2512.03573 Lemma 1.25). *If $(\mathcal{A}, L)$ is a separable QLCMS with the (B)-axiom witnessed by (h, μ) , then $\{\phi : \text{mk}_L(\phi, \mu) < \infty\}$ is weak-* dense in $\mathcal{S}(\mathcal{A})$. In particular, the (B)-axiom of Definition 2.2 implies the density condition of Definition 2.3.*

Definition 2.5 (Latrémolière 2512.03573 Def 1.37 + 1.42: pointed proper QMS). A *pinned proper QMS* is a pinned QLCMS $(\mathcal{A}, L, \mu)$ containing an L -Lipschitz μ -pinned exhaustive sequence $(h_n) \subseteq \text{dom}(L)$ with $L(h_n) \rightarrow 0$, $\mu(h_n) \rightarrow 1$, $\|h_n\| \rightarrow 1$. A *topography* $\mathfrak{M} \subseteq \mathcal{A}$ is an Abelian C*-subalgebra containing such an exhaustive sequence. A *pointed proper QMS* is a 4-tuple $(\mathcal{A}, L, \mathfrak{M}, \mu)$ with $\mathfrak{M}$ a topography and μ restricting to a character of $\mathfrak{M}$.

The associated structures are M-isometries, M-tunnels, M-local metametric $\mathfrak{D}_M$, parameter-family metametric $\mathfrak{D}^K$, and a 4-point relaxed forward triangle:

$$\mathfrak{D}^K(\mathbb{A}, \mathbb{B}) \leq (1 + \mathfrak{D}^K(\mathbb{A}, \mathbb{D}))^2 \mathfrak{D}^K(\mathbb{D}, \mathbb{B}) + (1 + \mathfrak{D}^K(\mathbb{D}, \mathbb{B}))^2 \mathfrak{D}^K(\mathbb{A}, \mathbb{D}).$$

The Latrémolière coincidence theorem (Thm 3.6) characterizes $\mathfrak{D}^K = 0$ via the existence of a full topographic M-isometry.

2.2 Krein operator-system substrate from Papers 44/45/46

The Krein-side substrate is the compact-temporal Lorentzian Krein space

$$\mathcal{K}_{n_{\max}, N_t, T} = \mathcal{H}_{\text{GV}}^{n_{\max}} \otimes \mathbb{C}^{N_t}$$

of Paper 44 [48], with fundamental symmetry $J_L = J_{\text{spat}} \otimes I_{N_t}$ where $J_{\text{spat}}^2 = +I$ is the spatial chirality swap; the K^+ -restricted Hilbert space is $\mathcal{K}^+ = \{|\psi\rangle : J_L|\psi\rangle = +|\psi\rangle\}$. The Lorentzian Dirac is (van den Dungen 2016 Prop. 4.1 [39])

$$D_L = i(\gamma^0 \otimes \partial_t + D_{\text{GV}} \otimes I_{N_t}),$$

Krein-self-adjoint: $D_L^\times = D_L$ where $\dagger$ is the Krein adjoint $\gamma^0 \cdot * \cdot \gamma^0$. The natural Lorentzian operator system is

$$\mathcal{O}^L = \text{span}_{\mathbb{C}}\{M_{N, L, M}^{\text{spat}} \otimes M_p^{\text{temp}} : N \leq n_{\max}, L < N, |M| \leq L, 0 \leq p \leq N_t - 1\},$$

of finite dimension at finite cutoff, with propagation number $\text{prop}(\mathcal{O}^L) = 2$ under the Weyl-doubled achievable envelope (Paper 44 [48] Prop. prop=2).

Paper 45 [49] is the K^+ -restricted weak-form *degeneracy* theorem: the natural Lorentzian-propinquity seminorm Λ^L vanishes identically, so the would-be convergence

$$\Lambda^L(\mathcal{T}_{n_{\max}, N_t, T}^L, \mathcal{T}_{S^3 \times S_T^1}^L) \leq C_3^{\text{joint}} \cdot \gamma_{n_{\max}, N_t, T}^{\text{joint}} \rightarrow 0$$

is a closed-form *rate-formula* evaluation (panel values $\Lambda^L(2, 3) = 2.0746$, $\Lambda^L(3, 5) = 1.6101$, $\Lambda^L(4, 7) = 1.3223$), *not* a quantum-metric distance — the Lorentzian quantum-metric convergence is descoped. Paper 46 [50] carries the same rate formula to a strong-form (no K^+ restriction) on the natural chirality-doubled scalar-multiplier substrate ($\Lambda^{\text{strong}} = \Lambda^L$, also degenerate), and proposes a route on a further enlarged substrate (Appendix B) with chirality-flipping generators M^{flip} satisfying $\{J_L, M^{\text{flip}}\} = 0$ — also descoped.

2.3 Mondino–Sämman pre-length space framework

We recall the central definitions from Mondino–Sämman [30].

Definition 2.6 (Mondino–Sämman Def 2.3: Lorentzian pre-length space). A *Lorentzian pre-length space* (LPLS) is a pair (X, ℓ) with X a set carrying a topology finer than the chronological topology, and $\ell : X \times X \rightarrow \{-\infty\} \cup [0, \infty]$ satisfying $\ell(x, x) \geq 0$ and the *reverse triangle inequality*

$$\ell(x, y) + \ell(y, z) \leq \ell(x, z) \quad \forall x, y, z \in X \quad (1)$$

(with convention $\pm\infty + \mp\infty = 0$ on the LHS yielding $-\infty$ when undefined). Timelike, causal, and causal-diamond relations are derived in the standard way: $x \ll y \iff \ell(x, y) > 0$, $x \leq y \iff \ell(x, y) \geq 0$, $J(x, y) = \{z : x \leq z \leq y\}$, and the time separation $\tau := \max(0, \ell)$.

Definition 2.7 (Mondino–Sämman Def 3.2: ε -net). An ε -net for $A \subseteq X$ is a collection $\mathcal{S} = (J_i)_{i \in \Omega}$ of causal diamonds with $\tau(J_i) \leq \varepsilon$ and $A \subseteq \bigcup_i J_i$.

Definition 2.8 (Mondino–Sämman Def 3.6: LGH convergence of subsets). A sequence $A_n \rightarrow A$ converges in the Lorentzian Gromov–Hausdorff (LGH) sense if there exist ε -nets S_n for A_n and S for A with: (i) matching cardinality; (ii) correspondences R_n between vertices with $\text{dis}(R_n) \rightarrow 0$; (iii) extension property of correspondences; (iv) forward density.

Definition 2.9 (Mondino–Sämman Def 3.8: covered LPLS). A *covered LPLS* is a 4-tuple $(X, \ell, o, \mathcal{U})$ where (X, ℓ) is an LPLS, $o \in X$ is a basepoint event, and $\mathcal{U} = (U_k)_{k \in \mathbb{N}}$ is a countable cover satisfying: (i) $\bigcup_k U_k = X$; (ii) $U_k \subseteq U_{k+1}$ (nested); (iii) $o \in U_k$ for all k ; (iv) $\sup_{x, y \in U_k} \tau(x, y) < \infty$ (slab bounds).

Definition 2.10 (Mondino–Sämman Def 3.12: pLGH convergence). $(X_n, \ell_n, o_n, \mathcal{U}_n) \rightarrow (X, \ell, o, \mathcal{U})$ in the *pointed LGH* sense iff for each k , $U_{k,n} \rightarrow U_{k,\infty}$ in LGH sense.

The Mondino–Sämman pre-compactness theorem (Thm 6.2) gives that any sequence of covered LPLS with (i) uniform slab bounds, (ii) uniform ε -net cardinality bounds, (iii) cumulative ε -net nesting has a pLGH-convergent subsequence.

2.4 Bisognano–Wichmann vacuum and the four-witness Wick-rotation theorem

The Paper 43 [47] hemispheric wedge is

$$W_L := P_W^{\text{spat}} \otimes P_{t \geq 0}, \quad P_W^{\text{spat}} = \frac{1}{2}(I + R_{\text{polar}}),$$

where R_{polar} is the spatial m_j -reflection involution (Paper 42 [46] § 4) and $P_{t \geq 0}$ is the diagonal projection on $\mathbb{C}^{N_t}$ to positive temporal indices. The Bisognano–Wichmann vacuum state under the BW- α geometric Hamiltonian (with $\beta = 2\pi/\kappa_g$) is

$$\rho_W^L := e^{-K_\alpha^W} / Z, \quad K_\alpha^W = J_{\text{polar}}^W \text{ on the wedge,}$$

with integer spectrum $\text{two_m}_j \in \mathbb{Z}_{>0}$. The Tomita–Takesaki modular Hamiltonian $K_{\text{TT}} = -\log \Delta$ is constructed from the wedge KMS state via polar decomposition (Paper 42 [46] § 6).

The Paper 42 [46] four-witness Wick-rotation theorem states that six physical witnesses (Bisognano–Wichmann canonical with $\kappa_g = 1$, Hartle–Hawking at $M = 1, 2$, Sewell at $M = 1$, Unruh at $a = 1, 2$) all yield bit-identical modular operators Δ and Hamiltonians K_{TT} under the BW unit normalization $H_{\text{local}} := K_\alpha^W / \beta$. The Riemannian-side bit-exact identity $\sigma_{2\pi}^\alpha(O) = O$ holds at every finite cutoff $n_{\text{max}} \in \{1, 2, 3, 4, 5\}$, and the Tomita BW- γ flow σ_t^{TT} satisfies the conjugacy $\sigma_t^{TT}(a) = \sigma_{-t}^\alpha(a)$ bit-exact. Paper 43 [47] extends to Lorentzian signature via BBB Krein construction [4] and verifies the same six-witness collapse at every (n_{max}, N_t) panel cell.

2.5 Connes–Rovelli thermal-time hypothesis

Connes and Rovelli [10] (1994) proposed:

Thermal-time hypothesis. *In quantum statistical mechanics on a generally covariant system, the natural physical time evolution at thermal equilibrium with state ω is given by the modular automorphism group σ_t^ω of ω (a state on the observable algebra $\mathcal{A}$).*

Applied to a KMS state at inverse temperature β on a Type III von Neumann algebra arising as the local algebra of a region of spacetime (Bisognano–Wichmann setting), the thermal time of ω relates to the geometric proper time of the underlying spacetime by

$$t_{\text{thermal}} = \frac{\beta}{2\pi} \cdot t_{\text{geometric}} = \frac{1}{\kappa_g} \cdot t_{\text{geometric}}, \quad (2)$$

where κ_g is the surface gravity (equivalently the BW Hawking-temperature parameter). This is the four-witness Wick-rotation theorem of Paper 42 [46] at the kinematic level: modular flow IS Lorentz-boost orbit on the BW wedge, with proportionality factor $1/\kappa_g = \beta/(2\pi)$.

2.6 Notation and conventions

Throughout the paper: $\mathcal{K} = \mathcal{K}_{n_{\text{max}}, N_t, T} = \mathcal{H}_{\text{GV}}^{n_{\text{max}}} \otimes \mathbb{C}^{N_t}$ denotes the compact-temporal Krein space; $J_L = J_{\text{spat}} \otimes I_{N_t}$ the fundamental symmetry; $\mathcal{K}^+, \mathcal{K}^-$ the $K^\pm$ -eigenspaces of J_L ; $\mathcal{O}^L$ the natural Lorentzian operator system; D_L the Krein-self-adjoint Lorentzian Dirac; $\omega_W^L = e^{-K_\alpha^W} / Z$ the BW vacuum; $K_\alpha^W = J_{\text{polar}}^W$ the BW geometric Hamiltonian (with integer spectrum two_m_j); $\sigma_t^{\omega_W^L}$ the Tomita–Takesaki modular flow of ω_W^L ; $\tau_{\text{mod}}^{\omega_W^L}(\hat{\omega}, \hat{\omega}') := \inf\{t \geq 0 : \hat{\omega}' = \hat{\omega} \circ \sigma_t^{\omega_W^L}\}$ the modular-flow time.

We write $\text{Spec}(\mathfrak{B})$ for the Gelfand spectrum of an Abelian C*-algebra $\mathfrak{B}$ and $\hat{\omega}$ for the character induced by a state ω restricting to $\mathfrak{B}$.

3 Krein-pointed proper quantum metric spaces

We lift Latrémolière’s pinned proper QMS framework to the Krein-algebraic setting. Throughout this section, the substrate is the K^+ -restricted Hilbert space; the natural-substrate generators all commute with J_L , so the Krein operator norm coincides with the Hilbert operator norm on the K^+ restriction. *Descope note (read all of this section under it):* on the Paper 45 substrate the Krein Lipschitz seminorm L^K is degenerate (its kernel exceeds the scalars; Status note (i)), so the axiom verifications below establish a *categorical/structural* lift, not a metric (proper quantum metric space) in the cited sense; the metric-level reading is descoped pending repair.

3.1 The Krein-Leibniz Lipschitz seminorm

Definition 3.1 (Krein-Leibniz Lipschitz seminorm). On the natural Lorentzian operator system $\mathcal{O}^L$ (or its continuum limit on $\mathcal{T}_{S^3 \times S_T^1}^L$), define

$$L^K(a) := \|[D_L, a]\|_{\text{op}}, \quad a \in \text{dom}(L^K) \subseteq \mathcal{O}^L,$$

where $\|\cdot\|_{\text{op}}$ is the operator norm on $\mathcal{B}(\mathcal{K})$ restricted to $\mathcal{K}^+$. We call L^K the *Krein-Leibniz Lipschitz seminorm*.

Lemma 3.2 (Krein-Leibniz inequality). *For every a, b in the natural chirality-doubled scalar-multiplier substrate,*

$$L^K(ab) \leq \|a\|_{\mathcal{O}^L} L^K(b) + L^K(a) \|b\|_{\mathcal{O}^L}.$$

Proof. On the natural substrate, all elements of $\mathcal{O}^L$ commute with J_L [48], so both a, b restrict cleanly to $\mathcal{K}^+$. On $\mathcal{K}^+$, the standard Hilbert operator norm $\|\cdot\|_{\text{op}}|_{\mathcal{K}^+}$ agrees with the algebra norm $\|\cdot\|_{\mathcal{O}^L}$. The standard operator-algebra Leibniz identity $[D_L, ab] = [D_L, a]b + a[D_L, b]$ holds in $\mathcal{B}(\mathcal{K})$; restricting to $\mathcal{K}^+$ and applying submultiplicativity yields the Krein-Leibniz inequality. $\square$

Remark 3.3 (Krein vs Hilbert operator norm on the enlarged substrate). The natural-substrate version of Lemma 3.2 uses the Hilbert operator norm on $\mathcal{K}^+$. On the Paper 46 [50] Appendix B *enlarged* substrate with chirality-flipping generators M^{flip} satisfying $\{J_L, M^{\text{flip}}\} = 0$, the $\mathcal{K}^+$ restriction is no longer well-defined as a multiplier (such generators map $\mathcal{K}^+$ to $\mathcal{K}^-$), and the two natural operator norms (Krein operator norm with respect to the indefinite Krein product vs. Hilbert operator norm on the full Krein space) differ. The $\mathcal{K}^+$ -weak-form route of this paper makes the distinction invisible; the strong-form route makes it visible at the level of the gradient-norm absorption mechanism of Paper 46 [50] Appendix B. For Q1' (Section 8.1) this distinction becomes load-bearing.

3.2 The Krein-separable QLCMS and Krein-pinned QLCMS

Definition 3.4 (Krein-separable QLCMS). A *Krein-separable QLCMS* is a pair $(\mathcal{A}^K, L^K)$ where $\mathcal{A}^K$ is the natural Lorentzian operator system on $\mathcal{K}$ (at finite cutoff: the C^* -algebra generated by $\mathcal{O}^L$ in $\mathcal{B}(\mathcal{K}^+)$, a finite-dimensional matrix algebra; in the continuum: $\mathcal{A}^K = C(\mathcal{T}_{S^3 \times S_T^1}^L)$ acting on $\mathcal{K}_{\text{continuum}}^+$), and L^K is the Krein-Leibniz Lipschitz seminorm on a dense $*$ -subalgebra $\text{dom}(L^K) \subseteq \mathcal{A}^K$, satisfying:

- (FM-K) The Krein-positive Fortet–Mourier distance metrizes the weak- $*$ topology on $\mathcal{S}(\mathcal{A}^K)$;
- (CB-K) The unit ball $\{a \in \text{dom}(L^K) : L^K(a) \leq 1\}$ is closed in the operator-norm topology;
- (B-K) There exists strictly positive $h^L \in \mathcal{A}_{\text{sa}}^K$ and a state $\mu^L \in \mathcal{S}(\mathcal{A}^K)$ with $\mu^L(h^L) = 1$ such that

$$\{h^L a h^L : a = b + t\mathbf{1}, L^K(b) \leq 1, \mu^L(b) = -t\}$$

is bounded.

Lemma 3.5. *The pair $(\mathcal{A}^K, L^K)$ with the natural Lorentzian operator system substrate is a Krein-separable QLCMS, with witnessing $(h^L, \mu^L) = (K_\alpha^W/Z, \omega_W^L)$.*

Proof sketch. The (FM-K), (CB-K), (B-K) axioms transport from Papers 42–45 substrate: (FM-K) via Paper 44 [48] § 6 Krein-positive Wasserstein–Kantorovich SDP framework, with weak- $*$ metrization following Latrémolière [25] Thm 1.9; (CB-K) via Banach–Alaoglu plus lower-semicontinuity of L^K ; (B-K) via Paper 42 [46] § 5 integer spectrum of K_α^W (h^L positive definite, $\|h^L\| = 1$, $\omega_W^L(h^L) = 1$ by BW pinning) combined with Paper 44 [48] propagation number = 2 (bounded sets in finite-dimensional subspaces are bounded). See A.2' § 2.3 of the synthesis memo for full detail. $\square$

Definition 3.6 (Krein-pinned QLCMS). A *Krein-pinned QLCMS* is a triple $(\mathcal{A}^K, L^K, \omega_W^L)$ with $\omega_W^L \in \mathcal{S}(\mathcal{A}^K)$ the BW vacuum, such that $\{\phi \in \mathcal{S}(\mathcal{A}^K) : \text{mk}_{L^K}(\phi, \omega_W^L) < \infty\}$ is weak-* dense in $\mathcal{S}(\mathcal{A}^K)$.

Lemma 3.7 (Continuum tightness via Lemma 2.4). $(\mathcal{A}^K, L^K, \omega_W^L)$ is a *Krein-pinned QLCMS*. Equivalently: the density condition of Definition 3.6 holds.

Proof. By Lemma 3.5, the (B-K) axiom holds with $(h^L, \mu^L) = (K_\alpha^W/Z, \omega_W^L)$. By Lemma 2.4 (Latrémolière 2512.03573 Lemma 1.25), the density of finite-Monge–Kantorovich states follows mechanically. $\square$

Remark 3.8. Lemma 3.7 dissolves a caveat that the diagnostic phase of the program had flagged as a bounded sub-sprint: the continuum tightness condition is a *corollary* of the (B-K) boundedness axiom via Latrémolière’s own Lemma 1.25. The Krein lift requires no separate verification beyond the (B-K) ingredients.

3.3 The Krein L-Lipschitz ω_W^L -pinned exhaustive sequence and the approximate-unit theorem

Definition 3.9 (Krein L-Lipschitz ω_W^L -pinned exhaustive sequence). A *Krein L-Lipschitz ω_W^L -pinned exhaustive sequence* is a sequence $(h_n)_{n \in \mathbb{N}} \subseteq \text{dom}_{\text{sa}}(L^K)$ with:

1. $\lim_{n \rightarrow \infty} L^K(h_n) = 0$,
2. $\lim_{n \rightarrow \infty} \omega_W^L(h_n) = 1$,
3. $\lim_{n \rightarrow \infty} \|h_n\|_{\mathcal{A}^K} = 1$.

Theorem 3.10 (Krein approximate-unit, transport of Latrémolière 2512.03573 Thm 1.30). *If $(\mathcal{A}^K, L^K, \omega_W^L)$ is a Krein-pinned QLCMS and (h_n) is a Krein L-Lipschitz ω_W^L -pinned exhaustive sequence, then (h_n) is an approximate unit for $\mathcal{A}^K$.*

Proof sketch. Direct transport of Latrémolière 2512.03573 Thm 1.30 via Claims 1.31–1.34: weak-* convergence to 1 (using Lemma 3.7); h_n^2 pinning (via Lemma 3.2 Leibniz applied to $h_n \cdot h_n$); $(1 - h_n^2)$ vanishing on compact sets of states (via Arzelà–Ascoli on the Fortet–Mourier metric ball); convergence to 1 in strict topology (via Kadison functional calculus on the K^+ -restricted commutative envelope of h_n). $\square$

The explicit Krein-side exhaustive sequence. The natural Krein-side exhaustive sequence is the truncated Krein triple projector sequence:

$$h_n := P_{n_{\max}(n), N_t(n), T(n)},$$

the orthogonal projection onto the $\mathcal{H}_{\text{GV}}^{n_{\max}(n)} \otimes \mathbb{C}^{N_t(n)}$ truncation, with $(n_{\max}(n), N_t(n), T(n)) \rightarrow (\infty, \infty, T_\infty)$ along admissible scaling per Paper 47 [51]. Verification: (i) $L^K(h_n) \leq \gamma_{n_{\max}(n), N_t(n), T(n)}^{\text{joint}} \cdot O(1) \rightarrow 0$ by Paper 45 [49] Sub-Sprint C¹ approximate-identity property; (ii) $\omega_W^L(h_n) \rightarrow 1$ by support of ω_W^L on the wedge; (iii) $\|h_n\| = 1$ since h_n is a projector.

3.4 Krein topography and Krein-pointed proper QMS

Definition 3.11 (Krein topography). A *Krein topography* $\mathcal{M}^L$ of a Krein-separable QLCMS $(\mathcal{A}^K, L^K)$ is an Abelian C*-subalgebra of $\mathcal{A}^K$ containing a strictly positive $h^L \in \text{dom}(L^K)$ and admitting a character μ^L of $\mathcal{M}^L$ witnessing the (B-K) boundedness axiom restricted to $\mathcal{M}^L$.

¹Sprint/track designations are anchors into the project’s internal chronicle (CHANGELOG.md in the repository).

Lemma 3.12 (The natural Krein topography). *The natural Krein topography on the GeoVac substrate is the sub-operator-system*

$$\mathcal{M}^L := \text{span}_{\mathbb{C}} \{ M_{N,L,0}^{\text{spat}} \otimes I_{N_t} : N \leq n_{\max}, L < N \} \subseteq \mathcal{A}^K,$$

i.e. the chirality-doubled scalar multipliers with M -quantum-number $M = 0$ tensored with the temporal identity.

Proof sketch. Abelian: Each $M_{N,L,0}^{\text{spat}}$ is a multiplication operator by a spatial 3-Y function $Y_{N,L,0}^{(3)}$, hence the algebra is Abelian; tensoring with I_{N_t} preserves the structure.

Strictly positive h^L : Take $h^L = K_{\alpha}^W/Z$; the BW geometric Hamiltonian is M-diagonal and positive on the wedge by Paper 42 [46] § 5.

(B-K) restricted to $\mathcal{M}^L$: Restrict Lemma 3.5 to $\mathcal{M}^L$; the boundedness condition is preserved.

Contains approximate unit: By Latrémolière Remark 1.41, this follows from the Leibniz property applied to the truncated projector sequence h_n , which is M-block-diagonal hence in $\mathcal{M}^L$. $\square$

Definition 3.13 (Krein-pointed proper QMS). A *Krein-pointed proper quantum metric space* (Krein PPQMS) is a 4-tuple

$$\mathbb{X}^K = (\mathcal{A}^K, L^K, \mathcal{M}^L, \omega_W^L)$$

where $(\mathcal{A}^K, L^K, \omega_W^L)$ is a Krein-pinned QLCMS, $\mathcal{M}^L$ is a Krein topography of $(\mathcal{A}^K, L^K)$, ω_W^L restricts to a character of $\mathcal{M}^L$, and there exists a Krein L-Lipschitz ω_W^L -pinned exhaustive sequence contained in $\mathcal{M}^L$.

Lemma 3.14 (The merged substrate is a Krein PPQMS). *The 4-tuple $(\mathcal{A}^K, L^K, \mathcal{M}^L, \omega_W^L)$ with $\mathcal{A}^K$ the natural Lorentzian operator system, L^K the Krein-Leibniz Lipschitz seminorm, $\mathcal{M}^L$ the natural Krein topography of Lemma 3.12, and ω_W^L the BW vacuum, satisfies all axioms of Definition 3.13.*

Proof sketch. The (FM-K), (CB-K), (B-K) axioms hold by Lemma 3.5; the continuum tightness holds by Lemma 3.7; the topography axioms hold by Lemma 3.12; $\omega_W^L|_{\mathcal{M}^L}$ is a character because $\mathcal{M}^L$ is Abelian (states on Abelian C*-algebras are characters by Gelfand); the truncated projector sequence $h_n = P_{n_{\max}(n), N_t(n), T(n)}$ lies in $\mathcal{M}^L$ (M-block-diagonal) and is a Krein L-Lipschitz ω_W^L -pinned exhaustive sequence. $\square$

3.5 Krein M-tunnels with extent element

Definition 3.15 (Krein M-tunnel). Let $\mathbb{X}_i^K = (\mathcal{A}_i^K, L_i^K, \mathcal{M}_i^L, \mu_i)$ for $i = 1, 2$ be two Krein PPQMS and $M \geq 1$. A *Krein M-tunnel* from $\mathbb{X}_1^K$ to $\mathbb{X}_2^K$ is a 6-tuple

$$\tau^L = (\mathcal{D}^L, L_{\mathcal{D}^L}^K, \mathfrak{M}_{\mathcal{D}^L}^L, \pi_1^K, \pi_2^K, e^L)$$

where $(\mathcal{D}^L, L_{\mathcal{D}^L}^K, \mathfrak{M}_{\mathcal{D}^L}^L, \mu_1 \circ \pi_1^K)$ is a pointed topographic separable Krein-side QLCMS, $e^L \in \text{dom}_{\text{sa}}(L_{\mathcal{D}^L}^K) \cap \mathfrak{M}_{\mathcal{D}^L}^L$ is the *extent element*, and $\pi_i^K : \mathcal{D}^L \rightarrow \mathcal{A}_i^K$ are topographic Krein M-isometries.

The natural Krein M-tunnel. For comparison of the truncated Krein triple $\mathcal{T}_{n_{\max}, N_t, T}^L$ with the continuum $\mathcal{T}_{S^3 \times S_T^1}^L$:

- $\mathcal{D}^L = \mathcal{O}_{n_{\max}, N_t, T}^L \oplus C(\mathcal{T}_{S^3 \times S_T^1}^L)$, direct sum of truncated operator system and continuum function algebra;
- $L_{\mathcal{D}^L}^K((a, f)) = \max\{L_1^K(a), L_2^K(f), \|a - B^{\text{joint}}(f)\|_{\text{op}}/\varepsilon_0\}$, the tunnel Lipschitz combining intrinsic seminorms with Berezin discrepancy at scale $\varepsilon_0 = \gamma^{\text{joint}}$;

- $\mathfrak{M}_{\mathcal{D}^L}^L = \mathcal{M}_1^L \oplus \mathcal{M}_2^L$;
- $\pi_1^K(a, f) = a, \pi_2^K(a, f) = f$;
- $e^L = (h_n, \mathbf{1})$, the natural extent element pairing the truncation projector with the continuum unit.

Lemma 3.16. *The natural Krein M-tunnel satisfies Definition 3.15.*

Proof sketch. The topography $\mathfrak{M}_{\mathcal{D}^L}^L$ is Abelian by direct sum of Abelian sub-algebras; e^L is in $\text{dom}_{\text{sa}}(L_{\mathcal{D}^L}^K)$ because both h_n and $\mathbf{1}$ are; $\pi_{1,2}^K$ are topographic M-isometries by Paper 45 [49] Sub-Sprint D Berezin/projection construction; e^L is K^+ -positive because both factors commute with J_L trivially. $\square$

4 Krein-pointed propinquity hypertopology

4.1 Extent of a Krein M-tunnel

Definition 4.1 (Krein extent of an M-tunnel). The *Krein extent* of an M-tunnel $\tau^L = (\mathcal{D}^L, L_{\mathcal{D}^L}^K, \mathfrak{M}_{\mathcal{D}^L}^L, \pi_1^K, \pi_2^K, e^L)$ is

$$\chi^K(\tau^L) := \max\{\chi_1^K, \chi_2^K, \chi_3^K\},$$

where χ_1^K is the reach (Hausdorff distance of pullback quasi-state spaces), χ_2^K is the height (pin-state alignment via Monge–Kantorovich on K^+ Wasserstein–Kantorovich), and χ_3^K is the extent-element Lipschitz-and-normalization:

$$\chi_3^K(\tau^L) = \max\{2ML^K(e^L), |1 - \mu_1 \circ \pi_1^K(e^L)|, |1 - \mu_2 \circ \pi_2^K(e^L)|\}.$$

Lemma 4.2. *For the natural Krein M-tunnel from $\mathcal{T}_{n_{\max}, N_t, T}^L$ to $\mathcal{T}_{S^3 \times S_T^1}^L$,*

$$\chi^K(\tau^L) \leq \gamma_{n_{\max}, N_t, T}^{\text{joint}} \rightarrow 0,$$

bit-identical to the Paper 45 [49] Sub-Sprint D propinquity bound.

Proof sketch. $\chi_1^K = \text{reach}_B^K \leq \gamma^{\text{joint}}$ by Paper 45 [49] Sub-Sprint D § 3.1; $\chi_2^K = K^+$ pin-state distance, bounded by γ^{joint} ; χ_3^K has three components, each bounded by γ^{joint} times a constant via the approximate-identity property of h_n applied to the extent element. $\square$

4.2 The Krein M-local quantum metametric

Definition 4.3 (Krein M-local quantum metametric). For $M \geq 1$ and Krein PPQMS $\mathbb{X}^K, \mathbb{Y}^K$:

$$\mathfrak{d}_M^K(\mathbb{X}^K, \mathbb{Y}^K) := \inf\{\chi^K(\tau^L) : \tau^L \text{ is a Krein M-tunnel from } \mathbb{X}^K \text{ to } \mathbb{Y}^K\}.$$

Theorem 4.4 (4-point relaxed forward triangle inequality). *For any $M \geq 1$ and Krein PPQMS $\mathbb{X}^K, \mathbb{Y}^K, \mathbb{Z}^K$:*

$$\mathfrak{d}_M^K(\mathbb{X}^K, \mathbb{Y}^K) \leq (1 + \mathfrak{d}_M^K(\mathbb{X}^K, \mathbb{Z}^K))^2 \mathfrak{d}_M^K(\mathbb{Z}^K, \mathbb{Y}^K) + (1 + \mathfrak{d}_M^K(\mathbb{Z}^K, \mathbb{Y}^K))^2 \mathfrak{d}_M^K(\mathbb{X}^K, \mathbb{Z}^K).$$

(Structural/categorical: $\mathfrak{d}_M^K$ is built from the degenerate Krein Lipschitz seminorm of Paper 45, so this is a property of the construction; its metric-level reading is descoped pending substrate repair — see the Status note.)

Proof sketch. Direct transport of Latrémolière 2512.03573 Thm 2.24 via Krein-side tunnel composition. The natural-substrate Krein structure is closed under tunnel composition (direct sums of operator systems give operator systems, Berezin composition preserves K^+ -positivity, the extent-element construction generalizes to triple composition). $\square$

4.3 Coincidence

Theorem 4.5 (Coincidence). *For Krein PPQMS $\mathbb{X}^K, \mathbb{Y}^K$, there exists a full topographic Krein M-isometry $\pi^K : \mathbb{X}^K \rightarrow \mathbb{Y}^K$ with $\mu_{\mathbb{Y}^K} \circ \pi^K = \mu_{\mathbb{X}^K}$ if and only if $\mathfrak{D}_M^K(\mathbb{X}^K, \mathbb{Y}^K) = 0$.* (Descoped at the metric level: with the degenerate Krein seminorm of Paper 45 the “only if” direction $\mathfrak{D}_M^K = 0 \Rightarrow$ isometry is conditional on substrate repair; as it stands the statement is categorical/structural — see the Status note.)

Proof sketch. The “if” direction is contained in Theorem 4.4. The “only if” direction transports Latrémolière 2512.03573 Thm 3.6 via target-set machinery: given a sequence of tunnels τ_n^K with $\chi^K(\tau_n^K) \rightarrow 0$, the target sets $\mathfrak{t}_{\tau_n^K}^K(a|l) := \{\pi_2^K(d) : \pi_1^K(d) = a, \|d\|_{L_{\mathcal{D}L}, M} \leq l\}$ are well-defined on the Krein side because of the K^+ -positivity of natural-substrate generators, and the four almost-properties (Latrémolière Lemmas 3.2–3.5) transport mechanically via Krein-Leibniz and Krein-approximate-unit. The diagonal extraction with Banach–Alaoglu on the K^+ -restricted state space produces a limit *-isomorphism preserving topography and BW pin state. $\square$

4.4 The Krein quantum metametric and hypertopology

Definition 4.6 (Krein quantum metametric). For Krein PPQMS $\mathbb{X}^K, \mathbb{Y}^K$:

$$\mathfrak{D}^K(\mathbb{X}^K, \mathbb{Y}^K) := \max\left\{\inf\{\varepsilon > 0 : \sup_{r \in [1, 1/\varepsilon]} \mathfrak{D}_r^K(\mathbb{X}^K, \mathbb{Y}^K) < \varepsilon\}, |e^{-\text{qdiam}(\mathbb{X}^K)} - e^{-\text{qdiam}(\mathbb{Y}^K)}|\right\}.$$

Theorem 4.7 (Inframetric structure). *$\mathfrak{D}^K$ is a K^+ -positive inframetric: symmetric, satisfying the 4-point relaxed triangle, and $\mathfrak{D}^K(\mathbb{X}^K, \mathbb{Y}^K) = 0$ iff a full topographic Krein M-isometry exists between $\mathbb{X}^K$ and $\mathbb{Y}^K$.* (Structural/categorical: built on the degenerate Krein seminorm of Paper 45; the non-degeneracy (metric-level) reading is descoped pending substrate repair — see the Status note.)

Proof sketch. Symmetry holds because Krein M-tunnels are symmetric in their roles; the 4-point relaxed triangle follows from Theorem 4.4; the coincidence statement is Theorem 4.5. $\square$

Definition 4.8 (Krein hypertopology). The *Krein hypertopology* on the class $p\mathcal{P}^K$ of Krein PPQMS is the topology with Kuratowski closure operator

$$\text{cl}^K(\mathcal{A}) := \{\mathbb{Y}^K \in p\mathcal{P}^K : \mathfrak{D}^K(\mathbb{Y}^K, \mathcal{A}) = 0\}$$

where $\mathfrak{D}^K(\mathbb{Y}^K, \mathcal{A}) := \inf\{\mathfrak{D}^K(\mathbb{Y}^K, \mathbb{X}^K) : \mathbb{X}^K \in \mathcal{A}\}$.

4.5 Compact-case agreement at $N_t = 1$

Theorem 4.9 (Compact-case agreement). *When restricted to $N_t = 1$, the Krein operator-system substrate and the $SU(2)$ rate factor reduce bit-exactly to the Paper 38 [43] $SU(2)$ state-space Gromov–Hausdorff construction. The joint $L1$ rate carries an additive $U(1)$ temporal offset $T/4$ (the Riemannian-limit temporal factor), so the bit-exact reduction is on the spatial substrate and its $SU(2)$ rate, not on the joint rate.*

Proof sketch. At $N_t = 1$: $\mathcal{K}_{n_{\max}, 1, T} = \mathcal{H}_{\text{GV}}^{n_{\max}}$ (spatial Krein space alone); $J_L = J_{\text{spat}}$; $\mathcal{K}^+ = \mathcal{H}_{\text{GV}}^{n_{\max}, +}$; $D_L|_{N_t=1} = iD_{\text{GV}}$ (temporal direction contributes nothing); $\mathcal{O}^L|_{N_t=1}$ is the chirality-doubled spinor lift of Paper 38 [43]’s scalar substrate. By Paper 45 [49] Sub-Sprint D § 5 Riemannian-limit recovery, the $SU(2)$ rate factor $\gamma_{SU(2)}(n_{\max}) = \Lambda_{n_{\max}}^{\text{Paper 38}}$ bit-exact; the joint rate is $\gamma_{L1}(n_{\max}, 1, T) = \gamma_{SU(2)}(n_{\max}) + T/4$, the $T/4$ being the $U(1)$ temporal-factor offset (uniform-circle expected $|\theta|$). The operator-system / spatial-substrate reduction is therefore bit-exact; the joint rate matches in its $SU(2)$ factor up to that $T/4$ offset. $\square$

Remark 4.10 (The compact-case agreement object). The compact-case object is the chirality-doubled spinor lift of Paper 38’s Riemannian $SU(2)$ Lipschitz spectral triple, pinned at the BW vacuum on the spatial wedge. This explicit identification ensures the Krein-pointed hypertopology is a strict extension of Paper 38’s state-space Gromov–Hausdorff construction, not a parallel one.

4.6 Numerical panel

The Krein-side propinquity bound inherits the bit-exact numerical panel from Paper 45 [49]:

$(n_{\max}, N_t)$	$\mathfrak{D}_M^K$ (qualitative)	Λ^L (Paper 45)
$(2, 3)$	≤ 2.0746	$= 2.0746$
$(3, 5)$	≤ 1.6101	$= 1.6101$
$(4, 7)$	≤ 1.3223	$= 1.3223$

Monotone decreasing, with ratio $\mathfrak{D}^K(4, 7)/\mathfrak{D}^K(2, 3) = 0.6374$ matching Paper 38 [43] single-factor bit-identical.

5 The Mondino–Sämman bridge: Wick-rotation functor

5.1 The F2 mismatch and three candidate resolutions

The structural mismatch between the Krein metametric $\mathfrak{D}^K$ (4-point relaxed forward, codomain $[0, \infty]$) and the Mondino–Sämman time separation ℓ (3-point reverse, codomain $\{-\infty\} \cup [0, \infty]$) is the **F2 mismatch**. We test three candidate resolutions.

5.1.1 R1: Negate time

Candidate. Define $\ell_{R1}^L(\hat{\omega}, \hat{\omega}') := -\mathfrak{D}^K(\mathbb{X}_{\hat{\omega}}^K, \mathbb{X}_{\hat{\omega}'}^K)$.

Verdict: NEGATIVE. R1 fails for two independent reasons: (i) Codomain mismatch: $\mathfrak{D}^K \in [0, \infty]$ gives $\ell_{R1}^L \in [-\infty, 0]$, while Mondino–Sämman requires $\{-\infty\} \cup [0, \infty]$. The MS reverse triangle encodes the Lorentzian convention that "longer causal curves accumulate more proper time" and the negation gives the opposite convention. (ii) Triangle structural mismatch: negating both sides of the 4-point relaxed forward inequality does NOT collapse to the 3-point reverse triangle, and the coefficient factors $(1 + \mathfrak{D}^K)^2$ have no MS-side analog.

5.1.2 R2: Functorial composite of Wick-rotation + modular-flow

Candidate. A bridge functor $F_a \circ F_c$ composed of: (a) Wick rotation F_a giving the right codomain shape (3-point reverse triangle on boost orbits via analytic continuation), and (c) modular flow F_c giving the right basepoint (BW vacuum) and the right cover scales ($\beta_k = 2\pi/\kappa_g$).

Verdict: PARTIAL. (a) gives the right structural shape but does not determine basepoint or cover scales. (c) gives basepoint and scales but inherits the F2 triangle-direction mismatch (modular flow is additive, not super-additive). The composite $F_a \circ F_c$ has all needed ingredients but the F2 mismatch is not resolved *by R2 alone*; the resolution requires R3.

5.1.3 R3: Connes–Rovelli thermal time as unifying framework

Candidate. Identify the Krein metametric $\mathfrak{D}^K$ as measuring *thermal time* of the wedge KMS state and the Mondino–Sämman ℓ as measuring *geometric proper time* of the same wedge KMS state. The Wick rotation between them, via equation (2), effects the analytic continuation

$t_{\text{thermal}} = -it_{\text{geometric}}/\kappa_g$ mapping additive thermal-time structure to super-additive geometric-time structure.

Verdict: POSITIVE. R3 supplies the unifying interpretation that resolves F2. Modular flow $\sigma_{t_1}^\omega \circ \sigma_{t_2}^\omega = \sigma_{t_1+t_2}^\omega$ is *additive* in thermal time (forward-triangle), while geometric proper time on a Lorentzian manifold is *super-additive* on causal curves (the "twin paradox"). The two triangle directions are not "mismatched": they encode the same wedge KMS state under two distinct projection mechanisms (operator-algebraic modular flow vs. synthetic causal geometry), connected by the Wick rotation.

5.2 Definition of the Wick-rotation functor

Definition 5.1 (The category $\mathbf{KreinMetaMet}_{\text{pp}}$). Let $\mathbf{KreinMetaMet}_{\text{pp}}$ be the category whose:

- Objects are Krein PPQMS $\mathbb{X}^K = (\mathcal{A}^K, L^K, \mathcal{M}^L, \omega_W^L)$ per Definition 3.13;
- Morphisms are topographic Krein M-isometries $\pi^K : \mathbb{X}_1^K \rightarrow \mathbb{X}_2^K$, i.e. proper *-epimorphisms preserving topography and pin state.

Definition 5.2 (The category $\mathbf{LorPLG}_{\text{cov}}$). Let $\mathbf{LorPLG}_{\text{cov}}$ be the category whose:

- Objects are covered LPLS $(X, \ell, o, \mathcal{U})$ per Definition 2.9;
- Morphisms are ℓ -preserving maps preserving basepoint o and cover $\mathcal{U}$.

Definition 5.3 (The Wick-rotation functor). The *Wick-rotation functor* $\mathbf{W} : \mathbf{KreinMetaMet}_{\text{pp}} \rightarrow \mathbf{LorPLG}_{\text{cov}}$ is defined on objects by

$$\mathbf{W}(\mathcal{A}^K, L^K, \mathcal{M}^L, \omega_W^L) := (\hat{\mathcal{M}}^L, \ell^L, \hat{\omega}_W^L, \hat{\mathcal{U}}^L),$$

where:

1. $\hat{\mathcal{M}}^L := \text{Spec}(\mathcal{M}^L)$ is the Gelfand spectrum of the M-diagonal Abelian topography;
2. $\hat{\omega}_W^L$ is the character of $\mathcal{M}^L$ induced by the BW vacuum;
3. $\hat{\mathcal{U}}^L = (\hat{U}_k)_{k \in \mathbb{N}}$ with $\hat{U}_k := \text{Spec}(\mathcal{M}^L|_{(n_{\max}(k), N_t(k), T(k))})$ along admissible scaling per Paper 47 [51];
4. The time separation ℓ^L on $\hat{\mathcal{M}}^L$ is

$$\ell^L(\hat{\omega}, \hat{\omega}') := \begin{cases} \kappa_g \cdot \tau_{\text{mod}}^{\omega_W^L}(\hat{\omega}, \hat{\omega}') & \text{if } \hat{\omega} \preceq \hat{\omega}' \text{ along modular flow,} \\ -\infty & \text{otherwise.} \end{cases}$$

On morphisms, $\mathbf{W}(\pi^K) := \hat{\pi}^K$ is the dual Gelfand spectrum map (contravariant by Gelfand duality: $\mathbf{KreinMetaMet}_{\text{pp}} \rightarrow \mathbf{LorPLG}_{\text{cov}}$ on objects sends $\mathbb{X}^K$ forward but on morphisms the arrow direction reverses).

5.3 The Krein–Mondino–Sämann Bridge Theorem

Theorem 5.4 (Krein–Mondino–Sämann Bridge). *The Wick-rotation functor $\mathbf{W}$ of Definition 5.3 is well-defined, with the following four properties:*

- (B1) Structural correspondence. $\mathbf{W}$ sends the Krein PPQMS substrate to the Mondino–Sämann covered LPLS structure row-by-row as in Table 1.

(B2) Reverse triangle inequality. For every $\mathbb{X}^K$ and every $\hat{\omega}_x, \hat{\omega}_y, \hat{\omega}_z \in \mathbf{W}(\mathbb{X}^K)$,

$$\ell^L(\hat{\omega}_x, \hat{\omega}_y) + \ell^L(\hat{\omega}_y, \hat{\omega}_z) \leq \ell^L(\hat{\omega}_x, \hat{\omega}_z).$$

(B3) Pre-compactness inheritance. The Mondino–Sämman pre-compactness conditions (Thm 6.2 of MS) hold on $\mathbf{W}(\mathbb{X}_n^K)$ for any sequence of truncated Krein PPQMS $(\mathbb{X}_n^K)$ at admissible scaling cutoffs.

(B4) Convergence transport (*descoped*). If $\mathcal{D}^K(\mathbb{X}_n^K, \mathbb{X}_\infty^K) \rightarrow 0$, then $\mathbf{W}(\mathbb{X}_n^K) \rightarrow \mathbf{W}(\mathbb{X}_\infty^K)$ in the Mondino–Sämman pLGH sense. (The premise $\mathcal{D}^K \rightarrow 0$ is built from the degenerate Krein metric of Paper 45, so this transport is conditional pending substrate repair, not an established metric convergence — see Theorem 5.8.)

5.4 Structural correspondence (B1)

The structural correspondence holds row-by-row per Table 1:

Table 1: The structural correspondence (B1) between Mondino–Sämman covered LPLS and Krein PPQMS structures, exhibited by the Wick-rotation functor $\mathbf{W}$.

#	Mondino–Sämman (synthetic)	Krein-algebraic-side analog
1	Underlying set X	$\hat{\mathcal{M}}^L = \text{Spec}(\mathcal{M}^L)$
2	Topology finer than chronological	weak-* topology on $\hat{\mathcal{M}}^L$
3	Time separation ℓ	Wick-rotated $\ell^L = \kappa_g \cdot \tau_{\text{mod}}$
4	Reverse triangle (Eq. (1))	on orbit: equality; off orbit: trivial $-\infty$
5	Timelike $\ll$	modular precedence with $t > 0$
6	Causal $\leq$	modular precedence with $t \geq 0$
7	Causal diamond	$\{\hat{\omega}_x \circ \sigma_s : 0 \leq s \leq t\}$
8	Basepoint event o	BW vacuum $\hat{\omega}_W^L$
9	Cover $\mathcal{U}$	truncated topography spectra $(\hat{U}_k)$
10	Cover scales β_k	BW canonical period $\beta_k = 2\pi/\kappa_g$
11	Slab bound	uniform 2π via modular periodicity
12	ε -net	modular causal diamond ε -net at scale $\gamma^{\text{joint}}(k)$
13	LGH convergence	truncation-projector correspondence
14	pLGH convergence	per-cover LGH convergence
15	Pre-compactness	cardinality bound from propagation = 2

5.5 Reverse triangle inequality (B2): Decomposition O on the M-diagonal topography

The substantive content of (B2) is the resolution of F2 via R3 plus the structural simplification that off-orbit cases are trivial at the K^+ -weak-form bridge.

Lemma 5.5 (Modular flow acts trivially on the topography). For every topographic generator $a = M_{N,L,0}^{\text{spat}} \otimes I_{N_t} \in \mathcal{M}^L$,

$$\sigma_t^{\omega_W^L}(a) = a \quad \forall t \in \mathbb{R}.$$

Consequently, the dual action on $\hat{\mathcal{M}}^L$ is trivial: $\hat{\sigma}_t^{\omega_W^L}(\hat{\omega}) = \hat{\omega}$ for all $\hat{\omega}$.

Proof. $[K_\alpha^W, M_{N,L,0}^{\text{spat}}] = 0$ because K_α^W is M-diagonal and the topographic generator has $M = 0$, both diagonal in the same (N, L, M) -block decomposition. Hence $\sigma_t^{\omega_W^L}(a) = e^{-itK_\alpha^W} a e^{itK_\alpha^W} = a$, and the dual Gelfand action is trivial. $\square$

Lemma 5.6 (Decomposition O for triples of characters). *For $\hat{\omega}_x, \hat{\omega}_y, \hat{\omega}_z \in \hat{\mathcal{M}}^L$ at finite cutoff, indexed by topographic eigenvalue labels $(N_x, L_x), (N_y, L_y), (N_z, L_z)$ with $M = 0$, the off-orbit case decomposes into four sub-cases:*

- (i) *All three on the same modular-flow orbit: $(N_x, L_x) = (N_y, L_y) = (N_z, L_z)$. On-orbit case;*
- (ii) *$\hat{\omega}_x, \hat{\omega}_z$ on the same orbit, $\hat{\omega}_y$ on a different orbit. Then $\ell^L(\hat{\omega}_x, \hat{\omega}_y) = -\infty$ (or $\ell^L(\hat{\omega}_y, \hat{\omega}_z) = -\infty$);*
- (iii) *$\hat{\omega}_x, \hat{\omega}_y$ on one orbit, $\hat{\omega}_y, \hat{\omega}_z$ on a different orbit, $\hat{\omega}_x, \hat{\omega}_z$ on yet a third. **Empty** (this would require $(N_x, L_x) = (N_y, L_y) \neq (N_z, L_z) = (N_x, L_x)$, a contradiction);*
- (iv) *All three on different orbits. At least one $\ell^L = -\infty$.*

Proof sketch. The orbit-preservation of modular flow (Lemma 5.5) means modular flow on $\hat{\mathcal{M}}^L$ preserves the (N, L) labels. Hence two characters are causally related ($\hat{\omega} \preceq \hat{\omega}'$ along modular flow with finite τ_{mod}) only if they share the same (N, L) label. Cases (i)–(iv) are then a complete enumeration of label-tuple configurations, and (iii) is structurally empty. $\square$

Proof of Theorem 5.4 (B2). By Lemma 5.6, we analyze (B2) case by case.

Case (i), on-orbit (equality). If $\hat{\omega}_x \preceq \hat{\omega}_y \preceq \hat{\omega}_z$ along modular flow with $\hat{\omega}_y = \hat{\omega}_x \circ \sigma_{t_1}^{\omega_W^L}$ and $\hat{\omega}_z = \hat{\omega}_y \circ \sigma_{t_2}^{\omega_W^L}$, then $\hat{\omega}_z = \hat{\omega}_x \circ \sigma_{t_1+t_2}^{\omega_W^L}$ by additivity of modular flow. Hence $\ell^L(\hat{\omega}_x, \hat{\omega}_y) + \ell^L(\hat{\omega}_y, \hat{\omega}_z) = \kappa_g(t_1 + t_2) = \ell^L(\hat{\omega}_x, \hat{\omega}_z)$.

Cases (ii), (iv), trivial $-\infty$ inheritance. At least one of $\ell^L(\hat{\omega}_x, \hat{\omega}_y)$ or $\ell^L(\hat{\omega}_y, \hat{\omega}_z)$ is $-\infty$, so the LHS is $-\infty$ by MS convention, and $-\infty \leq \ell^L(\hat{\omega}_x, \hat{\omega}_z)$ trivially.

Case (iii), empty. No contribution.

Hence (B2) holds in all cases. $\square$

Remark 5.7 (The super-additivity that DOES exist lives in $\mathcal{O}^L$, not in $\mathcal{M}^L$). The substantive super-additivity property (off-axis Lorentz-boost composition accumulating more proper time than direct composition) exists on the wedge but lives in the *full* Krein operator system $\mathcal{O}^L$, not in the M-diagonal topography $\mathcal{M}^L$. The Paper 46 [50] Appendix B enlarged substrate (chirality- flipping generators M^{flip} with $\{J_L, M^{\text{flip}}\} = 0$) supports such off-axis composition; on this enlarged substrate, the bridge **W** would extend to a richer LPLS structure with strict super-additivity off-orbit. This is the open question Q1' (Section 8.1).

5.6 Pre-compactness inheritance (B3)

Proof of Theorem 5.4 (B3). The Mondino–Sämman pre-compactness Thm 6.2 has three conditions: (i) uniform slab bounds; (ii) uniform ε -net cardinality bounds; (iii) cumulative ε -net nesting.

(i) *Uniform slab bound.* The timelike diameter of $\hat{U}_k$ equals $\sup \tau_{\text{mod}}^{\omega_W^L}$ on the wedge. Under BW canonical $\kappa_g = 1$, modular flow is 2π -periodic (Paper 42 [46] § 5 integer spectrum of K_α^W), so $\tau_{\text{mod}} \leq 2\pi$ uniformly: $T_k = 2\pi$ for all k .

(ii) *Uniform ε -net cardinality bound.* At finite cutoff $(n_{\text{max}}(k), N_t(k), T(k))$, $|\hat{U}_k| = O(n_{\text{max}}(k)^4 \cdot N_t(k))$ (dimension of the topography), and the ε -net cardinality is bounded above by $|\hat{U}_k|$ itself and by the standard discrete-to-continuous refinement bound $N(k, \varepsilon) = \min(|\hat{U}_k|, \text{Vol}(\hat{U}_k)/\varepsilon^{d_k})$. Both bounds are uniform via Paper 44 [48] propagation number = 2.

(iii) *Cumulative ε -net nesting.* By the admissible-scaling sequence $(n_{\text{max}}(k), N_t(k), T(k)) \rightarrow (\infty, \infty, T_\infty)$, the cover $\hat{U}_k \subseteq \hat{U}_{k+1}$ is nested by construction, and ε -nets of $\hat{U}_k$ extend to ε -nets of $\hat{U}_{k+1}$ by adding finitely many points.

By Mondino–Sämman Thm 6.2, the bridge image inherits pre-compactness. $\square$

5.7 Convergence transport (B4) via Plancherel-nested Berezin

Theorem 5.8 (Convergence transport — *descoped*). *If $\mathcal{D}^K(\mathbb{X}_n^K, \mathbb{X}_\infty^K) \rightarrow 0$ in the Krein hypertopology of Definition 4.8, then $\mathbf{W}(\mathbb{X}_n^K) \rightarrow \mathbf{W}(\mathbb{X}_\infty^K)$ in the Mondino–Sämman pLGH topology of Definition 2.10, with the strong (timelike forward density) sense. (Descoped: $\mathcal{D}^K$ is built from the degenerate Krein Lipschitz seminorm L^K of Paper 45, so the premise $\mathcal{D}^K \rightarrow 0$ is not a non-degenerate metric convergence; this transport is conditional pending substrate repair, not an established metric-level result.)*

Proof sketch. By the Krein hypertopology coincidence (Theorem 4.5) plus the bit-exact panel inheritance from Paper 45 [49], the Krein-side propinquity convergence implies that for each cover level k , the truncated topography spectra $\hat{U}_{k,n} \rightarrow \hat{U}_{k,\infty}$ in the LGH sense (per Definition 2.8). The Plancherel-nested Berezin/projection pair from Paper 45 [49] Sub-Sprint C/D supplies a Mondino–Sämman correspondence with vanishing distortion. Per-axiom verification:

Cardinality matching (MS Def 3.6(i)). At fixed admissible cutoff, the correspondence pair has $|S_n| = |S_\infty|$ on the M-diagonal topography (the Berezin map preserves the (N, L) shell structure).

Correspondence with vanishing distortion (MS Def 3.6(ii)). The Berezin/projection pair has $\text{reach} \leq \gamma^{\text{joint}} + \text{height} \leq \gamma^{\text{joint}} \cdot C_3^{\text{joint}}$, vanishing as $n \rightarrow \infty$.

Extension property (MS Def 3.6(iii)). The Plancherel-nested structure of Berezin maps IS the MS extension property: Berezin maps at finer cutoff extend Berezin maps at coarser cutoff because the Plancherel weights nest by construction (Paper 45 [49] Sub-Sprint C Lemma 5.1).

Forward density (MS Def 3.6(iv)). The modular-flow orbit discretization in the chronological topology provides the timelike forward density: for any $\hat{\omega}' \succ \hat{\omega}$ on a modular orbit, the truncation projects $\hat{\omega}'$ to a sequence $\hat{\omega}'_n$ converging to $\hat{\omega}'$ in the chronological topology with $\hat{\omega} \prec \hat{\omega}'_n$ for all sufficiently large n .

Per-cover LGH convergence at each k implies pLGH convergence per Definition 2.10. $\square$

5.8 Riemannian-limit recovery cross-check

Corollary 5.9. *At $N_t = 1$, the spatial/ $SU(2)$ factor of the Krein-side propinquity bound equals $\Lambda_{n_{\max}}^{\text{Paper 38}}$ bit-exact (Paper 45 Sub-Sprint D Lemma 5.1; the joint rate carries the $U(1)$ $T/4$ offset of Theorem 4.9), so the bridge image $\mathbf{W}(\mathbb{X}^K|_{N_t=1})$ is the synthetic-side counterpart of Paper 38's $SU(2)$ state-space Gromov–Hausdorff convergence: a trivial LPLS with no Lorentzian content (modular flow on the Riemannian limit collapses to identity, and $\ell^L|_{N_t=1} = 0$ along the Gelfand spectrum).*

5.9 Honest scope of the bridge theorem

The bridge functor $\mathbf{W}$ is constructed at the K^+ -weak-form level on the natural chirality-doubled scalar-multiplier substrate of Papers 44/45. The strong-form extension of $\mathbf{W}$ to the Paper 46 [50] Appendix B enlarged substrate (chirality-flipping generators) is **not** constructed in this paper; it is the open question Q1' (Section 8.1).

The bridge is **functorial**, **not isometric**: $\mathcal{D}^K$ and ℓ^L measure different physical quantities (thermal vs. geometric time) of the same wedge KMS state. The two triangle directions are not "mismatched" — they encode different projections of the same physical object, connected by the Wick rotation under R3 (Connes–Rovelli thermal time).

6 Application to the GeoVac hemispheric wedge

6.1 The GeoVac wedge as a Krein PPQMS

Lemma 6.1 (The wedge as Krein PPQMS). *The Paper 43 [47] hemispheric wedge $W_L = P_W^{\text{spat}} \otimes P_{t \geq 0}$ at every finite cutoff $(n_{\max}, N_t, T)$ defines a Krein PPQMS*

$$\mathbb{X}_{\text{wedge}}^K := (\mathcal{A}^L|_{W_L}, L^K|_{W_L}, \mathcal{M}^L|_{W_L}, \omega_W^L),$$

with all five Krein PPQMS axioms of Definition 3.13 verified by direct construction.

Proof sketch. (A1) $\mathcal{A}^L|_{W_L}$ is the wedge-restricted Lorentzian operator system (Paper 44 [48] § 2). (A2) $L^K|_{W_L}$ is the K^+ -restricted Krein-Leibniz Lipschitz seminorm of Definition 3.1. (A3) $\mathcal{M}^L|_{W_L}$ is the wedge-restricted M-diagonal Abelian topography of Lemma 3.12. (A4) $\omega_W^L = e^{-K_\alpha^W}/Z$ is the wedge KMS state (Paper 43 [47] § 4.2), diagonal in the M-eigenbasis hence a character of $\mathcal{M}^L|_{W_L}$. (A5) The Krein M-tunnel structure with $\chi^K \rightarrow 0$ is the wedge restriction of Lemma 4.2. $\square$

6.2 The seven newly accessible theorems

We apply the bridge functor $\mathbf{W}$ to the wedge PPQMS:

$$\mathbf{W}(\mathbb{X}_{\text{wedge}}^K) = (\hat{\mathcal{M}}^L|_{W_L}, \ell_{\text{wedge}}^L, \hat{\omega}_W^L, \hat{\mathcal{U}}_{\text{wedge}}^L),$$

where at panel cells the cardinality is $|\hat{\mathcal{M}}^L|_{W_L}| = \dim W_L = 16, 60, 160$ at $(n_{\max}, N_t) \in \{(2, 3), (3, 5), (4, 7)\}$ respectively.

6.2.1 T1: The wedge as Mondino–Sämann covered LPLS

Theorem 6.2 (T1: GeoVac wedge as Mondino–Sämann pointed pre-length space). *For every finite cutoff $(n_{\max}, N_t, T)$ with $T = 2\pi$ canonical, the GeoVac hemispheric wedge W_L defines a Mondino–Sämann covered Lorentzian pre-length space*

$$(\hat{\mathcal{M}}^L|_{W_L}, \ell_{\text{wedge}}^L, \hat{\omega}_W^L, \hat{\mathcal{U}}_{\text{wedge}}^L) \in \mathbf{LorPLG}_{\text{cov}},$$

with uniform timelike diameter 2π across all cover levels.

Proof sketch. Direct application of Theorem 5.4 (B1) plus Lemma 6.1. The uniform timelike diameter follows from Paper 43 [47] Thm bw_alpha_lorentz: modular flow is 2π -periodic on the wedge. $\square$

6.2.2 T2: Synthetic compactness of the wedge family

Theorem 6.3 (T2: GeoVac wedge synthetic compactness). *Any sequence $(\mathbb{X}_{\text{wedge}}^K(k_n))_{n \in \mathbb{N}}$ of GeoVac wedge Krein PPQMS at admissible-scaling cutoffs $(n_{\max}(k_n), N_t(k_n), T(k_n)) \rightarrow (\infty, \infty, T_\infty)$ has a subsequence whose bridge images $\mathbf{W}(\mathbb{X}_{\text{wedge}}^K(k_n))$ converge in the Mondino–Sämann pLGH sense to a covered LPLS limit. (The pre-compactness leg survives via the propagation-number cardinality bound; the **convergence** of the subsequence to its limit inherits the metric-level descope of Theorem 5.8 — it is conditional pending substrate repair.)*

Proof sketch. By Theorem 5.4 (B3), the three pre-compactness conditions of MS Thm 6.2 hold for the wedge sequence. The MS Thm 6.2 yields the pLGH-convergent subsequence. $\square$

6.2.3 T3: Synthetic bit-exact panel inheritance

[Descoped (see the Status note): the inherited values are evaluations of the closed-form rate formula, not measurements of a metric; the panel transports formula values, not metric data.]

Theorem 6.4 (T3: Synthetic pLGH-convergence rate panel for the GeoVac wedge). *The Mondino–Sämman pLGH-convergence rate of the GeoVac wedge sequence at the Paper 45 [49] panel cells satisfies:*

$(n_{\max}, N_t)$	$d_{\text{pLGH}}^{\text{wedge}} \text{ (Krein-side } \Lambda^L)$
(2, 3)	≤ 2.0746
(3, 5)	≤ 1.6101
(4, 7)	≤ 1.3223

*bit-identical to the Paper 45 Krein-side Λ^L values. These are the descoped Paper 45 **rate-formula** evaluations (not metric-convergence measurements), transported conditionally through the bridge: the Lorentzian quantum-metric convergence is descoped, so this is a rate panel, not an established pLGH-convergence panel.*

Proof sketch. The bridge **W** sends the Krein-side propinquity bound Λ^L to the synthetic-side pLGH rate d_{pLGH} . The functor is not isometric ($\mathfrak{D}^K$ and ℓ^L measure different physical quantities), but the rate parameter $\gamma^{\text{joint}} = O(\log n_{\max}/n_{\max} + T/N_t)$ is the same object on both sides, hence the numerical panel transports bit-identically. $\square$

Remark 6.5 (The substantive new content). All extant Mondino–Sämman-style examples (arXiv:2504.10380 v4 § 10) are constructed geometrically from continuous spacetimes: Chruściel–Grant approximations of globally hyperbolic spacetimes, sequences of warped products, etc. Theorem 6.4 is the first quantitative pLGH-convergence rate panel in the MS literature derived from *operator-algebraic* input via the bridge.

6.2.4 T4: Synthetic four-witness readings

Theorem 6.6 (T4: Synthetic four-witness readings). *The Paper 42 [46] four-witness Wick-rotation theorem identifies six physical thermal-time witnesses (Bisognano–Wichmann canonical, Hartle–Hawking at $M = 1, 2$, Sewell at $M = 1$, Unruh at $a = 1, 2$) on the operator-algebraic side, all giving bit-identical modular operators on the wedge KMS state. Via **W**, these six physical witnesses give six geometric-time readings of the same synthetic-Lorentzian wedge pre-length space, distinguished only by the rapidity-rate normalization κ_g :*

Witness	κ_g	Synthetic time scale
BW canonical	1	$\ell^L = \tau_{\text{mod}}$
Hartle–Hawking $M = 1$	1/4	$\ell^L = \tau_{\text{mod}}/4$
Hartle–Hawking $M = 2$	1/8	$\ell^L = \tau_{\text{mod}}/8$
Sewell $M = 1$	1/4	same as HH $M = 1$
Unruh $a = 1$	1	same as BW canonical
Unruh $a = 2$	2	$\ell^L = 2\tau_{\text{mod}}$

*Witnesses with the same κ_g produce identical synthetic LPLS: the six physical instantiations on the operator side reduce to **two synthetic-side equivalence classes**.*

Proof sketch. The bridge time separation $\ell^L = \kappa_g \cdot \tau_{\text{mod}}$ rescales linearly with κ_g . The bit-exact algebra-action collapse of the six witnesses on the Krein side (Paper 42 [46] § 8 Cor 8.1) translates to a bit-exact metric-rescaling identification on the synthetic side. $\square$

6.2.5 T5: Synthetic Pythagorean orthogonality with $1/\pi^2$ M1 signature

Theorem 6.7 (T5: Synthetic-side Pythagorean orthogonality). *Under the bridge $\mathbf{W}$ applied to the GeoVac hemispheric wedge at the Riemannian limit $N_t = 1$, the BW-aligned thermal-time generator $H_{\text{local}} = K_\alpha^W/\beta$ and the wedge-restricted Lorentzian Dirac D_W^L correspond to two structurally orthogonal foliation structures on the synthetic wedge pre-length space: a boost-orbit foliation $\mathcal{F}_\alpha$ (H_{local} generator, Π_W -even per Paper 43 [47] Lem parity_HS input (I1)) and a spinor-bundle foliation $\mathcal{F}_D$ (D_W^L generator, mixed Π_W -parity — temporal γ^0 odd, spatial D_{GV} even — per the corrected Paper 43 [47] input (I2)).*

The Paper 43 [47] § 10.2 Pythagorean closed-form residual

$$\|H_{\text{local}} - D_W^L\|_F^2 = \frac{\kappa_g^2 \cdot S(n_{\text{max}})}{4\pi^2} + D(n_{\text{max}})$$

translates to a synthetic-side decomposition of the wedge’s geometric data into a κ_g^2 -scaling Hopf-base-measure contribution ($S(n_{\text{max}})/(4\pi^2)$) and a κ_g -independent spinor-bundle contribution ($D(n_{\text{max}})$). The $1/\pi^2 = \text{Vol}(S^1)^{-2}$ master Mellin engine M1 signature (Paper 32 [42] § VIII rem:pythagorean_m1_closure) identifies as the inverse-squared measure of the bridge cover’s slab-period.

Proof sketch. The algebraic identity is Paper 43 [47] Thm pythagorean_formal. The synthetic-side reading is by application of $\mathbf{W}$: the Π_W -grading on the Krein side translates to the orthogonal foliation structure on the synthetic side via Gelfand duality, and the $1/\pi^2$ prefactor on the operator-algebraic Pythagorean residual IS the synthetic-side wedge’s slab-period inverse-squared measure ($\beta = 2\pi$, $1/\beta^2 = 1/(4\pi^2)$). $\square$

6.2.6 T6: G2 metric-level closure for the GeoVac wedge

[Descoped (see the Status note): the “metric-equivalent level” claimed below quantifies over the degenerate Paper 45 seminorm; the metric-level closure is open pending the upstream repair. The norm-resolvent ingredients from Paper 47 are unaffected.]

Theorem 6.8 (T6: G2 metric-level route for the GeoVac hemispheric wedge (descoped — conditional on the Paper 45 degenerate seminorm)). *For the GeoVac hemispheric wedge specifically, the de-compactification limit $T \rightarrow \infty$ closes at the propinquity-equivalent level on the synthetic side: the synthetic wedge images $\mathbf{W}(\mathbb{X}_{\text{wedge}}^K(k))$ at admissible-scaling cutoffs converge in the Mondino–Sämman pLGH topology to the synthetic non-compact wedge object $\mathbf{W}(\mathbb{X}_{\text{wedge}}^K(\infty))$ (where $\mathbb{X}_{\text{wedge}}^K(\infty)$ is the bridge image of the Paper 47 [51] non-compact carrier $S^3 \times \mathbb{R}_t$ wedge).*

Equivalently, $d_{\text{pLGH}}(\mathbf{W}(\mathbb{X}_{\text{wedge}}^K(k)), \mathbf{W}(\mathbb{X}_{\text{wedge}}^K(\infty))) \rightarrow 0$ as $T(k) \rightarrow \infty$.

*This is a **GeoVac-wedge-specific proposed route toward G2** (de-compactification limit, Paper 45 [49] § 1.4) at the **metric-equivalent level on the synthetic side** — descoped, conditional on the descoped Krein seminorm (Paper 45); it is not an established closure.*

Proof sketch. By Paper 47 [51] § 6, the three Krein-side carriers (periodic compact, bounded Dirichlet, non-compact $\mathbb{R}_t$) all converge in norm-resolvent sense to the same physical Lorentzian Dirac as $T \rightarrow \infty$. Under the bridge $\mathbf{W}$: $\mathbf{W}(\mathbb{X}_{\text{per}}^K(T))$, $\mathbf{W}(\mathbb{X}_{\text{Dir}}^K(T))$, and $\mathbf{W}(\mathbb{X}_{\mathbb{R}}^K)$ are well-defined Mondino–Sämman pre-length spaces (T1 applies). Norm-resolvent convergence on the Krein side plus the bridge functoriality plus the (B2)+(B4) bridge properties give pLGH convergence on the synthetic side.

The substantive content is that the synthetic-side bridge image carries the structural content of metric-level convergence even though the Krein-side propinquity Λ^L is not defined on non-compact carriers (the Paper 45 propinquity requires compact temporal radius). $\square$

6.2.7 T6a: Synthetic three-carrier identification

Corollary 6.9 (T6a: Synthetic three-carrier identification). *The Paper 47 [51] § 6 three-carrier identification on the operator-algebraic side translates via the bridge $\mathbf{W}$ to a synthetic-side three-carrier identification: the three synthetic wedge images $(\mathbf{W}(\mathbb{X}_{\text{per}}^K), \mathbf{W}(\mathbb{X}_{\text{Dir}}^K), \mathbf{W}(\mathbb{X}_{\mathbb{R}}^K))$ are isometric as Mondino–Sämman covered LPLS in the limit $T \rightarrow \infty$.*

Proof sketch. The Paper 47 isometric embeddings $\tilde{J}_T^{\text{per}}, \tilde{J}_T^{\text{Dir}} : \mathcal{K}_T^\bullet \hookrightarrow \mathcal{K}_{\mathbb{R}}$ commute with the M-diagonal topographies (which are time-independent), hence descend via Gelfand duality to maps on the synthetic side. The synthetic three-carrier identification is the dual of the Krein-side identification. $\square$

6.3 Honest scope of the wedge application

The bridge construction itself applies cleanly to every load-bearing GeoVac structure (Paper 42 four-witness, Paper 43 wedge + Pythagorean, Paper 45 panel, Paper 47 three-carrier). The headline content (T6 G2 metric-equivalent route for the wedge) is a *conditional* design — it inherits the descoped Paper 45 degeneracy, so the metric-level closure is *not* established (descoped), only a proposed route; the general non-compact Latrémolière problem remains open. The strong-form bridge to the enlarged substrate is Q1' (Section 8.1).

7 Synthetic-Lorentzian compactness theorems

7.1 Synthetic compactness of the GeoVac wedge family

Theorem 6.3 (T2) provides the synthetic compactness statement for the GeoVac wedge family. We restate it in the context of synthetic-Lorentzian compactness theory.

Theorem 7.1 (Synthetic compactness restated). *The class $\mathfrak{X}_{\text{GeoVac wedge}}$ of bridge images of GeoVac wedge Krein PPQMS at admissible-scaling cutoffs is pre-compact in the Mondino–Sämman pLGH topology of Definition 2.10.*

This is a substantive new content for the Mondino–Sämman literature: the wedge family is the first operator-algebraically derived class to satisfy the Mondino–Sämman pre-compactness criteria.

7.2 The first quantitative pLGH panel from operator-algebraic input

By Theorem 6.4 (T3), the GeoVac wedge family furnishes the first quantitative pLGH-convergence panel derived from operator-algebraic input. The bit-exact numerical values $\{2.0746, 1.6101, 1.3223\}$ across $(n_{\text{max}}, N_t) \in \{(2, 3), (3, 5), (4, 7)\}$ provide a concrete benchmark for future Mondino–Sämman-style synthetic-Lorentzian numerical work.

7.3 G2 metric-equivalent closure for the GeoVac wedge (proposed/descoped)

The deepest *proposed* content of the paper is Theorem 6.8 (T6) — a conditional route, *descoped* (it rests on Paper 45's degenerate Krein-side Λ^L ; see the Status note). We unpack the structural significance:

- **The non-compact Latrémolière propinquity is not in the published literature.** The original Latrémolière propinquity [22, 24] and its 2025 hypertopology extension [25] operate on compact (or proper, locally compact) quantum metric spaces. A non-compact extension to the non-compact carrier $S^3 \times \mathbb{R}_t$ remains open. This is the published-open problem G2 metric-level of Paper 45 [49] § 1.4.

- **The bridge supplies a workaround at the wedge level.** The Wick-rotation functor $\mathbf{W}$ converts the operator-algebraic non-compact propinquity question to a synthetic-Lorentzian pLGH convergence question (Mondino–Sämman pLGH on non-compact LPLS is well-defined). On the repaired (non-degenerate) seminorm the synthetic-side closure *would* exist and the bridge *would* transport it back to the operator side — but as it stands this rests on Paper 45’s degenerate Λ^L , so T6 is a *proposed/conditional* route, not an established closure (Status note).
- **The closure is wedge-specific.** The bridge applies to any Krein PPQMS, but the headline content (G2 closure) requires the 2π -periodicity of modular flow on the wedge and the bridge’s ability to extend along the admissible-scaling sequence to the non-compact carrier. For more general (non-wedge) carriers, the 2π -periodicity does not hold and the closure does not extend.

7.4 Synthetic three-carrier identification

Corollary 6.9 (T6a) extends the Paper 47 [51] three-carrier identification from the operator side to the synthetic side. This resolves an apparent tension between Papers 43 and 45’s different temporal-carrier choices (bounded Dirichlet vs. periodic compact) at the synthetic level: both, plus the non-compact carrier, are isometric as covered LPLS in the de-compactification limit.

7.5 Why this is genuinely new content for Mondino–Sämman pLGH theory

The Mondino–Sämman pLGH theory [30] develops the synthetic side independently of operator algebras. All extant examples in arXiv:2504.10380 v4 § 10 are constructed geometrically: Chruściel–Grant approximations of globally hyperbolic spacetimes, warped products, etc. The bridge construction of this paper introduces a categorically new source of MS-style examples: operator-algebraic truncations transported to the synthetic side via the Wick-rotation functor. The bridge construction itself is a genuinely new *source* of MS-style examples; the downstream headlines (T3 quantitative panel, T6 G2 route) are *conditional* on the bridge and descoped at the metric level (they inherit the Paper 45 degeneracy), so they are proposed content, not established new results, for the Mondino–Sämman literature.

7.6 Related synthetic-Lorentzian convergence frameworks

We position the bridge construction vs. adjacent synthetic-Lorentzian convergence frameworks. The Sormani–Vega null distance [36] and Sakovich–Sormani timed Gromov–Hausdorff [34] operate via null-distance metric-GH rather than via Mondino–Sämman’s ε -net-of-causal-diamonds. Minguzzi–Suhr bounded Lorentzian metric spaces [28] extend a complementary distinction metric. Allen–Burtscher metric GH [1] and Müller’s Cauchy slabs [31] provide alternative frameworks. The bridge of this paper targets specifically the Mondino–Sämman pLGH framework on covered LPLS; a parallel bridge to null-distance or timed-Hausdorff frameworks would be a separate construction.

8 Open questions

8.1 Q1’: Strong-form bridge on the enlarged substrate

The most natural follow-on to the K^+ -weak-form bridge of this paper is the strong-form bridge extending $\mathbf{W}$ to the Paper 46 [50] Appendix B enlarged substrate (chirality-flipping generators M^{flip} with $\{J_L, M^{\text{flip}}\} = 0$).

Open question Q1' (strong-form Krein–MS bridge). *Does the Wick-rotation functor $\mathbf{W}$ extend to the enlarged substrate of Paper 46 [50] Appendix B, and if so, does the off-orbit super-additivity close via direct transport of the Paper 42 [46] four-witness theorem to MS time separation?*

The substantive question that A.4'-A surfaced

On the K^+ -weak-form bridge of this paper, off-orbit super-additivity is structurally vacuous via Lemma 5.6 (Case (iii) is empty under the M-diagonal topography). On the strong-form bridge, the enlarged topography makes Case (iii) non-empty: there are triples $(\hat{\omega}_x, \hat{\omega}_y, \hat{\omega}_z)$ such that all three pairs are on different orbits with non-trivial chirality-flipping content.

In this regime, the operator-algebraic super-additivity from Paper 42's four-witness theorem becomes load-bearing: the on-orbit equality $\ell^L(x, y) + \ell^L(y, z) = \ell^L(x, z)$ generalizes to a strict super-additivity $\ell^L(x, y) + \ell^L(y, z) \leq \ell^L(x, z)$ via the analog of the special-relativistic twin paradox at the operator-algebraic level.

Q1' three-step decomposition

- (Q1'.A) Topography enlargement.** Extend the topography $\mathcal{M}^L$ to include M^{flip} generators; verify the topography axioms (Definition 3.11) hold for the enlarged structure. Substantive risk: the enlargement may break Abelianness (chirality-flipping generators do not commute with each other in general), in which case the Gelfand-spectrum construction does not apply directly.
- (Q1'.B) Gelfand spectrum.** Verify $\hat{\mathcal{M}}_{\text{enlarged}}^L$ is well-defined. If Abelianness breaks: generalize to non-commutative Mondino–Sämman pre-length spaces (a substantial NCG-research target not currently in the literature).
- (Q1'.C) Off-orbit super-additivity.** Prove $\ell^L(x, y) + \ell^L(y, z) \leq \ell^L(x, z)$ at strict-inequality level via Paper 42 four-witness theorem transport to off-axis modular flow.

Q1' effort estimate and recommendation

Estimated effort: a substantial open program, well beyond sprint scale. If Q1'.B forces the non-commutative MS extension route, the program deepens further (the non-commutative MS pre-length space concept does not exist in the published literature; constructing it would itself be a major NCG-mathematics target). Q1' is named here as the principal follow-on; it is the natural target for a Sprint L3e-Q1' or for an independent NCG-research program.

8.2 G2 metric-level closure on the general (non-wedge) carrier

Theorem 6.8 (T6) proposes a descoped, conditional route to G2 metric-level for the GeoVac wedge via the bridge image on the synthetic side. The general non-compact Latrémollière propinquity problem (Paper 45 [49] § 1.4 G2-metric on arbitrary non-wedge carriers) remains the named NCG-mathematics frontier. Paper 47 [51] establishes the analog closure at the norm-resolvent / spectral level on the general carrier; the metric-level analog requires either:

- A non-compact Latrémollière propinquity extension (a substantial NCG-mathematics target, beyond sprint scale); or
- A bridge to a non-wedge synthetic-Lorentzian framework that admits metric-level pLGH convergence on the general carrier (structurally distinct from the wedge bridge here).

8.3 G3 cross-manifold $\mathcal{T}_{S^3} \otimes \mathcal{T}_{\text{Hardy}}(S^5)$

The cross-manifold extension of the Lorentzian Krein spectral triple program is blocked at the NCG-framework level by the Paper 24 [40] § V four-layer Coulomb/HO asymmetry, including the (layer iv) modular-Hamiltonian structure of the wedge KMS state surfaced by Sprint L2-F.1. This obstruction is orthogonal to the present bridge: G3 is a structural extension target for the NCG framework itself, not for the bridge to Mondino–Sämann.

8.4 G4 inner-factor calibration data

The W3 spectral-zeta candidate for inner-factor calibration data was falsified (CLAUDE.md Sprint W3, 2026-05-08: CKM Wolfenstein candidate; three follow-up tracks ruled out PMNS, lepton masses, and the mechanically-generated basis). No concrete proposals for inner-factor calibration data remain in the literature. G4 is orthogonal to the present bridge.

8.5 Q2': Non-commutative Mondino–Sämann pre-length spaces

If Q1'.B (Section 8.1) forces breaking the Abelianness of the enlarged topography, the bridge construction would naturally generalize to non-commutative MS pre-length spaces. This is a substantial NCG-research target not currently in the literature.

Acknowledgments

This work is part of the GeoVac math.OA arc (Papers 29, 32, 38–47). The bridge construction was opened by the Latrémolière 2512.03573 [25] and Mondino–Sämann arXiv:2504.10380 v4 [30] December 2025 publications, with the Connes–Rovelli thermal-time hypothesis [10] supplying the unifying interpretation. Phase A formalization was conducted via six sub-sprint memos (Tier 3-Light diagnostic, A.2' Krein-lift, A.3' bridge construction + concurrent-work re-check, A.4'-A G-B2 closure, A.4'-B G-B4 closure, A.4'-C GeoVac wedge application) and synthesized at Phase A.5'. The paper draft is Phase B of the Sprint L3e-P3 program.

References

- [1] B. Allen, A. Burtscher, “Properties of the null distance and spacetime convergence,” *Int. Math. Res. Not. IMRN* (2022), no. 10, 7729–7808.
- [2] P. Bertozzini, R. Conti, W. Lewkeeratiyutkul, “Modular Theory, Non-Commutative Geometry and Quantum Gravity,” *SIGMA* 6 (2010), 067; arXiv:1007.4094.
- [3] J. J. Bisognano, E. H. Wichmann, “On the duality condition for quantum fields,” *J. Math. Phys.* **17** (1976), 303–321.
- [4] N. Bizi, C. Brouder, F. Besnard, “Space and time dimensions of algebras with applications to Lorentzian noncommutative geometry and quantum electrodynamics,” *J. Math. Phys.* **59** (2018), 062303. arXiv:1611.07062.
- [5] M. Braun, C. Sämann, “Gromov’s reconstruction theorem and measured Gromov–Hausdorff convergence in Lorentzian geometry,” arXiv:2506.10852 (June 2025).
- [6] R. Camporesi, A. Higuchi, “On the eigenfunctions of the Dirac operator on spheres and real hyperbolic spaces,” *J. Geom. Phys.* **20** (1996), 1–18.
- [7] A. H. Chamseddine, A. Connes, “The spectral action principle,” *Comm. Math. Phys.* **186** (1997), 731–750.

- [8] M. Che, R. Perales, C. Sormani, “Gromov’s compactness theorem for the intrinsic timed-Hausdorff distance,” *Differential Geometry and its Applications* **103** (2026), arXiv:2510.13069.
- [9] A. Connes, *Noncommutative Geometry*, Academic Press, San Diego, 1994.
- [10] A. Connes, C. Rovelli, “Von Neumann algebra automorphisms and time-thermodynamics relation in generally covariant quantum theories,” *Class. Quantum Grav.* **11** (1994), 2899–2918.
- [11] A. Connes, W. D. van Suijlekom, “Spectral truncations in noncommutative geometry and operator systems,” *Comm. Math. Phys.* **383** (2021), 2021–2067. arXiv:2004.14115.
- [12] C. Farsi, F. Latrémolière, “Collapse in noncommutative geometry and spectral continuity,” arXiv:2404.00240 (2024).
- [13] C. Farsi, F. Latrémolière, “Continuity for the spectral propinquity of the Dirac operators associated with an analytic path of Riemannian metrics,” arXiv:2504.11715 (April 2025).
- [14] N. Franco, M. Eckstein, “An algebraic formulation of causality for noncommutative geometry,” *Class. Quantum Grav.* **30** (2013), 135007 (and follow-ups).
- [15] J. B. Hartle, S. W. Hawking, “Path-integral derivation of black-hole radiance,” *Phys. Rev. D* **13** (1976), 2188–2203.
- [16] E.-M. Hekkelman and E. McDonald, “A noncommutative integral on spectrally truncated spectral triples, and a link with quantum ergodicity,” arXiv:2412.00628 (2024).
- [17] C. Ketterer, “Convergence of Lorentzian spaces and curvature bounds for generalized cones,” arXiv:2605.11271 (May 2026).
- [18] H. Kubota, “A Lorentzian coarea inequality,” arXiv:2605.09101 (May 2026).
- [19] M. Kunzinger, C. Sämann, “Lorentzian length spaces,” *Ann. Global Anal. Geom.* **54** (2018), 399–447. arXiv:1711.08990.
- [20] B. Kostant, “On the existence and irreducibility of certain series of representations,” *Bull. Amer. Math. Soc.* **75** (1969), 627–642.
- [21] F. Latrémolière, “The dual Gromov–Hausdorff propinquity,” *J. Math. Pures Appl.* **103** (2015), 303–351.
- [22] F. Latrémolière, “The quantum Gromov–Hausdorff propinquity,” *Trans. Amer. Math. Soc.* **368** (2016), 365–411. arXiv:1302.4058.
- [23] F. Latrémolière, “Convergence of Cauchy sequences for the covariant Gromov–Hausdorff propinquity,” *J. Math. Anal. Appl.* **469** (2019), 378–404.
- [24] F. Latrémolière, “The Gromov–Hausdorff propinquity for metric spectral triples,” *Adv. Math.* **404** (2022), Paper No. 108393. arXiv:1811.10843.
- [25] F. Latrémolière, “The quantum Gromov–Hausdorff hypertopology on the class of pointed proper quantum metric spaces,” arXiv:2512.03573 (December 2025).
- [26] M. Leimbach, “Convergence of Peter–Weyl truncations of compact quantum groups,” to appear, *J. Noncommut. Geom.* (2024); arXiv:2409.16698.
- [27] M. Marcolli, W. D. van Suijlekom, “Gauge networks in noncommutative geometry,” *J. Geom. Phys.* **75** (2014), 71–91. arXiv:1301.3480.

- [28] E. Minguzzi, S. Suhr, “Lorentzian metric spaces and their Gromov–Hausdorff convergence,” *Lett. Math. Phys.* **114** (2024), Paper No. 73, arXiv:2209.14384.
- [29] A. Mondino, V. Ryborz, C. Sämann, “Stability of synthetic timelike Ricci bounds under C^0 -limits and applications to impulsive gravitational waves,” arXiv:2605.03172 (May 2026).
- [30] A. Mondino, C. Sämann, “Lorentzian Gromov–Hausdorff convergence and pre-compactness,” arXiv:2504.10380 v4 (December 2025).
- [31] O. Müller, “Lorentzian Gromov–Hausdorff theory and finiteness results,” *Gen. Relativ. Gravit.* **54** (2022), Art. 117; arXiv:1912.00988.
- [32] G. Nieuviarts, “Emergence of Time from a Twisted Spectral Triple in Almost-Commutative Geometry,” arXiv:2512.15450 (December 2025, May 2026 v2).
- [33] M. Reed, B. Simon, *Methods of Modern Mathematical Physics Vol. IV: Analysis of Operators*, Academic Press, New York, 1978.
- [34] A. Sakovich, C. Sormani, “Introducing various notions of distances between space-times,” (2024). arXiv:2410.16800.
- [35] G. L. Sewell, “Quantum fields on manifolds: PCT and gravitationally induced thermal states,” *Ann. Phys. (NY)* **141** (1982), 201–224.
- [36] C. Sormani, C. Vega, “Null distance on a spacetime,” *Class. Quantum Grav.* **33** (2016), 085001. arXiv:1508.00531.
- [37] A. Strohmaier, “On noncommutative and pseudo-Riemannian geometry,” *J. Geom. Phys.* **56** (2006), 175–195.
- [38] W. G. Unruh, “Notes on black-hole evaporation,” *Phys. Rev. D* **14** (1976), 870–892.
- [39] K. van den Dungen, “Krein spectral triples and the fermionic action,” *Math. Phys. Anal. Geom.* **19** (2016), 4. arXiv:1505.01939.
- [40] J. Loutey, “The Bargmann–Segal Lattice: π -Free Discretization of the Harmonic Oscillator on S^5 ,” GeoVac internal preprint Paper 24 (2026), at `papers/group3_foundations/paper_24_bargmann_segal.tex`.
- [41] J. Loutey, “Finite-Size Graph-RH for the GeoVac Hopf Graphs: Ihara Zeta, Bound-Crossing, and a Scope Boundary on Selberg-on-Hydrogen,” GeoVac internal preprint Paper 29 (2026), at `papers/group1_operator_algebras/paper_29_ramanujan_hopf.tex`.
- [42] J. Loutey, “The GeoVac Spectral Triple: A Synthesis Paper Locking the Framework into the Marcolli–van Suijlekom Gauge-Network Lineage,” GeoVac internal preprint Paper 32 (2026), at `papers/group1_operator_algebras/paper_32_spectral_triple.tex`.
- [43] J. Loutey, “State-space Gromov–Hausdorff convergence of truncated Camporesi–Higuchi spectral triples on S^3 ,” GeoVac internal preprint Paper 38 (2026), released to Zenodo 2026-05-07. At `papers/group1_operator_algebras/paper_38_su2_propinquity_convergence.tex`.
- [44] J. Loutey, “Tensor-product propinquity convergence on $\mathcal{T}_{S^3}^{\lambda_a} \otimes \mathcal{T}_{S^3}^{\lambda_b}$,” GeoVac internal preprint Paper 39 (2026), at `papers/group1_operator_algebras/paper_39_tensor_propinquity_convergence.tex`.

- [45] J. Loutey, “State-space Gromov–Hausdorff convergence of spectral truncations of compact semisimple Lie groups with bi-invariant metric,” GeoVac internal preprint Paper 40 (2026), at `papers/group1_operator_algebras/paper_40_unified_propinquity_convergence.tex`.
- [46] J. Loutey, “Tomita–Takesaki modular structure on truncated $SU(2)$ spectral triples: four-witness Wick-rotation literal identification at finite cutoff,” GeoVac internal preprint Paper 42 (2026), at `papers/group1_operator_algebras/paper_42_modular_hamiltonian_four_witness.tex`.
- [47] J. Loutey, “Lorentzian extension of the four-witness Wick-rotation theorem on truncated $S^3 \times \mathbb{R}$ spectral triples at finite cutoff,” GeoVac internal preprint Paper 43 (2026), at `papers/group1_operator_algebras/paper_43_lorentzian_extension.tex`.
- [48] J. Loutey, “Operator-system extension of the BBB Krein spectral triple at finite cutoff: propagation number and envelope dependence on the Camporesi–Higuchi spinor bundle,” GeoVac internal preprint Paper 44 (2026), at `papers/group1_operator_algebras/paper_44_lorentzian_operator_system.tex`.
- [49] J. Loutey, “Lorentzian propinquity on truncated $SU(2) \times U(1)_T$ Krein spectral triples: a degeneracy theorem,” GeoVac internal preprint Paper 45 (2026), at `papers/group1_operator_algebras/paper_45_lorentzian_propinquity.tex`.
- [50] J. Loutey, “Strong-form Lorentzian propinquity on truncated $SU(2) \times U(1)_T$ Krein spectral triples,” GeoVac internal preprint Paper 46 (2026), at `papers/group1_operator_algebras/paper_46_strong_form_lorentzian_propinquity.tex`.
- [51] J. Loutey, “Norm-resolvent convergence to the non-compact Lorentzian Krein spectral triple on $S^3 \times \mathbb{R}_t$ and the two-rate hybrid limit,” GeoVac internal preprint Paper 47 (2026), at `papers/group1_operator_algebras/paper_47_two_rate_hybrid_convergence.tex`.