

Norm-resolvent convergence to the non-compact Lorentzian Krein spectral triple on $S^3 \times \mathbb{R}_t$ and the two-rate hybrid limit

J. Loutey*

May 2026 (preprint)

Abstract

We close the de-compactification gap (G2) of the Lorentzian propinquity program [27, 28] at the norm-resolvent / spectral level. Combining the parametric stability of Paper 45 [27] (and its strong-form extension Paper 46 [28]) with a standard compact-to-non-compact resolvent comparison, we establish a two-rate hybrid convergence

$$\mathcal{T}_{n_{\max}, N_t, T(N_t)}^L \xrightarrow{\Lambda^L\text{-rate } O(\log n_{\max}/n_{\max} + T(N_t)/N_t)} \mathcal{T}_{S^3 \times S_T^1(N_t)}^L \xrightarrow{\text{nr-rate } O(e^{-|\text{Im } z| T(N_t)/2})} \mathcal{T}_{S^3 \times \mathbb{R}_t}^L$$

along any admissible scaling $T(N_t) \rightarrow \infty$ with $T(N_t)/N_t \rightarrow 0$. The inner arrow is propinquity convergence (Paper 45 main theorem verbatim, at coupled cells); the outer arrow is norm-resolvent convergence of the Lorentzian Dirac on the compactified spacetime $S^3 \times S_T^1$ to the Lorentzian Dirac on the genuine spacetime $S^3 \times \mathbb{R}_t$, with an explicit exponential tail bound $O(e^{-|\text{Im } z| T/2})$ on compact-temporal-support subspaces. The construction preserves the BBB [1] Krein fundamental symmetry, the Krein-positive cone $\mathcal{K}^+$, and the Krein-self-adjointness of the Lorentzian Dirac across the entire two-rate diagram.

Honest scope. The outer arrow is at the norm-resolvent / spectral level, not at the propinquity / metric level. Propinquity-level convergence to the genuine non-compact Lorentzian Krein triple would require a non-compact extension of Latrémolière’s propinquity [13, 14], which is not in the published literature as of May 2026. We do not undertake this extension here; G2 closure at the metric level remains an open NCG-math problem.

Three-carrier identification. The norm-resolvent limit identifies three nominally distinct Krein-Lorentzian carriers as approximation schemes for the same physical operator: the periodic compact $S^3 \times S_T^1$ (Paper 45 substrate), the bounded Dirichlet interval $S^3 \times [-T_{\max}, T_{\max}]$ (Paper 43 [25] substrate), and the non-compact $S^3 \times \mathbb{R}_t$ (this paper’s limit object). All three converge to the same Lorentzian Dirac as the temporal scale grows.

Place in the math.OA standalone series. Ninth in the GeoVac math.OA arc (siblings: Papers 38 [21], 39 [22], 40 [23], 42 [24], 43 [25], 44 [26], 45 [27], 46 [28]). The result composes Paper 45/46 verbatim on the inner arrow with standard Reed–Simon [17] § XIII.16 / Hekkelman–McDonald 2024 [10] compact-to-non-compact-domain machinery on the outer arrow.

Contents

1	Introduction	2
1.1	The strategic reframing	3
1.2	Two-rate convergence: what it is and what it is not	3
1.3	Outline	4

*Independent researcher. jloutey@gmail.com. This work was developed in an AI-augmented agentic workflow with Anthropic’s Claude. Implementation, internal memos, and bibliographic collation were performed collaboratively; all theorems, design choices, and editorial decisions are the author’s responsibility.

2	Setup	4
2.1	Krein spaces at finite, bounded, and non-compact temporal carrier	4
2.2	Isometric embeddings	4
2.3	The Lorentzian Dirac at each carrier	5
2.4	Admissible scaling and parametric setup	6
3	The inner arrow: parametric propinquity stability	6
4	The outer arrow: norm-resolvent convergence	7
4.1	Definition	7
4.2	Main theorem (outer arrow)	7
4.3	Proof of Theorem 4.2	8
4.4	Extension to the dense compact-support subspace	9
5	The two-rate hybrid theorem	10
5.1	Choice of scaling	10
5.2	Empirical confirmation: numerical panel along admissible scalings	11
6	Three-carrier identification	11
6.1	Three carriers, one limit	11
6.2	Why both Paper 43 and Paper 45 exist	12
7	Discussion and related work	13
7.1	Position vs Mondino–Sämman synthetic Lorentzian GH	13
7.2	Position vs Hekkelman–McDonald 2024	13
7.3	Position vs Latrémolière	13
7.4	The two-rate hybrid framing in NCG	13
8	Open questions	14

1 Introduction

The Lorentzian propinquity program of Papers 45 [27]–46 [28] establishes that the finite-cutoff Lorentzian Krein spectral triple $\mathcal{T}_{n_{\max}, N_t, T}^L$ on the compactified spacetime $S^3 \times S_T^1$ converges, in the Latrémolière quantum Gromov–Hausdorff propinquity [13, 14], to the continuum compact-temporal Lorentzian Krein spectral triple $\mathcal{T}_{S^3 \times S_T^1}^L$ as $(n_{\max}, N_t) \rightarrow (\infty, \infty)$ at fixed temporal radius T . The natural canonical choice is the Bisognano–Wichmann modular period $T = 2\pi$, in which case the construction realizes the modular-flow sector of the wedge KMS state.

The construction at fixed $T = 2\pi$ leaves four explicitly named gaps toward the full Krein-signature theory (Paper 45 § 1.4):

- (G1) strong-form Lorentzian propinquity without $\mathcal{K}^+$ -restriction (closed on the natural substrate in Paper 46 [28]; strict-strong-form on the truly-enlarged substrate remains open),
- (G2) the *de-compactification limit* $T \rightarrow \infty$, recovering the canonical Wightman-style time axis $\mathbb{R}_t$,
- (G3) cross-manifold extensions $\mathcal{T}_{S^3} \otimes \mathcal{T}_{\text{Hardy}(S^5)}$ (blocked by Paper 24 [19] § V four-layer Coulomb/HO asymmetry), and
- (G4) inner-factor Yukawa calibration data (orthogonal to the convergence theorem).

This paper closes G2 at a precisely identified level. We establish a *two-rate hybrid convergence statement* of the form

$$\mathcal{T}_{n_{\max}, N_t, T(N_t)}^L \xrightarrow[\text{Paper 45/46}]{\Lambda^L} \mathcal{T}_{S^3 \times S_T^1(N_t)}^L \xrightarrow[\text{this paper}]{\text{nr}} \mathcal{T}_{S^3 \times \mathbb{R}_t}^L, \quad (1)$$

along any admissible scaling

$$T(N_t) \rightarrow \infty \quad \text{and} \quad T(N_t)/N_t \rightarrow 0 \quad \text{as } N_t \rightarrow \infty. \quad (2)$$

The inner arrow is propinquity convergence at coupled cells (a parametric reformulation of Paper 45 / Paper 46 main theorems). The outer arrow is *norm-resolvent convergence* in the standard sense of Reed–Simon [17] Vol. IV § XIII.16, with an explicit exponential tail estimate.

1.1 The strategic reframing

A natural first attempt at closing G2 would be the *inductive-limit propinquity* framework of Latrémolière [14] § 5 for AF C*-algebras. The truncated Lorentzian Krein triples $\{\mathcal{T}_{n_{\max}, N_t, T_n}^L\}$ are finite-dimensional, so the system is AF, and refinement maps $\mathcal{T}_{S^3 \times S_{T_n}^1}^L \hookrightarrow \mathcal{T}_{S^3 \times S_{T_{n+1}}^1}^L$ exist when $T_n \mid T_{n+1}$.

This approach is *structurally wrong* for the present setting: AF C*-algebras have totally-disconnected Gelfand spectrum, while $\mathbb{R}$ is connected, so $C_0(\mathbb{R})$ cannot be an AF inductive limit. The AF inductive limit of $\{C(S_{T_n}^1)\}$ along $T_n = 2^n T_0$ refinement chains is a Bunce–Deddens-style algebra [2], not $C_0(\mathbb{R})$. Therefore the inductive-limit propinquity machinery cannot reach the genuine non-compact target.

The right tool is norm-resolvent convergence of spectral triples [4] § VI: a notion of convergence that captures spectral data and operator-algebraic structure (including self-adjointness, fundamental symmetry, and the C*-algebra of bounded functions of D_L) but does *not* capture the Lipschitz/Wasserstein metric structure encoded by the propinquity. For G2’s *physical content* (“the Lorentzian Dirac on the compactified spacetime converges to the Lorentzian Dirac on the genuine spacetime”), norm-resolvent is the right framework. For G2’s *metric content* (Latrémolière-distance convergence on a non-compact carrier), the framework would need to be extended; this is not addressed here.

1.2 Two-rate convergence: what it is and what it is not

The composite (1) is a *hybrid* convergence statement: two different convergence notions on two different arrows, composed into a joint statement about the limiting behavior. It is not a single propinquity bound; the two rates do not combine into a uniform Latrémolière distance. Rather:

- For *any fixed* compactly-supported smooth test data on $S^3 \times \mathbb{R}_t$, the composite (1) gives strong convergence of expectations $\langle \xi, F(D_{L; n_{\max}, N_t, T(N_t)}) \xi \rangle \rightarrow \langle \xi, F(D_{L, \mathbb{R}}) \xi \rangle$ for every continuous $F : \mathbb{R} \rightarrow \mathbb{C}$ vanishing at infinity, and every ξ with compact temporal support.
- The convergence rate is exponential in T on the outer arrow ($O(e^{-|\text{Im } z| T/2})$ for the resolvent) and propinquity-rate on the inner arrow ($O(\log n_{\max}/n_{\max} + T(N_t)/N_t)$).
- Choosing an admissible scaling $T(N_t)$ trades the two rates: sub-linear scaling $T(N_t) = N_t / \log N_t$ gives faster de-compactification but slower inner rate; square-root scaling $T(N_t) = \sqrt{N_t}$ gives slower de-compactification but faster inner rate. The choice is a convention question, not a structural one.

The composite (1) is the appropriate G2 closure at the level GeoVac currently supports. It identifies the sequence of finite-cutoff compact-temporal Lorentzian Krein triples with the

canonical non-compact Lorentzian Dirac as the limit object, preserves the Krein structure across the entire diagram, and supplies explicit, computable convergence rates on both arrows. Metric-level closure requires non-compact Latrémolière propinquity, which is a multi-month NCG-math problem and not addressed here (§ 8).

1.3 Outline

Section 2 sets up the Krein-space construction, the isometric embeddings $\tilde{J}_T : \mathcal{K}_T \hookrightarrow \mathcal{K}_{\mathbb{R}}$ that allow comparing operators on different temporal carriers, and the Lorentzian Dirac on $S^3 \times \mathbb{R}_t$. Section 3 records the inner arrow (Paper 45/46 propinquity at coupled cells; we state this as Theorem 3.1 but the proof is Paper 45 main theorem verbatim, included for completeness). Section 4 proves the outer arrow (norm-resolvent convergence of $D_{L,T} \rightarrow D_{L,\mathbb{R}}$ as $T \rightarrow \infty$); the proof uses standard compact-to-non-compact-domain machinery (Reed–Simon [17] § XIII.16) plus a Krein-structure preservation argument specific to the BBB construction. Section 5 states and proves the two-rate hybrid convergence theorem (the main result, Theorem 5.1). Section 6 establishes the three-carrier identification: the periodic compact, bounded Dirichlet, and non-compact non-periodic carriers all converge to the same Lorentzian Dirac in norm-resolvent sense. Section 7 positions the result against the Mondino–Sämman synthetic Lorentzian GH program [15, 16], Hekkelman–McDonald 2024 [10], and the Strohmaier [18] / Bizi–Brouder–Besnard 2018 [1] Lorentzian-NCG literature. Section 8 records the remaining open questions (non-compact metric-level closure G1-genuine, cross-manifold G3, and inner-factor G4).

2 Setup

2.1 Krein spaces at finite, bounded, and non-compact temporal carrier

We work in the BBB chiral basis at KO-dimension $(m, n) = (4, 6)$, following Paper 45 [27] § 3 and the Sprint L2-B construction of Paper 43 [25]. The Camporesi–Higuchi spinor space on S^3 at cutoff $n_{\max}$ is $\mathcal{H}_{\text{GV}}^{n_{\max}}$ of dimension $\frac{2}{3}n_{\max}(n_{\max} + 1)(n_{\max} + 2)$.

Definition 2.1 (Krein spaces at three temporal carriers). Let $n_{\max} \in \mathbb{N}$, $N_t \in \mathbb{N}$, and $T > 0$. Define:

- (i) (*Compact periodic, finite-cutoff.*) $\mathcal{K}_{n_{\max}, N_t, T} := \mathcal{H}_{\text{GV}}^{n_{\max}} \otimes \mathbb{C}^{N_t}$, identifying $\mathbb{C}^{N_t}$ with the finite-Fourier truncation of $L^2(S_T^1)$ at modes $\omega_k = 2\pi k/T$, $k = -K_{\max}, \dots, +K_{\max}$ with $N_t = 2K_{\max} + 1$. Fundamental symmetry $J_L = J_{\text{spat}} \otimes I_{N_t}$, with J_{spat} the spatial BBB Krein structure on $\mathcal{H}_{\text{GV}}^{n_{\max}}$ (Paper 44 § 2).
- (ii) (*Compact periodic, continuum.*) $\mathcal{K}_T := \mathcal{H}_{\text{GV}} \otimes L^2(S_T^1)$, with $\mathcal{H}_{\text{GV}} = \varinjlim_{n_{\max}} \mathcal{H}_{\text{GV}}^{n_{\max}}$ the full Camporesi–Higuchi spinor Hilbert space on S^3 . Same $J_L = J_{\text{spat}} \otimes I$.
- (iii) (*Non-compact.*) $\mathcal{K}_{\mathbb{R}} := \mathcal{H}_{\text{GV}} \otimes L^2(\mathbb{R})$. Same $J_L = J_{\text{spat}} \otimes I$.

Each $\mathcal{K}_{\bullet}$ is a Krein space: a Hilbert space with an indefinite inner product $\langle \cdot, J_L \cdot \rangle$, decomposing as $\mathcal{K}_{\bullet} = \mathcal{K}_{\bullet}^+ \oplus \mathcal{K}_{\bullet}^-$ via the eigenspaces of J_L . The eigenspace projections $P_{\pm}$ are trace-class-finite at every $n_{\max}, N_t$ (item (i)) and projections in $\mathcal{B}(\mathcal{K}_{\bullet})$ (items (ii), (iii)).

2.2 Isometric embeddings

The natural identification $L^2(S_T^1) \cong L^2([-T/2, T/2])$ (square-integrable functions on the fundamental domain $[-T/2, T/2] \subset \mathbb{R}$ extended periodically) gives an isometric embedding into $L^2(\mathbb{R})$ by zero-extension:

$$J_T : L^2(S_T^1) \rightarrow L^2(\mathbb{R}), \quad (J_T f)(t) := \begin{cases} f(t) & t \in [-T/2, T/2], \\ 0 & |t| > T/2. \end{cases} \quad (3)$$

Proposition 2.2. *The map J_T is an isometry: $\|J_T f\|_{L^2(\mathbb{R})} = \|f\|_{L^2(S_T^1)}$ for every $f \in L^2(S_T^1)$. Its image is $\text{Im } J_T = L^2([-T/2, T/2]) \subset L^2(\mathbb{R})$, and the adjoint $J_T^* : L^2(\mathbb{R}) \rightarrow L^2(S_T^1)$ acts as restriction followed by periodic identification: $(J_T^* g)(t) = g(t)$ for $t \in [-T/2, T/2]$, extended periodically to all of S_T^1 .*

Proof. The fundamental-domain identification preserves L^2 -norm by definition, and zero-extension into $L^2(\mathbb{R})$ also preserves L^2 -norm. The image and adjoint claims are routine. $\square$

Extend J_T to the Krein spaces via $\tilde{J}_T := I_{\mathcal{H}_{\text{GV}}} \otimes J_T$: $\tilde{J}_T : \mathcal{K}_T \hookrightarrow \mathcal{K}_{\mathbb{R}}$. Restricting to the finite-cutoff substrate gives $\tilde{J}_T^{n_{\text{max}}, N_t} : \mathcal{K}_{n_{\text{max}}, N_t, T} \hookrightarrow \mathcal{K}_{\mathbb{R}}$ via the obvious tensor-product structure.

Lemma 2.3 (Krein-isometric embedding). *The embeddings $\tilde{J}_T$ and $\tilde{J}_T^{n_{\text{max}}, N_t}$ are Krein-isometric:*

$$\tilde{J}_T^* J_{L, \mathbb{R}} \tilde{J}_T = J_{L, T}, \quad (\tilde{J}_T^{n_{\text{max}}, N_t})^* J_{L, \mathbb{R}} \tilde{J}_T^{n_{\text{max}}, N_t} = J_{L, T}^{n_{\text{max}}, N_t}. \quad (4)$$

Proof. Both J_L act trivially on the temporal factor ($J_L = J_{\text{spat}} \otimes I_{\text{temp}}$), and $\tilde{J}_T$ is identity on the spatial factor. Equation (4) reduces to $J_T^*(I_{\mathbb{R}})J_T = J_T^*J_T = I_{S_T^1}$ on the temporal factor, which holds because J_T is isometric. $\square$

Corollary 2.4 ($\mathcal{K}^+$ -cone preservation). *$\tilde{J}_T(\mathcal{K}_T^+) \subset \mathcal{K}_{\mathbb{R}}^+$. Equivalently, $\tilde{J}_T$ commutes with the $\mathcal{K}^+$ -projection P_+ on its image.*

2.3 The Lorentzian Dirac at each carrier

Following Paper 45 [27] § 3 (after Paper 43 § 3 and the Sprint L2-C construction), the Lorentzian Dirac operator on each Krein space is

$$D_{L, T}^{n_{\text{max}}, N_t} := i(\gamma^0 \otimes \partial_t^{S_T^1, N_t} + D_{\text{GV}}^{n_{\text{max}}} \otimes I_{N_t}), \quad (5)$$

$$D_{L, T} := i(\gamma^0 \otimes \partial_t^{S_T^1} + D_{\text{GV}} \otimes I), \quad (6)$$

$$D_{L, \mathbb{R}} := i(\gamma^0 \otimes \partial_t^{\mathbb{R}} + D_{\text{GV}} \otimes I). \quad (7)$$

The temporal derivative is in each case:

- $\partial_t^{S_T^1, N_t}$: anti-Hermitian, diagonal in the finite-Fourier basis with eigenvalues $i\omega_k$, $\omega_k = 2\pi k/T$, $k = -K_{\text{max}}, \dots, +K_{\text{max}}$;
- $\partial_t^{S_T^1}$: essentially anti-self-adjoint on $C^\infty(S_T^1)$, closure has discrete spectrum $\{i\omega_k : k \in \mathbb{Z}\}$;
- $\partial_t^{\mathbb{R}}$: essentially anti-self-adjoint on $C_c^\infty(\mathbb{R})$, closure has continuous spectrum $i\mathbb{R}$ with absolutely continuous spectral measure (the standard Fourier transform).

Lemma 2.5 (Krein-self-adjointness). *Each operator in (5)–(7) is Krein-self-adjoint with respect to its J_L :*

$$J_{L, \bullet} (D_{L, \bullet})^* J_{L, \bullet}^{-1} = D_{L, \bullet} \quad (\text{formally, on the appropriate dense domain}).$$

Proof. The Sprint L2-C derivation (Paper 43 § 3) of $D_{L, T}^{n_{\text{max}}, N_t}$ as Krein-self-adjoint via $\{\gamma^0, D_{\text{GV}}\} = 0$ and $\partial_t^* = -\partial_t$ transports verbatim to the continuum case (6) (the spectrum is discrete, the algebraic identity is preserved) and to the non-compact case (7) (the spectrum is continuous, the algebraic identity is preserved on the core $C_c^\infty(\mathbb{R}) \otimes \mathcal{H}_{\text{GV}}$). $\square$

2.4 Admissible scaling and parametric setup

Definition 2.6 (Admissible scaling). A function $T : \mathbb{N} \rightarrow \mathbb{R}_{>0}$ is an *admissible scaling* if

$$T(N_t) \rightarrow \infty \quad \text{and} \quad \frac{T(N_t)}{N_t} \rightarrow 0 \quad \text{as } N_t \rightarrow \infty. \quad (8)$$

Example 2.7. Canonical admissible scalings:

- Sub-linear logarithmic: $T(N_t) = N_t / \log N_t$.
- Square root: $T(N_t) = \sqrt{N_t}$.
- General power-law: $T(N_t) = N_t^{1-\epsilon}$ for any $\epsilon \in (0, 1)$.

All three are admissible; the choice trades de-compactification rate against inner-propinquity decay rate (§ 5.1).

3 The inner arrow: parametric propinquity stability

We restate the inner arrow of (1) for completeness. The proof is Paper 45 [27] / Paper 46 [28] main theorem applied at coupled cells; the only content here is verifying that the application is uniform along an admissible scaling sequence.

Theorem 3.1 (Inner arrow). *Let $T(\cdot)$ be any admissible scaling (Definition 2.6). On the natural substrate of Paper 45 / Paper 46 main theorem,*

$$\Lambda^L(\mathcal{T}_{n_{\max}, N_t, T(N_t)}^L, \mathcal{T}_{S^3 \times S_{T(N_t)}^1}^L) \leq C_3^{\text{joint}}(n_{\max}) \gamma^{\text{joint}}(n_{\max}, N_t, T(N_t)), \quad (9)$$

with $C_3^{\text{joint}}(n_{\max}) = \sqrt{1 - 1/n_{\max}} \rightarrow 1^-$ (Paper 46 § 4.3 envelope-aware form, T - and N_t -independent) and

$$\gamma^{\text{joint}}(n_{\max}, N_t, T(N_t)) = \max(O(\log n_{\max}/n_{\max}), O(T(N_t)/N_t)) \rightarrow 0 \quad \text{as } (n_{\max}, N_t) \rightarrow (\infty, \infty). \quad (10)$$

The strong-form analog (Paper 46 main theorem, natural chirality-doubled scalar substrate) gives the same bound bit-exact via the “free upgrade” reading.

Proof. At every cell $(n_{\max}, N_t)$, the temporal radius $T(N_t)$ is a positive real number, and the Paper 45/46 main theorem applies as a black-box bound at that cell. The five lemmas L1' / L2 / L3 / L4 / L5 transport without modification:

- **L1' (Paper 44):** operator-system substrate, structurally independent of T (the temporal multiplier algebra is the commutative diagonal subalgebra of $M_{N_t}(\mathbb{C})$, with the specific Fourier momenta $\omega_k = 2\pi k/T$ appearing as numerical entries but not affecting the algebraic structure).
- **L2 (cb-norm, Paper 45 § 5):** $\|S_{K^{\text{joint}}}\|_{\text{cb}} = 2/(n_{\max} + 1)$, depending only on $n_{\max}$ via Bożejko–Fendler on the amenable compact product group $\text{SU}(2) \times U(1)_T$. The $U(1)_T$ factor contributes cb-norm 1 at every T .
- **L3 (Lichnerowicz, Paper 45 § 4):** structural identity $[D_L, a_s \otimes a_t] = i[D_{\text{GV}}, a_s] \otimes a_t$ bit-exact for every $(n_{\max}, N_t, T)$, because $[\gamma^0, a_s] = 0$ on the natural substrate and $[\partial_t, a_t] = 0$ for momentum-diagonal a_t . The constant $C_3^{\text{joint}}(n_{\max}, N_t) \leq \sqrt{1 - 1/n_{\max}}$ is T - and N_t -independent.

- **L4 (Berezin, Paper 45 § 5):** the joint Berezin map $B^{\text{joint}} = B^{\text{SU}(2)} \otimes B^{U(1)T}$ has approximate identity with rate $\gamma^{\text{joint}} = \max(O(\log n_{\text{max}}/n_{\text{max}}), O(T/N_t))$. The temporal-factor rate $O(T/N_t)$ is the standard Fejér-on-circle rate from Katznelson [12] Ch. I, with T -independent implicit constant. Substituting $T = T(N_t)$ gives $\gamma^{\text{joint}}(n_{\text{max}}, N_t, T(N_t)) = \max(O(\log n_{\text{max}}/n_{\text{max}}), O(T(N_t)/N_t))$, which vanishes by the admissibility condition $T(N_t)/N_t \rightarrow 0$.
- **L5 (propinquity assembly, Paper 45 § 6):** the tunneling-pair construction $(B^{\text{joint}}, P^{\text{joint}})$ and the reach/height bookkeeping produce (9) verbatim at each cell.

□

Remark 3.2 (The natural substrate is necessary). The inner arrow (9) holds on the *natural* substrate (chirality-doubled scalar spatial multipliers) of Paper 45 / Paper 46 main theorem. On the *enlarged* substrate of Paper 46 Appendix B (chirality-flipping generators with $\{J_L, M^{\text{flip}}\} = 0$), the gradient norm picks up a chirality-flip time-piece term $2\|\partial_t\|_{\text{op}}\|a^{\text{flip}}\|_{\text{op}}$ which *does not* survive the joint limit $T \rightarrow \infty$: the $O(N_t - 1)$ scaling of $\|\partial_t\|_{\text{op}}$ is paid by the gradient itself, not absorbed by the rate. Therefore the inner arrow’s T -uniformity is structurally tied to the natural substrate; the enlarged substrate has a separable open question (“L3c- β ” in our internal taxonomy).

4 The outer arrow: norm-resolvent convergence

The outer arrow of (1) is norm-resolvent convergence in the standard sense of Reed–Simon [17] Vol. IV § XIII.16, modulo the isometric embedding $\tilde{J}_T : \mathcal{K}_T \hookrightarrow \mathcal{K}_{\mathbb{R}}$.

4.1 Definition

Definition 4.1 (Norm-resolvent convergence modulo isometry). Let $\{A_T\}_{T>0}$ be a family of Krein-self-adjoint operators on $\mathcal{K}_T$, and $A_{\mathbb{R}}$ a Krein-self-adjoint operator on $\mathcal{K}_{\mathbb{R}}$, with isometric embeddings $\tilde{J}_T : \mathcal{K}_T \hookrightarrow \mathcal{K}_{\mathbb{R}}$. We say $A_T \rightarrow A_{\mathbb{R}}$ in *norm-resolvent sense modulo $\tilde{J}_T$* as $T \rightarrow \infty$ if, for every $z \in \mathbb{C} \setminus \mathbb{R}$,

$$\|\tilde{J}_T (A_T - z)^{-1} \tilde{J}_T^* - P_T (A_{\mathbb{R}} - z)^{-1} P_T\|_{\mathcal{B}(\mathcal{K}_{\mathbb{R}})} \rightarrow 0, \quad (11)$$

where $P_T = \tilde{J}_T \tilde{J}_T^*$ is the projection onto the image $\text{Im } \tilde{J}_T \subset \mathcal{K}_{\mathbb{R}}$.

This is the standard “compact-to-non-compact-domain” norm-resolvent convergence (e.g., Reed–Simon § XIII.16, Theorem XIII.86): one compares the resolvent on the smaller carrier (after embedding into the larger) with the resolvent on the larger carrier (after projecting onto the embedded image). The two should agree up to a T -dependent error that vanishes as $T \rightarrow \infty$.

4.2 Main theorem (outer arrow)

Theorem 4.2 (Outer arrow). *The Lorentzian Dirac operators $D_{L,T} \rightarrow D_{L,\mathbb{R}}$ in norm-resolvent sense modulo $\tilde{J}_T$ as $T \rightarrow \infty$. Explicitly, for every $z \in \mathbb{C} \setminus \mathbb{R}$ there exists a constant $C(z) > 0$ such that*

$$\|\tilde{J}_T (D_{L,T} - z)^{-1} \tilde{J}_T^* \xi - (D_{L,\mathbb{R}} - z)^{-1} \xi\|_{\mathcal{K}_{\mathbb{R}}} \leq C(z, \xi) e^{-|\text{Im } z| T/2} \quad (12)$$

for every $\xi \in \mathcal{K}_{\mathbb{R}}$ of compact temporal support $\text{supp}_t(\xi) \subset [-A, A]$ with $A < T/2$, where $C(z, \xi)$ depends only on $\|\xi\|_{\mathcal{K}_{\mathbb{R}}}$, $\text{Im } z$, and A .

The constant $C(z, \xi)$ can be tracked explicitly through the proof; in particular, it is bounded above by $C(z, \xi) \leq c_0 \|\xi\| (1 + |z|)^{-1} |\text{Im } z|^{-2}$ for a universal constant c_0 .

4.3 Proof of Theorem 4.2

The proof has four steps.

Step 1: Spatial factorization reduces to temporal-only convergence. The Lorentzian Dirac decomposes as

$$D_L = A + B, \quad A := i\gamma^0 \otimes \partial_t, \quad B := i D_{\text{GV}} \otimes I,$$

where the spatial term B acts identically on $\mathcal{K}_T$ and $\mathcal{K}_{\mathbb{R}}$ (both have the same $\mathcal{H}_{\text{GV}}$ spatial factor; the difference is only in the temporal Hilbert space) and is bounded by $\|B\|_{\text{op}} \leq \|D_{\text{GV}}\|_{\text{op}} =: M_{\text{spat}}$, finite at each n_{max} (but unbounded as $n_{\text{max}} \rightarrow \infty$; for the outer arrow at fixed n_{max} , the bound is finite). The temporal term A is unbounded on both $\mathcal{K}_T$ and $\mathcal{K}_{\mathbb{R}}$ but bounded relative to B : in fact, $\|A(B+i)^{-1}\|_{\text{op}}$ is uniformly bounded in T on the dense compact-support subspace.

By the standard sum-of-resolvents argument (Reed–Simon § VIII.7), norm-resolvent convergence $A_T \rightarrow A_{\mathbb{R}}$ together with B uniformly bounded implies norm-resolvent convergence $D_{L,T} = A_T + B \rightarrow A_{\mathbb{R}} + B = D_{L,\mathbb{R}}$, with the same rate.

It therefore suffices to establish norm-resolvent convergence $\partial_t^{S_T^1} \rightarrow \partial_t^{\mathbb{R}}$ as $T \rightarrow \infty$.

Step 2: Temporal resolvent kernel on $\mathbb{R}$. The operator $\partial_t^{\mathbb{R}}$ on $L^2(\mathbb{R})$ is essentially anti-self-adjoint on $C_c^\infty(\mathbb{R})$, with closure having continuous spectrum $i\mathbb{R}$. For $z \in \mathbb{C} \setminus \mathbb{R}$, the resolvent $(i\partial_t^{\mathbb{R}} - z)^{-1}$ acts as convolution with the integral kernel

$$K_z^{\mathbb{R}}(t) = -i \mathbf{1}_{\text{sgn Im } z}(t) e^{-izt}, \quad (13)$$

where $\mathbf{1}_+(t) = \mathbf{1}_{[0,\infty)}(t)$ if $\text{Im } z > 0$ and $\mathbf{1}_-(t) = -\mathbf{1}_{(-\infty,0]}(t)$ if $\text{Im } z < 0$. The kernel has exponential decay in $|t|$:

$$|K_z^{\mathbb{R}}(t)| \leq e^{-|\text{Im } z| |t|}.$$

Step 3: Temporal resolvent kernel on S_T^1 . The operator $\partial_t^{S_T^1}$ on $L^2(S_T^1)$ has discrete spectrum $\{i\omega_k : k \in \mathbb{Z}\}$, $\omega_k = 2\pi k/T$. For $z \in \mathbb{C} \setminus \mathbb{R}$ with $\text{Im } z > 0$, the resolvent acts as convolution (on S_T^1) with the periodic kernel

$$K_z^{S_T^1}(t) = \sum_{n \in \mathbb{Z}} K_z^{\mathbb{R}}(t + nT). \quad (14)$$

The sum converges absolutely because $K_z^{\mathbb{R}}$ has exponential decay: $|K_z^{S_T^1}(t)| \leq \sum_{n \in \mathbb{Z}} e^{-|\text{Im } z| |t+nT|} \leq e^{-|\text{Im } z| \text{dist}(t, \mathbb{Z}T)} (1 - e^{-|\text{Im } z| T})^{-1}$.

Equation (14) is the standard *Poisson-summation form* of the periodic resolvent: the periodic Green's function on S_T^1 is the sum of translates of the free Green's function on $\mathbb{R}$. This is a classical identity (Bloch decomposition; see Reed–Simon § XIII.16, Example 4).

Step 4: Embedded comparison. Fix $\xi \in \mathcal{K}_{\mathbb{R}}$ with compact temporal support $\text{supp}_t(\xi) \subset [-A, A]$, and let $T > 2A$. Then $\tilde{J}_T^* \xi \in \mathcal{K}_T$ is well-defined (the periodic identification of ξ restricted to $[-T/2, T/2]$ is just ξ itself, with the periodic structure trivial because $\text{supp}_t \xi \subset [-A, A] \subsetneq [-T/2, T/2]$).

Compute, for $\text{Im } z > 0$:

$$\begin{aligned} [(\partial_t^{\mathbb{R}} - z)^{-1} \xi](t) &= \int_{\mathbb{R}} K_z^{\mathbb{R}}(t-s) \xi(s) ds, \\ [\tilde{J}_T(\partial_t^{S_T^1} - z)^{-1} \tilde{J}_T^* \xi](t) &= \mathbf{1}_{[-T/2, T/2]}(t) \int_{[-T/2, T/2]} K_z^{S_T^1}(t-s) \xi(s) ds \\ &= \mathbf{1}_{[-T/2, T/2]}(t) \sum_{n \in \mathbb{Z}} \int_{[-T/2, T/2]} K_z^{\mathbb{R}}(t-s+nT) \xi(s) ds. \end{aligned}$$

The $n = 0$ term reproduces the $\mathbb{R}$ -resolvent restricted to $[-T/2, T/2] \times [-T/2, T/2]$:

$$\begin{aligned} \mathbf{1}_{[-T/2, T/2]}(t) \int_{[-T/2, T/2]} K_z^{\mathbb{R}}(t-s) \xi(s) ds &= \mathbf{1}_{[-T/2, T/2]}(t) \int_{\mathbb{R}} K_z^{\mathbb{R}}(t-s) \xi(s) ds \quad (\text{since } \text{supp } \xi \subset [-A, A]) \\ &= \mathbf{1}_{[-T/2, T/2]}(t) [(\partial_t^{\mathbb{R}} - z)^{-1} \xi](t). \end{aligned}$$

The $n \neq 0$ terms in the sum (14) give the error:

$$E_T(t) := \mathbf{1}_{[-T/2, T/2]}(t) \sum_{n \neq 0} \int_{[-A, A]} K_z^{\mathbb{R}}(t-s+nT) \xi(s) ds. \quad (15)$$

For $t \in [-T/2, T/2]$ and $s \in [-A, A]$ with $T > 2A$, we have $|t-s+nT| \geq |nT| - |t| - |s| \geq |n|T - T/2 - A \geq (|n|-1)T/2$ for $|n| \geq 1$. Therefore

$$|K_z^{\mathbb{R}}(t-s+nT)| \leq e^{-|\text{Im } z|(|n|-1)T/2}.$$

The remaining tail (for $|t| > T/2$) vanishes because of the $\mathbf{1}_{[-T/2, T/2]}$ factor; the remaining tail of the $\mathbb{R}$ -resolvent (for $|t| > T/2$) decays as $|(\partial_t^{\mathbb{R}} - z)^{-1} \xi(t)| \leq e^{-|\text{Im } z|(|t|-A)} \|\xi\|_{L^2}$ by the exponential decay of $K_z^{\mathbb{R}}$.

Combining: for $T > 2A$,

$$\begin{aligned} \|\tilde{J}_T(\partial_t^{S_T^1} - z)^{-1} \tilde{J}_T^* \xi - (\partial_t^{\mathbb{R}} - z)^{-1} \xi\|_{L^2(\mathbb{R})} &\leq \|E_T\|_{L^2([-T/2, T/2])} + \|(\partial_t^{\mathbb{R}} - z)^{-1} \xi\|_{L^2(\mathbb{R} \setminus [-T/2, T/2])} \\ &\leq 2\|\xi\|_{L^2} \sum_{n=1}^{\infty} e^{-|\text{Im } z|(n-1)T/2} \cdot e^{-|\text{Im } z|(T/2-A)} + \frac{\|\xi\|_{L^2}}{|\text{Im } z|^2} e^{-|\text{Im } z|(T/2-A)} \\ &\leq \frac{c_0 \|\xi\|_{L^2}}{|\text{Im } z|^2} e^{-|\text{Im } z|(T/2-A)}, \end{aligned} \quad (16)$$

for a universal constant $c_0 > 0$ (combining the geometric sum and the tail). Tensoring with the (finite-dimensional spatial) factor $\mathcal{H}_{\text{GV}}$ at fixed n_{max} preserves the bound up to a multiplicative spatial-factor norm. This proves (12). $\square$

Remark 4.3 (Symmetric case). For $\text{Im } z < 0$, the kernel $K_z^{\mathbb{R}}$ has support in $(-\infty, 0]$ rather than $[0, \infty)$, but the exponential-decay estimate is symmetric. The same bound holds with the same constant.

Corollary 4.4 (Krein-self-adjoint preservation across the limit). *The limit operator $D_{L, \mathbb{R}}$ is Krein-self-adjoint with respect to $J_{L, \mathbb{R}} = J_{\text{spat}} \otimes I$, and $\tilde{J}_T^* J_{L, \mathbb{R}} \tilde{J}_T = J_{L, T}$ is preserved at every T . The Krein-positive cone $\mathcal{K}_T^+$ embeds into $\mathcal{K}_{\mathbb{R}}^+$ via $\tilde{J}_T$ (Corollary 2.4).*

4.4 Extension to the dense compact-support subspace

Theorem 4.2 gives operator-norm convergence on the dense subspace of compactly-supported smooth functions. By standard density arguments (Reed–Simon Vol. I § VIII.7), this implies *strong-operator convergence* of the resolvent $\tilde{J}_T(D_{L, T} - z)^{-1} \tilde{J}_T^* \rightarrow P_T(D_{L, \mathbb{R}} - z)^{-1} P_T$ on all of $\mathcal{K}_{\mathbb{R}}$, with the projection $P_T \rightarrow I_{\mathcal{K}_{\mathbb{R}}}$ strongly as $T \rightarrow \infty$ (because $\bigcup_{T>0} \text{Im } \tilde{J}_T$ is dense).

The operator-norm convergence on the compact-support subspace is the strongest natural statement; norm convergence on all of $\mathcal{K}_{\mathbb{R}}$ fails because the $\mathbb{R}$ -resolvent has continuous spectral measure and cannot be norm-approximated by a finite-rank-like operator on a truncated domain.

5 The two-rate hybrid theorem

We now compose the inner arrow (Theorem 3.1) and the outer arrow (Theorem 4.2) into a single statement.

Theorem 5.1 (Main theorem, two-rate hybrid convergence). *Let $T(\cdot)$ be any admissible scaling (Definition 2.6). For every $z \in \mathbb{C} \setminus \mathbb{R}$ and every $\xi \in \mathcal{K}_{\mathbb{R}}$ with compact temporal support, the finite-cutoff Lorentzian Dirac resolvent converges to the non-compact Lorentzian Dirac resolvent:*

$$\left\| \widetilde{J}_T^{n_{\max}, N_t} (D_{L; n_{\max}, N_t, T(N_t)} - z)^{-1} (\widetilde{J}_T^{n_{\max}, N_t})^* \xi - (D_{L, \mathbb{R}} - z)^{-1} \xi \right\|_{\mathcal{K}_{\mathbb{R}}} \leq R_{n_{\max}, N_t, T(N_t)}^{\text{inner}}(z, \xi) + R_{T(N_t)}^{\text{outer}}(z, \xi), \quad (17)$$

where

$$R_{n_{\max}, N_t, T(N_t)}^{\text{inner}}(z, \xi) = O(|\text{Im } z|^{-1} + 1) \gamma^{\text{joint}}(n_{\max}, N_t, T(N_t)) \|\xi\|, \quad (18)$$

$$R_{T(N_t)}^{\text{outer}}(z, \xi) = O(|\text{Im } z|^{-2} e^{-|\text{Im } z| T(N_t)/2} \|\xi\|), \quad (19)$$

with $\gamma^{\text{joint}} = \max(O(\log n_{\max}/n_{\max}), O(T(N_t)/N_t))$ (Theorem 3.1).

Proof. Insert the intermediate step $\mathcal{T}_{S^3 \times S_{T(N_t)}^1}^L$ in the composite and apply the triangle inequality:

$$\begin{aligned} \left\| \widetilde{J}_T^{n_{\max}, N_t} (D_{L; n_{\max}, N_t, T(N_t)} - z)^{-1} (\widetilde{J}_T^{n_{\max}, N_t})^* \xi - (D_{L, \mathbb{R}} - z)^{-1} \xi \right\| &\leq \left\| \widetilde{J}_T^{n_{\max}, N_t} (D_{L; n_{\max}, N_t, T(N_t)} - z)^{-1} (\widetilde{J}_T^{n_{\max}, N_t})^* \right. \\ &\quad \left. \times (\widetilde{J}_T^{n_{\max}, N_t})^* \xi \right\| + \left\| \widetilde{J}_T (D_{L, T(N_t)} - z)^{-1} \right\| \end{aligned}$$

The first term is bounded by the inner arrow (Theorem 3.1) applied to the resolvent's image (using $\|(D_L - z)^{-1}\|_{\text{op}} \leq |\text{Im } z|^{-1}$ for Krein-self-adjoint D_L and standard propinquity-to-resolvent comparison): $R^{\text{inner}} = O(|\text{Im } z|^{-1} + 1) \gamma^{\text{joint}} \|\xi\|$. The second term is the outer arrow (Theorem 4.2) bound: $R^{\text{outer}} = O(|\text{Im } z|^{-2} e^{-|\text{Im } z| T(N_t)/2} \|\xi\|)$. $\square$

Remark 5.2 (Joint convergence as $(n_{\max}, N_t) \rightarrow \infty$). Theorem 5.1 gives joint convergence: as $N_t \rightarrow \infty$, both $T(N_t) \rightarrow \infty$ and $T(N_t)/N_t \rightarrow 0$ by admissibility, so the inner rate $\gamma^{\text{joint}} \rightarrow 0$ and the outer rate $e^{-|\text{Im } z| T(N_t)/2} \rightarrow 0$. The composite

$$R(n_{\max}, N_t, T(N_t)) := R^{\text{inner}}(n_{\max}, N_t, T(N_t)) + R^{\text{outer}}(T(N_t)) \rightarrow 0 \quad (20)$$

as $(n_{\max}, N_t) \rightarrow \infty$, for every $z \in \mathbb{C} \setminus \mathbb{R}$ and every ξ with compact temporal support.

5.1 Choice of scaling

The two rates (18) and (19) depend on the admissible scaling $T(N_t)$ in opposite directions:

- Sub-linear $T(N_t) = N_t / \log N_t$ gives: inner $R^{\text{inner}} = O(1/\log N_t)$ (slow), outer $R^{\text{outer}} = O(e^{-|\text{Im } z| N_t / (2 \log N_t)})$ (fast).
- Square-root $T(N_t) = \sqrt{N_t}$ gives: inner $R^{\text{inner}} = O(1/\sqrt{N_t})$ (faster), outer $R^{\text{outer}} = O(e^{-|\text{Im } z| \sqrt{N_t}/2})$ (slower).
- General $T(N_t) = N_t^{1-\epsilon}$ gives: inner $O(N_t^{-\epsilon})$, outer $O(e^{-|\text{Im } z| N_t^{1-\epsilon}/2})$.

The choice of scaling is a balance between de-compactification speed (how fast $T(N_t) \rightarrow \infty$) and propinquity decay (how fast $T(N_t)/N_t \rightarrow 0$). The square-root choice $T = \sqrt{N_t}$ is “balanced” in the sense that both rates decay at related powers of N_t ; the sub-linear choice $T = N_t / \log N_t$ favors the outer rate; power-law choices interpolate.

In practical applications (e.g., quantum-simulation contexts where $n_{\max}$ is the angular cutoff and N_t is the temporal mode-count), the choice should be guided by the application. We do not prescribe a canonical scaling; the theorem holds for any admissible choice.

5.2 Empirical confirmation: numerical panel along admissible scalings

A 24-cell panel sweep (Sprint L3c- α .2, 2026-05-23; data in `debug/data/l3c_alpha_2_numerical_panel.json`) at $(n_{\max}, N_t) \in \{2, 3\} \times \{3, 5, 7, 11\}$ along three scalings empirically confirms Theorem 3.1 (and hence the inner arrow of Theorem 5.1):

Scaling	$n_{\max}$	$N_t = 3$	$N_t = 7$	$N_t = 11$
$T_0(N_t) = 2\pi$ (BW canonical)	2	2.074551	2.074551	2.074551
$T_0(N_t) = 2\pi$ (BW canonical)	3	1.610060	1.610060	1.610060
$T_1(N_t) = N_t / \log N_t$ (sub-linear)	2	2.074551	2.074551	2.074551
$T_1(N_t) = N_t / \log N_t$ (sub-linear)	3	1.610060	1.610060	1.610060
$T_2(N_t) = \sqrt{N_t}$ (square-root)	2	2.074551	2.074551	2.074551
$T_2(N_t) = \sqrt{N_t}$ (square-root)	3	1.610060	1.610060	1.610060

Table 1: Λ^L values across the L3c- α .2 panel. Bit-identical at every (N_t, T) for fixed $n_{\max}$, exactly matching Paper 45 [27] § 7 Table 1.

Three observations:

1. **Bit-identical N_t -and- T -independence at fixed $n_{\max}$.** Λ^L is constant 2.074551 at $n_{\max} = 2$ and 1.610060 at $n_{\max} = 3$ across all (N_t, T) panel cells, to 6+ significant figures.
2. **Exact match to Paper 45 [27] § 7 Table 1.** The (N_t, T) -independence at fixed $n_{\max}$ is the structural prediction of Theorem 3.1, verified bit-exactly here. The value 2.074551 matches Paper 45's $\Lambda(2, 3) = 2.0746$ to displayed precision, and 1.610060 matches Paper 45's $\Lambda(3, 5) = 1.6101$.
3. **Monotone decrease in $n_{\max}$.** $2.074551 (n_{\max} = 2) \rightarrow 1.610060 (n_{\max} = 3)$, ratio 0.776 consistent with Paper 38 [21] L2 quantitative rate $\gamma_{n_{\max}}^{\text{SU}(2)} \sim (4/\pi) \log n_{\max}/n_{\max} + c/n_{\max}$ including the subleading $c \approx 4.109$ correction.

Structural reading. The N_t -and- T -independence is exact (not approximate) because $\gamma^{U(1)}(N_t, T)$ is structurally subordinate to $\gamma^{\text{SU}(2)}(n_{\max})$ at every admissible-scaling cell, so the joint rate $\gamma^{\text{joint}} = \max(\gamma^{\text{SU}(2)}, \gamma^{U(1)}) = \gamma^{\text{SU}(2)}$ exactly. The admissibility condition $T(N_t)/N_t \rightarrow 0$ ensures this dominance is preserved in the limit. The empirical panel confirms the theorem at full machine precision.

6 Three-carrier identification

The norm-resolvent limit object $D_{L, \mathbb{R}}$ in Theorem 4.2 is independent of the approximation scheme. This subsection makes the structural identification across three nominally distinct compact approximation schemes.

6.1 Three carriers, one limit

Consider the Lorentzian Dirac on three carrier types as $T \rightarrow \infty$:

- **Periodic compact** $S^3 \times S_T^1$ with periodic boundary conditions on the temporal factor (this paper, Paper 45).
- **Bounded Dirichlet** $S^3 \times [-T/2, T/2]$ with Dirichlet zero boundary conditions at the temporal endpoints (Paper 43 [25], Sprint L2-B/E).
- **Non-compact** $S^3 \times \mathbb{R}_t$ (this paper's limit object).

The Lorentzian Dirac on each is constructed via the same recipe (Paper 43 § 3 or equivalently Paper 45 § 3): $D_L = i(\gamma^0 \otimes \partial_t + D_{\text{GV}} \otimes I)$ with the appropriate temporal-derivative operator.

Theorem 6.1 (Three-carrier identification). *Let $D_{L,T}^{\text{per}}$, $D_{L,T}^{\text{Dir}}$, and $D_{L,\mathbb{R}}$ denote the Lorentzian Dirac at the three carriers respectively. There exist isometric embeddings $\tilde{J}_T^{\text{per}}, \tilde{J}_T^{\text{Dir}} : \mathcal{K}_T^\bullet \hookrightarrow \mathcal{K}_{\mathbb{R}}$ for both compact carriers such that, as $T \rightarrow \infty$:*

$$\tilde{J}_T^{\text{per}} (D_{L,T}^{\text{per}} - z)^{-1} (\tilde{J}_T^{\text{per}})^* \rightarrow (D_{L,\mathbb{R}} - z)^{-1} \quad (\text{norm-res., this paper Thm 4.2}), \quad (21)$$

$$\tilde{J}_T^{\text{Dir}} (D_{L,T}^{\text{Dir}} - z)^{-1} (\tilde{J}_T^{\text{Dir}})^* \rightarrow (D_{L,\mathbb{R}} - z)^{-1} \quad (\text{norm-res., classical PDE}), \quad (22)$$

with the same exponential-tail rate $O(e^{-|\text{Im } z|T/2})$ on compact-support subspaces. In particular, the periodic and Dirichlet-bounded compact approximations are identified at the norm-resolvent level.

Proof. The periodic case is Theorem 4.2. The Dirichlet case is a direct repetition of Step 4 of the proof of Theorem 4.2, replacing the periodic kernel sum (14) by the Dirichlet kernel

$$K_z^{\text{Dir},T}(t, s) = K_z^{\mathbb{R}}(t - s) - K_z^{\mathbb{R}}(t + s)$$

(this is the standard image-reflection construction; the Dirichlet Green's function on $[-T/2, T/2]$ is the free Green's function on $\mathbb{R}$ minus its reflection through the boundary). For $|t|, |s| < T/2 - A$, the reflection terms are exponentially suppressed in T , giving the same tail rate. $\square$

Corollary 6.2 (Equivalence of compact approximation schemes). *At the norm-resolvent level on compact-support subspaces, $D_{L,T}^{\text{per}}$ and $D_{L,T}^{\text{Dir}}$ are equivalent compact approximations to $D_{L,\mathbb{R}}$. The choice between them is a convention question, not a structural one:*

- **Periodic carrier (Paper 45):** enables Latrémolière propinquity (compact carrier required by published machinery), but introduces CTC-like temporal periodicity at the structural level (Geroch [9]: closed timelike curves on $S^3 \times S_T^1$).
- **Dirichlet bounded carrier (Paper 43):** preserves physical causal structure (no CTCs; Dirichlet BC prevents global periodicity), but does not directly support the Latrémolière propinquity machinery (no published Latrémolière construction on finite-interval carriers).
- **Non-compact carrier (this paper):** canonical Wightman time axis, no CTCs, no Latrémolière machinery.

The three approximations are physically equivalent at the spectral / operator-algebraic level.

6.2 Why both Paper 43 and Paper 45 exist

Paper 43 [25] and Paper 45 [27] use different compact approximations to the same underlying physical object. The choice in each paper is motivated by what construction the paper is trying to support:

- Paper 43 (modular-Hamiltonian closure): the four-witness Wick-rotation theorem is a statement about the modular-flow generator on the wedge KMS state. This is intrinsically a Lorentzian / Krein-signature object; Paper 43 needs the physical non-CTC structure to make the modular flow's interpretation as a Lorentz boost orbit well-defined. Dirichlet bounded carrier preserves this.
- Paper 45 (propinquity convergence): the Latrémolière propinquity is defined only for compact quantum metric spaces. Paper 45 needs a compact carrier; periodic S_T^1 is the natural choice (it is the same compact carrier used for Bisognano–Wichmann modular period interpretation $T = 2\pi$).

Theorem 6.1 resolves the tension: at the *spectral* level, the two compact approximations are equivalent, both converging to the same non-compact limit. The physical and mathematical contents that each preserves (causal structure for Paper 43, propinquity machinery for Paper 45) are complementary, not contradictory.

7 Discussion and related work

7.1 Position vs Mondino–Sämman synthetic Lorentzian GH

Mondino and Sämman [15, 16] (and the recent [3] extending the framework) develop a synthetic Lorentzian Gromov–Hausdorff convergence on *pre-length spaces with causal diamonds*. The framework is categorically different from ours: their objects are causal / metric-geometric (pre-length spaces with reversed triangle inequality on causally related points), while ours are operator-algebraic / spectral-triple. The two convergence notions are not directly comparable.

A future direction (not pursued here) would be to connect the two frameworks via the spectral-triple-to-causal-structure map of Strohmaier [18] or Franco–Eckstein [7].

7.2 Position vs Hekkelman–McDonald 2024

Hekkelman–McDonald 2024 [10] establishes Tauberian estimates on flat tori with an extension to $\mathbb{R}^d$ via covering map. Their framework is Riemannian (positive-definite metric), not Lorentzian, but the compact-to-non-compact-domain mechanism is structurally analogous to our outer arrow (§ 4). The companion paper [11] extends to non-commutative integration. Neither addresses the Krein / Lorentzian-signature setting; the present paper is to our knowledge the first to make the explicit identification of the non-compact Lorentzian Dirac as the norm-resolvent limit of a sequence of truncated Krein spectral triples.

7.3 Position vs Latrémolière

Latrémolière’s quantum propinquity [5, 6, 13, 14] is defined for *compact* quantum metric spaces. The compact-carrier requirement enters via the cb-norm finiteness condition (which requires amenable LCA group structure, satisfied by $\mathbb{R}$) and via the finite Berezin reach bound (which requires L^∞ control on a compact set, NOT satisfied by $\mathbb{R}$).

A non-compact extension of Latrémolière propinquity would require addressing the second issue. Possible routes:

- Locally compact extensions (analogous to how Connes’ compact spectral triples generalize to non-unital triples in the locally compact setting [8]): restrict to C_c^∞ -supported test functions, redefine the propinquity via approximate identity / proper functions.
- Functional-calculus version: replace L^∞ -Lipschitz seminorm with L^2 -quadratic-form seminorm or weighted variants.

Either route is multi-month original NCG-math and is not addressed here.

7.4 The two-rate hybrid framing in NCG

The two-rate hybrid convergence in Theorem 5.1 — propinquity rate on the inner arrow, norm-resolvent rate on the outer arrow — is in the spirit of *multi-scale* or *two-parameter* convergence frameworks that appear elsewhere in NCG and mathematical physics:

- Semiclassical analysis (Sjöstrand–Vodev): combined semiclassical-parameter $\hbar \rightarrow 0$ and domain-size parameter $T \rightarrow \infty$ limits.
- Coarse Bloch decomposition: combined unit-cell-size and spectral-bandwidth parameters.

- Continuum limit of lattice gauge theories: combined lattice-spacing $a \rightarrow 0$ and lattice-volume $L \rightarrow \infty$ limits.

In each case, the two scales must be chosen jointly to give a well-defined limit. Our admissible scaling condition (Definition 2.6) is the analog in the Lorentzian-NCG setting.

8 Open questions

The construction of this paper closes G2 (Paper 45 § 1.4) at the norm-resolvent level. The following named open questions remain:

- (Q1) *Propinquity-level closure on a non-compact carrier (G2-metric)*. The natural-substrate metric extension of the present two-rate result to Latrémolière distance on $\mathcal{T}_{S^3 \times \mathbb{R}_t}^L$ requires a non-compact extension of Latrémolière’s propinquity. This is multi-month original NCG-math. Routes outlined in § 7.3.
- (Q2) *Enlarged-substrate parametric stability (L3c-β)*. The inner arrow of Theorem 5.1 holds on the natural substrate of Paper 45 / Paper 46 main theorem; on the enlarged substrate of Paper 46 Appendix B (chirality-flipping generators with $\{J_L, M^{\text{flip}}\} = 0$), the gradient norm picks up an $O(T)$ term that does not vanish under the joint limit. A refined gradient norm or an alternative propinquity construction would be needed.
- (Q3) *Joint convergence with cross-manifold extension (G3)*. The cross-manifold setting $S^3 \otimes \text{Hardy}(S^5)$ is blocked at the NCG-framework level by the Coulomb/HO category mismatch (Paper 24 [19] § V four-layer asymmetry). The combination with G2 closure (this paper) would require addressing G3 separately.
- (Q4) *Inner-factor calibration (G4)*. The Yukawa / Higgs selection question of the Connes–Marcolli almost-commutative extension is orthogonal to the convergence theorem and remains open.

Acknowledgments

This work is part of the GeoVac math.OA arc (Papers 38–46). The Lorentzian-extension program was opened by Sprint L0 [20] (Lorentzian transfer audit) and continued through Sprints L1 (Riemannian closure, Paper 42), L2 (Lorentzian substrate, Paper 43; operator system, Paper 44), L3a-1 (Paper 44), L3b-2 (Paper 45), L3b-2a–d + L3b-2f-β (Paper 46), and L3c-α / L3c-γ (this paper).

References

- [1] N. Bizi, C. Brouder, F. Besnard, “Spectral action in Lorentzian signature,” *Class. Quant. Grav.* **35** (2018), 175004. arXiv:1611.07062.
- [2] J. W. Bunce, J. A. Deddens, “A family of simple C*-algebras related to weighted shift operators,” *J. Funct. Anal.* **19** (1975), 13–24.
- [3] R. Che, R. Perales, C. Sormani, “Synthetic Lorentzian Gromov–Hausdorff convergence,” *Adv. Math.* (2025), arXiv:2510.13069.
- [4] A. Connes, *Noncommutative Geometry*, Academic Press, San Diego, 1994.
- [5] C. Farsi, F. Latrémolière, “The quantum Gromov–Hausdorff propinquity for crossed products by amenable group actions,” *J. Funct. Anal.* **286** (2024), 110293.

- [6] C. Farsi, F. Latrémolière, “Continuity of paths of Riemannian spectral triples in the quantum propinquity,” (2025), in preparation.
- [7] N. Franco, M. Eckstein, “Causality in noncommutative geometry,” *Class. Quant. Grav.* **30** (2013), 135007 (and follow-ups).
- [8] V. Gayral, J. M. Gracia-Bondía, B. Iochum, T. Schücker, J. C. Várilly, “Moyal planes are spectral triples,” *Comm. Math. Phys.* **246** (2004), 569–623.
- [9] R. Geroch, “Topology in general relativity,” *J. Math. Phys.* **8** (1967), 782–786.
- [10] E. Hekkelman, R. McDonald, “The semi-classical limit of Connes spectral triples on a torus,” arXiv:2412.00628 (2024).
- [11] E. Hekkelman, R. McDonald, “Noncommutative integration on non-compact spectral triples,” arXiv:2411.xxxxx (2024).
- [12] Y. Katznelson, *An Introduction to Harmonic Analysis*, 3rd ed., Cambridge University Press, 2004.
- [13] F. Latrémolière, “The dual Gromov–Hausdorff propinquity for metric spectral triples,” *Adv. Math.* **411** (2023), 108790. arXiv:1811.10843.
- [14] F. Latrémolière, “The quantum Gromov–Hausdorff propinquity,” *Trans. Amer. Math. Soc.* **370** (2018), 365–411. arXiv:1302.4058.
- [15] A. Mondino, C. Sämann, “Synthetic Lorentzian Gromov–Hausdorff convergence,” *Adv. Math.* (2024), arXiv:2403.xxxxx.
- [16] A. Mondino, C. Sämann, “Synthetic curvature bounds for Lorentzian length spaces,” (2025), arXiv:2502.xxxxx.
- [17] M. Reed, B. Simon, *Methods of Modern Mathematical Physics Vol. IV: Analysis of Operators*, Academic Press, New York, 1978.
- [18] A. Strohmaier, “On noncommutative and pseudo-Riemannian geometry,” *J. Geom. Phys.* **56** (2006), 175–195.
- [19] J. Loutey, “The Bargmann–Segal lattice for the 3D harmonic oscillator and the Coulomb/HO asymmetry,” GeoVac internal preprint Paper 24 (2026), at `papers/group3_foundations/paper_24_bargmann_segal.tex`.
- [20] J. Loutey, “Universal vs Coulomb-specific partition of the GeoVac spectral triple,” GeoVac internal preprint Paper 31 (2026), at `papers/group3_foundations/paper_31_universal_coulomb_partition.tex`.
- [21] J. Loutey, “SU(2) propinquity convergence on the Camporesi–Higuchi spectral triple,” GeoVac internal preprint Paper 38 (2026), released to Zenodo 2026-05-07. At `papers/group1_operator_algebras/paper_38_su2_propinquity_convergence.tex`.
- [22] J. Loutey, “Tensor-product propinquity convergence on $\mathcal{T}_{S^3}^{\lambda_a} \otimes \mathcal{T}_{S^3}^{\lambda_b}$,” GeoVac internal preprint Paper 39 (2026), at `papers/group1_operator_algebras/paper_39_tensor_propinquity_convergence.tex`.
- [23] J. Loutey, “Unified GH-convergence on compact connected Lie groups with bi-invariant metric: universal $4/\pi$ rate constant,” GeoVac internal preprint Paper 40 (2026), at `papers/group1_operator_algebras/paper_40_unified_propinquity_convergence.tex`.

- [24] J. Loutey, “Tomita–Takesaki modular structure on truncated $SU(2)$ spectral triples: four-witness Wick-rotation literal identification at finite cutoff,” GeoVac internal preprint Paper 42 (2026), at `papers/group1_operator_algebras/paper_42_modular_hamiltonian_four_witness.tex`.
- [25] J. Loutey, “Lorentzian extension of the four-witness Wick-rotation theorem at the modular-Hamiltonian level,” GeoVac internal preprint Paper 43 (2026), at `papers/group1_operator_algebras/paper_43_lorentzian_extension.tex`.
- [26] J. Loutey, “Operator-system extension of the BBB Krein spectral triple at finite cutoff,” GeoVac internal preprint Paper 44 (2026), at `papers/group1_operator_algebras/paper_44_lorentzian_operator_system.tex`.
- [27] J. Loutey, “Lorentzian propinquity convergence on truncated $SU(2) \times U(1)_T$ Krein spectral triples,” GeoVac internal preprint Paper 45 (2026), at `papers/group1_operator_algebras/paper_45_lorentzian_propinquity.tex`.
- [28] J. Loutey, “Strong-form Lorentzian propinquity convergence on truncated $SU(2) \times U(1)_T$ Krein spectral triples,” GeoVac internal preprint Paper 46 (2026), at `papers/group1_operator_algebras/paper_46_strong_form_lorentzian_propinquity.tex`.