Potential-Independent Angular Sparsity for Qubit Hamiltonians in Spherical Fermion Systems

J. Loutey[1]

[1] *Independent Researcher, Kent, Washington*

(Dated: April 9, 2026)

For any N-fermion system whose single-particle orbitals are products of spherical harmonics and arbitrary radial functions, the fraction of nonzero two-body matrix elements is determined solely by angular momentum selection rules and is independent of the radial potential. Exact enumeration of the Gaunt coefficients yields the angular ERI density $D(l_{\max})$: 7.81% at $l_{\max} = 1$, 2.76% at $l_{\max} = 2$, 1.44% at $l_{\max} = 3$, 0.90% at $l_{\max} = 4$, and 0.62% at $l_{\max} = 5$. At $l_{\max} = 2$, 97.24% of all possible two-body matrix elements are guaranteed to vanish by the Wigner–Eckart theorem before any property of the potential is invoked. The bound is independent of the number of radial quantum numbers retained at fixed $l_{\max}$, and therefore independent of qubit count. Numerical verification across five potentials (Coulomb, harmonic oscillator, Woods–Saxon, square well, Yukawa) confirms the theorem: angular sparsity patterns are bit-identical across all potentials at matched orbital counts. The radial-angular factorization and the Gaunt selection rules are standard results from Condon & Shortley and Slater; the contribution of this paper is the explicit quantitative bound at each $l_{\max}$, framed as a universal resource guarantee for qubit Hamiltonian sparsity that applies to electronic structure, nuclear shell models, quantum dots, and any other spherical-potential fermion system.

I. INTRODUCTION

Quantum simulation of fermionic systems requires encoding second-quantized Hamiltonians as sums of Pauli operators. The number of Pauli terms, the number of qubitwise-commuting (QWC) measurement groups, and the Pauli 1-norm λ are the dominant resource metrics for near-term variational [?] and fault-tolerant [?] quantum algorithms. All three metrics scale with the number of nonzero two-body matrix elements in the chosen orbital basis, making Hamiltonian sparsity the central structural resource for quantum simulation.

Existing sparsity analyses are potential-specific. In quantum chemistry, the Coulomb $1/r_{12}$ interaction produces sparsity governed by the multipole expansion and the basis set's spatial locality [? ?]. In nuclear structure physics, the nucleon–nucleon (NN) interaction in a harmonic oscillator (HO) basis produces sparsity governed by Talmi–Moshinsky brackets and center-of-mass factorization [?]. These analyses are carried out independently, obscuring a structural feature that is common to both domains: the angular momentum selection rules imposed by the spherical harmonic component of the basis are identical.

In this paper we state the resulting potential-independent sparsity bound explicitly. For any rotationally invariant (central) two-body interaction expanded in a spherical harmonic basis, the zero structure of the two-body matrix elements depends only on the angular quantum numbers (l, m) and is independent of the radial wavefunctions, the radial potential, and the radial quantum numbers. The underlying factorization of two-body integrals into angular (Gaunt) coefficients and radial (Slater-type) integrals has been standard since Condon and Shortley [?] and Slater [?], and we claim no mathematical originality for it. The contribution of this paper is the explicit quantitative ERI density at each $l_{\max}$, verified numerically across five potentials, and framed as a universal resource bound for the quantum computing community.

This result was identified in the course of extending the GeoVac spectral graph theory framework [? ?] from electronic to nuclear structure applications. In the companion paper [?] we show that a separate feature of the GeoVac framework—namely the reduction of the one-body graph Laplacian spectrum to pure integers on S^3—is special to the Coulomb potential, so that the universal–specific partition identified here is sharp: angular sparsity is universal across spherical fermion systems, while the S^3 Fock projection and the π-free graph structure are Coulomb-specific.

II. TWO-BODY MATRIX ELEMENT FACTORIZATION

Consider N fermions in a single-particle basis $\{|n, l, m\rangle\}$ where n labels the radial quantum number, l is the orbital angular momentum, and m is the magnetic quantum number. The spatial orbitals are

$$\phi_{nlm}(\mathbf{r}) = R_{nl}(r)\, Y_l^m(\hat{r}), \tag{1}$$

where $R_{nl}(r)$ are radial wavefunctions determined by the single-particle potential $V(r)$ and Y_l^m are spherical harmonics.

Let $\hat{V}_{12} = V(|\mathbf{r}_1 - \mathbf{r}_2|)$ be a rotationally invariant two-body interaction. We expand $\hat{V}_{12}$ in Legendre polynomials:

$$V(|\mathbf{r}_1 - \mathbf{r}_2|) = \sum_{k=0}^{\infty} v_k(r_1, r_2)\, P_k(\cos\theta_{12}), \tag{2}$$

where $v_k(r_1, r_2)$ are radial expansion coefficients that depend on the specific form of V. For the Coulomb interaction, $v_k = r_<^k / r_>^{k+1}$; for other potentials, v_k takes different forms but the angular structure is identical.

Theorem 1 (Radial–Angular Factorization). *For any rotationally invariant two-body interaction $V(|\mathbf{r}_1 - \mathbf{r}_2|)$, the two-body matrix element factorizes as*

$$\langle n_1 l_1 m_1,\, n_2 l_2 m_2 \,|\, \hat{V}_{12} \,|\, n_3 l_3 m_3,\, n_4 l_4 m_4 \rangle = \sum_k c^k(l_1 m_1;\, l_3 m_3)\; c^k(l_2 m_2;\, l_4 m_4)\; R^k(n_1 l_1, n_2 l_2;\, n_3 l_3, n_4 l_4), \tag{3}$$

where the Gaunt coefficients

$$c^k(l\, m;\, l'\, m') = (-1)^m \sqrt{(2l+1)(2l'+1)} \begin{pmatrix} l & k & l' \\ 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} l & k & l' \\ -m & q & m' \end{pmatrix} \tag{4}$$

with $q = m - m'$ depend only on angular quantum numbers, and the Slater-type radial integrals

$$R^k = \int_0^\infty \int_0^\infty R_{n_1 l_1}(r_1)\, R_{n_3 l_3}(r_1)\, v_k(r_1, r_2)\, R_{n_2 l_2}(r_2)\, R_{n_4 l_4}(r_2)\, r_1^2 r_2^2\, dr_1\, dr_2 \tag{5}$$

depend on the potential V through the radial wavefunctions and the expansion coefficients v_k. The sum over k is bounded: $\max(|l_1 - l_3|, |l_2 - l_4|) \le k \le \min(l_1 + l_3, l_2 + l_4)$, with the parity constraints $l_1 + l_3 + k$ even and $l_2 + l_4 + k$ even.

Proof. Substituting the Legendre expansion (**??**) into the two-body integral and applying the spherical harmonic addition theorem $P_k(\cos\theta_{12}) = \frac{4\pi}{2k+1} \sum_q Y_k^{q*}(\hat{r}_1) Y_k^q(\hat{r}_2)$ yields a product of two angular integrals (Gaunt integrals) and one radial integral for each k. The Gaunt integral $\int Y_{l_1}^{m_1 *} Y_k^q Y_{l_3}^{m_3}\, d\Omega$ evaluates to a product of Wigner $3j$ symbols by the Wigner–Eckart theorem [**?**], giving Eq. (**??**). The bounds on k follow from the triangle inequality of the $3j$ symbol: $|l - l'| \le k \le l + l'$, applied to both Gaunt coefficients. The parity constraint follows from the $3j$ symbol $(l\ k\ l';\ 0\ 0\ 0)$, which vanishes unless $l + k + l'$ is even [**?**]. □

The factorization (**??**) is a standard result of angular momentum algebra [**?** **?** **?**]. The key observation for quantum computing applications is the following:

Theorem 2 (Potential-Independent Sparsity). *A two-body matrix element $\langle n_1 l_1 m_1, n_2 l_2 m_2 | \hat{V}_{12} | n_3 l_3 m_3, n_4 l_4 m_4 \rangle$ vanishes identically for every central potential whenever $c^k(l_1 m_1; l_3 m_3) \cdot c^k(l_2 m_2; l_4 m_4) = 0$ for every allowed k. Conversely, whenever at least one Gaunt product is nonzero, the matrix element is generically nonzero. The zero condition depends only on $(l_1, m_1, l_2, m_2, l_3, m_3, l_4, m_4)$ and is independent of the radial potential $V(r)$, the radial wavefunctions $R_{nl}(r)$, and the radial quantum numbers (n_1, n_2, n_3, n_4).*

Proof. The symmetry direction is immediate: if all angular factors vanish in Eq. (**??**), every term in the sum is zero regardless of R^k. This is the statement that the zero pattern generated by angular selection rules is a lower bound on sparsity.

For the converse, suppose there exists at least one k^* for which $c^{k^*}(l_1 m_1; l_3 m_3) \cdot c^{k^*}(l_2 m_2; l_4 m_4) \neq 0$. The corresponding Slater integral R^{k^*} is a weighted inner product of the radial functions $R_{n_1 l_1} R_{n_3 l_3}$ and $R_{n_2 l_2} R_{n_4 l_4}$ mediated by $v_{k^*}(r_1, r_2) > 0$ on a set of positive measure (true for any physical central potential that does not vanish identically in its multipole component). For any Sturm–Liouville potential with a discrete spectrum—including Coulomb, harmonic oscillator, Woods–Saxon, square well, and Yukawa—the radial products $R_{nl}(r) R_{n'l'}(r)$ are linearly independent as a function of r for distinct index tuples, and the bilinear functional defined by v_{k^*} is nondegenerate on this family. Therefore $R^{k^*} = 0$ can hold only for a measure-zero subset of the space of central potentials—pathological choices in which two or more R^k contributions conspire to cancel accidentally. Any infinitesimal perturbation of the potential restores $R^{k^*} \neq 0$. □

Remark. The theorem classifies zero matrix elements into two types: *symmetry-protected zeros* (those forced by the Gaunt selection rules) and *accidental zeros* (those produced by fine-tuned cancellations in R^k for specific potentials). Only symmetry-protected zeros transfer across potentials; accidental zeros are fragile under perturbation. Below, when we refer to the "angular ERI density" $D(l_{\max})$, we mean the density after symmetry-protected zeros are removed.

III. ERI DENSITY BOUNDS

Definition. The *angular ERI density* $D(l_{\max})$ is the fraction of four-index combinations $(l_1 m_1, l_2 m_2, l_3 m_3, l_4 m_4)$ with $l_i \le l_{\max}$ for which the matrix element is not forced to vanish by Gaunt selection rules. Equivalently, $D(l_{\max})$ is the fraction of four-index combinations for which (i) m-conservation $m_1 + m_2 = m_3 + m_4$ holds, and (ii) at least one value of k satisfies the triangle and parity conditions for both Gaunt factors with both factors nonzero.

The total number of angular orbital labels at $l_{\max}$ is $N_{\text{orb}} = (l_{\max} + 1)^2$, so the total number of unrestricted four-index combinations is N_{orb}^4. The angular ERI density counts the fraction that survives Gaunt selection rules.

Theorem 3 (ERI Density at Low $l_{\max}$). *Exact enumeration of the Gaunt selection rules gives:*

$l_{\max}$	Shells	N_{orb}	N_{orb}^4	$D(l_{\max})$
0	s	1	1	100.00%
1	$s+p$	4	256	7.8125%
2	$s+p+d$	9	6,561	2.7587%
3	$s+p+d+f$	16	65,536	1.4404%
4	$s+p+d+f+g$	25	390,625	0.9024%
5	through h	36	1,679,616	0.6190%

These values are obtained by exact enumeration of the N_{orb}^4 four-index combinations, applying the conditions of the definition above. The enumeration is implemented in `geovac/nuclear/potential_sparsity.py` *as* `angular_zero_count`*, and the values in the table are reproduced by* `debug/verify_paper22_lmax3.py`*.*

The density decreases monotonically with $l_{\max}$: the number of orbitals grows as $(l_{\max}+1)^2$ and the number of four-index combinations as $(l_{\max}+1)^8$, while the number of Gaunt-allowed transitions grows more slowly (polynomially). The asymptotic behavior $D(l_{\max}) \to 0$ as $l_{\max} \to \infty$ reflects the fact that most high-l four-index combinations fail the triangle and parity conditions for every k.

Independence from radial quantum numbers. When the full orbital set includes $n_{\max}$ radial quantum numbers for each (l, m), the total number of spatial orbitals becomes $N_{\text{spatial}} = n_{\max} \times (l_{\max}+1)^2$. The total ERI count scales as $N_{\text{spatial}}^4 = n_{\max}^4 \times N_{\text{orb}}^4$. By Theorem 2, every angular-forbidden element remains zero for all values of the radial quantum numbers, and every angular-allowed element is generically nonzero. The ERI density at fixed $l_{\max}$ is therefore independent of $n_{\max}$:

$$D(l_{\max}, n_{\max}) = D(l_{\max}) \quad \forall\, n_{\max} \geq 1. \tag{6}$$

This means the angular sparsity guarantee is independent of the qubit count $Q = 2 \times n_{\max} \times (l_{\max}+1)^2$ at fixed $l_{\max}$.

IV. NUMERICAL VERIFICATION

To verify Theorems 1–3, we constructed single-particle bases and computed all two-body matrix elements for five distinct radial potentials at $n_{\max} = 2$, $l_{\max} = 2$ ($N_{\text{spatial}} = 18$ orbitals, $Q = 36$ qubits):

1. **Coulomb:** $V(r) = -Z/r$ (hydrogen-like, electronic structure)
2. **Harmonic oscillator:** $V(r) = \frac{1}{2}\omega^2 r^2$ (nuclear shell model, quantum dots)
3. **Woods–Saxon:** $V(r) = -V_0/(1+e^{(r-R_0)/a})$ (nuclear mean field)
4. **Square well:** $V(r) = -V_0\,\Theta(R_0 - r)$ (textbook benchmark)
5. **Yukawa:** $V(r) = -V_0\, e^{-\mu r}/r$ (screened Coulomb, meson exchange)

For each potential, we solved the radial Schrödinger equation numerically (or analytically where available) to obtain radial wavefunctions $R_{nl}(r)$, then computed all $N_{\text{spatial}}^4 = 104{,}976$ two-body matrix elements using the Coulomb interaction $1/|\mathbf{r}_1 - \mathbf{r}_2|$ as the two-body operator. (The single-particle potential determines the *basis*; the two-body operator is the standard Coulomb repulsion in all cases. This is the physically relevant scenario for electronic structure. For nuclear structure, the two-body NN force is itself expanded in Legendre polynomials, and the same factorization applies.)

The results (Table **??**) confirm Theorem 2 to machine precision:

- The angular zero pattern is *bit-identical* across all five potentials. Not merely close—every element that is zero for Coulomb is zero for all other potentials, and every nonzero element is nonzero for all potentials.
- Radial zeros contribute exactly 0.00% additional sparsity. No Slater integral R^k accidentally vanishes for any of the five potentials tested.
- The ERI density is 2.76% for all potentials, matching the angular-only prediction of Theorem 3 at $l_{\max} = 2$.

The bit-identical angular patterns are not a numerical coincidence: they are a direct consequence of the Gaunt coefficients being pure functions of (l, m) quantum numbers, evaluated identically regardless of what potential produced the radial wavefunctions.

TABLE I. ERI sparsity decomposition at $n_{\max} = 2$, $l_{\max} = 2$ ($N_{\text{spatial}} = 18$). Angular zeros are determined by Gaunt selection rules; radial zeros count additional vanishings from radial integral structure.

Potential	Angular zeros (%)	Radial zeros (%)	ERI density (%)
Coulomb	97.24	0.00	2.76
Harmonic osc.	97.24	0.00	2.76
Woods–Saxon	97.24	0.00	2.76
Square well	97.24	0.00	2.76
Yukawa	97.24	0.00	2.76

V. IMPLICATIONS FOR QUANTUM SIMULATION

A. Universal resource bound

Theorem 2 provides a *potential-independent* upper bound on the number of nonzero two-body Pauli terms in any qubit Hamiltonian constructed from a spherical harmonic basis at angular momentum cutoff $l_{\max}$. At $l_{\max} = 2$, at most 2.76% of all possible two-body terms are nonzero; at $l_{\max} = 3$, at most 1.44%. This bound applies identically to:

- *Electronic structure:* Coulomb repulsion in hydrogen-like or Slater-type bases [**?**].
- *Nuclear shell model:* NN interaction in the harmonic oscillator *j-j* basis [**?**]. The central component of the NN force respects the same Gaunt selection rules.
- *Quantum dots:* Electron–electron repulsion in parabolic confinement bases [**?**].
- *Trapped ions and cold atoms:* Any system where single-particle states are expanded in spherical harmonics.

The bound means that qubit Hamiltonian sparsity estimates derived for one physical domain transfer directly to others at matched $l_{\max}$, without re-analysis of the interaction potential.

B. Composed architectures: O($Q^{2.5}$) Pauli scaling for arbitrary spherical potentials

The angular sparsity of Theorem 2 applies to single-center orbital bases. In the GeoVac composed architecture [**?** **?**], a molecular Hilbert space is decomposed into block-diagonal orbital subspaces (core, bond, lone pair), each centered on a specific nucleus and each carrying its own spherical harmonic basis. Cross-block electron repulsion integrals are suppressed structurally by the spatial factorization, and within-block integrals are subject to the angular selection rules of Theorem 2. The resulting Pauli scaling measured across the $LiH/BeH_2/H_2O/HF/NH_3/CH_4$ library is $O(Q^{2.5})$ [**?**], with exponent spread less than 0.05 across molecules.

Corollary 1 (Potential-Independent $O(Q^{2.5})$ Composed Sparsity). *In any composed block architecture whose per-block single-particle bases are spherical harmonic products $R_{nl}(r)Y_l^m$ and whose inter-block couplings respect the block decomposition, the Pauli scaling $O(Q^{2.5})$ observed for the Coulomb interaction in Ref. [**?**] holds for any central two-body interaction whatsoever, with exactly the same scaling exponent. The radial potential affects only the magnitudes of the surviving ERIs, not which ERIs survive.*

Proof sketch. The composed Pauli scaling decomposes into a block-diagonal factor (same for all potentials, depending only on the fiber-bundle structure) and a within-block angular factor (identical for all central potentials by Theorem 2). The radial integrals R^k merely renormalize the coefficients of surviving Pauli strings. Since the Pauli count is determined by which strings have nonzero coefficient, the count is potential-independent; the Pauli 1-norm λ and the QWC group count may differ, but the number of terms and the scaling exponent match. □

This corollary means that the quantum resource advantages of the GeoVac composed architecture—51× to 1,712× fewer Pauli terms than published Gaussian baselines for electronic structure—apply equally to nuclear shell model Hamiltonians, quantum dot Hamiltonians, and any other composed block architecture built from spherical harmonic bases. The angular selection rule structure is a computational resource that belongs to the geometry of angular momentum, not to the Coulomb potential.

C. Non-central interactions

Theorems 1–3 apply to central (rotationally invariant) two-body interactions. Non-central interactions—tensor forces (S_{12}), spin-orbit coupling ($\mathbf{L} \cdot \mathbf{S}$), and velocity-dependent potentials—have modified selection rules. The tensor force, for example, couples states with $|\Delta L| = 0, 2$ and mixes partial waves, expanding the set of allowed transitions relative to the central-force case. For such interactions, the angular sparsity bound of Theorem 3 is a *lower* bound on the ERI density (i.e., the Hamiltonian may be denser than the central-force prediction). The Gaunt selection rules for the central component still apply to that component, but additional nonzero terms appear from the non-central components. A complete sparsity analysis for nuclear Hamiltonians with tensor and spin-orbit forces requires extending the angular classification to include spin-dependent Gaunt-type coefficients, which is beyond the scope of this paper.

VI. UNIVERSAL VS. COULOMB-SPECIFIC STRUCTURE

The GeoVac spectral graph theory framework exhibits two distinct kinds of structure that are sometimes conflated. Theorem 2 and its corollary allow us to draw a sharp line between them.

Universal across spherical fermion systems. The following properties depend only on the angular momentum content of the basis, not on the specific radial potential:

- Radial–angular factorization of two-body matrix elements (Theorem 1).
- Gaunt-selection-rule sparsity $D(l_{\max})$ (Theorem 3).

- $O(Q^4) \to O(Q^{2.5})$ reduction from single-center to composed block architectures (Corollary 1).
- Bit-identical zero patterns across Coulomb, harmonic, Woods–Saxon, square well, and Yukawa potentials (Table **??**).

These properties hold for any central two-body interaction and any radial single-particle potential that produces a complete orbital basis.

Coulomb-specific. The following properties are tied to the $SO(4)$ accidental degeneracy of the hydrogen atom and do not extend to generic central potentials:

- The Fock stereographic projection from momentum space to S^3 that maps the hydrogen Schrödinger equation to an integer eigenvalue problem on the three-sphere [**?**].
- The π-free graph structure in which all one-body eigenvalues, degeneracies, and coupling coefficients are integers or rationals before projection to continuous manifolds.
- The universal constant $K = -1/16$ that maps graph eigenvalues to the Rydberg spectrum.
- The hydrogen shell degeneracy sequence n^2 as $O(n^2)$-dimensional $SO(4)$ irreducible representations.

A companion paper [**?**] establishes that the S^3 Fock projection is rigid in the following sense: modifying the Coulomb potential $-Z/r$ to any other central form breaks the $SO(4)$ symmetry, removes the shell degeneracy, and destroys the integer spectrum. The Coulomb case is the unique central potential for which the one-body Hamiltonian admits a π-free graph Laplacian interpretation.

This partition clarifies the scope of each structural feature. The angular sparsity results of this paper apply broadly across quantum-simulation domains; the S^3 graph Laplacian machinery applies only to Coulomb systems. The two classes of result should be cited and deployed accordingly.

VII. CONCLUSION

The zero structure of two-body matrix elements in a spherical harmonic basis is determined entirely by the Wigner $3j$ symbols (Gaunt coefficients) that encode angular momentum selection rules. This structure is independent of the radial potential that defines the single-particle orbitals, independent of the specific form of the two-body interaction (provided it is rotationally invariant), and independent of the number of radial quantum numbers retained in the basis.

The explicit angular ERI densities obtained by exact enumeration are 7.81% at $l_{\max} = 1$, 2.76% at $l_{\max} = 2$, 1.44% at $l_{\max} = 3$, 0.90% at $l_{\max} = 4$, and 0.62% at $l_{\max} = 5$. Numerical verification across five potentials (Coulomb, harmonic oscillator, Woods–Saxon, square well, Yukawa) at $l_{\max} = 2$ confirms the theorem to machine precision: the angular zero patterns are bit-identical and radial zeros contribute no additional sparsity.

The universal–specific partition is sharp. Angular sparsity, radial–angular factorization, and the $O(Q^{2.5})$ Pauli scaling of composed block architectures are universal across spherical fermion systems. The S^3 Fock projection, the π-free graph Laplacian, and the integer spectrum of the unperturbed hydrogen problem are Coulomb-specific and break under any modification of the radial potential [**?**]. Sparsity-aware quantum simulation techniques developed for electronic structure (qubit tapering, Pauli grouping, tensor factorization) transfer directly to nuclear structure, quantum dots, and other spherical-potential systems without modification, because the angular structure of the spherical harmonic basis is a computational resource that belongs to the geometry of angular momentum, not to any particular physical potential.

[1] A. Peruzzo *et al.*, "A variational eigenvalue solver on a photonic quantum processor," Nat. Commun. **5**, 4213 (2014).

[2] J. Lee *et al.*, "Even more efficient quantum computations of chemistry through tensor hypercontraction," PRX Quantum **2**, 030305 (2021).

[3] J. L. Whitten, "Coulombic potential energy integrals and approximations," J. Chem. Phys. **58**, 4496 (1973).

[4] B. I. Dunlap, N. Rösch, and S. B. Trickey, "Robust and variational fitting," Mol. Phys. **98**, 1639 (2000).

[5] J. Suhonen, *From Nucleons to Nucleus: Concepts of Microscopic Nuclear Theory* (Springer, Berlin, 2007).

[6] E. U. Condon and G. H. Shortley, *The Theory of Atomic Spectra* (Cambridge University Press, Cambridge, 1935).

[7] J. C. Slater, *Quantum Theory of Atomic Structure*, Vol. I (McGraw-Hill, New York, 1960).

[8] A. Messiah, *Quantum Mechanics*, Vol. II (North-Holland, Amsterdam, 1962).

[9] J. Loutey, "The Dimensionless Vacuum: Spectral Graph Theory and Conformal Equivalence on S^3," GeoVac Technical Report (2025).

[10] J. Loutey, "Structurally Sparse Qubit Hamiltonians from Spectral Graph Theory," GeoVac Technical Report (2025).

[11] J. Loutey, "Composed Natural Geometries for Core-Valence Molecular Electronic Structure," GeoVac Technical Report (2025).

[12] J. Loutey, "Exchange Constants and the Transcendental Taxonomy of Observables," GeoVac Technical Report (2026).

[13] J. Loutey, "Fock Rigidity: The S^3 Projection is Coulomb-Specific," GeoVac Technical Report (2026).

[14] S. M. Reimann and M. Manninen, "Electronic structure of quantum dots," Rev. Mod. Phys. **74**, 1283 (2002).