The GeoVac Spectral Triple: A Synthesis Paper Locking the Framework into the Marcolli–van Suijlekom Gauge-Network Lineage

GeoVac Project

May 2026

Abstract

The GeoVac framework has, over Papers 0–31, accumulated a body of structural results—the S^3 Fock-projection equivalence (Paper 7 [30]), the angular sparsity theorem (Paper 22 [33]), the Hopf-graph lattice-gauge structure (Paper 25 [36]), QED on S^3 (Paper 28 [37]), the Bargmann–Segal S^5 HO lattice (Paper 24 [35]), the SU(2) Wilson gauge (Paper 30 [38]), and the universal/Coulomb partition (Paper 31 [39])—each self-contained but together suggesting one parent structure. **[INTERNAL THEOREM]** This paper names and constructs that structure: a finite spectral triple $(\mathcal{A}_{\mathrm{GV}}, \mathcal{H}_{\mathrm{GV}}, D_{\mathrm{GV}})$ on the Fock-projected S^3 graph, with $\mathcal{A}_{\mathrm{GV}}$ the algebra of complex functions on graph nodes, $\mathcal{H}_{\mathrm{GV}}$ the Camporesi–Higuchi spinor space restricted to the truncation, and D_{GV} the diagonal Camporesi–Higuchi Dirac operator [22] on the truncated spinor space of unit S^3 (no graph form is attached to D_{GV}; Remark 6). This is not a new derivation; it is a structural map. Three results follow. **[OBSERVATION]** (1) *Lineage.* The construction sits inside the published Marcolli–van Suijlekom gauge-network framework [6] (with the Perez-Sanchez correction [11] that the continuum limit is Yang–Mills without Higgs), and inside the Connes–van Suijlekom spectral-truncation program [12, 14, 15, 16]. Every structural ingredient of the GeoVac construction has published precedent; what is not duplicated is the α observation ($K = \pi(B + F - \Delta)$ matches α^{-1} to 4.8×10^{-7}; the root of the cubic $1/\alpha = K - \alpha^2$ matches to 8.8×10^{-8}; Paper 2 [29]) from a discrete spectral construction. **[OBSERVATION]** (2) *Sub-sector identification.* Paper 28's QED on S^3 is the spectral-action computation on the spectrum of D_{GV}; Paper 31's universal/Coulomb partition is the $\mathcal{A}/D$ split of the present triple's spectral data. Paper 25's Hodge-1 Laplacian and Paper 30's SU(2) Wilson action are read as the Cartan-torus and full non-abelian slices of an almost-commutative extension by $M_2(\mathbb{C})$, but through the gauge-network (edge-hopping) Dirac of those papers, *not* through D_{GV}: the diagonal D_{GV} commutes with every $\pi(f)$ and so admits no inner fluctuations. The four papers are read as four projections of one structure, at the level of spectral and algebraic data. **[INTERNAL THEOREM]** (3) *Axiom audit.* $\mathcal{A}_{\mathrm{GV}}$ is finite, involutive, faithfully represented on $\mathcal{H}_{\mathrm{GV}}$; D_{GV} is self-adjoint, and since D_{GV} and every $\pi(a)$ are diagonal in the same basis, $[D_{\mathrm{GV}}, \pi(a)] = 0$ identically: the diagonal triple is a *spectral*, not a metric, object (the metric role belongs to the non-diagonal multipliers of the truncated operator system on the *true* Camporesi–Higuchi window — the span of complete Dirac eigenspaces, Paper 38 [61] — whose state spaces are proved to converge under the left-translation Lipschitz seminorm; convergence in the Dirac-commutator seminorm is **[OPEN]**, that seminorm being non-degenerate on the window only as measured at $n_{\max} \leq 3$). A real structure J_{GV} exists on the Dirac sector with $J_{\mathrm{GV}}^2 = -\mathbb{1}$ (the quaternionic/pseudoreal sign for the 2-dimensional Dirac spinor representation of $\mathrm{Spin}(3) = \mathrm{SU}(2)$) and $J_{\mathrm{GV}} D_{\mathrm{GV}} = +D_{\mathrm{GV}} J_{\mathrm{GV}}$, the signs $(\epsilon, \epsilon') = (-1, +1)$ of the continuum S^3 factor (KO-dimension 3 mod 8 [1]). The continuum factor is odd, and so is the true Camporesi–Higuchi window (measured, $n_{\max} \leq 3$: the commutant of its operator system is the scalars), the truncation that converges; only the chirality-doubled Fock/Weyl *model* truncation is even, via

the chirality-doubling grading $\gamma_{\mathrm{GV}} = \sigma_x \otimes \mathbb{1}$, with sign triple $(-,+,+)$ and no KO label attached at finite cutoff. **[MEASURED]** On the Fock/Weyl *model* of the operator system (the doubled computational object of this paper) the order-zero and order-one conditions fail at finite $n_{\max}$, with residuals that *grow* over the measured cutoffs (order zero at $n_{\max} = 2, 3, 4$, order one at $n_{\max} = 2, 3$; mechanism open). **[MEASURED]** On the true Camporesi–Higuchi window that converges, the charge conjugation $J\psi = (i\sigma_y)\overline{\psi}$ satisfies $J^2 = -\mathbb{1}$, $JD = +DJ$ and $J\mathcal{O}J^{-1} = \mathcal{O}$ exactly, while order zero and order one fail from $n_{\max} = 1$ with residuals that grow toward the sup-norm scale of the generators; there every residual is a Toeplitz boundary term of the compression, confined to the top Dirac levels of the window. **[OBSERVATION]** The four-way S^3 coincidence—Fock projection image, Hopf bundle total space S^3 (base S^2), Camporesi–Higuchi spin carrier, SU(2) gauge manifold—is the statement that the manifold reached by the operator-system truncations in the continuum limit, the spinor bundle of the Camporesi–Higuchi Dirac, and the gauge group of the SU(2) almost-commutative extension (Papers 25/30) all coincide on the unique rank-1 non-abelian compact simply-connected Lie group.

Scope. This paper does not derive new physics. It synthesizes existing GeoVac results into the axiomatic spectral-triple language. The synthesis lifts each result from a free-standing observation to a sub-sector of a single named structure, and makes explicit which axioms hold rigorously, which hold up to the standard finite-truncation caveats, and which remain open. **[PANEL-VERIFIED]** The real-structure question—whether the GeoVac Dirac operator D_{GV} (the diagonal spectral form) carries a real structure matching the signs of S^3 exactly at finite $n_{\max}$ or only in the continuum limit—is resolved in favour of finite-$n_{\max}$ exactness *for the three J-relations* $J^2 = -\mathbb{1}$, $JD = +DJ$, $J\mathcal{O}J^{-1} = \mathcal{O}$ only, measured on the Fock/Weyl model objects and, independently, on the true Camporesi–Higuchi window. **[MEASURED]** The order-zero and order-one conditions fail at finite $n_{\max}$ with residuals that grow with the cutoff, on the Fock/Weyl model of the operator system and on the true window alike (on the true window as a boundary term of the compression). The Fock/Weyl model truncation is even (chirality doubling; sign triple $(-,+,+)$, no KO label); the true window is odd (measured, $n_{\max} \leq 3$), like the continuum (its operator system generates all of $B(\mathcal{H})$, so it admits no grading), and there the pair $(\varepsilon, \varepsilon') = (-,+)$ is forced by equivariance; the residual open items are catalogued in Sec. 10.

1 Introduction

1.1 Why a spectral triple, and why now

Across the thirty-two papers (0–31) preceding this one, the GeoVac framework has carried four distinct strands. Strand 1 is the Fock projection $p_0 = \sqrt{-2E_n}$ that maps the hydrogenic Hamiltonian onto a function of the Laplace–Beltrami operator Δ on the unit three-sphere, $H = -\frac{Z^2}{2}(1-\Delta)^{-1}$, where Δ has continuum eigenvalues $\lambda_n = -(n^2-1)$; the graph nodes carry the quantum-number labels (n, l, m_l), not that spectrum (Paper 7). Strand 2 is the Hopf bundle $S^1 \to S^3 \to S^2$ that organizes the discrete edge structure into ladder operators with $U(1)$ phase content (Papers 2, 25). Strand 3 is the Camporesi–Higuchi Dirac spectrum $|\lambda_n^{\mathrm{D}}| = n + 3/2$ with degeneracy $g_n^{\mathrm{D}} = 2(n+1)(n+2)$ that controls QED on S^3 (Paper 28). Strand 4 is the Wilson lattice gauge structure with SU(2) link variables on the same graph (Paper 30).

Each strand uses the unit three-sphere in a different role:

(a) S^3 as the *conformal image* of the hydrogenic spectrum (Paper 7);

(b) S^3 as the *total space* of the Hopf bundle on which Paper 2's α Observation is phrased;

(c) S^3 as the *spin carrier* of the Camporesi–Higuchi Dirac operator (Paper 28);

(d) S^3 as the *gauge manifold* of SU(2) Wilson links (Paper 30).

[**OBSERVATION**] Working hypothesis WH4 in the project register flags this as the strangest and most consequential coincidence in the framework: $S^3 = \mathrm{SU}(2)$ is the unique rank-1 non-abelian compact simply connected Lie group where (a)–(d) all coincide; in any other dimension or any other rank, the four roles separate. WH4 takes the position, in its current (deflated) form, that the four strands are not four coincidences but one Fock-projection statement plus three forced consequences of $S^3 = \mathrm{SU}(2)$ (its parallelizability and maximal-torus structure).

This paper makes WH4 explicit. We give a concrete construction of a finite spectral triple $(\mathcal{A}_{\mathrm{GV}}, \mathcal{H}_{\mathrm{GV}}, D_{\mathrm{GV}})$ (Sec. 3) — a spectral, not metric, object, since its diagonal D_{GV} commutes with the diagonal algebra, the metric role belonging to the operator-system representative on the true Camporesi–Higuchi window (Paper 38); audit the standard Connes axioms one by one, separating what holds at finite cutoff (the three J-relations; oddness on the true window, which the chirality-doubled model replaces by an even grading) from what holds only in the continuum limit (order zero and order one, which fail at finite cutoff on the Fock/Weyl model of the operator system and on the true window) (Sec. 4); read Papers 25, 28, 30, and 31 as structural sub-sectors of the resulting structure (Sec. 5); read the Paper 2 combination formula $K = \pi(B + F - \Delta)$ inside the spectral-action language (Sec. 6); and place the Coulomb/HO asymmetry of Paper 31 in spectral-triple terms (Sec. 7).

1.2 What this paper is and is not

What it is. A synthesis paper. The construction is explicit, finite, and reproducible from existing GeoVac modules (`geovac/lattice.py`, `geovac/dirac_matrix_elements.py`, `geovac/fock_graph_hodge.py`); the lineage citations are to published mathematical work; the audit of the Connes axioms uses standard textbook definitions [1, 2, 3].

What it is not. A derivation of α. [**OBSERVATION**] Paper 2 stays where it is: a structural observation with three independent spectral homes for the three pieces B, F, Δ but no first-principles derivation of the combination rule. This paper does not change that status; it gives the spectral-action language in which the open question is phrased without promoting it beyond that observation.

What it is not, second. A claim of the Connes–Chamseddine Standard Model. The GeoVac construction is a *rank-1* non-abelian almost-commutative extension; the SM spectral triple of [5] is rank ≥ 3 with specific finite-algebra content $\mathbb{C} \oplus \mathbb{H} \oplus M_3(\mathbb{C})$. GeoVac is not the Standard Model. It is a finite spectral triple that is an *instance* of Marcolli–van Suijlekom 2014 [6] (with Perez-Sanchez 2025 continuum-limit correction [11]) at the data level and a GeoVac-original *extension* at the structural level (forced graph from Bertrand × Fock projection, master Mellin engine, explicit Hopf algebra $\mathcal{H}_{\mathrm{GV}}$, Tannakian dual U^*), [**OBSERVATION**] with the empirically interesting property that a combination of three invariants drawn from different sectors of the triple, $K = \pi(B+F-\Delta)$, satisfies, after subtraction of the leading Sprint-A correction α^2, a residual $K - 1/\alpha - \alpha^2 \approx 1.2 \times 10^{-5}$ corresponding to a post-correction relative residual of 8.8×10^{-8} that matches $\pi^3\alpha^3$ to within 0.25% (the raw match $|K - 1/\alpha|/(1/\alpha)$ is 4.77×10^{-7}).

Internal register. This paper makes explicit the spectral-triple framing already operating informally in the working hypothesis register (kept as §1.7 of the repository's `CLAUDE.md`; the sprint, track and door codenames used throughout this paper are chronicle anchors into that file and the repository `CHANGELOG.md`; WH1's commutative-factor leg — state-space convergence of the round-S^3 truncations on the true window — proved unconditionally as of 2026-06-10, Paper 38 [61];

convergence of the almost-commutative structure is [**OPEN**]). Papers continue under the rhetoric rule of §1.5: dual-description framing, no ontological priority claims, lead with computational results.

2 The Connes Data Recalled

A spectral triple is a triple $(\mathcal{A}, \mathcal{H}, D)$ where [1]

- $\mathcal{A}$ is an involutive (unital) algebra,
- $\mathcal{H}$ is a Hilbert space carrying a faithful $*$-representation $\pi : \mathcal{A} \to \mathcal{B}(\mathcal{H})$,
- $D : \mathcal{H} \to \mathcal{H}$ is an unbounded self-adjoint operator with compact resolvent, such that the commutators $[D, \pi(a)]$ are bounded for every $a \in \mathcal{A}$.

The triple is called *even* if there is a self-adjoint involution γ commuting with $\pi(\mathcal{A})$ and anticommuting with D; otherwise *odd*. A *real structure* on the triple is an antilinear isometry $J : \mathcal{H} \to \mathcal{H}$ with $J^2 = \epsilon \mathbb{1}$, $JD = \epsilon' DJ$, $J\gamma = \epsilon'' \gamma J$ (the signs $(\epsilon, \epsilon', \epsilon'')$ are determined by the KO-dimension mod 8 [1]); for the canonical Riemannian spin manifold of dimension n, the KO-dimension is n mod 8 and the signs are tabulated.

The two structural axioms of central interest below are:

(**O1**) *Order one:* $[[D, \pi(a)], \pi(b)^\circ] = 0$ for all $a, b \in \mathcal{A}$, where $b^\circ = JbJ^{-1}$ is the right action induced by the real structure;

(**R**) *Reality:* J exists with the signs above.

For an *almost-commutative* spectral triple $(\mathcal{A}_M \otimes \mathcal{A}_F, \mathcal{H}_M \otimes \mathcal{H}_F, D_M \otimes \mathbb{1} + \gamma_M \otimes D_F)$ [3, 5], the manifold sector $\mathcal{A}_M = C^\infty(M)$ is replaced in our finite setting by the algebra of complex-valued functions on a finite graph, with the convergence to $C^\infty(M)$ carried by the true-window operator-system representative of Theorem 5 (the diagonal algebra itself is the gauge-network convention; see Remark 1). This is exactly the move made by Marcolli and van Suijlekom in their gauge networks construction [6] and continued by Perez-Sanchez [11]: replace the manifold algebra by an algebra on a discrete vertex set, with edges carrying connection data. The GeoVac construction sits in this lineage by design.

3 The GeoVac Triple: Explicit Construction

Fix a truncation $n_{\max} \geq 1$. All objects below are finite.

3.1 The algebra $\mathcal{A}_{\mathrm{GV}}$

Let V_{Fock} denote the set of Fock-graph nodes at truncation $n_{\max}$, i.e. quantum-number triples (n, l, m_l) with $1 \leq n \leq n_{\max}$, $0 \leq l \leq n-1$, $-l \leq m_l \leq l$. The total node count is $N_{\mathrm{Fock}}(n_{\max}) = \sum_{n=1}^{n_{\max}} n^2$.

Definition 1 (Function algebra on the Fock graph)**.** $\mathcal{A}_{\mathrm{GV}}^{(n_{\max})} := \mathbb{C}^{V_{\mathrm{Fock}}}$*, the commutative* $*$*-algebra of complex-valued functions on* V_{Fock} *under pointwise multiplication, with involution* $f^*(v) = \overline{f(v)}$*.*

This is precisely the manifold algebra of the gauge networks of [6] (their $C(V)$, with V the vertex set), with the GeoVac-specific choice that V is the Fock-projected S^3 graph of Paper 7 rather than a generic graph. It is finite-dimensional ($\dim_{\mathbb{C}} \mathcal{A}_{\mathrm{GV}} = N_{\mathrm{Fock}}$), unital (1 is the constant function 1), and abelian.

Remark 1 ($\mathcal{A}_{\mathrm{GV}}$ as the diagonal representative of an operator system)**.** ***[OBSERVATION]*** *In the parallel reading of GeoVac through the spectral-truncation framework of Connes and van Suijlekom [12], Definition 1 describes $\mathcal{A}_{\mathrm{GV}}^{(n_{\max})} = \mathbb{C}^{V_{\mathrm{Fock}}}$ as the diagonal (gauge-network) representative attached to the operator system (its C^*-envelope is $M_N(\mathbb{C})$, since* prop $= 2$ *gives $\mathcal{O}^2_{n_{\max}} = M_N(\mathbb{C})$; corrected 2026-09-03)*

$$\mathcal{O}_{n_{\max}} := P_{n_{\max}}\, C^\infty(S^3)\, P_{n_{\max}},$$

where $P_{n_{\max}}$ is the index-truncation projecting onto the span of the first $n_{\max}$ Fock shells (in the spinor-harmonic basis of $L^2(S^3)$ adapted to the SO(4) decomposition). This is structurally identical to the Connes–van Suijlekom worked example on S^1 (their P_n onto $\mathrm{span}_{\mathbb{C}}\{e_1, e_2, \ldots, e_n\}$*, [12, §3.1]): a basis-index cutoff, not a Borel functional-calculus projection $\chi_{[-\Lambda,\Lambda]}(D)$. The genuine truncated algebra-image $\mathcal{O}_{n_{\max}}$ is $*$-closed but* not *multiplicatively closed; it carries nontrivial off-diagonal matrix elements*

$$\langle (n,l,m_l)|\, f\, |(n',l',m_l')\rangle = \int_{S^3} \overline{Y_{nlm_l}}\, f\, Y_{n'l'm_l'}\, dV$$

between distinct n-shells. These overlap integrals are exactly the Gaunt / Wigner-3j angular machinery and the hypergeometric Slater radial integrals already implemented throughout the GeoVac pipeline (Paper 22 angular sparsity; Paper 14 ERI structure). The off-diagonal data therefore already exists in the framework; $\mathcal{A}_{\mathrm{GV}}$ as written in Definition 1 is its diagonal restriction.

The two readings agree on diagonal elements and on the spectral data of D_{GV}; they differ in what is taken as the algebraic content surviving the truncation. Connes and van Suijlekom note that taking the C^*-envelope of $\mathcal{O}_\Lambda$ typically collapses to a "non-informative full matrix algebra" – the algebraic-structure information lives in the operator-system data thrown away by the envelope construction. Constructing $\mathcal{O}_{n_{\max}}$ explicitly in $\mathcal{B}(\mathcal{H}_{\mathrm{GV}})$ and computing its propagation number and Connes distance on node-evaluation states is the deliverable of the WH1 round-2 sprint (see `debug/wh1_connes_vs_mapping_memo.md`*, 2026-05-03). The present subsection retains the $\mathcal{A}_{\mathrm{GV}} = \mathbb{C}^{V_{\mathrm{Fock}}}$ definition because (a) it matches the Marcolli–van Suijlekom gauge-network convention used throughout Papers 25 and 30, and (b) it carries the spectral data (spectrum, degeneracies, real structure). The two representatives are* not *interchangeable for axiom-level or metric content: on the diagonal representative $[D_{\mathrm{GV}}, \pi(a)] = 0$, so it carries no metric and the order-zero/one conditions hold vacuously, whereas on $\mathcal{O}_{n_{\max}}$ the commutators are nonzero and the order-zero/one conditions fail at finite $n_{\max}$ (Sec. 4).*

Which operator system converges (2026-09-28). *The $\mathcal{O}_{n_{\max}}$ defined above — the first $n_{\max}$ Fock shells, and its Weyl and chirality-doubled spinor lifts (`SpinorTruncatedOperatorSystem`, `FullDiracTruncatedOperatorSystem`) — is a* computational model*, and every finite-cutoff axiom measurement in this paper (propagation number, Connes distances, the three J-relations, the order-zero/one residuals) is made on it, except the true-window real-structure audit (item 6 of Sec. 4). The object whose continuum limit is proved (Theorem 5, Paper 38 [61]) is a different one: the truncation to the* true *Camporesi–Higuchi window, the span of the complete Dirac eigenspaces with $|\lambda| \le n_{\max} + \frac{1}{2}$, whose operator system $\mathcal{O}^{\mathrm{CH}}_{n_{\max}}$ has dimension $5, 30, 91$ at $n_{\max} = 1, 2, 3$ (`tests/test_p38_true_ch_window.py`). The doubled model has the same labels and Hilbert-space dimensions but is not $\mathrm{SU}(2)_L$-invariant, omits the blocks coupling positive and negative Dirac levels,*

and reaches only the bands $N \le 2n_{\max} - 1$; its quantities are model quantities, not window quantities (Paper 38, § Truncations). On the true window the Dirac-commutator kernel is exactly the scalars at $n_{\max} \le 3$ (Paper 38, remark on the Dirac-commutator seminorm), and the real-structure audit is repeated on it in item 6 of Sec. 4; its propagation number is not measured.

Proposition 1 (Propagation number of $\mathcal{O}_{n_{\max}}$)**.** ***[PANEL-VERIFIED]*** *Let $\mathcal{O}_{n_{\max}} = P_{n_{\max}} C^\infty(S^3) P_{n_{\max}}$ be the truncated operator system of Remark 1, viewed as a $*$-closed subspace of $M_N(\mathbb{C})$ with $N = N_{\mathrm{Fock}}(n_{\max})$. The propagation number in the sense of Connes & van Suijlekom ([12], Definition 2.39) satisfies*

$$\mathrm{prop}(\mathcal{O}_{n_{\max}}) = 2 \qquad \text{for every } n_{\max} \in \{2, 3, 4\}, \tag{1}$$

matching the Toeplitz operator system $C(S^1)^{(n)}$ ([12], Proposition 4.2: $\mathrm{prop}(C(S^1)^{(n)}) = 2$ independently of n) verbatim. Equivalently, $\dim_{\mathbb{C}}(\mathcal{O}_{n_{\max}}^2) = N^2$ while $\dim_{\mathbb{C}}(\mathcal{O}_{n_{\max}}) < N^2$ strictly:

$n_{\max}$	N	N^2	$\dim(\mathcal{O}_{n_{\max}})$	$\dim(\mathcal{O}_{n_{\max}}^2)$
2	5	25	14	25
3	14	196	55	196
4	30	900	140	900

The dimension fraction $\dim(\mathcal{O}_{n_{\max}})/N^2$ falls $0.560 \to 0.281 \to 0.156$ across the three truncations: the operator system is a vanishing fraction of its C^*-envelope as $n_{\max}$ grows, but its second power fills the envelope completely. (This is the scalar Fock-shell model of Remark 1, not the true Camporesi–Higuchi window $\mathcal{O}^{\mathrm{CH}}_{n_{\max}}$ of the convergence theorem, whose propagation number is not measured here.)

Computational verification. The full construction and verification is documented in `geovac/operator_system.py` (the `TruncatedOperatorSystem` class) and `tests/test_operator_system.py` (24 tests, all passing; CPU time ∼20 s). In summary: a basis of $\mathcal{O}_{n_{\max}}$ is built from the multiplier matrices M_{NLM} corresponding to the hyperspherical harmonics $Y^{(3)}_{NLM}$, with selection rules $|n-n'|+1 \le N \le n+n'-1$ (SO(4) triangle), $|l-l'| \le L \le l+l'$ with $l+l'+L$ even (S^2 Gaunt parity), $M = m-m'$, and the angular S^2 part evaluated symbolically via `sympy.physics.wigner.wigner_3j`. Linear independence of $\{M_{NLM}\}$ is verified at machine precision; the propagation number is read off as the smallest k with $\dim(\mathcal{O}^k_{n_{\max}}) = N^2$. The result prop = 2 is robust: it is invariant under the choice of SO(4) radial-overlap placeholder (verified across three placeholders including a pure indicator that retains only the support pattern), so (1) is a structural invariant of the SO(4) selection rules at finite cutoff and does not depend on the radial value convention. See `debug/wh1_round2_propagation_memo.md` (2026-05-03) for the full computational record, including the explicit witness pair $a = M_{2,1,0}$ at $n_{\max} = 2$ for which $a^2 \notin \mathcal{O}_{n_{\max}}$ (Frobenius residual $\approx 14.9\%$), with the deficit concentrated on the top-shell diagonal entries. □

Remark 2 (Significance of prop = 2)**.** *Connes & van Suijlekom prove $\mathrm{prop}(C(S^1)^{(n)}) = 2$ for the Toeplitz operator system on the circle by showing that products of two Toeplitz shift generators reach every matrix unit e_{ij} [12]. The companion dual operator system $C(\mathbb{Z})^{(n)}$ has prop $= \infty$ (their Proposition 4.3) — so the value prop $= 2$ is non-trivial even at this level: it separates the Toeplitz construction from its dual. Eq. (1) states that the Fock-projected S^3 truncation is in the same propagation class as the Toeplitz S^1 at the level of a quantitative Connes–van Suijlekom invariant.* ***[OBSERVATION]*** *The value should not be over-read:* prop $= 2$ *is the* generic *value for a $*$-closed unital subspace of $M_N(\mathbb{C})$ of the same complex dimension (random such subspaces of dimension* 14 *in M_5 and* 55 *in M_{14} all have* prop $= 2$*; 2026-09-03), so the invariant certifies that $\mathcal{O}_{n_{\max}}$ is not an*

abelian or otherwise degenerate comparator, not that it is special among truncations. The strongest quantitative alignment between GeoVac and the published Connes–van Suijlekom framework is the convergence theorem in van Suijlekom's own distance (Theorem 5), not this invariant.

The mechanism producing $\mathcal{O}_{n_{\max}} \subsetneq M_N(\mathbb{C})$ is the "raise-then-lower at the top shell escapes the truncation" boundary correction: a multiplier $M_{N\geq 2,L,M}$ acting at the top shell $n = n_{\max}$ would, in the continuum, be balanced by intermediate sums over $n' > n_{\max}$, but the truncation $P_{n_{\max}}$ kills these contributions. This is structurally the same boundary effect as the pendant-edge theorem of Paper 28 [37], $\Sigma_{\text{graph}}(\text{GS}) = 2(n_{\max}-1)/n_{\max}$ for the graph-native ground-state self-energy, viewed from the operator-system side rather than the QED side.

Corollary 1 (prop = 2 separates $\mathcal{O}_{n_{\max}}$ from abelian comparators)**.** ***[PANEL-VERIFIED]*** *The propagation value* $\text{prop}(\mathcal{O}_{n_{\max}}) = 2$ *in Proposition 1 separates the Connes–van Suijlekom spectral truncation of S^3 from the abelian (point-evaluation) discretizations of S^3; it does* not *single it out among non-abelian truncations, for which* prop = 2 *is generic (retitled 2026-09-03 from "structural specificity"). The natural circulant-style comparator on S^3 at N points*

$$A_{\text{circ},N} := \text{span}\{E_{i,i} : i = 0, \ldots, N-1\} \subset M_N(\mathbb{C}),$$

i.e., the commutative C^*-subalgebra of diagonal matrices in the "point-evaluation" basis of $\mathbb{C}^N$, satisfies

$$\text{prop}(A_{\text{circ},N}) = \begin{cases} 1 & \text{(intrinsic envelope: } A_{\text{circ},N} \text{ is its own } C^*\text{-algebra)}, \\ \infty & \text{(ambient envelope: } M_N(\mathbb{C}) \text{ is non-abelian; no power of an abelian subalgebra reaches it).} \end{cases} \tag{2}$$

Verified at $N \in \{2, 5, 14, 30, 55, 140\}$, matching both dimension-matching conventions (envelope-match $N(\text{circ}) = N(\text{GV})$ and operator-system-match $N(\text{circ}) = \dim(\mathcal{O}_{n_{\max}})$). Either reading of (2) *is categorically different from the spectral-truncation value* 2*.*

Computational verification. The construction lives in `geovac/circulant_s3.py` (the `CirculantS3Truncation` class) with verification in `tests/test_circulant_s3.py` (35 tests, all pass; runtime ∼1.4 s). Algorithmically, $\text{prop}(A_{\text{circ},N})$ is computed via the same vec-stack rank algorithm used in the proof of Proposition 1, applied to the diagonal-matrix basis of $A_{\text{circ},N}$. The intrinsic-envelope reading exits at $k = 1$ since $\dim(A^1_{\text{circ},N}) = N = \dim_{\mathbb{C}}(A_{\text{circ},N})$ already saturates the intrinsic C^*-envelope. In the ambient-envelope reading, the $M_N(\mathbb{C})$ envelope has dimension N^2 for $N \geq 2$ but $\dim(A^k_{\text{circ},N}) = N$ for every k (commutativity is preserved under self-products), so no finite k achieves saturation; prop $= \infty$ by Connes–van Suijlekom Definition 2.39. See `debug/wh1_round3_falsification_memo.md` (2026-05-03) for the full record, including the verbatim Connes–van Suijlekom Outlook §6 contrast (lines 2611–2618 of arXiv:2004.14115v2) between spectral and circulant truncations on S^1, of which Eq. (2) is the verbatim S^3 analog. □

Remark 3 (Connes distance on $\mathcal{O}_{n_{\max}}$: SDP framework and structural findings)**.** ***[MEASURED]*** *The Connes distance on operator systems ([12], Eq. (2))*

$$d_D(\phi, \psi) = \sup_{x \in \mathcal{O}} \big\{ |\phi(x) - \psi(x)| : \|[D, x]\|_{\text{op}} \leq 1 \big\}$$

restricted to pure node-evaluation states $\phi_v(x) := \langle v|x|v\rangle$ on the truncated operator system has been computed by semidefinite programming on $\mathcal{O}_{n_{\max}}$ at $n_{\max} = 2, 3$. The implementation lives in `geovac/connes_distance.py` *(a* `cvxpy` *SDP with the operator-norm constraint lifted to the LMI $\left[\begin{smallmatrix} I & [D,x] \\ [D,x]^* & I \end{smallmatrix}\right] \succeq 0$, and a gauge-fixing constraint orthogonalizing x to $\ker([D, \cdot]) \cap \mathcal{O}_h$); 17 tests in* `tests/test_connes_distance.py` *(all pass; runtime ∼9 s). Three structural findings emerged at the placeholder convention:*

1. ***SO(3) m-reflection forces zero distance.*** *For every $|n, l, -m\rangle$ and $|n, l, +m\rangle$ pair, $d_D(\phi_{|n,l,-m\rangle}, \phi_{|n,l,+m\rangle})$ 0 exactly (one such pair at $n_{\max} = 2$, four at $n_{\max} = 3$). This is the SO(3) m-reflection symmetry of the operator system showing through, not a pathology: the symmetry identifies the two states as members of the same orbit, and any point-discriminating function must be constant on the orbit.*

2. ***Rational structure at $n_{\max} = 2$ under placeholder multipliers (retracted as a placeholder artifact, Remark 4).*** *Under the placeholder values the five non-zero entries of the 5×5 distance matrix at $n_{\max} = 2$ grouped into ratios $1 : 2 : 3$ (specifically $28.87 : 57.73 : 86.60$), with $86.60 \approx 50\sqrt{3} = 86.6025$. This rational/algebraic structure is lost at $n_{\max} = 3$ as the SDP gains degrees of freedom. Under physical values the 1:2:3 grouping does not persist, though the $50\sqrt{3}$ entry does (Remark 4).*

3. ***Non-monotonicity in the Fock-graph distance.*** *Pearson correlation between $d_D(\phi_v, \phi_w)$ and the combinatorial Fock-graph distance $d_{\text{graph}}(v, w)$ (counting $\Delta n = \pm1$ and $\Delta l = \pm1$ ladder steps) is $\rho \approx -0.04$ at $n_{\max} = 2$, growing to $+0.14$ at $n_{\max} = 3$ (Spearman $0.09 \to 0.18$) under placeholder values; under physical values the sign flips (Remark 4), and the absolute level is far from a discretized-geodesic metric at our reachable cutoffs.*

These findings should be read with three explicit caveats, the first two of which are closed in Remark 4.

- *The SO(4) radial overlap integral $I^{lLl'}_{nNn'}$ in the construction of $\mathcal{O}_{n_{\max}}$ (Remark 1) is currently a placeholder rather than the genuine Avery–Wen–Avery 3-Y integral on S^3. Round 2's propagation-number result (Proposition 1) is invariant under this choice; the Connes distance is not. The absolute distance values reported above are convention-dependent until the placeholder is replaced.*

- *The construction is on the scalar Fock basis $\mathcal{H}_{\text{GV}}$ only; the spinor lift to $\mathcal{H}_{\text{GV}} \otimes \mathbb{C}^4$ with the native Camporesi–Higuchi Dirac in the $(n_{\text{D}}, \kappa, m_j)$ basis is not yet incorporated. The diagonal scalar Dirac on $\mathcal{H}_{\text{GV}}$ has a large $\ker([D, \cdot]) \cap \mathcal{O}$ that makes the SDP unbounded on most state pairs; the metric values reported here use an off-diagonal Dirac proxy as a finite-distance regulator, which is a second convention choice.*

- *No comparison to the round-S^3 continuum geodesic distance is attempted. Such a comparison is the actual content of Connes–vS Gromov–Hausdorff convergence (Sec. 4 discussion of Row 9 of the round-1 mapping). The scalar case is published: Gaudillot-Estrada–van Suijlekom [17] prove spectral-truncation convergence for all compact metric groups, including $\text{SU}(2) \cong S^3$; the spinor case — on the true Camporesi–Higuchi window of complete Dirac eigenspaces, not the chirality-doubled Fock model used in this remark — is the contribution of Theorem 5 below and Paper 38.*

The SO(3) m-reflection forced zeros (and the $50\sqrt{3}$ fingerprint) are robust across the placeholder choice; the rational grouping at $n_{\max} = 2$, the absolute distance values and the non-monotonicity diagnostic are not (Remark 4). See `debug/wh1_round3_connes_distance_memo.md` for the full record. Caveats (1) and (2) above are subsequently closed in Remark 4, where two of the three structural findings (the 1:2:3 grouping and the trend-toward-correlation reading) are retracted as placeholder artifacts.

Remark 4 (R3.1+R3.2 closure of round-3 caveats)**.** ***[OBSERVATION]*** *Two of the three caveats above have been closed in subsequent sub-sprints (continuation of the same dispatch on 2026-05-03), and two of the three structural findings (the* 1:2:3 *grouping and the trend-toward-correlation reading) have been retracted as placeholder artifacts.*

R3.1 (Avery–Wen–Avery integral, placeholder → physical). *The placeholder for* $I^{lLl'}_{nNn'}$ *is replaced with the closed-form Avery–Wen–Avery* 3-Y *integral on* S^3*, factorized as* $I_{\rm rad} \cdot G_{\rm Gaunt}$ *with* $I_{\rm rad}$ *the Gegenbauer triple integral on* χ *and* $G_{\rm Gaunt}$ *the standard* S^2 *Gaunt integral. Implemented in* `geovac/so4_three_y_integral.py` *(sympy-exact,* `lru_cache`*), verified by hyperspherical orthonormality at* $n_{\max} \le 3$ *in exact arithmetic.* $\mathrm{prop}(\mathcal{O}_{n_{\max}}) = 2$ *holds at* $n_{\max} = 2, 3, 4$ *under the physical default with dim sequences* $14 \to 25,\ 55 \to 196,\ 140 \to 900$ bit-identical *to the placeholder values—confirming the round-2 invariance claim by replacement, not by sampling.*

The Connes distance recomputed under physical values retains the m*-reflection forced zeros and the* $50\sqrt{3}$ *fingerprint, but* retracts *the* $1:2:3$ *rational structure as a placeholder artifact: under Avery the distinct nonzero values at* $n_{\max} = 2$ *are* $\{1.27, 86.60, 87.18, 174.93, 175.53, 262.11\}$ *with no clean integer grouping. The* $M_{2,0,0}$ *multiplier rescales by* $\sim 3\times$ *between placeholder and physical values, propagating as* $\sim 9\times$ *on pairs that use it twice. More importantly, the non-monotonicity* flips sign *at* $n_{\max} = 3$*: Pearson goes from* $+0.14$ *(placeholder) to* -0.36 *(physical), Spearman from* $+0.18$ *to* -0.33*. The optimistic "slow trend toward correlation" reading is retracted; under physical values the metric is* anti-correlated *with the Fock-graph distance. Memo:* `debug/wh1_r31_avery_wen_avery_memo.md`*.*

R3.2 (spinor lift, scalar Fock → Weyl spinor on S^3**).** *The construction extends to the Weyl sector of the Camporesi–Higuchi spinor bundle on* S^3 *via Clebsch–Gordan decomposition* $|\psi_{l,j=l+1/2,m_j}\rangle = \alpha_+ |Y_{l,m_j-1/2}\rangle \otimes |\uparrow\rangle + \alpha_- |Y_{l,m_j+1/2}\rangle \otimes |\downarrow\rangle$ *with* $\alpha_\pm = \sqrt{(l \pm m_j + 1/2)/(2l+1)}$*. Each scalar Avery multiplier* M_{NLM} *lifts to a CG-weighted sum of two scalar matrix elements; the resulting* $\widetilde{\mathcal{O}}_{n_{\max}}$ *is API-compatible with the existing SDP machinery. Implemented in* `geovac/spinor_operator_system.` $\dim \widetilde{\mathcal{O}} = \dim \mathcal{O}$ *at every* $n_{\max}$ *(the CG decomposition introduces no new linear independencies); identity,* $*$*-closure, and* m_j*-reflection forced zeros all generalize naturally.*

Two clean R3.2 findings. ***(i)*** The truthful Camporesi–Higuchi Dirac on the spinor bundle is too n-degenerate *for the Connes distance to be well-defined on cross-shell pure-state pairs: at* $n_{\max} = 2$*,* 24 *of* 28 *cross-pairs give* $+\infty$ *because the* n*-shell-diagonal multipliers* $M_{N,L=\mathrm{even},M=0}$ *commute with* $D = \mathrm{diag}(n_{\rm Fock} + 1/2)$*, making the SDP unbounded. This is the spinor analog of the round-3* `shell_scalar` *diagnostic. It is a property of the Fock/Weyl spinor* model*, not of the truncated Camporesi–Higuchi triple: on the true window (complete Dirac eigenspaces) the Dirac-commutator kernel is exactly the scalars at* $n_{\max} \le 3$ *(Paper 38, remark on the Dirac-commutator seminorm;* `tests/test_p38_true_ch_window.py`*). An earlier version called the* n*-degeneracy "structural" (withdrawn 2026-09-28).* ***(ii)*** Under a CH+offdiag perturbation that lifts the n-degeneracy, the metric remains anti-correlated *with the Fock-graph distance: Pearson nz* -0.36 *at* $n_{\max} = 2$*,* -0.26 *at* $n_{\max} = 3$*. The non-monotonicity is therefore robust under both the placeholder → Avery upgrade (R3.1)* and *the scalar → spinor lift (R3.2). Memo:* `debug/wh1_r32_spinor_lift_memo.md`*.*

Disambiguation settled. *The round-3 §5.4 two-reading question ("placeholder artifact" vs "pure-state-distance pathology") lands on the latter. R3.1 rules out the placeholder reading (anti-correlation* strengthens *under physical values). R3.2 rules out the scalar-Dirac-proxy reading (anti-correlation persists under the spinor lift). The Connes distance on* $\mathcal{O}_{n_{\max}}$ *or its spinor lift, with pure node-evaluation states, is structurally not a discretization of the round-*S^3 *geodesic distance at any cutoff*

we can reach. The third caveat above (no GH comparison on the full state space) remains the natural next move; whether the metric becomes monotone in the GH limit via a Kantorovich–Wasserstein lift to mixed states requires Peter–Weyl on SU(2) *(genuinely new NCG mathematics; deferred by Connes & van Suijlekom 2021 three times). The S^3 GH convergence itself is no longer unpublished: the scalar case is closed by Gaudillot-Estrada–van Suijlekom [17] for all compact metric groups including* SU(2)*, and the spinor case, on the true Camporesi–Higuchi window (not the chirality-doubled Fock/Weyl model of this remark), by Theorem 5 and Paper 38; the Kantorovich–Wasserstein monotonicity question remains open.*

Remark 5 (Two-sided structural alignment)**.** ***[OBSERVATION]*** *Connes & van Suijlekom's worked S^1 example exhibits the spectral-vs-circulant distinction along two axes simultaneously: the Toeplitz operator system $C(S^1)^{(n)}$ has* prop $= 2$ *(their Proposition 4.2), while the circulant matrix algebra $C(C_m)$ has* prop $= 1$ *trivially (it is its own C^*-algebra). Eqs.* (1) *(Proposition 1) and* (2) *(Corollary 1) reproduce both axes verbatim on S^3:*

	spectral truncation	*circulant analog*
S^1 *(Connes–vS)*	$C(S^1)^{(n)}$, prop $= 2$	$C(C_m)$, prop $= 1$
S^3 *(this work)*	$\mathcal{O}_{n_{\max}}$, prop $= 2$	$A_{\mathrm{circ},N}$, prop $= 1$ *or* ∞

The two-sided match separates spectral truncation from abelian discretization on both manifolds; it does not foreclose the generic non-abelian value (Remark above). Together with the SO(4)-isometry-preservation property of Sec. 4 below (the property Connes & van Suijlekom themselves identify as distinguishing spectral truncation from circulant discretization), it is a consistency check on the construction; the load-bearing alignment is Theorem 5.

3.2 The Hilbert space $\mathcal{H}_{\mathrm{GV}}$

Let V_{Dirac} denote the set of Dirac-graph nodes at the matching truncation, labeled by Camporesi–Higuchi triples $(n_{\mathrm{D}}, \kappa, m_j)$ with $0 \le n_{\mathrm{D}} \le n_{\max} - 1$ (the standard convention $n_{\mathrm{Fock}} = n_{\mathrm{CH}} + 1$ from `geovac/dirac_matrix_elements.py`), Dirac–Coulomb relativistic angular momentum $\kappa \in \{-(l+1), +(l+1)\}$ for each $0 \le l \le n_{\mathrm{D}}$ — equivalently the fixed $j = l + \frac{1}{2}$ carrying a ± 1 chirality label — and total angular momentum projection $m_j \in \{-j, \ldots, j\}$ where $j = |\kappa| - 1/2$. *Corrected 2026-09-03:* this read $\kappa \in \{-l-1, +l\}$, which is arithmetically inconsistent with the node count below — that set gives $2, 10, 28$ at $n_{\max} = 1, 2, 3$ against the formula's $4, 16, 40$ — and is not what `geovac/full_dirac_operator_system.py` builds; its labels carry an explicit chirality, and its own documentation rejects the $j = l - \frac{1}{2}$ reading. The total node count is

$$N_{\mathrm{Dirac}}(n_{\max}) = \sum_{n=0}^{n_{\max}-1} g_n^{\mathrm{D}} = \sum_{n=0}^{n_{\max}-1} 2(n+1)(n+2) = \tfrac{2}{3} n_{\max}(n_{\max}+1)(n_{\max}+2).$$

Definition 2 (Spinor space on the Dirac graph)**.** $\mathcal{H}_{\mathrm{GV}}^{(n_{\max})} := \mathbb{C}^{V_{\mathrm{Dirac}}}$*, with the standard Hermitian inner product on* $\mathbb{C}^{N_{\mathrm{Dirac}}}$*.*

The faithful $*$-representation of $\mathcal{A}_{\mathrm{GV}}$ on $\mathcal{H}_{\mathrm{GV}}$ is by the natural pullback: for $f \in \mathcal{A}_{\mathrm{GV}}$ and $\psi \in \mathcal{H}_{\mathrm{GV}}$,

$$(\pi(f)\psi)(n_{\mathrm{D}}, \kappa, m_j) := f(n_{\mathrm{Fock}}(n_{\mathrm{D}}), l(\kappa), m_l(\kappa, m_j)) \cdot \psi(n_{\mathrm{D}}, \kappa, m_j), \tag{3}$$

where the maps $n_{\mathrm{Fock}}, l, m_l$ are the standard Dirac-to-Fock quantum-number bridges: $n_{\mathrm{Fock}}(n_{\mathrm{D}}) = n_{\mathrm{D}} + 1$, $l(\kappa) = |\kappa| - 1$ for both signs (corrected 2026-09-04 together with the label set: the earlier

"$-1-\kappa$ if $\kappa<0$, κ if $\kappa>0$" returns $l+1$ for $\kappa=+(l+1)$, which maps top-shell nodes outside V_{Fock} and breaks the order-zero argument as printed), and m_l via the standard $|j,m_j\rangle = \langle l,m_l;\tfrac{1}{2},m_s|j,m_j\rangle|l,m_l\rangle|s,m_s\rangle$ Clebsch–Gordan decomposition. Each $\pi(f)$ acts diagonally; in the spinor basis it is the diagonal matrix $\mathrm{diag}(f(v_{\mathrm{Fock}}(\cdot)))$. The representation is faithful: $\pi(f)=0 \Leftrightarrow f\equiv 0$ (every Fock node has at least one Dirac node above it for $n_{\max}\geq 1$).

3.3 The Dirac operator D_{GV}

The continuum Camporesi–Higuchi Dirac operator on the unit three-sphere has spectrum $|\lambda_n^{\mathrm{D}}| = n+3/2$ with degeneracy $g_n^{\mathrm{D}} = 2(n+1)(n+2)$ [22]. We promote this to a finite operator on $\mathcal{H}_{\mathrm{GV}}$ by truncation of its spectral decomposition.

Definition 3 (D_{GV})**.** *D_{GV} is the diagonal operator on $\mathcal{H}_{\mathrm{GV}}$ with eigenvalues $\lambda_n^{\mathrm{D},\pm} = \pm(n_{\mathrm{D}}+3/2)$ on the eigenstate $|n_{\mathrm{D}},\kappa,m_j\rangle$ with sign $+$ for $\kappa<0$ and $-$ for $\kappa>0$ (a GeoVac Dirac–Coulomb labelling convention for the two signs of the Camporesi–Higuchi spectrum; Camporesi–Higuchi [22] supply the spinor bundle and the spectrum, not a κ labelling).*

This is the operator used everywhere in the paper: every axiom verification below, the real structure J_{GV} of Proposition 4, and every finite-truncation spectral-action computation are carried out on this diagonal D_{GV} (`geovac/full_dirac_operator_system.py`, `geovac/real_structure.py`; backing tests `tests/test_full_dirac_operator_system.py`, `tests/test_real_structure.py`). The π-free certificate of the Dirac sector (Paper 28 §graph_native_qed.6) is a statement about this spectrum: every eigenvalue is a half-integer rational, every degeneracy is a positive integer.

Remark 6 (No separate "graph form" of D_{GV})**.** *Earlier drafts of this section defined a second, "graph form" $D_{\mathrm{GV}}^{\mathrm{graph}}$ – a first-order hopping operator on a Camporesi–Higuchi edge set "with weights chosen so that the spectrum matches" Definition 3 – and stated their unitary equivalence as a proposition. That proposition was tautological (two Hermitian operators with the same spectrum and multiplicities are unitarily equivalent by the spectral theorem, and the weights had been chosen to make the spectra agree), and the module it cited, `geovac/dirac_matrix_elements.py`, holds closed-form angular (Szmytkowski) and radial (hydrogenic) matrix elements in the (κ,m_j) basis, not a hopping operator. The paper therefore carries one Dirac operator, the diagonal one above. The Dirac-sector* graph *that appears elsewhere in the corpus (`geovac/dirac_lattice.py`, the E1-dipole adjacency of Paper 28's graph-native QED) is a different operator, $H=D+W$, whose adjacency encodes the photon-vertex selection rules rather than D_{GV} itself. Withdrawn at the 2026-09-02 trunk certification run (`CHANGELOG.md`, v5.3.0); the module contents asserted here are pinned by `tests/test_paper32_dirac_module_contents.py`.*

3.4 Bounded commutators

Proposition 2 (Bounded commutators)**.** *For every $a\in\mathcal{A}_{\mathrm{GV}}$, the commutator $[D_{\mathrm{GV}},\pi(a)]$ is a bounded operator on $\mathcal{H}_{\mathrm{GV}}$.*

Proof. D_{GV} has finite-dimensional domain $\mathcal{H}_{\mathrm{GV}}$ at every $n_{\max}$, hence bounded. $\pi(a)$ is a diagonal multiplication operator, also bounded. Any commutator of bounded operators is bounded. In fact more is true: D_{GV} and $\pi(a)$ are both diagonal in the basis $|n_{\mathrm{D}},\kappa,m_j\rangle$, so $[D_{\mathrm{GV}},\pi(a)]=0$ identically. □

[SYMBOLIC PROOF] The check is therefore trivial in a strong sense: the commutator seminorm $a\mapsto\|[D_{\mathrm{GV}},\pi(a)]\|$ vanishes on all of $\mathcal{A}_{\mathrm{GV}}$, the Connes distance it would induce is

degenerate, the first-order differential forms $\Omega^1_{D_{\mathrm{GV}}}$ are zero, and D_{GV} admits no inner fluctuations. The diagonal triple $(\mathcal{A}_{\mathrm{GV}}, \mathcal{H}_{\mathrm{GV}}, D_{\mathrm{GV}})$ is a *spectral*, not a metric, object: it carries the Camporesi–Higuchi spectrum, the degeneracies and the real structure, but no Lipschitz geometry. The metric role belongs instead to the *non-diagonal* multipliers of the truncated operator system on the true Camporesi–Higuchi window (Remark 1; Paper 38 [61]), whose state spaces converge under the left-translation Lipschitz seminorm (Theorem 5); the theorem uses no Dirac commutator, and convergence in the Dirac-commutator seminorm is **[OPEN]** (that seminorm is non-degenerate on the window as measured at $n_{\max} \leq 3$, `tests/test_p38_true_ch_window.py`). Both halves of the contrast — $[D_{\mathrm{GV}}, \pi(f)] = 0$ identically on the diagonal representative, and nonzero commutators on the non-diagonal multipliers of the chirality-doubled model $\mathcal{O}_{n_{\max}}$ — are regression-pinned in `tests/test_paper32_diagonal_triple.py`.

3.5 The triple, summarized

Putting Definitions 1, 2, 3 and Proposition 2 together:

Theorem 1 (The GeoVac spectral triple)**.** *For every $n_{\max} \geq 1$, the data $(\mathcal{A}^{(n_{\max})}_{\mathrm{GV}}, \mathcal{H}^{(n_{\max})}_{\mathrm{GV}}, D^{(n_{\max})}_{\mathrm{GV}})$ is a finite spectral triple in the sense of [1]: $\mathcal{A}_{\mathrm{GV}}$ is unital and involutive, $\pi : \mathcal{A}_{\mathrm{GV}} \to \mathcal{B}(\mathcal{H}_{\mathrm{GV}})$ is a faithful $*$-representation, D_{GV} is self-adjoint with bounded commutators. **[INTERNAL THEOREM]** As $n_{\max} \to \infty$ the state spaces converge to that of the canonical spectral triple of the unit Riemannian spin three-sphere S^3 with the Camporesi–Higuchi Dirac operator* for the operator-system representative on the true Camporesi–Higuchi window $\mathcal{T}_{n_{\max}} = (\mathcal{O}^{\mathrm{CH}}_{n_{\max}}, \mathcal{H}_{n_{\max}}, D_{n_{\max}})$ *of Theorem 5 (van Suijlekom's state-space Gromov–Hausdorff distance, translation-seminorm metrization, limit identified up to isometry; the target's spectral-triple properties are inputs, not outputs, of the convergence); neither the Fock/Weyl model $\mathcal{O}_{n_{\max}}$ of the axiom audit nor the diagonal (gauge-network) representative $\mathcal{A}_{\mathrm{GV}} = \mathbb{C}^{V_{\mathrm{Fock}}}$ of Remark 1, used for the axiom audit above, is the object that converges (Corollary 1 places the diagonal representative on the comparator side of the Connes–van Suijlekom distinction; scope clause added 2026-09-03).*

The continuum-limit clause is the spectral-truncation statement of Connes–van Suijlekom [12, 14, 15] for the true window $\mathcal{O}^{\mathrm{CH}}_{n_{\max}}$; no Cauchy-sequence statement for the diagonal algebra is made (a sentence to that effect was withdrawn 2026-09-03).

4 Axiom Audit

We check each Connes axiom in turn, distinguishing what holds rigorously at finite $n_{\max}$ from what holds only in the continuum limit.

4.1 Order zero (involutive faithful representation)

[SYMBOLIC PROOF] Holds by construction from Definitions 1–2 and Eq. (3): $\pi(f)$ acts diagonally in the $(n_{\mathrm{D}}, \kappa, m_j)$ basis, so π is multiplicative and $\pi(f^*) = \pi(f)^\dagger$, and it is faithful because the quantum-number bridges of Eq. (3) map V_{Dirac} onto V_{Fock}. No numerical certificate is involved (an earlier citation of `tests/test_dirac_matrix_elements.py` here covered the $\kappa \leftrightarrow (l, j)$ bridges, not the representation axiom, and is withdrawn).

4.2 Bounded commutators (regularity)

Holds rigorously at every $n_{\max}$ by Proposition 2.

4.3 Compact resolvent / finite summability

In the finite-$n_{\max}$ truncation $\mathcal{H}_{\mathrm{GV}}$ is finite-dimensional so this is automatic. The continuum statement is that D_{GV} has compact resolvent and is 3^+-summable (Dixmier-trace summable in dimension $3+\epsilon$); this is the Camporesi–Higuchi summability of the continuum operator on S^3 and is a standard result [22, 1].

4.4 Order one

Proposition 3 (Order one, diagonal representative)**.** *For every $a, b \in \mathcal{A}_{\mathrm{GV}}$ (the diagonal representative), $[[D_{\mathrm{GV}}, \pi(a)], \pi(b)^\circ] = 0$ holds trivially, because already $[D_{\mathrm{GV}}, \pi(a)] = 0$ (Proposition 2).*

This is vacuous rather than informative: the diagonal triple has no nonzero commutators to constrain. It is *not* the statement for the Fock/Weyl model $\mathcal{O}_{n_{\max}}$ of the operator-system representative, whose multipliers do not commute with D and for which the order-zero and order-one conditions (with the real structure) *fail* at finite $n_{\max} \geq 2$ (item 5 of the finite-$n_{\max}$ audit below; Table 1 labels the two representatives separately). Order-one becomes non-trivial only when one extends to an almost-commutative triple $\mathcal{A}_{\mathrm{GV}} \otimes M_n(\mathbb{C})$ with non-abelian fiber, at which point it constrains the form of fluctuations of D. This is exactly the setting where Papers 25 ($n=1$, $U(1)$ fiber) and 30 ($n=2$, SU(2) fiber) live; we treat them in Sec. 5.

4.5 Real structure

S^3 is odd-dimensional; in dimension n the canonical KO-dimension is n mod 8 for a Riemannian spin manifold, and the sign table [1, 3] for $n=3$ gives

$$\epsilon = -1, \quad \epsilon' = +1, \quad \epsilon'' = \text{ (none)} \quad \text{(no } \gamma \text{ in odd dimension)}.$$

The sign $\epsilon = -1$ reflects the fact that the 2-dimensional spinor representation of $\mathrm{Spin}(3) = \mathrm{SU}(2)$ is quaternionic (pseudoreal): the standard charge-conjugation operator on this representation squares to $-\mathbb{1}$, equivalently $\mathbb{H} \otimes_{\mathbb{R}} \mathbb{C} \cong M_2(\mathbb{C})$ acts on a complex 2-dimensional space with antiunitary J satisfying $J^2 = -\mathbb{1}$. This is the same sign that appears in the canonical Lorentzian Dirac formalism in $3+1$ dimensions for the charge-conjugation matrix C.

Proposition 4 (Real structure on $\mathcal{H}_{\mathrm{GV}}$)**.** *There exists an antilinear isometry $J_{\mathrm{GV}} : \mathcal{H}_{\mathrm{GV}} \to \mathcal{H}_{\mathrm{GV}}$ with $J_{\mathrm{GV}}^2 = -\mathbb{1}$ and $J_{\mathrm{GV}} D_{\mathrm{GV}} = +D_{\mathrm{GV}} J_{\mathrm{GV}}$, namely the charge-conjugation operator on the Camporesi–Higuchi spinor bundle restricted to the truncation.*

Construction. The continuum charge conjugation J_{S^3} on the spinor bundle of S^3 satisfies the listed signs by construction [1] (the KO-dimension-3 row; Camporesi–Higuchi [22] supply the spinor bundle and spectrum, not a charge conjugation); the finite-cutoff signs are verified directly by `tests/test_real_structure.py`. Restriction to the truncation $\mathcal{H}_{\mathrm{GV}}^{(n_{\max})}$ commutes with the spectral truncation because D_{GV} acts diagonally in the spectral basis and J_{S^3} permutes the spectral basis (sending $|n_{\mathrm{D}}, \kappa, m_j\rangle \mapsto |n_{\mathrm{D}}, \kappa, -m_j\rangle$ up to a phase — *m_j only, with the chirality label held fixed* — whose antilinear-squaring gives $-\mathbb{1}$ on the quaternionic 2-spinor representation). *Corrected 2026-09-03:* the construction printed here sent $\kappa \mapsto -\kappa$, which under the label set above flips chirality; since $D_{\mathrm{GV}} = \chi\,(n + \frac{1}{2})$ that would give $\varepsilon' = -1$ and contradict this proposition's own $JD = +DJ$. The implemented J (`geovac/real_structure.py`) maps m_j alone, as §IV describes and as the real-structure tests verify. The restriction therefore inherits the signs. □

Finite-$n_{\max}$ audit (Sprint WH1-Connes Step 2, 2026-05). The qualitative argument above is now backed by a quantitative computational audit at $n_{\max} \in \{1, 2, 3\}$ on both the Weyl sector (`geovac/spinor_operator_system.py`, $\dim_{\mathcal{H}} = 2, 8, 20$) and the full-Dirac sector (`geovac/full_dirac_operator_s` $\dim_{\mathcal{H}} = 4, 16, 40$), with J_{GV} realized as an antilinear permutation-phase operator $J = U\mathcal{K}$ (with $\mathcal{K}$ complex conjugation) and phase convention $\sigma(l, m_j) = i^{2m_j}(-1)^l$ giving $\sigma_a \overline{\sigma_{J(a)}} = -1$ on every basis label. Implementation in `geovac/real_structure.py`; verification in `tests/test_real_structure.py` (44 tests passing); raw audit data `debug/data/wh1_real_structure_audit.json`. The audit verdict for the four Connes-axiom conditions involving J at finite $n_{\max}$ on the truthful Camporesi–Higuchi Dirac is as follows; items 1–5 are measured on the Fock/Weyl and chirality-doubled *model* objects of Remark 1, and item 6 repeats the audit on the true window whose continuum limit Theorem 5 proves:

1. $J^2 = -\mathbb{1}$ **HOLDS EXACTLY** at every $n_{\max} \in \{1, 2, 3\}$ on both sectors (bit-exact zero residual). The phase formula $\sigma_a \overline{\sigma_{\pi(a)}} = -1$ is verified pair-by-pair across the spinor and full-Dirac bases.

2. $J_{\mathrm{GV}} D_{\mathrm{GV}}^{\mathrm{truthful}} = +D_{\mathrm{GV}}^{\mathrm{truthful}} J_{\mathrm{GV}}$ **HOLDS EXACTLY** at every $n_{\max}$ (bit-exact zero residual). Mechanism: the truthful CH Dirac is diagonal in $(n_{\mathrm{Fock}}, l, m_j, \text{chirality})$ with eigenvalue chi $\cdot$ $(n_{\mathrm{Fock}} + 1/2)$ depending only on n and chi; J_{GV} permutes only m_j, leaving n and chi fixed, so the matrix relation $U\overline{D} = DU$ is diagonal-by-diagonal.

3. $J_{\mathrm{GV}} \mathcal{O}_{n_{\max}} J_{\mathrm{GV}}^{-1} = \mathcal{O}_{n_{\max}}$ **HOLDS EXACTLY** at every $n_{\max}$ (bit-exact zero residual on each generator a, with 0 of $1, 14, 55$ generators failing the $\mathcal{O}$-membership test). Mechanism: each multiplier matrix $\widetilde{M}_{NLM}$ has J-conjugate $\widetilde{M}_{NL(-M)}$ via the $m_l \to -m_l$ symmetry of the Wigner-3j coupling coefficients (Edmonds Eq. 3.7.5 / Avery–Wen–Avery 3-Y integral); the operator system is closed under this involution.

4. $J_{\mathrm{GV}} D_{\mathrm{GV}}^{\mathrm{offdiag}} = +D_{\mathrm{GV}}^{\mathrm{offdiag}} J_{\mathrm{GV}}$ **FAILS** structurally (max residual 2×10^0 at $n_{\max} \in \{2, 3\}$; 10^{-2} at $n_{\max} = 1$ from the m-linear lifter alone). This is the natural counter-example: the round-3 `offdiag` Dirac is a hand-crafted Hermitian perturbation with E1-type hopping $D[a, b] = \alpha$ on the rule $|\Delta n| = |\Delta l| = 1, |\Delta m_j| \le 1$, and the permutation-phase σ with $(-1)^l$ parity does not commute with this perturbation because the l-parity sign flips under bra/ket $l \to l \pm 1$. The genuine 4-component Camporesi–Higuchi spinor Dirac (which couples large/small components via the proper $\mathrm{SU}(2)_L \times \mathrm{SU}(2)_R$ angular gradient with Clifford-algebra γ matrices) would resolve this, but lies outside the present construction; the offdiag CH is best read as an SDP-bounding device (round-3 R3.5 Connes distance), not a spectral-action-foundational object.

5. **Order-zero** $[a, JbJ^{-1}] = 0$ **and order-one** $[[D, a], JbJ^{-1}] = 0$ **FAIL at finite** $n_{\max} \ge 2$ on the Weyl-sector model $\widetilde{\mathcal{O}}_{n_{\max}}$ of the operator system (not the true window of Theorem 5), with residuals of order 0.05–0.2 (largest matrix entry in absolute value, not operator norm). At $n_{\max} = 2$: 124 of 196 pairs fail order-zero (max residual 5.5×10^{-2}); 43 of 196 pairs fail order-one (max residual 1.0×10^{-1}); at $n_{\max} = 3$: 2,594 of 3,025 and 1,337 of 3,025 respectively (max residuals 7.9×10^{-2} and 2.0×10^{-1}). The conditions hold trivially at $n_{\max} = 1$ (single-generator operator system). **[MEASURED]** The order-zero failure *grows* with the cutoff (Weyl sector, 2026-09-28 re-measurement): max residual $5.5 \times 10^{-2} \to 7.9 \times 10^{-2} \to 9.6 \times 10^{-2}$ and failing fraction 63% → 86% → 92% (17,970 of 19,600 pairs) at $n_{\max} = 2, 3, 4$; on a *fixed* set of generators (the degree-1 harmonics, $N = 2$) the operator-norm residual runs $0.051, 0.068, 0.076, 0.081$ at $n_{\max} = 2$–5, and its ratio to the (growing) generator norm runs $0.225, 0.212, 0.209, 0.208$ — still drifting down, so whether it tends to a nonzero limit is

not established (measured through $n_{\max} = 5$). The order-zero counts/residuals, the model order-one counts at $n_{\max} = 2, 3$ and the fixed-generator growth are regression-pinned in `tests/test_paper32_order_condition_growth.py`.

6. **True Camporesi–Higuchi window (2026-09-28).** **[MEASURED]** The audit repeated on the object of Theorem 5: spinors $L^2(\mathrm{SU}(2)) \otimes \mathbb{C}^2$ in the left-invariant frame, basis $e^j_{mm's} = \sqrt{2j+1}\, D^j_{mm'} \otimes e_s$, $D = 2\sigma \cdot J_R + \frac{3}{2}$, window $|\lambda| \le n_{\max} + \frac{1}{2}$, $\mathcal{O}^{\mathrm{CH}}_{n_{\max}} = P\, C(S^3)\, P$ of dimension $5, 30, 91, 204$ at $n_{\max} = 1$–4, spanned by the compressed matrix coefficients $D^J_{MM'}$ of bands $N = 2J + 1 \le 2n_{\max}$ (sup norm 1). The charge conjugation is $J = UK$, $U\, e^j_{mm's} = (-1)^{m-m'} (i\sigma_y)_{s's}\, e^j_{-m,-m',s'}$: pointwise complex conjugation of the spinor ($\overline{D^j_{mm'}} = (-1)^{m-m'} D^j_{-m,-m'}$) twisted by $i\sigma_y$, which reverses J_R and σ and so fixes $\sigma \cdot J_R$. It commutes with D, hence preserves the window, and $J^2 = -\mathbb{1}$, $JD = +DJ$ hold exactly (test-pinned at $n_{\max} = 1$–4). The signs are not a phase convention: the antiunitaries intertwining D and the isometry generators $(J_L,\ J_R + \sigma/2)$ form a family of one phase per irreducible block (dimension $2n_{\max}$), every member has $J^2 = -\mathbb{1}$, and no equivariant antiunitary anticommutes with D. $J\mathcal{O}^{\mathrm{CH}}J^{-1} = \mathcal{O}^{\mathrm{CH}}$ holds exactly; indeed $J(PfP)^*J^{-1} = PfP$. Order zero $[a, Jb^*J^{-1}]$ and order one $[[D, a], Jb^*J^{-1}]$ *fail*, already at $n_{\max} = 1$ (where the model holds trivially), over all basis pairs:

$n_{\max}$	order 0 failing	max	order 1 failing	max entry	op. norm
1	12/25 (48%)	1/2	4/25 (16%)	3/4	0.750
2	572/900 (64%)	5/9	513/900 (57%)	5/3	1.863
3	5842/8281 (71%)	11/16	5856/8281 (71%)	21/8	2.902
4	30462/41616 (73%)	19/25	31021/41616 (75%)	18/5	3.923

(max entry in the Peter–Weyl basis; at order zero it equals the operator norm). On the fixed degree-1 generators the order-zero operator-norm residual runs $1/2, 5/9, 11/16, 19/25, 29/36, 41/49$ and the order-one residual $3/4, 7/9, 13/16, 21/25, 31/36, 43/49$ at $n_{\max} = 1$–6, increasing toward 1, the product of the generators' sup norms (the circle analogue: on $\mathrm{span}\{e^{ik\theta}\}_{|k|\le n}$ the commutator of the compressions of $e^{\pm i\theta}$ is a difference of two rank-one projections, of norm 1 at every n). Relative to the generator norm the true-window residual (0.72 at $n_{\max} = 2$, 0.86 at $n_{\max} = 5$) is *larger* than the model's (≈ 0.21). *Mechanism (exact).* Since $fg = gf$, $[PfP, PgP] = Pg(1-P)fP - Pf(1-P)gP$; at order one $[D, f] = c(df)$ commutes with g pointwise, so the same identity holds with f replaced by $c(df)$. A band-N multiplier moves the Dirac level by at most $N - 1$ (Paper 38's shell-difference rule), so every residual vanishes on all levels more than $\max(N_a, N_b) - 1$ below the top (measured: largest entry outside that strip $\le 3 \times 10^{-15}$, $n_{\max} = 2$–5). The degree-1 residual lives on the top Dirac level alone, whose share of the window is $3/(n_{\max} + 2) \to 0$, and its normalised Hilbert–Schmidt norm $\|C\|_2/\sqrt{\dim \mathcal{H}}$ falls $0.30, 0.29, 0.28, 0.26$ at $n_{\max} = 2$–5. The failure is a boundary term of the compression — full operator norm, shrinking weight — not a bulk non-commutativity. Two independent constructions (Clebsch–Gordan multipliers with the closed-form J; sampled Wigner-D functions with J obtained from pointwise conjugation of the sampled basis) agree pair by pair to 10^{-10} at $n_{\max} = 1, 2$ (`tests/test_true_window_real_structure.py`). Test-pinned there: the J-relations (with $\dim \mathcal{O} = 5, 30, 91, 204$) at $n_{\max} = 1$–4, the order tables at $n_{\max} = 1$–3, the degree-1 residuals at $n_{\max} = 1$–4, and the boundary strip at $n_{\max} = 2, 3$; the $n_{\max} = 4$ order-zero/one table row, the degree-1 values at $n_{\max} = 5, 6$, the Hilbert–Schmidt weights and the strip check at $n_{\max} = 4, 5$ are measured in the audit driver, not test-pinned.

Reading of the order-condition failures. The order-zero condition $[a, JbJ^{-1}] = 0$ is the abelian-classical statement *multiplications by classical functions commute*, automatic for $f, g \in C^\infty(S^3)$ in the continuum. At finite $n_{\max}$, the truncated objects $P_{n_{\max}} f P_{n_{\max}}$ and $J P_{n_{\max}} g P_{n_{\max}} J^{-1}$ are no longer multiplications by classical functions; they are operator-system elements ([12, Sec. 2]), and their commutator is generically nonzero. That explains why the residual is nonzero; it does *not* explain its size or its trend. An earlier version of this paragraph inferred from the GH convergence of Theorem 5 that the residuals decay in the $n_{\max} \to \infty$ limit, and called the failures finite-resolution artifacts. The measurement above refutes the decay at the cutoffs reached: the residuals grow, and on fixed low-degree generators they do not decay through $n_{\max} = 5$. The same growth is measured on the true window that converges (item 6), where it is identified exactly as a Toeplitz boundary term of the compression. No contradiction with Theorem 5 arises: GH convergence of state spaces does not by itself control the commutator $[a, JbJ^{-1}]$ in norm, and on the true window the residual, though of full operator norm, is confined to a boundary strip of vanishing weight. **[MEASURED]** status, regression-pinned in `tests/test_paper32_order_condition_growth.py` (model) and `tests/test_true_window_real_structure.py` (true window); the mechanism on the model (the boundary-term identity of item 6 is the plausible reading there too), and whether the continuum identity is recovered in any weaker (e.g. weak-operator or state-space) sense, are open.

Net status of the real structure at finite $n_{\max}$. The introductory abstract flagged as open "whether the GeoVac Dirac operator D_{GV} (the diagonal spectral form) carries a real structure matching the signs of S^3 exactly at finite $n_{\max}$ or only in the continuum limit." The Sprint WH1-Connes Step 2 audit closes this question *positively for the three J-relations on the truthful CH Dirac*: $J^2 = -\mathbb{1}$, $JD = +DJ$, and $J\mathcal{O}J^{-1} = \mathcal{O}$ all hold with bit-exact zero residual at every finite $n_{\max} \in \{1, 2, 3\}$ on both the Weyl and full-Dirac (chirality-doubled) model sectors, and on the true Camporesi–Higuchi window (test-pinned at $n_{\max} = 1$–4), where the signs are forced by equivariance (item 6). It does *not* close it for the order-zero and order-one conditions (involving the abelian-multiplication identity that defines the classical limit of the algebra): on the model those fail at finite $n_{\max} \geq 2$ with residuals that grow over the measured cutoffs (item 5 above; **[MEASURED]**, regression-pinned in `tests/test_paper32_order_condition_growth.py`, mechanism open), and on the true window they fail from $n_{\max} = 1$ with residuals growing toward the generators' sup-norm scale, as a Toeplitz boundary term of the compression (item 6; **[MEASURED]**, `tests/test_true_window_real_structure.py`). The offdiag CH Dirac fails $JD = +DJ$ structurally and is therefore not a spectral-action-foundational object; the truthful CH is the natural domain for the present audit. (On the diagonal representative $\mathcal{A}_{\mathrm{GV}}$ the truthful CH Dirac has no nonzero commutators and so no inner fluctuations, Proposition 2; an inner-fluctuation calculus for the Track 3 almost-commutative extension needs the operator-system representative or a non-diagonal Dirac, Sec. 6 discussion.)

4.6 $\mathbb{Z}_2$-grading

Proposition 5 (Odd continuum; even model truncation; odd true window)**.** *(a) The continuum spectral triple of the unit Riemannian spin S^3 with the Camporesi–Higuchi Dirac operator is* odd*: it carries no $\mathbb{Z}_2$-grading commuting with $C^\infty(S^3)$ and anticommuting with the Dirac operator. (b) The* chirality-doubled model *truncation $(\mathcal{A}_{\mathrm{GV}}, \mathcal{H}_{\mathrm{GV}}, D_{\mathrm{GV}})$ is* even*: the chirality-doubling operator $\gamma_{\mathrm{GV}} = \sigma_x \otimes \mathbb{1}$ (exchanging the two chirality copies of each (n, l, m_j) label) is a self-adjoint involution that commutes with $\pi(\mathcal{A}_{\mathrm{GV}})$ and anticommutes with D_{GV}. (c) The* true *Camporesi–Higuchi window of Theorem 5 (complete Dirac eigenspaces, the object that converges) is* odd *(measured, $n_{\max} \leq 3$), like the continuum: at $n_{\max} = 1, 2, 3$ the commutant of its operator system $\mathcal{O}^{\mathrm{CH}}_{n_{\max}}$ is the scalars, so*

no $\mathbb{Z}_2$-grading commutes with $\mathcal{O}^{\mathrm{CH}}$ and anticommutes with D. The evenness in (b) is a property of the chirality-doubled model, not of the converging truncation.

Proof. (a) S^3 is odd-dimensional; this is the standard statement for odd Riemannian spin manifolds [1]. (b) $\pi(f)$ does not depend on the chirality label (Eq. (3)), so it commutes with any operator acting on the chirality index alone, in particular with $\sigma_x \otimes \mathbb{1}$; $D_{\mathrm{GV}} = \chi\,(n + \frac{1}{2})$ is $\sigma_z \otimes (n+\frac{1}{2})$ in the chirality index, and σ_x anticommutes with σ_z. Measured bit-exactly at $n_{\max} \leq 3$ together with $J\gamma = +\gamma J$ (`geovac/chirality_grading.py`; `tests/test_almost_commutative.py`, §8.2; `tests/test_paper32_diagonal_triple.py`, which also contrasts the diagonal $\sigma_z \otimes \mathbb{1}$ candidate, which commutes with D_{GV} instead of anticommuting). **[SYMBOLIC PROOF + MEASURED]** (c) **[MEASURED]** At $n_{\max} = 1, 2, 3$ ($\dim \mathcal{H} = 4, 16, 40$) the commutant of $\mathcal{O}^{\mathrm{CH}}_{n_{\max}}$ (the null space of $X \mapsto [X, a]$ over a basis of $\mathcal{O}^{\mathrm{CH}}$) has dimension 1 (Peter–Weyl construction, `tests/test_true_window_real_structure.py`). Two further routes in the same test file corroborate it at $n_{\max} = 1, 2$: the sampled-function construction of $\mathcal{O}^{\mathrm{CH}}$, whose commutant is again the scalars and whose generated unital algebra is all of $B(\mathcal{H})$ (dimension $16,256$, so by Burnside's theorem its commutant is the scalars; dimension 1600 at $n_{\max} = 3$ is computed outside the test), and an independent polynomial construction on $S^3 \subset \mathbb{R}^4$ (no Wigner-D, Clebsch–Gordan or sampling code). At $n_{\max} = 3$ the oddness rests on the Peter–Weyl route. A scalar cannot anticommute with the invertible D, so the grading space is $\{0\}$. On the true window only the pair $(\varepsilon, \varepsilon') = (-, +)$ is defined, and it is forced by equivariance (audit item 6). □

The chirality-doubled model triple of (b) therefore carries the sign triple $(\varepsilon, \varepsilon', \varepsilon'') = (-, +, +)$, and, per the rule stated in §8.2, *no* KO-dimension label is attached at finite cutoff: the continuum factor is KO-3 (odd), the model's grading reads as the KO-4 column, and any single label at finite cutoff is a convention. The grading is a doubling artefact of the chirality-labelled model basis, not a chirality of S^3; it is absent on the true window (c) (measured, $n_{\max} \leq 3$) and in the continuum limit, where the odd description is the correct one. Odd-dimensional spectral triples construct their spectral action from $|D|$ rather than from D^2 projected onto chirality eigenspaces, which is the structure that Paper 28's QED-on-S^3 computations use.

4.7 Summary

Finite-cutoff Krein-carrier sign audit at $(m, n) = (4, 6)$ (Sprint L2-B/C/D, 2026-05-16). The Riemannian audit above sits at signature $(s, t) = (3, 0)$ (KO-dim 3 for the continuum factor). A signature-blind sign audit of a Krein carrier at $(s, t) = (3, 1)$ — not a Lorentzian geometry in any metrically visible sense; the framework's Lorentzian programme is descoped (Paper 45) — via the Bizi–Brouder–Besnard 2018 [50] Krein-space spectral-triple classification translates to the BBB index pair $(m, n) = (4, 6)$ (via $m \equiv t + s$, $n \equiv t - s$ (mod 8)). Sprints L2-B (Krein space, `geovac/krein_space_construction.py`), L2-C (Lorentzian Dirac, `geovac/lorentzian_dirac.py`), and L2-D (Connes axiom audit, `geovac/connes_axiom_audit_31.py`) **[PANEL-VERIFIED]** verify four BBB-predicted-sign axioms bit-exactly at $n_{\max} \in \{1, 2, 3\}$ and $N_t \in \{1, 11, 21\}$ on the truthful Camporesi–Higuchi spatial Dirac. Per BBB Table 1 at $(4, 6)$: $\varepsilon = +1$, $\varepsilon'' = -1$, $\kappa = -1$, $\kappa'' = +1$ (verified directly against [50] v2 PDF). Sub-deliverables and verdicts are summarised in Sec. 8.6 and the subsections § 8.7, § 8.8, § 8.9.

Structural finding on χD. [PANEL-VERIFIED] The universal BBB Sec 5(v) axiom $\chi D = -D\chi$ fails bit-exactly on the pair (γ_K^5, D_L) built in Sprints L2-B/C, with Frobenius residual $2\|D_{\mathrm{GV}}\|_F$ at $N_t = 1$ (numerical values $6.00, 18.33, 38.88$ at $n_{\max} = 1, 2, 3$ respectively). The mechanism is structural and follows from two locked Riemannian-side conventions: (i) the Sprint L2-B

convention identifies `FullDiracLabel.chirality` with the γ^5 eigenvalue (the chiral-basis Peskin–Schroeder convention which keeps the Riemannian limit identification with $\mathcal{H}_{\mathrm{GV}}$ transparent); (ii) the truthful Camporesi–Higuchi Dirac is diagonal in chirality with eigenvalue chi $\cdot\,(n_{\mathrm{Fock}} + 1/2)$, so D_{GV} *commutes* with γ^5 rather than anticommuting with it. [**SYMBOLIC PROOF**] These two facts together force $\{\gamma^5, D_L\} = 2i\,(\gamma^5 D_{\mathrm{GV}}) \otimes I_{N_t} \neq 0$ in general. This is a structural property of GeoVac's chirality-diagonal D_{GV} in the Peskin–Schroeder chiral basis, not a basis-convention error and not a defect of the Lorentzian lift. Three resolutions are documented: **(R1) accept the structural finding** (truthful D_{GV} is Krein-self-adjoint with the BBB-predicted (4, 6) signs on the J-relations and is the right object for axioms requiring $JD = +DJ$, even though it does not form a full BBB indefinite spectral triple); **(R2) use the offdiag Camporesi–Higuchi Dirac** for axioms that require $\{\chi, D\} = 0$ (the same offdiag CH used as an SDP-bounding device in the Riemannian Connes-distance round 3 R3.5); **(R3) redefine γ^5 on $\mathcal{H}_{\mathrm{GV}}$** as an off-diagonal grading anticommuting with truthful D_{GV} (the chirality-doubling $\gamma_{\mathrm{GV}} = \sigma_x \otimes \mathbb{1}$ of Proposition 5(b) is such a grading), which breaks Sprint L2-B's chirality-as-γ^5 identification and is essentially a basis-convention rebuild. The recommended path for Sprint L2-E (Krein-level Paper 42 redo) is R1 + R2 in parallel, matching the Riemannian-side practice of using truthful CH for spectral-action and reality-condition tests while using offdiag CH for SDP-bounding Connes-distance computations.

Theorem 2 (State-space limit identification)**.** *The state spaces of the true-window truncations (Theorem 5) converge, in van Suijlekom's state-space Gromov–Hausdorff distance under the translation seminorm, to the state space $(P(S^3), W_d)$ of the canonical spectral triple of the unit Riemannian spin three-sphere with the Camporesi–Higuchi Dirac operator, up to isometry (Paper 38, limit-identification proposition). That target triple is real, odd, and of* KO*-dimension* 3*; these are classical properties of the target, not outputs of the convergence.*

[**INTERNAL THEOREM**] This is a structural theorem; no fitting parameters or empirical inputs. It is a statement about the continuum limit only: at finite cutoff the true-window truncation is odd like its limit (measured, $n_{\max} \leq 3$; Proposition 5(c)), only the chirality-doubled model being even (Proposition 5(b)), and the Fock/Weyl model of the operator system fails order zero and order one, as does the true window, by a boundary term of the compression (Table 1, audit items 5–6). The audit in Table 1 establishes that the spectral-triple framing is not metaphorical at the level of spectral data and the three J-relations.

5 Sub-sector Identification: Papers 25, 28, 30, 31

We now read the major framework papers as projections of the GeoVac triple.

5.1 Paper 28 (QED on S^3): the spectral action

Paper 28 [37] computes Tr $f(D^2/\Lambda^2)$ and related spectral sums on the Dirac operator D_{GV} in the continuum limit, with SO(4) vertex selection rules from the Hopf-bundle decomposition. In the language of Sec. 2:

- Theorem T9 (Paper 28 Sec. T9): the spectral zeta of D^2 is a two-term polynomial in π^2 with rational coefficients at every integer s. In spectral-triple language: the heat-kernel coefficients $a_k(D^2)$ on S^3 saturate at $k = 2$ (Paper 28's "SD two-term exactness theorem").
- Self-energy structural zero (Paper 28 Theorem 4): $\Sigma(n_{\mathrm{ext}} = 0) = 0$ on the continuum spectral sum, identically by vertex parity. This is the standard heat-kernel bosonic sector cancellation specialized to the Camporesi–Higuchi spectrum.

- Vector-photon QED on S^3 (Paper 28 §vector_photon_qed): adding photon (q, m_q) labels with Wigner $3j$ vertex coupling recovers 7/8 selection rules for scalar electrons and 8/8 for Dirac electrons. The per-loop calibration is the S^2 Weyl constant $1/(4\pi)$ (M1 volume slot).

In the spectral-triple language Paper 28 is the spectral-action computation $\mathrm{Tr}\, f(D_{\mathrm{GV}}^2/\Lambda^2)$ at one loop; the Camporesi–Higuchi spectrum is the input data of D_{GV}; the SO(4) selection rules are the harmonic decomposition of the spinor bundle. Paper 28 is not a free-standing observation about S^3; it is the spectral action of the present triple.

5.2 Paper 25 (Hopf graph as lattice gauge): the Cartan torus

Paper 25 [36] introduces the signed node-edge incidence matrix B of the Fock graph, with $L_0 = BB^\top$ the node Laplacian (its earlier reading as the matter propagator is withdrawn: L_0 is spectrally inert on S^3 and S^5, 2026-09-25) and $L_1 = B^\top B$ the discrete Hodge-1 Laplacian. **[OBSERVATION]** The spectral-triple reading of L_1 is *not* an inner fluctuation of D_{GV}: $[D_{\mathrm{GV}}, \pi(f)] = 0$ (Proposition 2), so $\Omega^1_{D_{\mathrm{GV}}} = 0$ and D_{GV} has no fluctuations at all. The reading goes through the gauge-network (edge-hopping) Dirac of Papers 25/30 — the Marcolli–van Suijlekom setting of Sec. 9 — whose $U(1)$ link variables fluctuate the edge data and whose quadratic kinetic term is L_1. In that setting L_1 is the Cartan-subalgebra (maximal-torus) sector, and its assignment as "photon propagator" in Paper 25 is the gauge-network spectral-action reading. A statement of the same identification on the non-diagonal multipliers of $\mathcal{O}_{n_{\max}}$, with a backing test, is open.

5.3 Paper 30 (SU(2) Wilson gauge): the full non-abelian sector

Paper 30 [38] builds the SU(2) Wilson lattice gauge theory on the same graph and proves: (a) the maximal-torus reduction reproduces Paper 25's $U(1)$ action exactly; (b) the weak-coupling kinetic term equals L_1 acting on $\mathfrak{su}(2)$-valued 1-cochains. **[OBSERVATION]** In the spectral-triple language the natural setting is the almost-commutative algebra $\mathcal{A}_{\mathrm{GV}} \otimes M_2(\mathbb{C})$ (a rank-1 non-abelian M_2 fiber). With the diagonal $D_{\mathrm{GV}} \otimes \mathbb{1}$ this extension still has no inner fluctuations ($[D_{\mathrm{GV}} \otimes \mathbb{1}, a \otimes m] = [D_{\mathrm{GV}}, a] \otimes m = 0$); the identification of Paper 30's $\mathfrak{su}(2)$-valued link variables as the fluctuations, and of the Wilson plaquette action as the lattice-gauge implementation of the spectral-action curvature term, is made in the Marcolli–van Suijlekom gauge-network setting with its edge-hopping Dirac (Sec. 9), where the order-one condition is non-trivial.

Read that way, Paper 30 is the rank-1 non-abelian almost-commutative gauge-network extension over the GeoVac label set; Paper 25 is the abelian projection of Paper 30; together they exhibit the standard Cartan torus / full group structure of that gauge sector. They are not fluctuations of the diagonal D_{GV}.

5.4 Paper 31 (universal/Coulomb partition): the $\mathcal{A}/D$ split

Paper 31 [39] partitions GeoVac results into a universal sector that depends only on the algebra $\mathcal{A}$ (angular sparsity, composed scaling, selection rules) and a potential-specific sector that depends on the Dirac operator D (spectrum, calibration κ, transcendental content, rigidity, α observation). In the spectral-triple language this is exactly the $\mathcal{A}/D$ split: properties of $\mathcal{A}_{\mathrm{GV}}$ alone (rank, dimension, representation content) transfer to any spectral triple with the same algebra; properties of D_{GV} specifically depend on the choice of Dirac operator. The D-dependent side is a statement about *spectral* data (spectrum, degeneracies, the transcendentals of spectral sums): on the diagonal representative D_{GV} commutes with $\mathcal{A}_{\mathrm{GV}}$ (Proposition 2), so the split carries no metric content of its own.

Paper 31 is the structural reading of Paper 22 (angular sparsity theorem, an $\mathcal{A}$-statement) and Paper 23 (nuclear shell model, a different D-choice on the same $\mathcal{A}$): when one keeps $\mathcal{A}$ fixed and varies D, the universal results transfer; when one varies $\mathcal{A}$ (say, replace S^3 with S^5, Paper 24), the entire structural picture reorganizes.

5.5 Summary of sub-sector identification

Track NI as an explicit composed real-space spectral triple. The "Different D, same $\mathcal{A}$" row of Table 3 (Paper 23 [34], nuclear shell model) covers a second class of construction in Paper 23 §VI: the composed nuclear–electronic deuterium Hamiltonian, a 26-qubit Hilbert-space tensor product of a 16-qubit deuteron register (harmonic-oscillator basis) with a 10-qubit hydrogenic electronic register (Fock-projected S^3 lattice), coupled through a cross-register hyperfine $\mathbf{I}\cdot\mathbf{S}$ operator. This is structurally distinct from both the SM almost-commutative geometry (which couples a Riemannian factor with a finite *internal* factor) and from the standard spectral-truncation convergence machinery (which handles single triples or one-infinite-plus-one-finite products); **[PANEL-VERIFIED]** it is constructed as an explicit Connes-style real-space multi-particle spectral triple, and its Hamiltonian reproduces the singlet–triplet hyperfine pattern $(-3A/4, +A/4)$ for the input coupling A_{hf} set to the 21 cm value (1.62×10^{-7} Ha for the $3A/4$ shift) — a structural check of the Clebsch–Gordan pattern, not a measurement against an external reference, since the coupling is the input (2026-09-03). **[OBSERVATION]** The positioning observation (that the canonical NCG / spectral-triple literature does not contain a directly comparable construction) is formalized in the standalone Track NI Zenodo memo (`papers/group4_quantum_computing/track_ni_spectral_triple_zenodo.m`). The W2b cross-manifold structural blocker of Sec. 8.5 is exactly the convergence-theorem gap that surrounds this construction.

Paper 39 closes the same-manifold tensor-product case. The same-manifold sub-case of W2b — the tensor product $\mathcal{T}_{S^3}^{\lambda_a} \otimes \mathcal{T}_{S^3}^{\lambda_b}$ of two truncated Camporesi–Higuchi spectral triples on S^3 at distinct focal lengths $\lambda_a \neq \lambda_b$ — is closed in a companion standalone paper (Paper 39 [43]). The result extends Paper 38's convergence to the KO-dim-6 graded tensor product via the *lifted-state* construction (leading constant 1): Paper 38's unconditional-theorem facts, lifted factor-wise and metrized by the translation seminorm, with the limit the graded (Clifford-doubled) tensor product of the two odd Camporesi–Higuchi real spectral triples — the naive $\gamma_a \otimes D_b$ being unrealisable ($\gamma_a = 0$ on S^3). (An earlier (B, P)-pair route via a Pythagorean refinement of the triangle inequality gave only a weaker conditional bound with constant 2 (its internal Lipschitz comparison constant being $C_3^{(2)} \leq \sqrt{2}$); its height leg was refuted and it is superseded by the lifted-state route [retracted 2026-09-05: p39-pythagorean-height-route].) This is the mathematical foundation for the GeoVac multi-focal-composition program, where two electrons in distinct hydrogenic shells live on S^3 at distinct focal lengths $\lambda = n/Z$; the convergence theorem of Paper 39 says that the cross-register joint operator system on $\mathcal{O}_{n_{\max,a}} \otimes \mathcal{O}_{n_{\max,b}}$ converges qualitatively to the continuum tensor algebra $C^\infty(S^3 \times S^3)$ in van Suijlekom's state-space Gromov–Hausdorff distance (the Latrémolière propinquity strengthening is the same open gap as for the single S^3 factor; Remark 44). The two remaining cross-manifold variants ($\mathcal{T}_{S^3} \otimes \mathcal{T}_{S^5}^{\mathrm{Bargmann}}$ and $\mathcal{T}_{S^3} \otimes \mathcal{T}_{S^5}^{\mathrm{Riem}}$) remain the W2b content of Sec. 8.5.

Remark 7 (Multi-center composition: the reducible–irreducible transition)**.** *The multi-focal program above concerns two electrons at distinct focal lengths on a single S^3; the complementary* multi-center *composition concerns the orthogonal projections P_A onto the single-center orbital subspaces of a polyatomic molecule in the shared Hilbert space with overlap metric G. The natural*

setting is the $*$-representation *theory of orthogonal projections. For a* diatomic *the pair* $\{P_A, P_B\}$ *is* tame *(Type I): by the two-projection theorem (Halmos [67], with the principal-angle norm and block calculus catalogued in Böttcher–Spitkovsky [68]) the generated* $*$*-algebra decomposes into blocks of dimension* ≤ 2*, completely classified by the principal angles between the two subspaces—* ***[MEASURED]*** *for LiH at* R_{eq} *the three angles* $7.6°/44.7°/67.3°$*, the middle one within* $0.3°$ *of* $45°$*, so that* $\|[P_A, P_B]\| = 0.49997$ *sits essentially at the ceiling* $\frac{1}{2}$ *(the ceiling itself, and the reducibility of any pair, are forced by the two-projection theorem* ***[SYMBOLIC]****; the molecular data supply only the angles). The* $*$*-algebra generated by* three or more *projections is, by contrast,* $*$*-wild in general; the passage from two to three centers is precisely this tame-to-wild threshold.* ***[PANEL-VERIFIED]*** *What the molecular data adds is that the three-center configuration lands on an* irreducible *(Schurian) point of that wild family: for linear* $\mathrm{BeH_2}$ *(the* σ *pair* $\{1s, 2p_0\}$ *per center) the three center-projections generate the* full *matrix algebra* M_6 *and act irreducibly (commutant* $= \mathbb{C}\mathbb{1}$*), welding the whole space into a single indecomposable block that admits no decomposition; every pair remains reducible, as it must by the two-projection theorem.*

Tameness is not composability, and conflating the two misreads the obstruction. The diatomic pair is completely classified, but it emphatically does not commute—$\|[P_A, P_B]\| = 0.49997$ *is essentially the maximal possible value* $\frac{1}{2}$*—and it is that* two*-projection non-commutativity, not the three-projection wildness, which constitutes the multi-focal composition wall recorded above. The obstruction accordingly has two distinct rungs. At two centers the projections fail to commute while remaining completely classified by their principal angles: the obstruction is quantitative, and the cost of removing it is known in closed form. At three centers the classification itself fails, and no complete list of invariants exists. Forcing commutation at the first rung—Löwdin orthogonalization—is the framework's sparsity-destroying move; at the second rung there is no corresponding normal form available to force.*

The wildness is specific to the subspace configuration and is not *a generic consequence of non-commutativity. The framework's other non-commutative angular structure—*$SO(3)/SU(2)$ *coupling, the Gaunt and* $6j$ *machinery—is the representation theory of a* compact *group, hence type I by Peter–Weyl, hence tame however non-abelian it may be.* ***[MEASURED]*** *The contrast is quantitative: on the menu* $l = 0, 1, 2$ *the* $\mathfrak{su}(2)$ *generators satisfy* $\|[J_a, J_b]\| = 2$ *yet their commutant is three-dimensional (one dimension per irreducible constituent, and* $\sum_\lambda m_\lambda^2$ *in general—four for* $l=1$ *at multiplicity two, five for* $l=1 \oplus l=2 \oplus l=2$*), whereas the three center-projections, at the* smaller *non-commutativity* $\|[P_A, P_B]\| \approx 0.49$*, have commutant* $\mathbb{C}\mathbb{1}$*. Non-abelianness and wildness are independent properties. This is the structural reason the framework's angular sparsity is graded, finite and potential-independent (Paper 22), and the reason no Gaunt-style graded reduction should be expected to rescue multi-center composition: arbitrary configurations of subspaces are not group representations and escape Peter–Weyl.*

Two invariants must be kept separate. ***[PANEL-VERIFIED]*** *(i) The* categorical *statement—pairs reducible, triple irreducible—is robust: it survives enriching the per-center* σ *menu* $\{1s, 2p_0\} \to \{1s, 2s, 2p_0\}$ *(*$M_6 \to M_9$*, irreducible at the physical* $Z = 2$ *charge, regression-pinned; the structural* $Z = 1$ *case is likewise regression-pinned,* `tests/test_paper32_composition_pins.py`*) and holds at real molecular geometries. It is, however, a statement about center* count*: three coupled projections in general position are irreducible at essentially any bonding-relevant separation, the commutant reverting to reducible only once residual overlaps fall below resolution.* ***[MEASURED]*** *(ii) The bonding-dependent content is instead the* magnitude *of the irreducible three-body coupling, measured by the nested commutator* $\|[[P_A, P_B], P_C]\|$ *(which cannot even be formed for a diatomic):* 0.32 *at bonding (regression-pinned,* `tests/test_paper32_composition_pins.py`*), decaying smoothly to zero as the centers separate.*

[PANEL-VERIFIED] *Geometry sharpens the picture. At* exactly *linear geometry the in-*

plane basis $\{s, p_x, p_z\}$ per center decouples by axial symmetry into a σ and a π sector and the nine-dimensional algebra is reducible; any *bend mixes the sectors and welds the full space into one irreducible M_9 block (regression-pinned at the* H_2O *geometry* $104.5°$*; the sweep across* $160° \to 90°$ *and apex charges $Z = 1, 2, 4$ is likewise regression-pinned,* `tests/test_paper32_composition_pins.py`*)—the operator-algebra face of the Renner–Teller mechanism.* ***[OBSERVATION]*** *As an internal observation the configuration operator $F = \sum_i P_i$ carries, over molecular geometry, a lattice of conical intersections each with a π ($\mathbb{Z}_2$) Berry phase and, under bending, a structural Renner–Teller promotion; this is the* generic *topology of a symmetric-triatomic degeneracy, distinct from the molecule's physical electronic conical intersection, not an identity with it. (Computed and regression-pinned in* `tests/test_paper32_composition.py` *on validated two-center overlaps, with the commutant routine self-validated on cases of known answer; chronicle in CHANGELOG v5.1.0.)*

Remark 8 (The configuration operator's Berry curvature: a flat $\mathbb{Z}_2$ bundle over nuclear geometry)**.** *The configuration operator $F = \sum_i P_i$ of the previous remark varies over the nuclear-geometry manifold $\mathcal{M}$, and its geometric-phase structure is the finished form of the "lattice of conical intersections" noted there.* ***[SYMBOLIC PROOF]*** *Because the σ orbitals are real, F is real-symmetric at every geometry, so its Berry connection is* flat *with only a $\mathbb{Z}_2$ ($\pm$) holonomy: the curvature is not a smooth field but a sum of π-flux Dirac deltas at the conical intersections,*

$$\Omega \;=\; \sum_{CI} \pi\, \delta^2(m - m_{CI}),$$

the Longuet–Higgins sign-change / molecular Aharonov–Bohm structure (Longuet-Higgins; Berry; Mead–Truhlar) realized here for F rather than the electronic Hamiltonian. ***[SYMBOLIC PROOF]*** *On the linear-stretch plane (d_1, d_2) the Fukui–Hatsugai–Suzuki lattice flux is exactly* 0 *or* π *on every plaquette (forced by the reality of F; the machine-exact regression check is a consistency check, not independent evidence).* ***[MEASURED]*** *There are three π-sources—the central intersection at* $(2.45, 2.45)$ *and an antisymmetric-stretch mirror pair— carried by the two crossing bands and absent on the other four (the central source and the mirror pair are both regression-pinned,* `tests/test_paper32_composition_pins.py`*). The holonomy is the $\mathbb{Z}_2$ representation of $\pi_1(\mathcal{M} \setminus \mathrm{CI})$ sending a loop to $\pi \cdot (\#\, \text{enclosed} \bmod 2)$, cross-checked by direct parallel transport (π around one intersection and around all three).*

[MEASURED] *Each intersection is the real (equatorial) section of a charge-$\frac{1}{2}$ Berry monopole: in a fixed real frame the local 2×2 model is $F_0 + a_z\sigma_z + a_x\sigma_x$ with the σ_y coefficient* identically zero *(****[SYMBOLIC]****: forced by F real-symmetric)—the real family lives on the sphere's equator—and branching vectors $g = \nabla a_z$ along the symmetric stretch (the tuning mode) and $h = \nabla a_x$ along the antisymmetric stretch (the coupling mode), with nonzero winding.* ***[SYMBOLIC PROOF]*** *The $\mathbb{Z}_2$ (rather than $U(1)$) character is* forced by time-reversal symmetry *—the reality of the Coulomb overlaps—not by planarity: real orbitals give real overlaps in any geometry, so a non-coplanar real center leaves F real-symmetric and does not lift the phase. Breaking time reversal does.* ***[PANEL-VERIFIED]*** *A magnetic (Peierls) phase ϕ threading the* H–Be–H *loop makes F complex-Hermitian, opens the empty σ_y axis, and turns each intersection into a genuine $U(1)$ Berry monopole of integer charge: the first Chern number over a small sphere enclosing $(d^*, d^*, 0)$ in (d_1, d_2, ϕ) is ± 1 (opposite on the two crossing bands), and the real $\phi = 0$ plane cuts the monopole at its equator, recovering the π holonomy.* ***[OBSERVATION]*** *This is an internal observation about the geometry of the composition, distinct from the molecule's physical electronic conical intersection. (Regression-pinned in* `tests/test_paper32_berry_curvature.py`*; chronicle in CHANGELOG v5.1.0.)*

Remark 9 (Four centers: a continuous four-body modulus (the $\tilde{D}_4$ analog))**.** *The three-center geometric phase of Remark 8 is discrete ($\mathbb{Z}_2$ conical intersections). A* fourth *center adds a genuinely new,* continuous *four-body invariant. With $S_{ij} = Q_i^\top Q_j$ the metric-orthonormal overlaps between center subspaces, the loop holonomy of this overlap connection around the* 4*-cycle, $W = \mathrm{Tr}(S_{12}S_{23}S_{34}S_{41}) = \mathrm{Tr}(P_1P_2P_3P_4)$, is invariant under the per-center gauge $Q_i \mapsto Q_iO_i$ ($O_i \in O(r)$) and is therefore a physical configuration invariant. Two facts make it the four-center analogue of an elliptic modulus.* ***[PANEL-VERIFIED]*** *First, a* threshold*: for four rank-one (single-orbital) subspaces W is a function of the pairwise and triple traces—the 4×4 Gram is the entire configuration, so there is no four-body degree of freedom—whereas for rank ≥ 2 it is not (a local Jacobian-nullspace test gives residual $\gtrsim 0.3$). A genuine four-body invariant appears exactly when each center carries a σ–π pair, mirroring the rank-one→rank-two step in the tame four-subspace ($\tilde{D}_4$) problem.* ***[MEASURED]*** *Second, over molecular shape W is a* smooth continuous modulus*: for a planar rhombus with $\{1s, 2p\}$ per center it runs monotonically (from ≈ 0.8 to ≈ 0.04 over the sprint's diagonal-ratio range; the monotonicity and the sprint-range endpoints ($0.79 \to 0.042$) are both regression-pinned,* `tests/test_paper32_composition_pins.py`*) as the diagonal ratio varies —a one-parameter family of inequivalent four-center configurations, the continuous freedom the three-center case lacks. For real orbitals W is real (its phase is $\mathbb{Z}_2$); a magnetic (Peierls) flux through the* 4*-cycle makes* $\arg W$ *grow continuously from zero—the four-body $U(1)$ lift.* ***[OBSERVATION]*** *The object is the $*$-representation (non-abelian holonomy) analogue of the $\tilde{D}_4$ cross-ratio: four* orthogonal *projections are $*$-wild rather than tame, so the modulus is a matrix loop holonomy, not a single scalar cross-ratio. It is a geometry-axis invariant—a function of nuclear shape—distinct from the density-axis cross-ratio of the three-center integral in Paper 59 [66]. (Regression-pinned in* `tests/test_paper32_four_center_modulus.py`*; chronicle in CHANGELOG v5.1.0.)*

Remark 10 (The configuration operator is the composed basis's overlap metric)**.** *The three preceding remarks treat $F = \sum_i P_i$ as an abstract configuration operator. It is not abstract. Let G be the Gram matrix of the union of the per-center bases, with blocks $G_{ij} = \langle \chi_i | \chi_j \rangle$, and suppose each* intra*-center block is the identity—automatic when each center's own functions are orthonormal, and exact for the Shibuya–Wulfman metric of Paper 60 [57]. Put $X = \mathrm{chol}(G)^\mathsf{T}$, so that $X^\mathsf{T}X = G$ and the columns of X are the basis vectors expressed in an orthonormal ambient frame; let W_k collect the columns belonging to center k. Then $W_k^\mathsf{T}W_k = G_{kk} = I$, so W_k has orthonormal columns, the pseudoinverse in $P_k = W_kW_k^+$ collapses to W_k^T, and*

$$F \;=\; \sum_k W_k W_k^\mathsf{T} \;=\; XX^\mathsf{T} \;=\; X\,(X^\mathsf{T}X)\,X^{-1} \;=\; X\,G\,X^{-1}.$$

[SYMBOLIC PROOF] *Hence $F \simeq G$—*similar*, not merely cospectral. Every spectral statement in the preceding remarks (the conical intersections, the band gaps, the $\mathbb{Z}_2$ holonomy of the eigenvectors) is therefore a statement about the* overlap metric of the composed basis *over nuclear geometry.* ***[MEASURED]*** *Verified across the whole construction: linear* BeH_2*, bent* BeH_2 *(including the* H_2O *angle* $104.5°$*), and the four-center configuration, with* $\max|\mathrm{spec}\,F - \mathrm{spec}\,G| \leq 3 \times 10^{-15}$ *and* $\|FX - XG\|_{\max} \leq 9 \times 10^{-16}$*. The intra-center hypothesis is load-bearing rather than decorative: perturbing one intra-center block off the identity by* 0.3 *breaks the identity outright (spectral discrepancy* 6.7×10^{-2}*). The four-center holonomy of Remark 9 is the same statement read blockwise,* $\mathrm{Tr}(P_1P_2P_3P_4) = \mathrm{Tr}(G_{12}G_{23}G_{34}G_{41})$ *(residual* 3×10^{-16}*).*

Two consequences follow, one positive and one negative. ***[MEASURED]*** (i) An orthogonalization criterion. *Since the intersections are degeneracies of the metric, the eigenvectors of G are double-valued around one: parallel transport of the near-degenerate eigenvector around a*

small loop enclosing the central intersection returns it with overlap -1.0000*, against* $+1.0000$ *on a CI-free control. Any orthogonalization that* selects *metric eigenvectors—canonical orthogonalization with a drop threshold, the standard remedy for near-linear-dependence—is therefore path-dependent along a geometry scan through such a point, whereas symmetric (Löwdin) orthogonalization* $G^{-1/2}$*, being a function of the matrix rather than of its eigenvectors, returns bit-exactly (**[SYMBOLIC]**: single-valuedness of any matrix function; the numerical return is a consistency check). **[MEASURED]*** (ii) It is not a conditioning statement. *The intersections are* mid-*spectrum degeneracies (crossing eigenvalue* ≈ 0.71 *against* $\lambda_{\min} \approx 0.13$*), and* $\mathrm{cond}(G)$ *passes through* d^* *monotonically (*$19.9 \to 16.2 \to 15.0$*); the metric-conditioning growth of Paper 60 is a different functional of the same object, and no kink at* d^* *should be expected or claimed. The distinction drawn in the preceding remarks between* F *and the molecule's* physical electronic *conical intersection is unchanged; what changes is that* F *is an invariant of the framework's own encoding rather than a generic operator introduced for the analysis. (Regression-pinned in* `tests/test_paper32_config_operator_metric.py`*; chronicle in CHANGELOG v5.1.2.)*

6 The $K = \pi(B + F - \Delta)$ Formula in Spectral-Action Language

Paper 2 [29] observed the numerical coincidence $K = \pi(B + F - \Delta) \approx 1/\alpha$ at 4.77×10^{-7} (raw; 8.8×10^{-8} after the Sprint-A α^2 correction), with

$$B = 42, \quad F = \tfrac{\pi^2}{6}, \quad \Delta = \tfrac{1}{40},$$

and the Phase 4B–4I sprint series identified three independent spectral homes for the three pieces (project development-frontier log):

- B is the finite Casimir trace at $m = 3$: $B = \sum_{n=1}^{3}(2l+1)l(l+1)|_{n,l} = 42$ (Paper 2 Eq. 17).
- $F = D_{n^2}(d_{\max} = 4) = \zeta_R(2)$, the Fock-degeneracy Dirichlet series at the packing exponent (Phase 4F α-J).
- $\Delta^{-1} = g_3^{\mathrm{D}} = 2(3+1)(3+2) = 40$, the total (both-sign) Camporesi–Higuchi Dirac degeneracy at level $k = 3$, $|\lambda| = k + 3/2 = 9/2$ (Paper 7's g_3^{Dirac}; Phase 4H SM-D).

In spectral-triple language, B, F, Δ live in three different sectors of the same triple:

- B is a *finite scalar Casimir trace* on the truncated algebra: in the language of Sec. 4, it is $\mathrm{tr}_{\mathcal{A}_{\mathrm{GV}}}(C \cdot \not{K})$ at $n_{\max} = 3$, either for $C = L^2$ the SO(3) Casimir $l(l+1)$ (the sum printed above) or for $C = (n^2-1)/2$, half the SO(4) Casimir $L^2 + A^2 = n^2 - 1$; with the unhalved SO(4) Casimir the same trace is $84 = 2B$ (both routes' agreement at 42, and the un-halved trace at 84, are regression-pinned in `tests/test_paper32_B_normalisation.py`).
- F is an *infinite scalar spectral zeta*: it equals $\zeta_{1-\Delta_{S^3}}(2) = \sum_{n\geq 1} n^2 \cdot n^{-4} = \zeta(2)$ in the continuum limit, where Δ_{S^3} is the scalar Laplace–Beltrami operator on S^3 and $1 - \Delta_{S^3}$ has eigenvalue n^2 with the Fock degeneracy n^2. It is *not* $\zeta_{-\Delta_{S^3}}(2) = \sum_{n\geq 2} n^2/(n^2-1)^2 \approx 0.885$ (corrected 2026-09-30).
- Δ^{-1} is a *single-level Dirac degeneracy*: it equals g_3^{D}, the total multiplicity (both signs) of the Camporesi–Higuchi level $k = 3$, $|\lambda_3^{\mathrm{D}}| = 9/2$.

These three are categorically different objects: a finite combinatorial trace, an infinite analytic Dirichlet series, and a single-level multiplicity. They sit in different sectors of the spectral triple

(scalar/spinor; finite/infinite; trace/zeta/dimension) and **[PANEL-VERIFIED]** Phases 4B–4I exhausted nine attempts to find a common spectral generator that produces all three simultaneously, without success (twelve counting Sprint A and Sprint K-CC; see the E6 no-single-mechanism observation, Observation 3).

Observation 1 (Three-sector composition)**.** *The Paper 2 combination rule $K = \pi(B + F - \Delta)$ is a sum across three structurally distinct sectors of the GeoVac spectral triple. Each piece has been individually identified as a clean spectral invariant; the combination rule itself remains a numerical observation.*

This rephrasing is what working hypothesis WH5 in the project register is built on: α is the conversion factor between three projection regimes that do not share a generator. The spectral-triple framing does not derive α; it makes precise where the obstruction is.

Remark 11 (Sprint A, $\pi^3\alpha^3$ residual)**.** ***[MEASURED]*** *Sprint A (April 2026, working-hypothesis register) verified at 80 dps that the post-cubic residual $K - 1/\alpha - \alpha^2 = 1.2079 \times 10^{-5}$ matches $\pi^3\alpha^3 = 1.2049 \times 10^{-5}$ to within 0.25%. The structural reading is $\pi^3 = \mathrm{Vol}(S^5)$, hinting at a contribution from the Bargmann–Segal S^5 HO sector (Paper 24 [35]) at higher order. This is a structural hint, not a derivation; the cross-check (Paper 24's HO triple is a distinct spectral triple, not a sub-sector of the GeoVac triple) shows that the $\pi^3\alpha^3$ identification, if real, is a* cross-triple *relation and not internal to the GeoVac spectral-action expansion.*

7 The Coulomb/HO Asymmetry in Spectral-Triple Language

Paper 31 [39] formalized a layered asymmetry (Paper 24 §5 catalogues seven layers, six standing after the 2026-09-25 withdrawal of the spectrum-computing role of L_0 [retracted 2026-09-26: coulomb-ho-layer1-and-three-discriminators]; the table below tabulates structural rows, including that withdrawn one, marked; the CM Hodge structure is Paper 24 §5) between the Coulomb spectral triple of the present paper and the harmonic oscillator spectral triple of Paper 24 [35], summarized here:

[OBSERVATION] In spectral-triple language the asymmetry is structural: the Coulomb case is (in its continuum limit) a Riemannian-spin spectral triple of KO-dimension 3, the HO case is a spectral triple of complex Euler type on the holomorphic Hardy space of S^5. These are different objects even though both are "almost-commutative" in the broad sense. Calibration π enters in the Coulomb case through the $k = 0$ volume slot $\mathrm{Vol}(S^2)/4$ in the spectral-action coefficients; it does not enter the HO case because the HO spectral action is built from a linear (first-order) operator whose heat kernel does not produce π at the rational level of the truncation [35].

Observation 2 (Coulomb-specificity of π)**.** *The presence of π in the spectral-action coefficients of QED is specific to the Coulomb-projected S^3 triple of this paper, not a universal feature of GeoVac-style discretizations. The HO triple is π-free at the discrete level. This is the structural answer to "why does QED need π"—it needs π because it lives on the Coulomb-projection S^3 triple, where calibration π enters through the $k = 0$ volume slot $\mathrm{Vol}(S^2)/4$ (M1) and the spectral action's π^{2k} (M2).*

Theorem 3 (G3 closure: Bertrand-rigidity category obstruction)**.** ***[PROOF-BY-ARGUMENT; scoped to published constructions]*** *Within standard Connes spectral triples and Berezin–Toeplitz quantization, there is no tensor-product spectral triple $\mathcal{T}_{S^3} \otimes \mathcal{T}^{\mathrm{BL}}_{S^5}$ that simultaneously preserves:*

(a) the Riemannian-Dirac character of $\mathcal{T}_{S^3}$ (second-order conformal projection and spectral action built from D^2, spinor bundle, first-order Camporesi–Higuchi Dirac with spectrum $|\lambda_n| = n + 3/2$); and

(b) the Bargmann character of $\mathcal{T}^{\mathrm{BL}}_{S^5}$ (π-free certificate, Hardy-sector tower structure, first-order Euler spectrum $E_N = N + 3/2$, Paper 24).

The obstruction is categorical and rests on three premises:

- Bertrand's theorem (1873): *the only central potentials with all bounded orbits closed are $V = -Z/r$ (Coulomb) and $V = \omega^2 r^2$ (harmonic oscillator).*
- Fock rigidity (Paper 23 [34], Thm. 4): *the S^3 conformal projection is unique to $-Z/r$; the projection is second-order (stereographic / conformal), yielding a Riemannian spectral triple with the Laplace–Beltrami operator.*
- HO rigidity (Paper 24 [35], Thm. 3): *the S^5 Bargmann–Segal projection is unique to $\omega^2 r^2$; the projection is first-order (complex-analytic / holomorphic), yielding a Berezin–Toeplitz construction with the Euler number operator on the Hardy space.*

Proof sketch. Step 1 (category identification). Fock rigidity and HO rigidity place the two potentials in different mathematical categories: Coulomb → second-order conformal → Riemannian spectral triple; HO → first-order holomorphic → Berezin–Toeplitz quantization.

Step 2 (exhaustion of tensor-product options). Three constructions are available: (i) the standard graded tensor product with the round-S^5 Riemannian Dirac—this preserves (a) but violates (b) (π re-enters via the Seeley–DeWitt coefficients on S^5); (ii) the tensor product with the Bargmann–Euler operator—this preserves (b) but violates (a) (E_{BL} is not a Dirac-type operator on S^5, being the radial part of the Laplacian restricted to the holomorphic sector, not an elliptic differential operator on the total space); (iii) a hypothetical framework-extending construction beyond standard Connes spectral triples and Berezin–Toeplitz quantization—we know of no such framework in the published mathematics (a literature-absence premise, which is why the theorem is scoped to published constructions).

Step 3 (irreducibility). Bertrand exhausts the potentials; the two rigidity theorems fix the projection type for each. The category mismatch is a structural feature of central-force quantum mechanics, not an artifact of the GeoVac construction. The asymmetry layers of Paper 24 § V (seven catalogued, six standing; corrected 2026-09-03 from "five"; five are listed here, the remaining two, including the CM Hodge structure, are in Paper 24 § V) — spectrum-computing role of L_0 (*withdrawn* 2026-09-25: L_0 does not converge to Δ_{S^3} and is spectrally inert on both spheres); calibration π; Wilson gauge with natural matter coupling (a representation-theoretic layer, not an L_0 one); modular-Hamiltonian Pythagorean orthogonality; spectral-action gravity termination — are, the withdrawn one excepted, read here as downstream of this single categorical obstruction; the proof sketch argues the obstruction itself, not each layer's derivation from it. □

Remark 12 (G3 scope)**.** *Theorem 3 does not assert that S^3 and S^5 are unrelated — they are the two Bertrand geometries, deeply connected through the angular sparsity theorem (Paper 22). It asserts that their* spectral-triple-type structures *live in different mathematical categories, and no published framework bridges the gap while preserving both structural contents. Cross-manifold unification, if achievable, requires framework-extending mathematics beyond the scope of this paper and the published NCG literature.*

8 The π-Source Case-Exhaustion Theorem

[PANEL-VERIFIED] The empirical principle of Paper 35 [41] (Prediction 1, graduated to load-bearing status on a cumulative 208/208 confirmation across sprints KG and TX-B, May 2026) states that a GeoVac observable contains π if and only if its evaluation includes a continuous integration over a temporal or spectral parameter promoted from the discrete label/Dirac spectrum. Read in the spectral-triple language of §3–§4, this principle has a structural underpinning: the only places in the GeoVac framework where such continuous integrations occur are anchored to three named mechanisms of the present triple. **[CONDITIONAL]** We state the case-exhaustion below and give its proof sketch; the (i)⇔(ii) leg rests on the empirical Paper 35 verification rather than on a proof, and the converse (iii)⇒(i) is withdrawn (2026-09-30): no mechanism is known to put π into every one of its outputs (see the proof sketch).

Theorem 4 (π-Source Case-Exhaustion)**.** *Let $\mathcal{C} = (P_{i_k} \circ \cdots \circ P_{i_1})$ be any finite composition of projections drawn from the Paper 34 list [40] § III.1–§ III.15, applied to a Layer-1 datum on $(\mathcal{A}_{\mathrm{GV}}, \mathcal{H}_{\mathrm{GV}}, D_{\mathrm{GV}})$ or its Bargmann–Segal sibling triple. The following hold: (i)⇔(ii) (the empirical leg) and (ii)⇒(iii). The converse (iii)⇒(i) is withdrawn: it is not established (see the proof sketch):*

- *(i) The converged value $\mathcal{C}(\mathrm{datum})$ contains a transcendental of the form π, $\sqrt{\pi}$, π^k for $k \in \mathbb{Q}$, or π in a denominator.*
- *(ii) The chain $\mathcal{C}$ contains at least one projection P_{i_j} whose evaluation pipeline includes a continuous integration over a parameter promoted from the discrete label/Dirac spectrum (heat-kernel time, Matsubara frequency, Mellin contour, S^2 angular (volume-slot) measure, $\mathrm{SU}(N)$ Haar measure, or analytic-continuation parameter).*
- *(iii) The chain $\mathcal{C}$ engages at least one of three spectral-triple mechanisms:*
  - ***M1*** *(the $k = 0$ volume slot): introduces $\pi = \mathrm{Vol}(S^2)/4$ via the Hopf fibration $S^1 \to S^3 \to S^2$ (corrected 2026-09-03 from "$S^3 \to S^2 \times S^1$", which misstates the fibration and asserts a product S^3 is not), or its higher-rank analogues [36, 38].*
  - ***M2*** *(Seeley–DeWitt heat-kernel coefficient): introduces $\sqrt{\pi}$ via heat-kernel coefficients on the Camporesi–Higuchi Dirac spectrum on unit S^3 (Paper 28 Theorem T9 [37]: $\zeta_{D^2}(s)$ is a two-term polynomial in π^2 with rational coefficients at every integer $s \geq 1$; at half-integer $s \geq 5/2$ it carries odd zeta values instead). The sub-mechanism for the spinor sector is the half-integer Hurwitz $\zeta(s, 3/2)$.*
  - ***M3*** *(vertex-topology Dirichlet L-content): introduces Catalan $G = \beta(2)$ and Dirichlet $\beta(s)$ via quarter-integer Hurwitz shifts $\zeta(s, 3/4)$, $\zeta(s, 5/4)$ at two-loop sunset vertices (Paper 28 Theorem 3: $D_{\mathrm{even}}(s) - D_{\mathrm{odd}}(s) = 2^{s-1}(\beta(s) - \beta(s-2))$).*

Proof sketch. (i) ⇔ (ii) is Paper 35 Prediction 1, verified across 208 individual checks (sprints KG, TX-B, May 2026 [41]).

(ii) ⇒ (iii): the proof body that follows enumerates the fifteen Paper 34 projections in §III.1–§III.15 (the corpus state at this theorem's drafting; Paper 34 has since grown to twenty-eight projections [40], and the theorem's claim is here restricted to those first fifteen pending an extension of the case-by-case enumeration to projections 16–28). Each of the fifteen Paper 34 projections is either π-free at the projection step (projections 1, 3, 4, 5, 8, 9, 12, 13, 14: Fock conformal, Bargmann–Segal, Stereographic, Sturmian, Wigner $3j$, Wigner D, Molecule-frame, Drake–Swainson, Rest-mass) or π-bearing through a continuous integration that reduces to one of M1, M2, M3 by direct identification (Hopf bundle, Spectral action, Camporesi–Higuchi spinor lift, Wilson plaquette, Vector-photon promotion, Observation/temporal-window). The nine π-free projections live in either Paper 31's

universal sector (algebra-only, transfers to any spherical-fermion system) or Paper 22's potential-independent angular sparsity tier; neither sector contains π. The six π-bearing projections are tagged in `debug/track_ts_e1_theorem_promotion_memo.md` §2 to one or more of M1/M2/M3, with closed-form identification for each (Paper 25 for M1, Paper 28 T9 theorem for M2, Paper 28 Theorem 3 for M3).

The converse (iii) ⇒ (i) is withdrawn (2026-09-30): engaging a mechanism is not known to force π into the converged value. Paper 55 tags the rational Casimir trace $B = 42$ as M1's "rational core" (a tagging convention, not a proof that B is the converged value of a chain engaging M1); the spinor spectral zeta of M2's sub-mechanism at half-integer s carries no known π, e.g. $\zeta_{D^2}(5/2) = 14\,\zeta(3) - \frac{31}{2}\,\zeta(5)$ (Paper 28 T9; this lies outside the pure-Tate M2 ring, which is scoped to Seeley–DeWitt coefficients and integer s); M3's convergent closed form at $s = 4$ gives $2^3(\beta(4) - \beta(2)) = 8(\beta(4) - G)$, with no known π (none of $\zeta(3)$, $\zeta(5)$, G, $\beta(4)$ is known to lie in $\mathbb{Q}[\pi]$, nor proved not to). What remains closed-form is the forward identification per mechanism: M1's $\mathrm{Vol}(S^2)/4 = \pi$ and M2's heat-kernel $a_0(S^3) = \sqrt{\pi}$ carry π where those specific quantities enter. (Earlier versions asserted the converse for every mechanism, then for M1 and M2 only; both are withdrawn.) □

Corollary 2 (M1 sub-mechanism lies in the localised pure-Tate sub-ring generated by π)**.** *The M1 sub-mechanism (the $k = 0$ volume slot of the master Mellin engine, historically given a Hopf-base name after the factor $\mathrm{Vol}(S^2)/4 = \pi$ of Paper 25's $S^3 \to S^2$ bundle; that name is a misnomer for Paper 38's $4/\pi$, which is* twice *the unit-radius sphere-volume quotient $\mathrm{Vol}(S^2)/\mathrm{Vol}(S^3) = 2/\pi$ and not a base-to-total ratio (corrected 2026-09-03; the "$4/\pi$ is a quotient of unit-radius sphere volumes" wording printed here until then was itself wrong, since $\mathrm{Vol}(S^3) = 2\pi^2$, and it contradicted Paper 38)) produces period values in the localised pure-Tate sub-ring $\mathbb{Q}[\pi, \pi^{-1}]$ of mixed Tate periods over $\mathbb{Q}$, at depth 0 and Tate weight $\in \mathbb{Z}$. **[PANEL-VERIFIED]** Every M1 closed form witnessed across the corpus (Paper 25 Hopf base $\mathrm{Vol}(S^2)/4 = \pi$; Paper 38 §L2 asymptotic state-space GH rate $4/\pi = \mathrm{Vol}(S^2)/\pi^2$; Paper 40 [62] Thm 1 universal $4/\pi$ across compact connected semisimple Lie groups (rigorous at rank 1, with an analytical universality mechanism at proof-sketch tier and numerical support at rank 2 plus a noisier rank-3 datapoint; a circle factor carries $2/\pi$); and the M1 factor π^{-2} of Paper 50 [63] §3's F-theorem value $-3\zeta(3)/(16\pi^2)$, an M1×M3 cross-product whose $\zeta(3)$ is M3 content — a continuum Klebanov–Pufu–Safdi value re-derived there, not a finite-cutoff output) is a $\mathbb{Q}$-linear combination of π^n for $n \in \mathbb{Z}$, with $|n| \le 2$ on the empirical witnesses logged so far. Higher integer powers of π enter only through M2 (Seeley–DeWitt) or M3 (vertex-parity sums). The M1 ring does not engage M2's $\sqrt{\pi}$ vs π^2 volume-normalisation choice, because the volume slot is itself a volume integral and does not engage the heat-kernel asymptotic; it is* not *free of the Killing-form normalisation, however: under Kac's basic form the Paper 38 constant reads $2\sqrt{2}/\pi$ (discussion of Theorem 5; Paper 38), which lies in $\mathbb{Q}(\sqrt{2})[\pi, \pi^{-1}]$ rather than $\mathbb{Q}[\pi, \pi^{-1}]$, so the ring statement holds in the corpus's dual-Coxeter convention. Sprint M1 Pure-Tate (June 2026, memo `debug/sprint_m1_pure_tate_memo.md`).*

Corollary 3 (M2 sub-mechanism inherits Fathizadeh–Marcolli mixed-Tate classification, with pure-Tate refinement)**.** *At the standard volume-normalized Connes–Chamseddine convention, the M2 sub-mechanism Seeley–DeWitt coefficients on S^3 are mixed-Tate periods over $\mathbb{Q}$ by direct inheritance from Fathizadeh–Marcolli [7] (the static-R–W sub-case $a(t) \equiv 1$ on $\mathbb{R} \times S^3$). **[PANEL-VERIFIED]** The GeoVac S^3 M2 sector is in fact the strictly smaller pure-Tate sub-ring $\bigoplus_k \pi^{2k} \cdot \mathbb{Q}$ of even-weight Tate periods, with no $\zeta(3), \zeta(5)$, or MZV content appearing at the SD level. This refines F–M's continuum classification in two structural ways: (i) the Dirac SD expansion is two-term exact ($a_k^{D^2} = 0$ for $k \ge 2$, Paper 51 [54] Cor. 2.1), not generically infinite; (ii) the scalar Laplacian SD coefficients have the closed-form geometric-series collapse $a_k^{\Delta} = 2\pi^2/k!$. The M3*

sub-mechanism (vertex-parity Hurwitz, Catalan G, $\beta(s)$) lives in cyclotomic mixed Tate at level 4 *(mixed Tate over $\mathbb{Z}[i]$, not over $\mathbb{Q}$) and does* not *contaminate the SD coefficients, by the operator-order partition of Paper 18 [32] § III.7 ($k = 2$ for SD vs $k = 1$ for vertex parity). Sprint Mixed-Tate Test (June 2026, memo* `debug/sprint_mixed_tate_test_memo.md`*).*

Corollary 4 (M3 sub-mechanism inherits Deligne–Glanois cyclotomic mixed-Tate classification at level ≤ 4)**.** *The M3 sub-mechanism (vertex-parity Hurwitz / Dirichlet L / depth-k loop tower) on the discrete Camporesi–Higuchi S^3 spectral triple lies in the cyclotomic mixed-Tate period ring at level ≤ 4 over $\mathcal{O}_4[1/4] = \mathbb{Z}[i, 1/2]$ at depth equal to the loop order, by direct application of Deligne [8] (proven for $N \in \{2, 3, 4, 6, 8\}$) and the explicit basis of Glanois [9]. Three sub-rings are realised:*

$$\mathcal{MT}(\mathbb{Z}) \;\subset\; \mathcal{MT}(\mathbb{Z}[1/2]) \;\subset\; \mathcal{MT}(\mathbb{Z}[i, 1/2]),$$

selected by (i) un-restricted vs vertex-restricted summation, and (ii) loop order = motivic depth. Specifically:

- *Un-restricted Dirac Dirichlet $D(s)$ at integer $s \geq 2$ (Paper 28 [37] Thm 2) lies in $\mathcal{MT}(\mathbb{Z})$ at depth* 1*;*
- *Vertex-restricted $D_{\text{even}} - D_{\text{odd}} = 2^{s-1}(\beta(s) - \beta(s-2))$ at integer $s \geq 2$ (Paper 28 [37] Thm 3) lies in $\mathcal{MT}(\mathbb{Z}[i, 1/2])$ at level* 4*, depth* 1*, via Hurwitz at quarter-integer shifts; Catalan $G = \beta(2)$ and $\beta(4)$ are level-4 Dirichlet L-value periods in this ring;*
- *Two-loop irreducible $S_{\min}$ (Paper 28 §$S_{\min}$) lies in $\mathcal{MT}(\mathbb{Z}[1/2])$ at depth* 2*; the depth-k tower (Paper 28 Prop. depth_k) generates new irreducible periods at depth k = loop order in $\mathcal{MT}$ at level ≤ 4.*

The vertex-parity-as-χ_{-4} Dirichlet character mechanism IS the Deligne–Glanois Galois descent from level 4 *to level* 2 *(the $N' = 2 \mid N = 4$ instance of Glanois [9] Cor. 1.2) realised on the framework's natural QED observables. Combined with Corollary 3, the master Mellin engine of Theorem 4 now has a complete period-theoretic stratification: M1 in $\mathbb{Q}[\pi, 1/\pi]$, M2 (Seeley–DeWitt coefficients and integer-s values) in the pure-Tate sub-ring $\bigoplus_k \pi^{2k} \cdot \mathbb{Q}$, and M3 in $\mathcal{MT}(\mathbb{Z}[i, 1/2])$ at level ≤ 4. Specific motivic identification of irreducible constants at depth ≥ 2 ($S_{\min}$, $S^{(3)}$, …) within the Deligne–Glanois basis is an open question at each depth. Sprint M3 Cyclotomic Mixed-Tate (June 2026, memo* `debug/sprint_m3_cyclotomic_mixed_tate_memo.md`*).*

Remark 13 (Joint engagement)**.** *M1, M2, M3 are not set-theoretically disjoint as constructions: the spectral action invokes both M1 (the $k = 0$ volume slot on the spinor bundle) and M2 (Seeley–DeWitt coefficients) simultaneously, and the vacuum-polarization coefficient $\Pi = 1/(48\pi^2)$ (Paper 28 § V) contains one factor of π from each mechanism. The theorem requires only that* at least one *of the three be engaged; multiple engagement is the typical case at higher loop order. The mechanisms are role-disjoint in the Paper 18 § III.7 Mellin-engine sense (M1 = packing-tier ω_d, M2 = heat-kernel via Mellin/Gaussian, M3 = vertex-parity quarter-integer Hurwitz character mechanism). The Track TS-E3 sprint (May 2026) further sharpened this by showing that M1, M2, M3 are all sub-cases of a single master Mellin-engine mechanism $\mathrm{Tr}(D^k \cdot e^{-tD^2})$ on a compact Riemannian manifold, with $k \in \{0, 1, 2\}$ selecting the sub-mechanism.*

Remark 14 (Status of $K = \pi(B + F - \Delta)$)**.** ***[OBSERVATION]*** *Paper 2's combination $K = \pi(B + F - \Delta) \approx 1/\alpha$ matches its target at 4.77×10^{-7} (raw; 8.8×10^{-8} through the cubic's root after the α^2 correction). Under Theorem 4, K's chain (Fock ∘ Hopf∘ spinor) engages M1 (the explicit prefactor π, identified by Sprint A as $\mathrm{Vol}(S^2)/4$ [29]) and M2 (in $F = \pi^2/6 = D_{n^2}(d_{\max})$ via the*

Mellin transform of the scalar Fock-degeneracy). The π-content is therefore consistent with case-exhaustion. The remaining anomaly is at the depth-prediction *layer of Paper 34 § VIII.3 (depth predicts $\sim 3\%$ residual at three projections; K matches at 5×10^{-7}, the cubic's root at 10^{-7}): a quantitative-error puzzle, distinct from the qualitative case-exhaustion of the present theorem, and unaffected by it.*

Remark 15 (Falsification target)**.** ***[OPEN]*** *A counter-example to Theorem 4 would be a finite chain of the first fifteen projections introducing π via a fourth mechanism not in {M1, M2, M3}; a sixteenth projection doing so would fall outside the theorem's scope and would instead block its extension. Three natural candidates were investigated in Track TS-E3 (May 2026) and each REDUCED to combinations of the three existing mechanisms: continuum-gravity coupling (M1+M2), anomaly-class projections (M1+M2+M3 via the eta-invariant's first-order Mellin), and non-trivial principal-bundle projections (M1 normalisations against integer integrands). The strongest remaining computational falsification target is the discrete first-Chern-class invariant on the Hopf S^3 graph at $n_{\max} = 3$, expected π-free by the case-exhaustion reading; if a π appears unexpectedly, a fourth mechanism M4 would need to be named.*

Remark 16 (Mechanism-as-domain sharpening (Sprint MR-A/B/C, May 2026))**.** *A three-track follow-up sprint (MR-A: Dirac state-space GH rate; MR-B: spectral-action modular residual; MR-C: L2 next-order constant) tested whether the master Mellin engine of Paper 18 § III.7 is also* predictive *for convergence rates, not only classifying for π-sources. The combined verdict sharpens the case-exhaustion theorem into a mechanism-as-domain partition.*

- *M1 ($k = 0$) is the natural domain of state-space GH rates. The L2 sprint (2026-05-06) measured the asymptotic constant of the SU(2) scalar central Fejér kernel as $4/\pi$ (dual-Coxeter normalisation; $2/\pi$ on the unit S^3, Paper 38, 2026-09-03). The constant is the sphere-volume quotient $\mathrm{Vol}(S^2)/\mathrm{Vol}(S^3) = 2/\pi$ for unit-radius spheres, twice that in the dual-Coxeter normalisation; it is not the Hopf fibration's base-to-total ratio, whose base carries half the radius (Paper 38, volume-ratio note).*

- *M2 ($k = 2$) is the natural domain of heat-kernel and spectral-action convergence. Sprint MR-B derived a closed form for the Camporesi–Higuchi modular residual:*

$$\varepsilon(t) \;=\; \sum_{m \geq 1} (-1)^m \sqrt{\pi}\, e^{-m^2\pi^2/t} \left[t^{-3/2} - 2m^2\pi^2\, t^{-5/2} - \tfrac{1}{2}\, t^{-1/2} \right].$$

  Verified at > 100 digits at every tested t; every coefficient is $\sqrt{\pi}$ times an element of $\bigoplus_k \pi^{2k}\,\mathbb{Q}$; modular exponent exactly π^2 via Jacobi ϑ_2.

- *M3 ($k = 1$) is the natural domain of vertex-restricted parity-character observables (Paper 28 § QED-vertex: Catalan G, $\beta(4)$). Sprint MR-A demonstrated that asking for M3 from a half-integer-only Peter–Weyl truncation (the state-space GH machinery of Theorem 5) is a* category error*: the half-integer-spin truncation cannot span the constant function χ_0 on* SU(2)*, so the kernel does not localize and the moment is identically $\gamma_n^{\mathrm{Dirac}} = \pi$ for every $n_{\max} \geq 1$. The L2 scalar rate's $(4/\pi)\log(n)/n$ decay (dual-Coxeter normalisation; $2/\pi$ on the unit S^3) is an integer×half-integer interference effect; removing the integer sector kills it. This is exactly what the M1 mechanism predicts when the sub-bundle lacks the constant function.*

The reading is therefore: the index $k \in \{0, 1, 2\}$ classifies both *the sub-mechanism and the natural domain of observable from which its transcendental signature can be extracted. The case-exhaustion*

theorem 4 classifies which *transcendentals appear; the domain partition refines to* which observables *can produce them. Sprint MR-C extracted the L2 next-order constant* $c = 4.10932146748987562042\ldots$ *to* 22 *verified decimal digits (an earlier Richardson-extrapolated* ≥ 80*-digit string was contaminated from digit* 12 *by float-precision panel points; see Paper 38 § Open questions) but found no PSLQ identification in* $M_1 \cup M_2 \cup M_3 \cup \{\gamma_E, \log 2, G, \zeta(3)\}$ *at coefficient ceiling* 10^4*. This is consistent with c being a downstream Stein–Weiss IBP derivative of the M1 mechanism rather than a direct Mellin output, and is a partial negative against extending the predictive engine to all asymptotic-expansion coefficients along subsequent analytical pipelines. The engine is predictive* within *each domain. Source:* `debug/mr_a_dirac_propinquity_*`, `debug/mr_b_spectral_action_rate_*`, `debug/mr_c_l2_subleading_*`

Remark 17 (CH-substrate-specificity of the chirality-parity factorization (Sprint Q5'-S5-falsifier, 2026-06-05))**.** *The bit-exact* $\mathbb{Z}/2$ *chirality-parity factorization of the master Mellin source on the truncated Camporesi–Higuchi spectral triple (Sprint Q5'-CH-1) is CH-specific: it requires both spectral* $\pm$*-symmetry and chirality alignment* $\gamma_{ii} = \chi_i$, $\Lambda_{ii} = \chi_i|\lambda_i|$ *of the CH Dirac. The Bargmann–Segal Hardy-on-*S^5 *triple (Paper 24) has neither at any finite* $N_{\max}$*: the Hardy spectrum is strictly positive (no* $\pm$*-partner) and the* $(N, 0)$ SU(3) *tower carries no canonical chirality grading (the two natural candidates* $\chi_N = (-1)^N$ *and* $\chi_l = (-1)^l$ *coincide because* $(N, 0)$ *branching forces l-parity* $\equiv N$*-parity, and neither reproduces the CH-1 even-j zeros). The* ω^{tri} *enrichment program of Paper 55 [64] § subsec:open_m2_m3 correctly targets the CH triple as substrate; the case-exhaustion classification 4 keeps its stated scope (conditional, first fifteen projections) but its bit-exact* finite-cutoff *witness form is CH-substrate-specific. This places a sixth layer on the Coulomb/HO asymmetry of Paper 24 § V. Source:* `debug/sprint_q5p_s5_falsifier_memo.md`.

Remark 18 (Stage-1 cochain-level witness of the master-Mellin partition (Sprint Q5'-Stage1-JLO-nmax2, 2026-06-05))**.** ***[PANEL-VERIFIED]*** *On the truncated Camporesi–Higuchi spectral triple at* $n_{\max} = 2$*, the master-Mellin partition* (M_1, M_2, M_3) *is bit-exactly visible at the cyclic-cochain level as the symbol* $\omega^{\mathrm{tri}}(\mathcal{T}_2) = (16, 84, 36) \in \mathbb{Z}^3$*. The first two slots live inside a single JLO degree-0 cochain:* M_1 *at* t^0 *of* $\phi_0^{\mathrm{odd}}(1; t) = \mathrm{Tr}\, e^{-tD^2}$*,* M_2 *at* $-t^1$ *of the same.* M_3 *lives outside the JLO framework on commutative* $\mathcal{A} = \mathbb{C}^5$ *and instead in the Connes–Moscovici residue framework as the leading coefficient of* $\mathrm{Tr}(\gamma\, D\, e^{-tD^2})$*. The first non-trivial JLO cochain degree on commutative* $\mathcal{A}$ *is* $n = 2$*; structural vanishing* $\phi_1(a_0, a_1; t) \equiv 0$ *is proved bit-exactly for all idempotent pairs* (a_0, a_1)*. The cutoff sweep (Sub-Sprint 2a) extends the symbol bit-exactly to* $n_{\max} \in \{1, 2, 3, 4\}$ *via the closed forms* $M_1(n) = 2n(n{+}1)(n{+}2)/3$*,* $M_2(n) = n(n{+}1)(n{+}2)(2n{+}1)(2n{+}3)/10$*,* $M_3(n) = n(n{+}1)^2(n{+}2)/2$*, all re-derived from the chirality-balanced shell degeneracy* $2n(n{+}1)$ *and the CH eigenvalue* $n{+}1/2$*. See Paper 55 § subsec:open_m2_m3 for the Stage-1 Q5' framing. Source:* `debug/sprint_q5p_jlo_nmax2_memo.md`, `debug/sprint_q5p_2a_jlo_nmax_sweep_memo.md`.

Remark 19 (Mellin extraction points of the master-Mellin partition (Sprint Q5'-Stage1-2b, 2026-06-05))**.** *The master Mellin partition* (M_1, M_2, M_3) *corresponds to three categorically different extraction points of the formal Mellin transform of the JLO degree-0 cochain on a compact 3-dimensional spectral triple:*

- M_1 *at the spectral-dimension pole* $s = d/2 = 3/2$ *of* $\Gamma(s)\,\zeta_{D^2}(s)$*: the Weyl-law short-time leading* $\mathrm{Tr}(e^{-tD^2}) \sim \sqrt{\pi}/2 \cdot t^{-3/2}$ *supplies the* $k = 0$ *volume-slot* π *residue.*
- M_2 *at integer-s regular points of the same* $\Gamma(s)\,\zeta_{D^2}(s)$*: Seeley–DeWitt pure-Tate continuum values (e.g.,* $\zeta_{D^2}(2) = \pi^2 - \pi^4/12$ *on unit CH) reproduce the Sprint Q5'-CH-2 panel bit-exactly.*
- M_3 *at the analogous extraction on a structurally different Mellin complex, the* η*-style pairing* $\Gamma(s)\,\mathcal{M}[\mathrm{Tr}(\gamma\, D\, e^{-tD^2})](s)$*: vertex-parity Hurwitz content at quarter-integer shifts.*

At finite $n_{\max}$ the spectral zeta is bit-exact rational (e.g., $\zeta^{(2)}_{D^2}(1) = 1\,978\,465\,196\,515\,232/532\,976\,619\,280\,625$ for the full Dirac at $n_{\max} = 2$, $\kappa = -1/16$); the $s \to 0$ residue is exactly $\dim \mathcal{H} \in \mathbb{Z}$ (no π). Master-Mellin transcendentals enter only in the $n_{\max} \to \infty$ continuum limit; the cosmic-Galois Stage 2 motivic-Galois action acts at continuum, not at finite cutoff. Source: `debug/sprint_q5p_2b_phi0_mellin_memo.md`

Remark 20 (Continuum residue identification of the master-Mellin M_1 volume-slot coefficient (Sprint Q5'-Stage1-2b-continuum, 2026-06-05))**.** *The Mellin extraction-points reading of Remark 19 is closed at theorem grade for the M_1 component on the truncated Camporesi–Higuchi spectral triple of dimension $d = 3$. The continuum spectral zeta $\zeta^{\mathrm{cont}}_{D^2}(s) = 2\,\zeta(2s-2, 3/2) - \frac{1}{2}\,\zeta(2s, 3/2)$ has a simple pole at the spectral-dimension value $s = d/2 = 3/2$ with bit-exact meromorphic residue 1 (verified two independent ways: from the Hurwitz pole at $u = 2s - 2 = 1$ with residue 1, and from direct Laurent expansion via the digamma identity). The Mellin residue of $\Gamma(s)\,\zeta^{\mathrm{cont}}_{D^2}(s)$ at $s = 3/2$ is therefore $\Gamma(3/2) \cdot 1 = \sqrt{\pi}/2$, which matches the standard heat-kernel Weyl-law leading coefficient $r\,\mathrm{Vol}(S^3)/(4\pi)^{3/2}\big|_{r=2}$ ([23, 24]) bit-exactly. The three published instances of the M_1 volume slot ($\mathrm{Vol}(S^2)/4 = \pi$ in Paper 18 § III.2; $\mathrm{Vol}(S^2)/\pi^2 = 4/\pi$ (dual-Coxeter, a convention-dependent constant; $2/\pi$ at unit radius) in Paper 38 (Lemma L2) as the L2 asymptote rate constant; $\Gamma(d/2) = \sqrt{\pi}/2$ here at the Mellin pole) are **[OBSERVATION]** sibling normalizations of the same spectral object, related by exact factors of $\pi^2/4$ and $\pi^{3/2}/8$ with no spectral-data input. The M2 continuum panel at integer $s \in \{1, 2, 3, 4, 5\}$ reproduces the Sprint Q5'-CH-2 Seeley–DeWitt panel bit-exactly (5/5 matches). Tauberian rate uniformity in a neighborhood of $s = 3/2$ is the Stage-2-relevant open question, reachable from Karamata [25] and Korevaar [26]. Source:* `debug/sprint_q5p_continuum_mellin_memo.md`.

Remark 21 ($\mathrm{HP}^{\mathrm{even}}$ class identification on the bicomplex of the truncated CH triple (Sprint Q5'-Stage1-2c-Bicomplex, 2026-06-05))**.** *The cyclic (b, B) bicomplex of the truncated Camporesi–Higuchi spectral triple at $n_{\max} = 2$ has been constructed bit-exactly. The load-bearing degree-1 entire-cyclic cocycle condition $b\phi_0 + B\phi_2 = 0$ holds bit-exactly on the full panel of 36 idempotent-pair inputs at three t-orders t^0, t^1, t^2, both even and odd flavors (216 bit-exact zero residuals). The explicit $\mathrm{HP}^{\mathrm{even}}$ class on the Morita-trivial baseline $\mathbb{Q}^5 = \mathrm{HP}_0(\mathbb{C}^5)$ is*

$$[\phi_{\mathrm{JLO}}]^{\mathrm{HP}^{\mathrm{even}}} \;=\; (+2, -2, +2, +2, -4) \in \mathbb{Q}^5,$$

the sector-resolved McKean–Singer index summing to $\mathrm{Tr}(\gamma) = 0 = \chi(S^3)$. The Sub-Sprint 1 symbol $(16, 84, 36)$ is therefore a JLO cocycle representative, not a coboundary. A structural finding (honest scope): at degree 3, $(b\phi_2 + B\phi_4)$ has bit-exact residual $\pm 1/(3 \cdot 2^{16}) = \pm 1/196\,608$ on specific palindromic 4-tuples, a JLO-tower truncation artifact tied to the structural vanishing of odd-degree cochains $\phi_1 \equiv \phi_3 \equiv 0$ on commutative $\mathcal{A}$. The CM-residue cocycle for M_3 is single-cochain and immune to this artifact; the JLO/CM-residue distinction is Stage-2-relevant. Source: `debug/sprint_q5p_2c_bicomplex_memo.md`.

Remark 22 (CM-residue η-class identification, Sprint Q5'-Stage1-CM-Bicomplex, June 2026)**.** *The second span of the Stage-1 cosmic-Galois U^* bridge — the Connes–Moscovici η-pairing residue cocycle on the truncated Camporesi–Higuchi spectral triple at $n_{\max} = 2$ — has been constructed bit-exactly in* sympy.Rational *(*`debug/sprint_q5p_cm_bicomplex_memo.md`*,* `debug/data/sprint_q5p_cm_bicomplex.js`*. The CM-η cochain $\psi^\eta_n(a_0, \ldots, a_n; t) := \int_{\Delta_n} \mathrm{Tr}(\gamma D\, a_0\, e^{-s_0 t D^2}\,[D, a_1] \cdots [D, a_n]\, e^{-s_n t D^2})\, ds$ replaces the JLO leftmost insertion γ by γD, and shares the bicomplex structure $(b + B)\psi = 0$ with JLO via the same formulas. The degree-1 cocycle condition $b\psi^\eta_0 + B\psi^\eta_2 = 0$ holds bit-exactly on the full panel of 36 idempotent-pair inputs at three t-orders both flavors (216 bit-exact zero residuals), with NO truncation artifact of the JLO $\pm 1/196608$ type. The CM-η class on the Morita-trivial baseline*

$\mathbb{Q}^5 = \mathrm{HP}_0(\mathbb{C}^5)$ *is explicitly*

$$[\psi_{\mathrm{CM}}^{\eta,\mathrm{even}}]^{\mathrm{HP}_\eta} = (3, 3, 5, 15, 10) \in \mathbb{Z}^5,$$

summing to $36 = M_3(n_{\max} = 2)$ *(the Sub-Sprint 1 master-Mellin M3 ground truth). The sector-resolved structure is the chirality-symmetrized eigenvalue magnitude per sector* $\eta_s = \dim_s \cdot (n_s + 1/2)$*, in clean dual to the JLO* $\chi_s = \dim(\gamma_+ e_s \mathcal{H}) - \dim(\gamma_- e_s \mathcal{H})$ *chirality balance per sector. The two classes share the same* K_0 *sector-decomposition* $\{[e_s]\}_{s=0..4}$ *but encode distinct cohomological content: JLO captures the zeroth heat-kernel supertrace coefficient (McKean–Singer index* $\chi(S^3) = 0$*), CM-*η *captures the leading* t^0 η*-density supertrace coefficient (*M_3 *mechanism leading). Cross-check at* $n_{\max} = 3$*:* $M_3(3) = 120$ *bit-exact, CM-*η *degree-1 cocycle bit-exact zero on all 9 diagonal sectors. See Paper 55 § subsec:open_m2_m3 for the Stage-1 Q5' construction context.*

Remark 23 (Q5' Stage 1 pro-system functoriality of the JLO and CM-η classes (Sprint Q5'-Stage1-Prosystem, June 2026))**.** *The Stage-1 cocycle classes form a strictly compatible inverse system across the cutoff pro-system* $\{\mathcal{T}_{n_{\max}}\}_{n_{\max}\geq 1}$ *(`debug/sprint_q5p_prosystem_memo.md`, `debug/data/sprint_q5p_prosystem.json`). Sector idempotents are indexed by* (n, l) *with* $1 \leq n \leq n_{\max}$*,* $0 \leq l \leq n$*, giving* $N(n_{\max}) = n_{\max}(n_{\max} + 3)/2$ *sectors at each cutoff (2, 5, 9, 14 at* $n_{\max} = 1, 2, 3, 4$*, with strict nesting); the truncation* $P_{n+1\to n} : \mathcal{O}_{n+1} \to \mathcal{O}_n$ *is sector projection (algebra homomorphism), the algebraic Berezin* $B_{n\to n+1}$ *is sector inclusion. Both the JLO* $\mathrm{HP}^{\mathrm{even}}$ *class* $\chi_{(n,l)}$ *and the CM-*η *class* $\eta_{(n,l)}$ *are* sector-local *— the per-sector value depends only on the sector label, not on the cutoff:*

$$\chi_{(n,l)} = \begin{cases} +2 & l < n \\ -2n & l = n \end{cases}, \qquad \eta_{(n,l)} = \begin{cases} (2l+1)(2n+1) & l < n \\ n(2n+1) & l = n \end{cases}.$$

Bit-exact pull-back identity $P_{n+1\to n}^*[\psi^{(n+1)}] = [\psi^{(n)}]$ *holds across all three consecutive cutoff pairs for both classes (six inverse-image identities, each with bit-zero residual); algebraic Berezin compatibility holds with the same bit-exactness on the common sectors. Total sums match the Sub-Sprint 2a polynomial closed forms* $M_1(n_{\max}) = 2n_{\max}(n_{\max}+1)(n_{\max}+2)/3$*,* $\sum_s \chi_s = 0$ *(McKean–Singer index* $\chi(S^3) = 0$ *preserved cutoff-uniformly),* $\sum_s \eta_s = M_3(n_{\max}) = n_{\max}(n_{\max}+1)^2(n_{\max}+2)/2$ *bit-exactly at all four cutoffs. The pro-system is the cleanest possible inverse-system lift of the cohomological Stage-1 data: strict at the class level (no twist, no coboundary modification), consistent with the sector-locality of the Camporesi–Higuchi shell decomposition. The strict-strong-form question — pro-system functoriality at the level of* cochain morphisms *in the entire-cyclic bicomplex (vs the class level closed here) — is sharpened to a bit-exact dichotomy in Remark 25; the deep NCG-mathematics sub-question on entire-cyclic functoriality under algebra truncations (which we believe open in the literature; we know of no treatment in Connes 1994 Ch. IV, but have not established this) remains open, beyond sprint scale, independent of the Stage-1 closure here. See Paper 55 § subsec:open_m2_m3 for the Q5' Stage 1 construction context.*

Remark 24 (Q5' Stage 2 candidate Hopf algebra substrate, Sprint Q5'-Stage2-Hopf, June 2026)**.** *The first scoping step of Stage 2 of the cosmic-Galois* U^* *program produces a concrete bit-exact finite-cutoff candidate Hopf algebra* $\mathcal{H}_{\mathrm{GV}}^{(n_{\max})} = \mathrm{Sym}_{\mathbb{Q}}(V_{n_{\max}})$ *on the pro-system substrate of Remark 23, where* $V_{n_{\max}}$ *is the* $\mathbb{Q}$*-vector space of primitive generators* $x_{(n,l),k}$ *indexed by* $(n, l) \in \mathcal{S}_{n_{\max}}$ *and Mellin slot* $k \in \{0, 1, 2\}$ *(with* $k = 0, 1, 2$ *corresponding to M1, M3, M2 of §8, respectively). With primitive coproduct* $\Delta(x_g) = x_g \otimes 1 + 1 \otimes x_g$*, counit* $\varepsilon(x_g) = 0$*, and antipode* $S(x_g) = -x_g$ *on generators (extended as algebra homomorphisms),* $\mathcal{H}_{\mathrm{GV}}$ *satisfies all five Hopf axioms bit-exactly at* $n_{\max} \in \{2, 3\}$ *(269 bit-exact zero residuals on the axiom panel: 52 coassociativity, 104 counit-left/right, 104 antipode-left/right, 9 bialgebra-compatibility), and the* k*-grading is preserved by* Δ *so that* $\mathcal{H}_{\mathrm{GV}} = \mathcal{H}_{\mathrm{GV}}^{[0]} \otimes \mathcal{H}_{\mathrm{GV}}^{[1]} \otimes \mathcal{H}_{\mathrm{GV}}^{[2]}$ *factors as a tensor product of three Hopf sub-algebras (42 bit-exact generator-level* k*-preservation checks). The pro-system truncation* $P_{n+1\to n}$ *of Remark 23 lifts*

bit-exactly to a Hopf-algebra homomorphism $\mathcal{H}_{\mathrm{GV}}^{(n+1)} \to \mathcal{H}_{\mathrm{GV}}^{(n)}$ (126 bit-exact zero residuals on the truncation panel at $(n_{\max}, n_{\max}-1) \in \{(2,1),(3,2)\}$). The candidate motivic Galois group at finite cutoff is the additive affine group $U_{\mathrm{GeoVac}}^{*(n_{\max})} = \mathbb{G}_a^{3N(n_{\max})}$ (dimensions 15 and 27 at $n_{\max} = 2, 3$), with the Mellin-slot factorisation $U_{\mathrm{GeoVac}}^{*(n_{\max})} = \prod_{k=0}^{2} \mathbb{G}_a^{N(n_{\max})}$ structurally enforcing the M1/M2/M3 partition of Theorem 4 at the Stage-2 cosmic-Galois level. The candidate is the abelian primitive-cospan *of the Connes–Kreimer renormalization Hopf algebra (rooted trees in Connes–Kreimer 1998, arXiv:hep-th/9808042; Feynman graphs in Connes–Kreimer 2000, arXiv:hep-th/9912092): same generator-and-grading shape, same pro-system truncation structure, but the sub-graph coproduct of CK is replaced by the primitive coproduct here because the CH Fock sector idempotents are sector-disjoint (no proper sub-sector nesting). The natural continuation toward Stage-2's full Tannakian construction (Connes–Marcolli 2004, arXiv:math/0409306) — a substantial open program — is to enrich the coproduct with a sub-sector or nested-Hopf-tower structure that introduces non-abelian pro-unipotent content. See Paper 55 § subsec:open_m2_m3 for the cosmic-Galois narrative.*

Remark 25 (Q5' Stage 1 strict-strong-form cochain-morphism drift, Sprint Q5'-Stage1-StrictStrong, June 2026)**.** *Sharpening the v3.60.0 honest-scope caveat of Remark 23: while pro-system functoriality at the level of cocycle CLASSES is bit-exactly strict across the cutoff pro-system $\{\mathcal{T}_{n_{\max}}\}_{n_{\max}\geq 1}$ on commutative $\mathcal{A}$, pro-system functoriality at the COCHAIN-MORPHISM level of the entire-cyclic bicomplex FAILS bit-exactly on the same algebra (`debug/sprint_q5p_strict_strong_memo.md`, `debug/data/sprint_q5p_strict_strong.json`). The degree-3 closure residual $(b\phi_2 + B\phi_4)$ on the load-bearing (e_2, e_3) palindromic 4-tuples carries bit-exact values*

$$n_{\max} = 2 : \pm 1/(3 \cdot 2^{16}) \quad \to \quad n_{\max} \geq 3 : \pm 1/2^{16} \quad (\textit{bit-exact fixed point at } n_{\max} = 3, 4),$$

with the pull-back cochain increment $\Delta(B\phi_4)_{3\to 2} = -1/(3 \cdot 2^{14})$ closing the residual identity bit-exactly: $1/196608 + (-1/49152) = -1/65536$. The CM-$\eta$ cocycle tower exhibits the same structural pattern at degree 3 with structurally distinct quantitative values (the v3.59.0 reading that the CM-η tower is the cochain-clean alternative is restricted to degree 1 and does not extend to degree 3). The pro-system functoriality verdict is therefore a sharp dichotomy: strict at the class level (Remark 23), bit-exact non-strict with $n_{\max} = 3$ fixed-point structural drift at the cochain-morphism level (this remark). The drift is consistent with a boundary-vs-interior shell partition: $n_{\max} = 2$ is the boundary cutoff (no shell above $n = 2$ in the ϕ_4 matrix-element channel), $n_{\max} \geq 3$ are interior cutoffs (stabilized). Empirical confirmation at bit-exact precision of a question we believe open in the literature (entire-cyclic functoriality under algebra truncations; no treatment known to us in Connes 1994 Ch. IV). See Paper 55 § subsec:open_m2_m3 for the Stage-1 Q5' construction context.

Remark 26 (Tauberian rate uniformity in a neighborhood of $s = 3/2$ (Sprint Q5'-Stage1-Tauberian, 2026-06-05))**.** *The pointwise rate of approach in Remark 20 is sharpened to a uniform rate on compact above-pole neighborhoods of the spectral-dimension pole $s = d/2 = 3/2$ via the standard uniform Tauberian theorem of Karamata [25] (foundational), Korevaar [26] § III.4 (modern treatment with boundary-saturation Remark 3), and Tenenbaum [27] § II.7 (Tauberian theorems; the specific theorem number and the claim of uniform rates with explicit constants are not verified against the text, and complex Tauberian theorems of Wiener–Ikehara type do not in general give rates, so the rate content here should be read as resting on the direct tail estimate rather than on this citation). The four hypotheses transport bit-exactly to the CH spectral zeta: (H1) positivity $a_n = 2n(n+1) > 0$; (H2) polynomial growth $a_n = O(n^2)$; (H3) meromorphic continuation with unique simple pole at $\sigma_0 = 3/2$ of residue 1 (the v3.59.0 Track 2 closure); (H4) vertical-strip boundedness from standard Hurwitz convexity. Conclusion: on the compact above-pole neighborhood $U_> = [3/2 + \epsilon,\, 3/2 + 1/4]$, $\sup_{s\in U_>} |\zeta_{D^2}^{(N)}(s) - \zeta_{D^2}^{\mathrm{cont}}(s)| = O(N^{-2\epsilon+\delta})$ uniformly in $U_>$, for any $\delta > 0$. Boundary-saturation:*

the supremum is attained at the boundary point closest to the pole, verified bit-exactly at $n_{\max} \in \{2,3,4,5\}$ *on the six-grid panel* $\{5/4, 11/8, 23/16, 25/16, 13/8, 7/4\}$ *(sup attained at* $s = 25/16$ *at every cutoff; empirical slope* -0.092 *at* $n_{\max} \in \{2,\dots,5\}$ *monotonically approaches the predicted* $-1/8$*, hitting* -0.123 *at* $n_{\max} = 100$*). Source:* `debug/sprint_q5p_tauberian_memo.md`.

Remark 27 (Continuum residue identification of the master-Mellin M_3 η-tower (Sprint Q5'-Stage1-M3-Continuum, 2026-06-05))**.** *The continuum* M_3 *η-Mellin object on the truncated Camporesi–Higuchi spectral triple of dimension* $d = 3$ *is closed at theorem grade with parity-of-s stratification. The continuum η-Dirichlet series* $\eta_D(s) := \sum_{n \geq 0} g_n |\lambda_n|^{1-s}$ *admits the Hurwitz reduction* $\eta_D(s) = 2\zeta(s-3, 3/2) - \frac{1}{2}\zeta(s-1, 3/2)$ *(equivalently* $\eta_D(s) = D(s-1)$ *via Paper 28's Dirac Dirichlet series), with TWO simple poles at* $s = 4$ *and* $s = 2$ *of bit-exact meromorphic residues* 2 *and* $-1/2$ *respectively (verified two independent routes: Hurwitz pole rule at* $u = 1$ *AND direct Laurent via digamma). The corresponding Mellin residues* $\mathrm{Res}_{s=4}\Gamma(s)\eta_D(s) = 12$ *and* $\mathrm{Res}_{s=2}\Gamma(s)\eta_D(s) = -1/2$ *are rational integer values, NOT in the M3 cyclotomic ring; the genuine M3 cyclotomic content lives in the CHIRALITY-RESOLVED η-Mellin* $(D_{\mathrm{even}} - D_{\mathrm{odd}})(s-1)$*, which inherits the Sprint Q5'-CH-3 even-s M3 stratification via the* $\mathbb{Q}(i) = \mathbb{Q}(\zeta_4)$ *Galois descent. Specifically at* $s = 3$ *the chirality discriminant equals* $-1 + 2G$ *(Catalan G; genuine M3 depth 1); at* $s = 5$ *it equals* $8\beta_4 - 8G$*; at* $s = 7$ *it equals* $-32\beta_4 + 32\beta_6$ *(both genuine M3 depth 1+). Odd-*$(s-1)$ *values collapse to* M_1 *via* $\beta(2k+1) \in \pi^{\mathrm{odd}}\mathbb{Q}$*. The discrete-side Karamata Tauberian signature at the* $s = 4$ *pole holds exactly: partial sums on the v3.60.0 sector-local substrate satisfy* $S(n_{\max})/\log(n_{\max}) \to 2$ *at* $n_{\max} \geq 200$ *(meromorphic residue; no Jacobian since the natural η-Mellin variable IS the spectral* $|\lambda| = n + 1/2$*). This closes the* M_3 *component of the continuum-residue identification triad started by Remark 20. Structural sharpening: unlike* M_1 *(which admits three exact-factor sibling normalizations* π*,* $4/\pi$*,* $\sqrt{\pi}/2$ *of a single generator),* M_3*'s ring* $\mathcal{MT}(\mathbb{Z}[i, 1/2], 4)$ *is depth-graded with generators* $\{1, G, \beta_4, \beta_6, \dots\}$ *at successive depths with no* π*-power reduction at any depth* ≥ 1 *(a transcendence independence of Catalan G from* $\pi^{2k}\mathbb{Q}$ *— conjectural: even the irrationality of G is open).* Conjecturally, this asymmetry is the operational content of the master-Mellin partition M_1 *at* $k = 0$ *depth* 0 *vs* M_3 *at* $k = 1$ *depth* $1+$*. Source:* `debug/sprint_q5p_m3_continuum_memo.md`.

Remark 28 (Q5' Stage 2 substrate enrichment via $J^*(S^3)$, Sprint Q5'-J-Star-S3, June 2026)**.** *A first scoping step of the nested-Hopf-tower substrate enrichment of Remark 24 (a substantial open program) replaces the disjoint sector-idempotent substrate* $\mathbb{C}^{N(n_{\max})}$ *with the SU(2) Peter–Weyl matrix-coefficient substrate*

$$J^*_{j_{\max}}(S^3) = \mathrm{span}_{\mathbb{Q}}\{\pi^j_{mn} : 0 \leq j \leq j_{\max},\ -j \leq m, n \leq j\},$$

with the matrix-coefficient coproduct $\Delta\pi^j_{mn} = \sum_p \pi^j_{mp} \otimes \pi^j_{pn}$*. This coproduct is bit-exactly* non-primitive *on every* $j > 0$ *generator (verified at* $j_{\max} \in \{1/2, 1, 3/2\}$ *on 29 generators with* $j > 0$*); coassociativity and counit hold bit-exactly in the free polynomial algebra (147 zero residuals); the antipode axiom requires passing to the standard* $\mathcal{O}(SU(2)) \cong \mathcal{O}(SL_2)$ *quotient by SU(2) unitarity relations (textbook construction; Klimyk–Schmüdgen 1997 §1.3.2); the pro-system truncation* $P_{j_{\max}+1/2 \to j_{\max}}$ *lifts bit-exactly to a Hopf-algebra homomorphism (132 zero residuals). The candidate motivic Galois group at the quotient is* $U^{*(j_{\max})}_{\mathrm{GeoVac},J^*} = SL_2$ *for every* $j_{\max} \geq 1/2$ *— a non-abelian semisimple algebraic group, achieving the qualitative goal of breaking the abelian primitive shape of Remark 24. The trade is axis-orthogonal:* $J^*(S^3)$ *has non-abelian content but sacrifices the Mellin-slot k-partition; v3.61.0's* $\mathcal{H}_{\mathrm{GV}}$ *has the Mellin partition but is abelian. The natural continuation, beyond sprint scale, is a synthesis combining both — either (a) tensor product* $\mathcal{H}_{\mathrm{v3.61}} \otimes \mathcal{H}^{J^*}$ *giving* $U^* = \mathbb{G}_a^{3N} \times SL_2$*, or (b) Mellin-slot-decorated Peter–Weyl* $\pi^{j,k}_{mn}$ *giving* $U^* = SL_2^3$*. The* $SL_2 \times \mathbb{G}_a^d$

form would be the Levi decomposition of a general algebraic group (semisimple times pro-unipotent), suggesting the synthesis substrate is the right Stage-2 target. See Paper 55 § subsec:open_m2_m3.

Remark 29 (Q5' Stage 2 cross-shell off-diagonal Dirac enrichment scoping, Sprint Q5'-OffDiag-Dirac, June 2026)**.** *The first scoping step of the cross-shell off-diagonal Dirac perturbation enrichment — itself a substantial open program — (second of three structural ingredients flagged in Remark 24) promotes the v3.61.0 pro-system substrate to track the off-diagonal Dirac perturbation κA via transition generators $T_{s' \to s} := e_s \cdot (\kappa A) \cdot e_{s'}$ (non-zero for E1-Gaunt-selected sector pairs). At $n_{\max} = 2$ on the truncated CH spectral triple, the enriched substrate breaks the sector-locality structural condition that v3.61.0 Track A identified as forcing the abelian primitive coproduct: bit-exact in* sympy.Rational*, 12 of 12 single-step transitions carry non-vanishing η-class values in $\kappa^2 \cdot \mathbb{Z}$ (the M3 cocycle slot non-vanishes on inter-sector content), 12 of 12 chain compositions land on idempotents $T_2 \cdot T_1 = c \cdot e_s$ with non-zero $2\eta(T_1)\eta(T_2)$ cross-term cocycle image, and an additional 18 chain compositions land on two-step transitions outside the basic single-step generator basis (algebra-closure failure). The lowest-order non-trivial commutators $[T_a, T_b] \neq 0$ appear at 24 of 66 = 36% of transition pairs — explicit non-abelian Lie generators. The non-primitivity content has denominator family $\kappa^2 \cdot \mathbb{Z} = 2^{-8} \cdot \mathbb{Z}$ times inverse path counts, structurally aligned with the bit-exact $\pm 1/65536$ closure drift residual of Remark 25 modulo a 2^3 JLO simplex normalization factor; the exact bit-exact bridge identification between the two phenomena is a feasible follow-on. The enriched substrate's candidate motivic Galois group $U^{*(n_{\max})}_{\text{GeoVac,enriched}}$ has both an abelian (idempotent-generated) part and a unipotent (transition-generated) part, structurally targeting the pro-unipotent factor of the Connes–Marcolli cosmic-Galois U^* (Connes–Marcolli 2004, arXiv:math/0409306; built on the Connes–Kreimer Hopf algebra) at the Stage-2 level. See Paper 55 § subsec:open_m2_m3 for the cosmic-Galois narrative.*

Remark 30 (Q5' Stage 2 Cuntz-extension enrichment: structural incompatibility, Sprint Q5'-Cuntz, June 2026)**.** *The first scoping step of the long-range Cuntz-extension enrichment ingredient flagged in Remark 24 returns a structural negative. Replace the commutative algebra $\mathcal{A}^{(n_{\max})}_{\text{GV}} = \mathbb{C}^{N(n_{\max})}$ of sector idempotents by the Cuntz algebra $\mathcal{O}_N$ generated by $N = 3 \cdot N(n_{\max})$ isometries $S_{(n,l),k}$ (one per sector (n,l) and Mellin slot $k \in \{0,1,2\}$). The two natural candidate coproducts on $\mathcal{O}_N$ are bit-exactly incompatible with the Cuntz relations: (i) the primitive coproduct $\Delta(S_i) = S_i \otimes 1 + 1 \otimes S_i$ extended as a $*$-algebra homomorphism fails $S_i^* S_j = \delta_{ij}$ on 0/225 pairs at $n_{\max} = 2$ with $N = 15$; (ii) the diagonal coproduct $\Delta(S_i) = S_i \otimes S_i$ passes the first Cuntz relation on 225/225 pairs but fails $\sum_i S_i S_i^* = 1$ by exactly $N(N-1) = 210$ missing cross-terms. Net: $\mathcal{O}_N$ is structurally a Hopf-*comodule (over the gauge $U(1)$ scaling each S_i by a phase) rather than a Hopf-algebra. The M1/M2/M3 partition survives only at the trivial comodule level via fixed-k generator labels, structurally weaker than the clean Hopf tensor product $\mathcal{H}_{\text{GV}} = \otimes_k \mathcal{H}^{[k]}_{\text{GV}}$ established for the commutative case in Remark 24. The JLO degree-1 cochain ϕ_1 is structurally non-zero on $\mathcal{O}_N$ (the commutative-case vanishing theorem requires sector orthogonality $e_s e_t = \delta_{st} e_s$ which Cuntz isometries violate), but extraction at theorem grade requires the full CH-adjacency-extended Dirac representation through the Pimsner–Voiculescu sequence (Pimsner–Voiculescu 1980,* J. Operator Theory *4; Pimsner 1997, Fields Inst. Commun. 12), genuinely long-range. The Cuntz route is therefore the structurally most disruptive of the three enrichment ingredients flagged in Remark 24: it does not enrich the Hopf substrate but shifts the framework into the categorically different Hopf-comodule-over-gauge-group setting (Cuntz–Pimsner extension). The remaining two ingredients (nested-Hopf-tower $J^*(S^3)$ closed at finite cutoff in Remark 28; cross-shell off-diagonal Dirac closed at finite cutoff in Remark 29) preserve the Hopf-algebra category at the substrate level and are correspondingly more tractable. See Paper 55 § subsec:open_m2_m3 for the cosmic-Galois program narrative.*

Remark 31 (Q5' Stage 2 substrate HEADLINE: Levi-decomposition tensor synthesis, Sprint Q5'-Levi-Synthesis, June 2026)**.** *The HEADLINE Stage-2 substrate construction combines the v3.61.0 abelian primitive substrate of Remark 24 with the v3.62.0 Peter–Weyl substrate of Remark 28 into the tensor-product Hopf algebra*

$$\mathcal{H}_{\mathrm{GV}}^{\mathrm{Levi},(n_{\max},j_{\max})} := \mathcal{H}_{\mathrm{GV}}^{(\mathrm{v3.61}),n_{\max}} \otimes_{\mathbb{Q}} \mathcal{H}_{\mathrm{GV}}^{J^*,j_{\max}}.$$

At finite cutoff $(n_{\max}, j_{\max}) \in \{(2,1/2),(2,1),(3,1/2)\}$*, all five Hopf axioms (coassociativity, counit, antipode), bialgebra compatibility, the M1/M3/M2 Mellin-slot k-grading preservation on the first factor, and the non-cocommutativity on the second factor (encoding non-abelian* SL_2 *content) are verified bit-exactly (882 zero residuals across the joint axiom + truncation panel: 636 axiom residuals + 246 truncation residuals). The Mellin partition lifts to* $\mathcal{H}^{\mathrm{Levi}} = \bigotimes_{k=0}^{2} \mathcal{H}_{\mathrm{v3.61}}^{[k]} \otimes \mathcal{H}^{J^*}$*. The candidate motivic Galois group at the standard* $\mathcal{O}(SL_2)$ *quotient is*

$$\boxed{U_{\mathrm{GeoVac,Levi}}^{*(n_{\max},j_{\max})} = \mathbb{G}_a^{3N(n_{\max})} \times SL_2}$$

by the Tannakian theorem (tensor product of Hopf algebras = direct product of affine group schemes; Waterhouse 1979 §1.4; Sweedler 1969 Ch. IV; Deligne 1990 Catégories tannakiennes*). The semidirect product* $\mathbb{G}_a^{3N(n_{\max})} \rtimes SL_2$ *reduces to a direct product because the Levi action on the unipotent radical is trivial: the v3.61.0 generators* $x_{(n,l),k}$ *carry no* $SU(2)$ *representation content (they are abelian primitives indexed by sector and Mellin slot), so any* SL_2 *element acts on them as the identity (Hochschild 1981 Ch. VII Levi-decomposition direct-product reduction). The pro-unipotent* $\mathbb{G}_a^{3N(n_{\max})}$ *factor carries the master-Mellin engine M1/M3/M2 partition (Remark 24); the semisimple* SL_2 *factor carries the Peter–Weyl non-abelian content (Remark 28). The Levi-decomposition shape is modelled on (but differs from) the published structural target for the cosmic-Galois* U^* *in the Connes–Marcolli motivic Galois machinery (Connes–Marcolli 2004 arXiv:math/0409306; Connes–Marcolli 2008 book [2] Ch. 1 §7.3, "Equivariant principal bundles and the group* $G^* = G \rtimes \mathbb{G}_m$*"): pro-unipotent times the multiplicative group (reductive, not semisimple). The long-range Stage-2 Tannakian construction reduces to (a) full Tannakian closure of the pro-unipotent factor, and (b) verification that the cosmic-Galois* U^* *of Connes–Marcolli 2004 acts on* $\mathcal{H}_{\mathrm{GV}}^{\mathrm{Levi}}$ *in the expected way. See Paper 55 § subsec:open_m2_m3.*

Remark 32 (Q5' Stage 2 substrate enrichment via Mellin-slot-decorated Peter–Weyl, Sprint Q5'-Decorated-PW, June 2026)**.** *The alternative synthesis flagged in Remark 28 option (b) — the Mellin-slot-decorated Peter–Weyl substrate*

$$\mathcal{H}_{\mathrm{dec}}^{(j_{\max})} = \mathrm{span}_{\mathbb{Q}}\{\pi_{mn}^{j,k} : 0 \le j \le j_{\max},\ -j \le m,n \le j,\ k \in \{0,1,2\}\}$$

with k-preserving coproduct $\Delta\pi_{mn}^{j,k} = \sum_p \pi_{mp}^{j,k} \otimes \pi_{pn}^{j,k}$ *— is constructed and verified bit-exactly at* $j_{\max} \in \{1/2, 1\}$ *with* $n_k = 3$ *Mellin slots. All Hopf axioms hold bit-exactly with the standard counit and per-slot* $SU(2) \to SL_2$ *quotient (411 zero residuals: coassoc 57, counit L+R 114, antipode at quotient 57, k-grading preservation 57, pro-system Hopf-hom 126). The substrate factorises as* $\mathcal{H}_{\mathrm{dec}} = \mathcal{O}(SL_2)^{\otimes 3}$ *with the three Mellin mechanisms M1, M2, M3 carried by three independent non-abelian semisimple* SL_2 *factors. The candidate motivic Galois group at the* $SU(2)^{\otimes 3}$ *quotient is* $U_{\mathrm{dec},(a)}^* = SL_2^3$ *for every* $j_{\max} \ge 1/2$*. The alternative k-summing coproduct* $\Delta\pi_{mn}^{j,k} = \sum_{k_1+k_2=k \pmod 3} \sum_p \pi_{mp}^{j,k_1} \otimes \pi_{pn}^{j,k_2}$ *passes coassociativity and a* $\mathbb{Z}/3$*-augmented counit* $\varepsilon_{\mathrm{aug}}(\pi_{mn}^{j,k}) = \delta_{mn}\delta_{k,0}$ *but FAILS the antipode axiom at the per-slot* $SU(2)$ *quotient by a structural*

factor of $n_k = 3$ — fixing the normalisation requires $1/n_k \in \mathbb{Q}(1/3)$ breaking the discrete-for-skeleton discipline. The two synthesis routes (this remark's SL_2^3 vs Remark 31's tensor synthesis $\mathbb{G}_a^{3N} \times SL_2$) capture different aspects of the Stage-2 substrate: the tensor synthesis is the Levi-decomposition shape with explicit v3.61.0 abelian primitive content; this remark's decorated-PW is the Mellin-decorated symmetry shape with the M1/M2/M3 partition realised as independent SL_2 factors. Both substrates are valid candidates; the strategic choice between them is the next sprint-scale decision in the long-range Stage-2 program. See Paper 55 § subsec:open_m2_m3.

Remark 33 (Q5' Stage 2 bridge identity closing the T3b umbrella, Sprint Q5'-Bridge-Id, June 2026)**.** *The bit-exact bridge identification flagged as a feasible follow-on in Remark 29 closes (`debug/sprint_q5p_br` `debug/data/sprint_q5p_bridge_id.json`). On the load-bearing (e_2, e_3, e_3, e_2) palindrome at the strict-strong-form cochain-morphism interior cutoff $n_{\max} \geq 3$, the closure drift residual is bit-exactly $-\kappa^4 = -1/2^{16}$, equivalent in four closed forms:*

$$\mathrm{drift}_{n_{\max}\geq 3} = -\kappa^4 = -\frac{1}{4!}\kappa^4\, T^{\mathrm{int}}_{\mathrm{path}} = -\frac{2\eta(T_{(2,0)\to(2,1)})\eta(T_{(2,1)\to(2,0)})}{40} = -\frac{5/8192}{40},$$

with $T^{\mathrm{int}}_{\mathrm{path}} = 24$ the chirality-weighted two-step E1 path count at $n_{\max} \geq 3$ and the integer $40 = 2\cdot 10\cdot 2$ the palindromic two-step path-count product on the OffDiag substrate $T_{(2,0)\to(2,1)}, T_{(2,1)\to(2,0)}$ of Remark 29. The boundary cutoff $n_{\max} = 2$ value $+\kappa^4/3 = +1/196608$ uses $T^{\mathrm{bdy}}_{\mathrm{path}} = 8$, with the bit-exact ratio $T^{\mathrm{int}}_{\mathrm{path}}/T^{\mathrm{bdy}}_{\mathrm{path}} = 24/8 = 3$ identifying the -3 boundary-vs-interior factor of Remark 25 as an explicit integer path-count ratio. The pullback increment closes the integer arithmetic: $32 = 24 + 8$. The denominator-scale mismatch 2^{13} (T3b $\eta \otimes \eta$) vs 2^{16} (Track B drift) is bit-exactly the 2^3 piece of the JLO simplex factor $1/4! = 1/(3\cdot 2^3)$; the leftover numerator 5 in 5/8192 is half the path-count product $40 = 5\cdot 8$. The T3b umbrella conjecture ("the two findings are aspects of one Stage-2 substrate-enrichment story with denominator scales aligned modulo a 2^3 JLO simplex factor") is therefore bit-exactly correct, and the bridge identity is constructive at finite cutoff. Long-range follow-ons (closed-form $T_{\mathrm{path}}(n_{\max})$, Lie-algebra structure constants, continuum-limit Mellin lift) remain open.

Remark 34 (Q5' Stage 2 OffDiag algebra-closure dimension at $n_{\max} = 3$, Sprint Q5'-OffDiag-Closure-nmax3, June 2026)**.** *The first long-range scoping step of the algebra-closure dimension at general $n_{\max}$ (named follow-on (1) of Remark 29) identifies the closure of the OffDiag-enriched substrate $\mathcal{A}_{\mathrm{OD}}^{(n_{\max})} = \mathbb{Q}\langle e_s, A_{\mathrm{off}}\rangle$ under matrix multiplication as the path algebra of the sector quiver realised in $M_{\dim H}(\mathbb{Q})$ via the canonical adjacency representation $A_{\mathrm{off}} = D - \Lambda = \kappa A_{\mathrm{graph}}$. Bit-exact in* sympy.Rational *at $n_{\max} \in \{2, 3\}$: $\dim \mathcal{A}_{\mathrm{OD}}^{(2)} = 84$ and $\dim \mathcal{A}_{\mathrm{OD}}^{(3)} = 592$, with closure decomposing as a sum of per-block ranks over all $N(n_{\max})^2$ sector pairs (all 25 of 25 blocks at $n_{\max} = 2$, all 81 of 81 at $n_{\max} = 3$ realised non-trivially — the sector quiver is fully connected via multi-step E1 dipole paths). Fill-fractions $84/256 = 21/64$ and $592/1600 = 37/100$ slowly increasing; asymptotic limit requires $n_{\max} = 4$ for honest identification. Lie subalgebra of single-step transitions does NOT close: 82 of 378 commutator pairs lie outside the 28-dim transition span at $n_{\max} = 3$, generalising the 24 of 66 at $n_{\max} = 2$. Named sprint-scale follow-on: extend bit-exact closure to $n_{\max} = 4$ for third data point + asymptotic fill-fraction identification. Long-range: closed-form per-block rank formula from quiver combinatorics, and embedding of $\{\mathcal{A}_{\mathrm{OD}}^{(n_{\max})}\}_{n_{\max}}$ into the inductive system whose limit hosts the conjectural pro-unipotent factor of the cosmic-Galois U^* Hopf algebra (Connes–Marcolli 2004, arXiv:math/0409306). See Paper 55 § subsec:open_m2_m3.*

Remark 35 (Q5' Stage-2 combined substrate L5 = L1 categorically, Sprint Q5'-Combined-Substrate, June 2026)**.** *The first long-range scoping step of combining the T3a Peter–Weyl substrate (Remark 28) and T3b cross-shell off-diagonal Dirac substrate (Remark 29) on a SINGLE combined*

substrate closes via categorical reduction. The combined substrate $\mathcal{A}^{\text{combined}} = \mathcal{A}^{J^*}_{j_{\max}} \otimes \mathcal{A}^{\text{OD}}_{n_{\max}}$ *reduces to the categorical TENSOR PRODUCT of Remark 31 at the basic substrate level, not a smash product. Two structural obstructions block the smash-product upgrade: (i) Peter–Weyl matrix coefficients* π^j_{mn} *are polynomial* functions *on* SL_2*, while OffDiag transitions* $T_{s'\to s}$ *are operators on* $\mathcal{H}$ *with* scalar *cocycle data (*η*-class values in* $\kappa^2\cdot\mathbb{Z}$*) — categorically different objects; no* SL_2 *group element evaluates the 4 spin-*1/2 *matrix coefficients to the 12 OffDiag* η*-rationals simultaneously. (ii) The natural* $SU(2)$ *z-rotation* U_θ *does NOT preserve the basic 5-sector OffDiag basis: bit-exact symbolic computation (`debug/compute_q5p_combined_substrate_part2.py`) gives three distinct phase factors* $\{1, e^{i\theta}, e^{-i\theta}\}$ *on the conjugate* $U_\theta T U_\theta^{-1}$ *— the result lives in the span of* m_J*-resolved subtransitions, not in the span of the basic basis. The candidate motivic Galois group at the quotient is therefore the Levi-decomposition product*

$$U^{*(n_{\max},j_{\max})}_{\text{combined}} = SL_2 \times \mathbb{G}_a^{N_{\text{OD}}(n_{\max})},$$

exactly the long-range synthesis target predicted by Remark 31. ***L5 reduces to L1 at the substrate level****; the genuinely new content (a non-trivial smash product) lives at the* m_J*-resolved refinement of the OffDiag basis, which loses the v3.61.0 sector structure as load-bearing data and constitutes a long-range continuation. See Paper 55 § subsec:open_m2_m3.*

Remark 36 (Q5' Stage-2 substrate decision: L1 is forced via the primitive-space invariant, Sprint Q5'-L1-vs-L2-Diagnostic, June 2026)**.** *The strategic question flagged in Remark 35 and in Sprint Q5'-Levi-Arc § 5 – whether the v3.61.0 abelian primitives* $x_{(n,l),k}$ *in* $\mathcal{H}^{(n_{\max})}_{\text{GV}}$ *are reducible to Peter–Weyl matrix coefficients in disguise (so that the simpler* SL_2^3 *substrate of Remark 32 would suffice) or carry independent content (so that the Levi-decomposition substrate of Remark 31 is forced) – is resolved bit-exactly at finite cutoff by the primitive-space invariant* $\dim \text{Prim}(\mathcal{H}) = \dim \text{Lie}(\text{Spec}\,\mathcal{H})$*, a Hopf-isomorphism invariant. For* $\mathcal{H}^{(n_{\max})}_{\text{GV}}$ *the Milnor–Moore structure theorem for connected cocommutative Hopf algebras over* $\mathbb{Q}$ *identifies* Prim *with the generating vector space* $V_{n_{\max}}$*, giving* $\dim \text{Prim}(\mathcal{H}_{\text{GV}}) = 3N(n_{\max}) \in \{6, 15, 27, 42, 60, 81\}$ *at* $n_{\max} \in \{1, \ldots, 6\}$*. For* $\mathcal{H}^{(j_{\max})}_{\text{dec}}$ *the standard identification* $\text{Prim}\,\mathcal{O}(SL_2)^{\otimes 3} = \mathfrak{sl}_2 \oplus \mathfrak{sl}_2 \oplus \mathfrak{sl}_2$ *gives* $\dim \text{Prim}(\mathcal{H}_{\text{dec}}) = 9$ *independent of* $j_{\max}$ *(higher* $j_{\max}$ *adds symmetric-square Clebsch–Gordan polynomials of degree* ≥ 2 *in the* $j = 1/2$ *generators, contributing zero new dimensions to* $\mathfrak{m}/\mathfrak{m}^2$ *at the augmentation ideal). At every* $n_{\max} \geq 2$ *the dim defect is strictly positive* $(6, 18, 33, 51, 72)$*, so no Hopf isomorphism exists between the two substrates; L1 is forced. The substrate-dim coincidences flagged in Remark 32 (*$15 = 15$ *at* $(n_{\max}, j_{\max}) = (2, 1/2)$ *and* $42 = 42$ *at* $(4, 1)$*) are small-cutoff artifacts breaking at* $n_{\max} \in \{1, 3, 5, 6\}$*. An independent Hopf-isomorphism invariant (cocommutativity) separates the substrates at every* $n_{\max} \geq 1$ *–* $\mathcal{H}_{\text{GV}}$ *is cocommutative (every generator primitive) while* $\mathcal{H}_{\text{dec}}$ *is not on* $j > 0$ *matrix coefficients. Bit-exact panel: 258 zero residuals / consistent checks (driver `debug/compute_q5p_l1_vs_l2_diagnostic.py`; data `debug/data/sprint_q5p_l1_vs_l2_diagnostic.json`). The L2* SL_2^3 *substrate of Remark 32 remains valid as a complementary semisimple-only reading of the M1/M2/M3 partition; the Levi-decomposition* $U^*_{\text{GeoVac,Levi}} = \mathbb{G}_a^{3N(n_{\max})} \times SL_2$ *of Remark 31 is the structurally-correct Stage-2 target, an analogue of (not a match to) the published Connes–Marcolli motivic-Galois shape* $U \rtimes \mathbb{G}_m$*, whose reductive factor is* $\mathbb{G}_m$*, not* SL_2*. The canonical long-range next step of the Q5' program is the Tannakian closure of the pro-unipotent factor* $\mathbb{G}_a^{3N(n_{\max})}$ *in the Connes–Marcolli machinery. See Paper 55 § subsec:open_m2_m3.*

Remark 37 (Q5' Stage-2 closed-form generating function for $T^{\text{abs}}_{\text{path}}(n_{\max})$ on the OffDiag substrate, with cutoff-independence + E1 selection + McKean-Singer-on-A^2 closures, Sprint Q5'-T-Path-Generating

(T2), June 2026)**.** *The chirality-weighted two-step path count*

$$N^{(2)}_{s'\to s} \;:=\; \mathrm{Tr}(\gamma\, A\, e_s\, A\, e_{s'})/\kappa^2$$

on the OffDiag substrate (Remark 29) is bit-exactly cutoff-independent*: every realised sector pair (s', s) has the same $N^{(2)}$ value at every $n_{\max}$ where both sectors are present. Hypothesis verified bit-exactly across $n_{\max} \in \{1, 2, 3, 4, 5\}$, including boundary pairs. The E1 selection rule $|\Delta n|, |\Delta l| \le 1$ holds for all realised transitions. The total signed sum vanishes identically:*

$$T^{\mathrm{signed}}_{\mathrm{path}}(n_{\max}) \;=\; \sum_{(s',s)} N^{(2)}_{s'\to s} \;=\; \mathrm{Tr}(\gamma A^2)/\kappa^2 \;\equiv\; 0,$$

extending McKean-Singer ($\chi(S^3) = 0$, McKean-Singer on $A^0 = I$) to the chirality-weighted A^2-trace. The total absolute path count admits the bit-exact closed form

$$\boxed{T^{\mathrm{abs}}_{\mathrm{path}}(n_{\max}) \;=\; \frac{(n_{\max}^2 + 9n_{\max} - 4)(n_{\max}^2 + 13n_{\max} - 6)}{6}}$$

verified at panel $(8, 72, 224, 496, 924)$ for $n_{\max} \in \{1, 2, 3, 4, 5\}$, predicting $T^{\mathrm{abs}}_{\mathrm{path}}(6) = 1548$. Bit-exact verification (driver `debug/compute_q5p_t_path_generating.py`*; data* `debug/data/sprint_q5p_t_path_generati`
memo `debug/sprint_q5p_t_path_generating_memo.md`*). Unlocks the continuum Mellin lift of Remark 38.*

Remark 38 (Q5' Stage-2 continuum Mellin lift of the discrete bridge identity to the spectral zeta side with M2/M3 decomposition, Sprint Q5'-Mellin-Lift-Bridge (T1), June 2026)**.** *Using the quartic closed form of Remark 37, the continuum Mellin lift of the v3.63.0 L3 discrete bridge identity* $\mathrm{drift}_{n_{\max}\ge 3} = -\kappa^4$ *(Remark 33) is the generating function*

$$F(s) \;:=\; \sum_{n\ge 1} T^{\mathrm{abs}}_{\mathrm{path}}(n)\cdot n^{-s} \;=\; \tfrac{1}{6}\zeta(s-4) + \tfrac{11}{3}\zeta(s-3) + \tfrac{107}{6}\zeta(s-2) - \tfrac{53}{3}\zeta(s-1) + 4\,\zeta(s).$$

$F(s)$ admits the bit-exact decomposition by shift parity

$$F(s) \;=\; F_{\mathrm{M2}}(s) + F_{\mathrm{M3}}(s),$$

$$F_{\mathrm{M2}}(s) = \tfrac{1}{6}\zeta(s-4) + \tfrac{107}{6}\zeta(s-2) + 4\,\zeta(s) \quad (\textit{even shifts, M2 Seeley–DeWitt}),$$
$$F_{\mathrm{M3}}(s) = \tfrac{11}{3}\zeta(s-3) - \tfrac{53}{3}\zeta(s-1) \quad (\textit{odd shifts, M3 vertex-parity-Hurwitz}).$$

Bit-exact verification: $F - F_{M2} - F_{M3} = 0$ symbolically; 5 simple poles at $s \in \{1, 2, 3, 4, 5\}$ with residues $\{4, -53/3, 107/6, 11/3, 1/6\}$; integer-s panel exhibits mixed M2 (π^{2k}) + M3 (ζ(odd)) content *at every test point (e.g. $F(6) = -53\zeta(5)/3 + \pi^2/36 + 4\pi^6/945 + 11\zeta(3)/3 + 107\pi^4/540$). F is structurally distinct from the v3.62.0 M3 $\eta_D(s)$ Hurwitz Mellin (shares only the odd-shift structure with $\mathbb{Q}$ vs $2^s\,\mathbb{Q}$ coefficients). This is an* algebraic-side realisation of the M2/M3 part of the master Mellin engine partition on a single Hopf-algebra-level invariant *(F has no M1 part; its M3 content is the odd-ζ sub-sector, $\zeta(\mathrm{odd}) \in M_3 \cap \mathrm{MT}(\mathbb{Z})$, Paper 55) — the Mellin engine extends from the trace-evaluation level (case-exhaustion theorem 4) to the substrate-side Mellin generating function. Driver* `debug/compute_q5p_mellin_lift_bridge.py`*; data* `debug/data/sprint_q5p_mellin_lift_bridge.json`*; memo* `debug/sprint_q5p_mellin_lift_bridge_memo.md`*.*

Remark 39 (Q5' Stage-2 m_J-resolved smash-product foundation: bit-exact $\Delta m_J \in \{-1, 0, +1\}$ grading and $U(1)$ z-rotation diagonalisation at $n_{\max} = 2$, Sprint Q5'-mJ-Smash-Product (T4), June 2026)**.** *The m_J-resolved OffDiag substrate at $n_{\max} = 2$ admits a bit-exact $\mathbb{Z}_3$-grading by $\Delta m_J = m_{J,\text{to}} - m_{J,\text{from}} \in \{-1, 0, +1\}$ (Wigner-Eckart selection for the vector operator $A = \kappa A_{\text{graph}}$) with symmetric dimension count*

$$\dim \mathcal{A}_{-1} = 30, \quad \dim \mathcal{A}_0 = 40, \quad \dim \mathcal{A}_{+1} = 30, \quad \textit{total} = 100,$$

over all 100 non-zero off-diagonal Dirac entries. The $U(1)$ z-rotation $U_\theta|n, \kappa, m_J\rangle = e^{im_J\theta}|n, \kappa, m_J\rangle$ acts diagonally: $U_\theta T_{m'_J \to m_J} U_\theta^{-1} = e^{i\Delta m_J \theta} T_{m'_J \to m_J}$, exactly reproducing the three phase factors $\{1, e^{i\theta}, e^{-i\theta}\}$ that Remark 35 identified as broken on the basic substrate — the m_J-resolved refinement diagonalises *the $U(1)$ action. Critical scope finding: the j-content at $n_{\max} = 2$ includes $j = 3/2$ states (from $\kappa = \pm 2$ in sectors $(2, 1)$ and $(2, 2)$), so the decorated-PW substrate of Remark 32 at $j_{\max} = 1/2$ is* insufficient *for the full $SU(2)$ action on the m_J-resolved OffDiag;* L2 must be extended to $j_{\max} \geq 3/2$ *as the sprint-scale prerequisite to the long-range smash-product construction $\mathcal{A}^{m_J\text{-}OD} \rtimes \mathcal{O}(SL_2)$ — the only path to a non-trivial Hopf extension beyond Levi-decomposition. Driver* `debug/compute_q5p_mj_smash_product.py`*; data* `debug/data/sprint_q5p_mj_smash_product.json`*; memo* `debug/sprint_q5p_mj_smash_product_memo.md`*.*

Remark 40 (Q5' L2 substrate extended to $j_{\max} = 3/2$: sprint-scale prerequisite for the long-range smash-product construction closed, Sprint Q5'-FO1 (v3.66.0), June 2026)**.** *The sprint-scale prerequisite flagged by Remark 39 (T4 of v3.65.0) for the long-range smash-product construction $\mathcal{A}^{m_J\text{-}OD} \rtimes \mathcal{O}(SL_2)^{\otimes 3}$ at $n_{\max} = 2$ is closed bit-exactly. The L2 decorated-PW substrate of Remark 32 extends to $j_{\max} = 3/2$ with all 7 Hopf axioms passing: dim $\mathcal{H}_{\text{dec}}^{(3/2)} = 90$ (per slot: $\sum_{2j \leq 3}(2j+1)^2 = 1 + 4 + 9 + 16 = 30$), 621 bit-exact zero residuals across coassociativity (90/90), counit L+R (180/180), antipode at SU(2)$^{\otimes 3}$ quotient (90/90 structural per slot), k-grading preservation (90/90), and two pro-system Hopf-homomorphism truncations $P_{3/2\to 1}$ (126/126) and $P_{3/2\to 1/2}$ (45/45). The substrate factorisation $\mathcal{O}(SL_2)^{\otimes 3}$ and candidate motivic Galois group SL_2^3 are unchanged from v3.63.0 $j_{\max} \in \{1/2, 1\}$ panel — the $j_{\max}$ parameter controls only which irrep matrix coefficients are explicitly included; the underlying group structure is $j_{\max}$-invariant. This unblocks the long-range smash-product construction at the sprint-scale prerequisite level. Driver* `debug/compute_q5p_fo1_l2_extension_jmax32.py`*; data* `debug/data/sprint_q5p_fo1_l2_extension_jmax32.js` *memo* `debug/sprint_q5p_fo1_l2_extension_jmax32_memo.md`*.*

Remark 41 (Q5' Interpretation C closure: $F(s)$ integer-s panel sits in MT period ring at level ≤ 4 over $\mathbb{Z}[i, 1/2]$, AND U^* acts on $\chi, \eta, F(s)$ cocycle classes via period-pairing, Sprint Q5'-FO2+FO3 (v3.66.0), June 2026)**.** *T5 of v3.65.0 identified three interpretations of "U^* acts on $\mathcal{H}_{\text{Levi}}$": (A) Hopf-coaction, (B) Hopf-automorphism, (C) period-pairing on the image $\pi(\mathcal{H}_{\text{Levi}}) \subset \mathbb{C}$ via motivic Galois conjugation. Interpretation C is the cleanest sprint-scale closure. This remark records its bit-exact closure for all named cocycle-class families. T5 SQ1+SQ2+SQ5 combined: (i) every $\chi_{(n,l)}$ value and every $\eta_{(n,l)}$ value (60 + 60 entries up to $n_{\max} = 4$) is an integer in $\mathbb{Z}$ at depth 0 in the MT period ring; (ii) every term of $F(s)$ (Remark 38) at integer $s \in \{6, 7, 8, 9, 10\}$ (5 cells $\times$ 5 terms = 25 term classifications) is either pure-Tate $\pi^{2k}\,\mathbb{Q}$ (M2 component of v3.65.0 T1) or odd-Riemann $\zeta(\text{odd})\,\mathbb{Q}$ (M3 component); both sit in* $\text{MT}(\mathbb{Q}, 1)$*, which embeds in* $\text{MT}(\mathbb{Z}[i, 1/2], 4)$ *trivially. The Paper 18 §III.7 M1/M2/M3 master Mellin engine partition aligns bit-exactly with the standard MT depth/weight grading at integer-shifted ζ values: $k = 0$ M1 (volume slot, depth 0, weight 1 in π), $k = 2$ M2 (Seeley–DeWitt, depth 0, weight $2k'$ in $\pi^{2k'}$), $k = 1$ M3 (vertex-parity-Hurwitz, depth 1, weight $2k' + 1$ in $\zeta(2k' + 1)$, its odd-ζ sub-sector; Paper 55). The cosmic-Galois*

U^* acts on χ, η trivially (depth 0 fixed-points), on the M2 component of $F(s)$ as the Tate subgroup (invariant up to π^{2k} scaling), and on the M3 component as the standard motivic Galois action on $\mathrm{MT}(\mathbb{Q}, 1)$ *odd-zeta classes (Brown 2012, Glanois 2015). Total bit-exact zero residuals: 145 (25 + 60 + 60). Driver* `debug/compute_q5p_fo2_fo3_mt_period_containment.py`*; data* `debug/data/sprint_q5p_fo2_fo3_mt_period.json`*; memo* `debug/sprint_q5p_fo2_fo3_mt_period_memo.md`*. Honest scope: Interpretation C closes for the named cocycle classes ($\chi, \eta, F(s)$ panel); Interpretations A (Hopf-coaction) and B (Hopf-automorphism) remain open as named sprint-scale (B via SQ3) and long-range (A) follow-ons. The L1 follow-on (a) Tannakian closure of the pro-unipotent factor in the Connes–Marcolli machinery remains the canonical long-range next step.*

Remark 42 (Rational-residue extension of the case-exhaustion theorem: the Bernoulli ladder and the functional-equation reading of the two-layer split (Sprint GB, 2026-05-29))**.** *The case-exhaustion theorem 4 classifies (conditionally, for chains of the first fifteen Paper 34 projections) the π-content as $\mathcal{M}[\mathrm{Tr}(D^k\, e^{-tD^2})]$. The companion question is where the framework's* rational *coefficients come from. A diagnostic on the gravity sector's small rationals places them on a single Bernoulli ladder, with each rung B_{2n} glued across two windows by the Riemann functional equation:*

$$\zeta(2n) = (-1)^{n+1}\frac{B_{2n}(2\pi)^{2n}}{2(2n)!} \in \pi^{2n}\,\mathbb{Q} \qquad \Longleftrightarrow \qquad \zeta(1-2n) = -\frac{B_{2n}}{2n} \in \mathbb{Q}.$$

The left (positive-even) window is the M2 transcendental ring of Layer 2 observables; the right (negative-integer) window is the π-free rational residue carried by Layer 1 skeleton coefficients. Verified placements (all exact in `sympy`*):*

- *Conical-defect tip coefficient $\frac{1}{12} = -\zeta(-1) = B_2/2$ (§4.15); per-t UV target $\frac{1}{24} = -\zeta(-1)/2 = B_2/4$ (the string $c/24$); scalar S^3 Casimir $\frac{1}{240} = +\zeta(-3)/2$ (B_4 rung); the α-observation's $F = \pi^2/6 = \pi^2 B_2 = \zeta(2)$ (Paper 2; the M2 window of the same B_2 whose skeleton window is the conical tip).*
- Non-trivial internal check*: the replica-entropy derivative is forced, not free — $\frac{d}{d\alpha}\big[\frac{1}{12}(\frac{1}{\alpha} - \alpha)\big]\big|_{\alpha=1} = -\frac{1}{6} = -B_2 = 2\zeta(-1)$, the α-derivative of the tip's $B_2/2$.*
- Controls behave*: $\kappa = -1/16$ (numerically coincident with the Fock Jacobian factor $1/16$; an Observation, Paper 0 [28]) has no clean ζ form; the Dirac Casimir $17/480$ misses the scalar B_4 rung by exactly $1/32$ and requires the half-integer Hurwitz shift $\zeta(-3, \frac{1}{2})$, i.e. the scalar/spinor distinction is the integer/half-integer (M2/M3) split.*

The reading is the rational-residue extension of Theorem 4: the rational coefficients are the negative-integer analytic continuation of the same *Mellin object whose positive-even values are the Layer-2 transcendentals, and the empirical two-layer split (Paper 34/35) is the Riemann functional-equation reflection $\zeta(s) \leftrightarrow \zeta(1-s)$ instantiated per Bernoulli order.*

Honest scope. *This is a structural correspondence at the same epistemic level as Theorem 4, not a forcing proof; it concerns ζ at* integer *arguments (special values), not the non-trivial zeros, so it makes no contact with the critical-line structure (cf. WH6 and the three classical-RH walls RH-N/O/M, which remain in force). The B_2, B_4,* and *B_6 rungs are now populated (Sprint GB-3): the B_6 rung's falsifiable target was met by the conformal-scalar Casimir on S^5 (a GeoVac manifold via the Bargmann–Segal lattice, Paper 24), which evaluates to $-31/60480 = \frac{1}{24}(\zeta(-5) - \zeta(-3))$ — the B_6 rung $\zeta(-5) = -1/252$ enters with clean coefficient $+\frac{1}{24}$, alongside B_4 at $-\frac{1}{24}$, with B_2* absent*. The $\zeta(-5) - \zeta(-3)$ spreading (rather than a pure $\zeta(-5)$) is the* same *quadratic-multiplicity mechanism that obstructs the functional equation in Paper 28's functional-equation obstruction remark (the S^5 harmonic degeneracy is quartic), now seen on the Casimir side from an independent*

observable; the method reproduces the S^3 control $1/240 = \frac{1}{2}\zeta(-3)$ (Paper 35). The integer/half-integer (M2/M3) split recurs: the conformal scalar gives the clean ladder value, the half-integer HO/Bargmann S^5 Casimir gives $-17/3840$ (the "17" echoing the Dirac-S^3 17/480). Next rung: S^7 conformal Casimir is predicted to spread across $\zeta(-7), \zeta(-5), \zeta(-3)$. The ladder is thus supported across three rungs and two observable types, but remains a structural correspondence, not a forcing theorem. Source: `debug/sprint_gb_bernoulli_ladder_memo.md`, `debug/sprint_gb3_b6_rung_memo.md`.

Remark 43 (Bisognano–Wichmann reading of the M1 sub-mechanism on temporal compactifications (Track D, May 2026))**.** *The M1 sub-mechanism of the master Mellin engine, $\mathcal{M}[\mathrm{Tr}(D^0\, e^{-tD^2})]$, produces a* Vol/Mellin-measure *factor on the spatial S^3 ($4/\pi = 2\,\mathrm{Vol}(S^2)/\mathrm{Vol}(S^3)$ at unit radius, numerically $\mathrm{Vol}(S^2)/\pi^2$, the L2 state-space GH rate constant in the dual-Coxeter normalisation* — not *a quantity on the Hopf base, whose radius is half the total space's; label corrected 2026-09-03; Sprint MR-A/B in Remark 16) and on a temporal S^1_β factor (the bare circumference β itself; Sprint TD Track 1, $T_{S^3} \otimes T_{S^1_\beta}$). Sprint TD Track 4 (*`debug/sprint_td_track4_memo.md`*, May 2026) extends the temporal-side instantiation to the Euclidean Schwarzschild cigar, where smoothness at $r = 2M$ forces the τ-circle period $\beta_{\mathrm{cigar}} = 8\pi M$, consistent with the Hawking temperature $T_H = 1/(8\pi M)$ (sympy residual zero): the M1 volume slot re-expresses the continuum value, it does not derive it—M is external (it comes from the Einstein equations) and β is posited at finite cutoff. Track D (*`debug/bisognano_wichmann_track_d_memo.md`*, May 2026) makes explicit the dual physical reading of M1 on temporal compactifications: under the standard Wick-rotation chain (Hartle–Hawking [47]; Sewell [46]; Bisognano–Wichmann [44, 45]; reviewed in [49]), the framework's M1 2π on S^1_τ matches the period of the modular-flow / Lorentz-boost orbit of a stationary observer outside the Schwarzschild horizon. Equivalently, the M1 sub-mechanism on temporal compactifications corresponds, by analogy, to the Wick-rotated Bisognano–Wichmann modular-flow mechanism; identifying the framework's generator with a boost is open (Paper 42's O0). This is a* structural correspondence *(an analogy), not a literal identification: two mathematical objects (a volume factor on the Riemannian side; a modular-automorphism period on the Lorentzian side) are matched by the published Wick-rotation prescription (Bisognano–Wichmann 1976; Sewell 1982; Hartle–Hawking 1976), not by an operator-level proof inside the truncated spectral triple $\mathcal{T}_{n_{\max}}$. The natural follow-up that would turn the correspondence into an operator-level identification — compute the modular Hamiltonian K on $\mathcal{T}_{n_{\max}}$ for a wedge-restricted Camporesi–Higuchi state and verify $\sigma_{i\cdot 2\pi}$ = identity at finite $n_{\max}$, then take the GH limit using the lemmas of Theorem 5 — is named as the principal falsifier in Track D §3.1; estimated beyond sprint scale; priority MEDIUM-HIGH. The case-exhaustion theorem 4 itself is unchanged: M1's 2π carries* two readings of one number *(the $k = 0$ volume slot, the framework's own reading; and the boost-orbit period via Bisognano–Wichmann, an analogy under Wick rotation), not two independent constraints on the master Mellin engine partition. The corollary on Rindler wedges (the Unruh temperature $T_U = a/(2\pi)$, with $\beta_{\mathrm{Rindler}} = 2\pi/a$ posited and re-expressed through the same M1 volume slot — a consistency check, not a derivation) is carried out in the Unruh paragraph below.*

Unruh pendant (May 2026). *The Track D §5.4 follow-up was executed as a pendant sprint (*`debug/unruh_pe`*s May 2026). Wick-rotating Rindler coordinates (η, ρ) for an observer at proper acceleration a produces Euclidean polar (η_E, ρ) with conical regularity at $\rho = 0$ forcing η_E-period 2π, hence proper-time period $\beta_U = 2\pi/a$ and Unruh $T_U = a/(2\pi)$ (Unruh [48]). Sprint TD Track 1's* `matsubara_spectrum()` *applied verbatim with $\beta = 2\pi/a$ reproduces the Unruh-thermal Matsubara spectrum bit-identically (sympy residual zero on $T_U = 1/\beta_U$, lowest bosonic mode $= a$, lowest fermionic mode $= a/2 = \pi T_U$). The single 2π in β_U is the same M1 ($k = 0$ volume slot) / $\mathrm{Vol}(S^1)$ signature that drove the cigar correspondence, instantiated on the Rindler η_E-circle instead of the cigar τ-circle. The reading is identical: under the Wick-rotation chain, the framework's M1 2π on*

$S^1_{\eta_E}$ *matches the period of the modular-flow / boost-orbit of the accelerated observer (an analogy; identifying the framework's generator with a boost is open, Paper 42's O0). Hawking, Bisognano–Wichmann, and Unruh are three faces of one continuum Wick-rotation theorem, whose shared* 2π *the framework's M1 (*$k = 0$*) volume slot matches by analogy (it re-expresses the continuum value and does not derive it); the published Wick-rotation prescription supplies the bridge to the Lorentzian-side modular-flow vocabulary in each case. The* operator-level falsifier *(modular Hamiltonian on* $\mathcal{T}_{n_{\max}}$ *for a wedge-restricted state has analytic period* 2π*) named in Track D §3.1 covers all three faces simultaneously, since the bridge runs through the same KMS-Tomita-Takesaki algebra in every case; lifting the period-closure check on any one face lifts the structural correspondence on Hawking, Bisognano–Wichmann, and Unruh together. Same scope (structural correspondence, not an operator-level identification); same falsifier; same MEDIUM-HIGH priority and scale.*

Sprint L1 (2026-05-16): a finite-cutoff period closure, which does not close the falsifier. [SYMBOLIC (trivial) + MEASURED] *Sprint L1 (`debug/l1_modular_hamiltonian_results_memo.m`) takes as modular generator on* $\mathcal{T}_{n_{\max}}$ *the rotation generator preserving the hemispheric wedge,* $K = J_{\text{polar}} = 2J_z$ *(`geovac/modular_hamiltonian.py`), and verifies* $\sigma_{2\pi}(O) = O$ *for four witnesses labelled Bisognano–Wichmann, Hartle–Hawking, Sewell and Unruh at every tested* $n_{\max} \in \{2, 3, 4, 5\}$ *(maximum residual* 1.8×10^{-16}*,* 4.8×10^{-16}*,* 9.3×10^{-16}*,* 1.6×10^{-15} *at* $\dim_H = 16, 40, 80, 140$*; the value* 40 *at* $n_{\max} = 3$ *is the cumulative node count* $4 + 12 + 24$*, numerically equal to but distinct from* $g_3^{\text{Dirac}} = \Delta^{-1}$*). The closure is exact, but it is exact for a reason that carries no Bisognano–Wichmann content:* $K = 2J_z$ *has integer spectrum (the odd integers* $2m_j$*), so* $e^{i \cdot 2\pi \cdot n} = 1$ *gives* $\sigma_{2\pi} = \text{id}$ *for* any *operator* O *(the module's own documentation says so), and the four "witnesses" are the same* K *with the surface-gravity label* κ_g *relabelled, so their bit-exact agreement (max consistency residual* 0.0*) holds by construction. Any other integer-spectrum generator passes the same checks, including one with no rotation meaning (regression-pinned in `tests/test_modular_period_closure_automatic.py`, which also shows a non-integer-spectrum generator fails the same closure). The computation therefore identifies* no *Bisognano–Wichmann modular flow, and the four-witness statement is* not *made framework-internal at finite cutoff by it.The falsifier named above —* derive K *as the modular Hamiltonian of a wedge-restricted Camporesi–Higuchi state, rather than choose it, and check its period — remains* ***OPEN****. The continuum four-witness statement (*$\beta = 2\pi/\kappa_g$ *from Hartle–Hawking, Sewell, Bisognano–Wichmann and Unruh) is prior art and is unaffected. The signature-blind Krein-carrier sign audit at* $(3, 1)$ *is §8.10; the Lorentzian programme is descoped (Paper 45). The L1-A architecture obstructions §§10a–10e of `debug/l1_modular_hamiltonian_architecture_memo.md` were addressed at the level of the chosen* K*: §10a operator-system leakage is zero (*$\sigma_{2\pi}(O) = O$ *exactly, so* $\sigma_{2\pi}(O)$ *trivially stays in* $\mathcal{O}_{n_{\max}}$*); §10b/c chirality / M3 trivialization consistency: the* 2π *in the period is the period* chosen *with the integer-spectrum generator, posited rather than derived; its tag as* $\text{Vol}(S^1)$ *(M1) under Sprint TS-E1's master Mellin engine case-exhaustion theorem 4 is a classification of the posited value, not a derivation; §10d Gibbs phenomenon: the hemispheric projector* P_W *uses a discrete* $m_j \to -m_j$ *reflection (no smooth cutoff needed, exact involution); §10e* J_{TT}*-vs-*J_{GV} *compatibility: the Tomita conjugation is constructed as a fresh class `geovac.modular_hamiltonian.TomitaConjugation` with verified* $J_{\text{TT}}^2 = +I$*, categorically distinct from the real structure* J_{GV} *of `geovac.real_structure` (which has* $J_{GV}^2 = -I$*).*

Sprint L1-tighten (2026-05-16 evening): the Tomita–Takesaki cross-check is circular. *Sprint L1-tighten (`debug/l1_tighten_tomita_results_memo.md`) builds the Tomita–Takesaki data on the GNS Hilbert–Schmidt space* $H_{\text{GNS}} = M_{\dim_W}(\mathbb{C})$ *for the state* $\rho_W = e^{-K}/Z$ *with the* same *chosen* K*, extracts* $K_{\text{TT}} = -\log \Delta$ *from the polar decomposition* $S = J_{\text{TT}} \Delta^{1/2}$*, and verifies* $\sigma_{2\pi}^{\text{TT}}(O) = O$ *for the four labels at* $n_{\max} \in \{2, 3, 4, 5\}$ *(maximum residual* 1.09×10^{-16}*,* 1.07×10^{-15}*,* 3.03×10^{-15}*,* 4.79×10^{-15}*), with* $\sigma_t^{\text{TT}}(a) = \sigma_{-t}^{\alpha}(a)$ *at* $t = 1$ *(residual* $< 10^{-16}$*) and* $J_{\text{TT}}^2 = +I$

to machine precision. Because ρ_W is built as e^{-K}/Z, the modular generator of that state is K by construction, so this step re-derives the input rather than testing it: it is a correct consistency check of the Tomita–Takesaki machinery, not an independent realization of Bisognano–Wichmann. The result is a compact-period closure of an integer-spectrum flow — signature-blind, not a metric statement — and the falsifier status above is unchanged.

Theorem 5 (Gromov–Hausdorff convergence on S^3)**.** *Let $\mathcal{T}_{S^3} = (C^\infty(S^3), L^2(S^3, \Sigma), D_{\mathrm{CH}})$ be the round-S^3 metric spectral triple with the Camporesi–Higuchi Dirac, and let $\mathcal{T}_{n_{\max}} = (\mathcal{O}^{\mathrm{CH}}_{n_{\max}}, \mathcal{H}_{n_{\max}}, D_{n_{\max}})$ be the Connes–van Suijlekom truncation at cutoff $n_{\max}$ on the* true *Camporesi–Higuchi window: $\mathcal{H}_{n_{\max}}$ is the span of the complete Dirac eigenspaces with $|\lambda| \le n_{\max}+\frac{1}{2}$, $\mathcal{O}^{\mathrm{CH}}_{n_{\max}} = P_{n_{\max}} C^\infty(S^3) P_{n_{\max}}$ (*$\dim = 5, 30, 91$ *at $n_{\max} = 1, 2, 3$), and the state spaces are metrized by the left-translation Lipschitz seminorm (Paper 38; this is* not *the Fock-shell operator system of sprint R2.1, a model object, Remark 1). Then the truncated triples converge to $\mathcal{T}_{S^3}$ in van Suijlekom's state-space Gromov–Hausdorff distance (the strengthening to the Latrémolière quantum propinquity remains an open named gap; see Remark 44):*

$$d_{\mathrm{GH}}(\mathcal{T}_{n_{\max}}, \mathcal{T}_{S^3}) \;\le\; \gamma_{n_{\max}} \;\xrightarrow[n_{\max}\to\infty]{}\; 0,$$

where (matching Paper 38's corrected unconditional display, whose bound is $\le \gamma$ outright; the abandoned-route Lipschitz constant was $C_3 = 1$) $\gamma_{n_{\max}}$ is the L2 mass-concentration moment of the central spectral Fejér kernel on SU(2). The asymptotic rate $\gamma_{n_{\max}} = (4/\pi)\log n_{\max}/n_{\max} + O(1/n_{\max})$ (dual-Coxeter normalisation, $\mathrm{Cas}(\mathrm{ad}) = h^\vee$*; equivalently $2/\pi$ on the unit S^3 this paper declares, and $2\sqrt{2}/\pi$ under the field-standard Kac form — the constant scales with the metric) is* ***[INTERNAL THEOREM + MEASURED]*** *in Paper 38 [61] (main theorem, translation-seminorm metrization, 2026-06-10, stated there as $(4/\pi + o(1))\log n_{\max}/n_{\max}$; derived, with its two logarithmic sub-coefficients pinned by doubling estimators in its appendix): what is proved unconditionally is the bound $d_{\mathrm{GH}} \le \gamma_{n_{\max}}$ and $\gamma_{n_{\max}} \to 0$, while the uniform bound $\gamma_{n_{\max}} \le 6\log n_{\max}/n_{\max}$ ($n_{\max} \ge 2$) is* ***[PANEL-VERIFIED]****; the small-n closed forms are consistent with it. Convergence in the Dirac-commutator seminorm, and of the almost-commutative (gauge-network, $\mathcal{A}\otimes M_n$) structure, is* ***[OPEN]****.*

Proof sketch. The theorem is established by Paper 38's [61] lifted-state unconditional route in the van Suijlekom state-space Gromov–Hausdorff distance (*not* a Latrémolière propinquity — that strengthening is the open named gap of Remark 44). The historical five-lemma UCP-pair assembly of WH1 sprint R2.5 (April–May 2026) is recorded below for provenance; its lemma L5 is withdrawn (2026-09-03, Remark 44):

- **L1′** (Sprint R3.5, 2026-05-04; `debug/wh1_r35_full_dirac_memo.md`): the offdiag Camporesi–Higuchi operator system substrate has every non-forced cross-shell Connes-distance pair finite. This is a statement about the doubled Fock computational model, and the unconditional theorem does not use it (Paper 38, Lemma L1′); the state-space distance of the theorem is defined by the translation seminorm on the true window (the strengthening to the Latrémolière propinquity remains the open named gap of Remark 44).

- **L2** (Sprint R2.5/L2, 2026-05-04; `debug/r25_l2_proof_memo.md`): the positive central spectral Fejér kernel $K_{n_{\max}} = Z^{-1}_{n_{\max}} |\sum_{j\le j_{\max}} \sqrt{2j+1}\,\chi_j|^2$ on SU(2) is normalized, central, with mass-concentration moment $\gamma_{n_{\max}} \to 0$ (closed-form algebraic values $\gamma_2, \gamma_3, \gamma_4$, monotone decrease), mass distribution $N/Z_{n_{\max}}$ on the Fock-shell index (the kernel's mass distribution, not its multiplier symbol $\sigma_{n_{\max}}(N)$; Paper 38, corrected 2026-09-30), and central-multiplier cb-norm

$\|T_K\|_{\mathrm{cb}} = 1$ on the central subalgebra (the smoothing map is unital completely positive; the value $2/(n_{\max}+1)$ recorded by the original sprint is the mass-distribution maximum, not a cb-norm, Remark 44(i); corrected 2026-09-28) (the Bożejko–Fendler transcription of the original sprint is withdrawn: that result concerns discrete groups; the cb-norm 1 follows from complete positivity).

L3 (Paper 38, Lemma L3): **[INTERNAL THEOREM]** the Lipschitz comparison $\|[D_{\mathrm{CH}}, P_{n_{\max}} M_f P_{n_{\max}}]\|_{\mathrm{op}} \leq C_3 \|\nabla f\|_{L^\infty}$ holds with $C_3 = 1$ for *every* f and every $n_{\max}$, by compression from the continuum identity $\|[D_{\mathrm{CH}}, M_f]\| = \|\nabla f\|_{L^\infty}$ (the projection $P_{n_{\max}}$ commutes with D_{CH} and contracts operator norms); the constant is sharp in the limit. The earlier Sprint R2.5 per-harmonic ratio bound behind this lemma is false and withdrawn in Paper 38 (2026-09-28), together with the reading of the Track-A "torsion-and-curvature correction" as spectral-gap quantization that rested on it. The lemma is not used by the unconditional theorem.

L4 (Sprint R2.5/L4, 2026-05-06; `debug/r25_l4_proof_memo.md`): the Berezin-type reconstruction map $B_{n_{\max}} : C(S^3) \to \mathcal{O}_{n_{\max}}$ defined, in Paper 38's corrected form, by $B_{n_{\max}}(f) = \sum_{N \leq 2n_{\max}-1} \sum_{L,M} \sigma_{n_{\max}}(N)\, c_{NLM}\, M_{NLM}$ with $\sigma_{n_{\max}}$ the L2 multiplier symbol and Avery–Wen–Avery multipliers from sprint R3.1 (the sprint's original weighting by the mass distribution $N/Z_{n_{\max}}$ over $N \leq n_{\max}$ is not unital, $B(\mathbf{1}) = Z^{-1} I$, and is withdrawn; this historical lemma is not used by Paper 38's unconditional theorem) is positive (preserves $f \geq 0$), contractive ($\|B_{n_{\max}}(f)\|_{\mathrm{op}} \leq \|f\|_{C(S^3)}$), an approximate identity ($\|B_{n_{\max}}(f) - P_{n_{\max}} M_f P_{n_{\max}}\|_{\mathrm{op}} \leq \gamma_{n_{\max}} \|\nabla f\|_{L^\infty}$), and L3-compatible ($\|[D_{\mathrm{CH}}, B_{n_{\max}}(f)]\|_{\mathrm{op}} \leq \|\nabla f\|_{L^\infty}$). The map is the SU(2) analog of the Fejér-kernel smoothing map on S^1 of van Suijlekom [13] (Connes–van Suijlekom 2021 §4.2 define the Fejér–Riesz operator system itself, and their Lemma 4.4 is the Fejér–Riesz factorization, not a reconstruction map), equivalently the non-Kähler analog of the Hawkins 1999 Berezin map for Kähler coadjoint orbits (Comm. Math. Phys. 202).

L5 (Sprint R2.5/L5, 2026-05-06; `debug/r25_l5_proof_memo.md`): the tunneling pair $(B_{n_{\max}}, P_{n_{\max}})$ between $\mathcal{T}_{S^3}$ and $\mathcal{T}_{n_{\max}}$ has reach $\leq \gamma_{n_{\max}}$ in both directions (L4(c) approximate identity + the dual estimate via L2(g) Bożejko–Fendler; *the dual estimate is a withdrawn named gap*), height contributions bounded constants (L4(b) contractivity + $\mathrm{height}_P = 0$ — *both withdrawn 2026-09-03: L4(b) is not the relevant height, and* $\mathrm{height}_P = 0$ *is false by the same finite-band witness as* height_B); the withdrawn assembly then gave $d_{\mathrm{GH}}(\mathcal{T}_{n_{\max}}, \mathcal{T}_{S^3}) \leq C_3 \cdot \gamma_{n_{\max}} \to 0$ (the theorem now rests on Paper 38's lifted-state route instead) — the state-space distance of the theorem statement, not the Latrémolière propinquity (the symbol Λ denotes van Suijlekom's distance in Paper 38), whose label remains an open named gap.

The module's test panel does *not* verify Lemma L5 (measured 2026-09-03): the Lipschitz-normalised panel height rises with the cutoff (0.667, 0.808, 0.878, 0.913, 0.937, 0.952 at $n_{\max} = 2, \ldots, 7$) while $\gamma_{n_{\max}}$ falls, so the margin shrinks from $+1.41$ to $+0.05$ and turns negative at $n_{\max} = 7$ (-0.07; measured 2026-09-03). Normalised as printed (height over $\|f\|_{\mathrm{Lip}}$) the panel entries rescale into the unit-Lipschitz ball, so they are a *subset* of the set L5's supremum ranges over and therefore a lower bound on it: an entry exceeding $\gamma_{n_{\max}}$ *contradicts* the bound rather than leaving it open. **Lemma L5 is accordingly withdrawn as stated** (2026-09-03): Paper 38 gives a one-line witness — $B_{n_{\max}}$ is finite-band, so the unit-Lipschitz ball contains f with $B_{n_{\max}}(f) = 0$ and $\mathrm{height}_B \equiv 1$ at every cutoff — and the crossing is simply where $\gamma_{n_{\max}}$ falls below 1. This paragraph read "this neither confirms nor contradicts L5… an open check" until then. **Scope:** Paper 38's unconditional theorem uses no height constituent and is untouched, so the convergence statement and the WH1 keystone's commutative-factor leg stand. What the module reports as the bound

is $C_3\gamma_{n_{\max}} = 2.075, 1.610, 1.322$ (these are the L2 moments $\gamma_2, \gamma_3, \gamma_4$, not independent measurements; ratio 0.637). Module `geovac/gh_convergence.py` packages the assembly (48 unit tests, of which the panel-inequality checks added 2026-09-03 are the ones that can fail — the earlier version reported the bound as γ by construction), integrating with the L1′–L4 modules verbatim. □

Remark 44 (Status and named gaps (2026-06-09))**.** *Four gaps in the five-lemma proof chain behind Theorem 5 are named here. (i) The L2 cb-norm statement conflated the kernel's Plancherel* mass distribution *with its convolution-*multiplier symbol*: the smoothing map is unital completely positive with cb-norm exactly* 1*, and the quantity* $2/(n_{\max}+1)$ *is the mass-distribution maximum, not a cb-norm; the rate content (the mass-concentration moment* $\gamma_{n_{\max}}$*, its closed forms, and the* $4/\pi$ *asymptote) is unaffected. (ii) The* reach_P *dual estimate in L5 is a* named gap*: a forward cb-norm controls no partial inverse; the correct mechanism is an antiderivative-transference lemma in the style of Leimbach–van Suijlekom (Adv. Math.* ***439*** *(2024) 109496); for scalar sectors on compact metric groups the conclusion is covered by Gaudillot-Estrada–van Suijlekom [17]. (iii) The distance proved is van Suijlekom's state-space Gromov–Hausdorff distance (arXiv:2005.08544 lineage); the Latrémolière propinquity label is an open named gap — the cited Latrémolière framework builds tunnels from quantum isometries, not UCP pairs. (iv) The kernel condition* $L(a) = 0 \iff a \in \mathbb{C}\mathbf{1}$ *for the Dirac-commutator seminorm. The counts* $10/14$ *(*$n_{\max} = 2$*) and* $26/55$ *(*$n_{\max} = 3$*) for the chirality-diagonal CH Dirac, and* $1/14$*,* $1/55$ *for the engineered offdiag Dirac, are counts on the* doubled Fock model *of Remark 1, not on the truncated Camporesi–Higuchi triple: on the true window the Dirac-commutator kernel is exactly the scalars at* $n_{\max} \le 3$ *(****[MEASURED]****, Paper 38, remark on the Dirac-commutator seminorm;* `tests/test_p38_true_ch_window.py`*). An earlier version of this item read the model counts as a failure of the Dirac seminorm on the truncated triple (withdrawn 2026-09-28). Items (ii) and (iii) are resolved (2026-06-10) by the translation-seminorm reframing of the companion* S^3 *paper, adopted because its compression and lifted-state estimates close, not because the Dirac seminorm fails; whether the convergence can be proved in the Dirac seminorm is* ***[OPEN]****. The truncated state space is metrized by the left-translation Lipschitz seminorm (equal to the Lipschitz constant on the continuum), whose kernel is exactly the scalars on the true window at every cutoff (Schur multiplicity-one plus per-band injectivity of band-limited compression); the dual direction is supplied by an exact-fit spinor lifted state (the Fejér vector* $h \otimes \chi$*, lying in the window by the* $V_j \otimes V_{j\pm 1/2}$ *block decomposition of the Camporesi–Higuchi eigenspaces), and both almost-inverse defects are Fejér smoothings at the common moment* $\gamma_{n_{\max}}$ *— no partial inverse or transference estimate remains.* ***[INTERNAL THEOREM]*** *The convergence theorem is unconditional in van Suijlekom's state-space framework, bounded by* $\gamma_{n_{\max}} \to 0$*;* ***[INTERNAL THEOREM + MEASURED]*** *its asymptotic rate is* $(4/\pi + o(1)) \log n_{\max}/n_{\max}$ *in the dual-Coxeter normalisation of* $\gamma_{n_{\max}}$ *(*$2/\pi$ *on the unit* S^3*), the tier Paper 38 gives the constant.*

Remark 45 (Limit identification)**.** ***[INTERNAL THEOREM]*** *Paper 38 (its limit-identification proposition) identifies the state-space limit,* up to isometry*, as* $(P(S^3), W_d)$*: the unconditional theorem gives GH convergence of the true-window state spaces, metrized by the Monge–Kantorovich metric of the left-translation seminorm* L*, to* $(S(C(S^3)), \mathrm{MK}_L)$*; a GH limit is unique up to isometry, and since* $L = \mathrm{Lip}_d$*, Kantorovich–Rubinstein duality gives* $\mathrm{MK}_L = W_d$*, the Wasserstein distance for the (dual-Coxeter) geodesic distance* d*. The identification is for the translation-seminorm metric, not for the Dirac Connes distance* $d_{D_{n_{\max}}}$*: convergence in the Dirac-commutator seminorm is* ***[OPEN]****. (An earlier version of this remark identified the limit of* $(\mathcal{T}_{n_{\max}}, d_{D_{n_{\max}}})$ *via the withdrawn L4/L5 assembly; withdrawn 2026-09-28.)*

Remark 46 (WH1 keystone: the commutative factor)**.** *Theorem 5 closes the commutative-factor leg of the WH1 keystone (working-hypothesis register).* ***[INTERNAL THEOREM]*** *The Connes–*

*van Suijlekom truncations of the commutative factor — the round-S^3 Camporesi–Higuchi triple — are proved to converge in state-space Gromov–Hausdorff distance ($d_{\mathrm{GH}} \le \gamma_{n_{\max}} \to 0$). **[OPEN]** The structural claim of WH1 as a whole — that GeoVac is an almost-commutative spectral triple whose truncations, including the gauge-network $\mathcal{A} \otimes M_n(\mathbb{C})$ structure, converge — is* not *proved: convergence of the almost-commutative structure is open, as is convergence in the Dirac-commutator seminorm. **[INTERNAL THEOREM + MEASURED]** The displayed rate is $(4/\pi)\log n_{\max}/n_{\max} + O(1/n_{\max})$ (dual-Coxeter normalisation,* $\mathrm{Cas}(\mathrm{ad}) = h^\vee$*; equivalently $2/\pi$ on the unit S^3 this paper declares, and $2\sqrt{2}/\pi$ under the field-standard Kac form — the constant scales with the metric) (Paper 38), on the true Camporesi–Higuchi window under the translation-seminorm metrization. The L3 constant $C_3 = 1$ holds for every f and every $n_{\max}$ by compression (Paper 38, Lemma L3, **[INTERNAL THEOREM]**); the earlier per-harmonic route to it, and the reading of the Track-A "torsion-and-curvature correction" as spectral-gap quantization, are withdrawn (2026-09-28). The remaining open questions (Higgs-gap analysis on the four-way S^3 coincidence; the order-zero and order-one conditions of the real structure at finite $n_{\max}$ — the KO sign pair itself is settled, Sec. 4) are downstream of this theorem. The proof chain was corrected in June 2026 (Remark 44): the theorem stands unconditionally in van Suijlekom's state-space framework under the translation-seminorm metrization.*

Remark 47 (Pythagorean orthogonality of the L2-E residual; a normalisation-dependent $1/\pi^2$)**.** *The case-exhaustion theorem 4 classifies (conditionally, for chains of the first fifteen Paper 34 projections) the transcendental ring in which the π-content must sit. A post-Sprint-L2-E PSLQ probe (`debug/h_local_residual_pslq_memo.md`, 2026-05-17) records the following operator-level identity; its π-content is fixed by a posited normalisation (below), so it is* not *an instantiation of that theorem. On the truncated Lorentzian Krein spectral triple (Paper 42 [53] §11.7), the local Hamiltonian $H_{\mathrm{local}} = K_L^{\alpha,W}/\beta$ (posited generator $2J_z$; BW identification open, Paper 42's O0) and the wedge-restricted Lorentzian Dirac D_W^L are Hilbert–Schmidt orthogonal,*

$$\langle H_{\mathrm{local}}, D_W^L \rangle_{\mathrm{HS}} \;=\; 0 \qquad \textit{(bit-exact at every tested } (n_{\max}, N_t)\textit{)},$$

and the squared residual decomposes as a Pythagorean sum:

$$\left\| H_{\mathrm{local}} - D_W^L \right\|_F^2 \;=\; \frac{\kappa_g^2 \cdot S(n_{\max})}{4\pi^2} \;+\; D(n_{\max}),$$

with $S, D \in \mathbb{Q}[n_{\max}]$ given by cumulative wedge Casimir traces (Paper 42 §11.7, Cor. 11.15 therein). The sole transcendental in the residual is $1/\pi^2$ (corrected 2026-09-03 from "$= 1/\mathrm{Vol}(S^1)^2$", which is false arithmetic: $\mathrm{Vol}(S^1) = 2\pi$, so that quantity is $1/(4\pi^2)$ — the displayed equation above carries the correct form), and it enters only *through the posited normalisation $H_{\mathrm{local}} = K/\beta$ with $\beta = 2\pi/\kappa_g$: it is the continuum KMS period imported as a normalisation and squared (a temporal-compactification π in the sense of Paper 35),* not *an M1 Mellin output. The orthogonality itself is a chirality-parity identity that holds for any rescaling of K and is independent of β (Paper 42, Thm. 11.14). An earlier version of this remark read the $1/\pi^2$ as "the squared Hopf-base-measure signature, exactly the M1 output the case-exhaustion theorem predicts"; thatreading is withdrawn (2026-09-28), since K is posited rather than derived (Paper 42, O0).*

The residual's ring $\mathbb{Q}[n_{\max}] \oplus \pi^{-2}\,\mathbb{Q}[n_{\max}]$ is therefore a statement about the chosen normalisation, and it does not add an instantiation of the mechanism-as-domain partition of Remark 16 (M1 owns state-space GH-rate observables, M2 heat-kernel residuals, M3 vertex-restricted parity observables); an earlier version counted it as a fourth M1 instantiation and as a prediction of the engine (withdrawn 2026-09-28).

The orthogonality holds for all six modular witnesses (Paper 42 [53], the corollary on the single operator-system construction; Paper 42 §11.7), via κ_g-linearity of H_{local} and κ_g-independence of D_W^L — by construction, since the witnesses share one posited K. The mechanism is a chirality-parity identity, proved in general (Paper 42, Thm. 11.14 and Cor. 11.15): in the unfolded wedge spinor basis H_{local} is real, diagonal and independent of the chirality χ (eigenvalues two_m_j*), while the wedge Dirac is chirality-diagonal with χ-paired eigenvalues of opposite sign, so the pairs cancel in the trace. (An earlier version described the wedge Dirac as off-diagonal in this basis and left the general proof as a named follow-on; both corrected 2026-09-28.) The regression test (`tests/test_paper43_pythagorean_hs_orth` checks 7 cells — Riemannian $n_{\max} = 2, 3, 4$ and Lorentzian $(n_{\max}, N_t) = (2,1), (3,1), (2,3), (3,3)$ — not the* 18 *an earlier version cited.*

Cross-references. *Paper 42 §11.7 Cor. 11.15 (closed form and witness-universality); Paper 42 §8, the Hilbert–Schmidt orthogonality remark (which states that the orthogonality does* not *underlie the six-witness agreement, which holds by construction); Paper 24 §5.3 Coulomb/HO asymmetry (seven catalogued, six standing; the structural finding that the Pythagorean orthogonality is S^3-specific to the construction — the S^5 Hardy sector has no clean analog, adding the Pythagorean orthogonality for the posited wedge generator as a further Coulomb/HO asymmetry layer alongside calibration π and Wilson matter coupling; the spectrum-computing-L_0 layer once listed here is withdrawn, 2026-09-25).*

Remark 48 (BH Phase 0 — clean negative on wedge KMS area-law)**.** *The modular-Hamiltonian construction of Sprint L1 (a finite-cutoff period closure for a chosen integer-spectrum generator, this section; it identifies no Bisognano–Wichmann flow) and Sprint L2-E (its signature-blind Krein-carrier counterpart) is compatible with the Hartle–Hawking and Unruh* temperatures *via the M1 mechanism (§ 43). A natural follow-up question is whether the same construction also reproduces the Bekenstein–Hawking* entropy $S_{BH} = A/(4G)$ *at finite cutoff via the von Neumann entropy of the wedge KMS state $\rho_W = e^{-K_\alpha^W}/Z$ (the unit-β Gibbs state of the posited integer-spectrum generator $K_\alpha^W = 2J_z$, not BW-normalised: that would be $e^{-\pi K_\alpha^W}/Z$, Paper 50 [63]). Sprint BH-Phase0 (`debug/bh_phase0_diagnostic_memo.md`, 2026-05-22) tests this directly at $n_{\max} \in \{2, 3, 4, 5, 6, 7\}$.*

Result. *$S(\rho_W) \approx 1.944 \log(n_{\max}) + 0.565$ at $R^2 = 0.99991$ across the panel. No polynomial scaling appears in any of the four candidate fits (log-linear, power-law, poly-2, poly-3). An s-deformation $\rho(s) = e^{-sK_\alpha^W}/Z$ across $s \in \{0.01, 0.05, 0.1, 0.5, 1, 2, 2\pi\}$ shows power-law exponents $\alpha \in [0.64, 0.72]$, never close to the $\alpha = 2$ that a naive Bekenstein–Hawking area-law $S \sim n_{\max}^2$ would predict.*

Mechanism. *The structural reading is*

$$S(\rho_W) \;\approx\; \log\bigl(n_{\text{equator}}\bigr), \qquad n_{\text{equator}} = n_{\max}\,(n_{\max}+1),$$

where n_{equator} is the count of wedge states with smallest $|m_j| = 1/2$ (the equator analog). The unit-β state ($s = 1$) is effectively the maximally mixed state on the lowest-K_α shell, with exponentially suppressed contributions $\sim e^{-(2k+1)}$ from shells $k = 1, 2, \ldots$. The two asymptotic limits saturate bit-exactly: $S(s \to 0) = \log(\dim W)$ (hot, maximally mixed) and $S(s \to \infty) = \log(n_{\text{equator}})$ (cold, lowest shell only). The unit-β state ($s = 1$) sits between, closer to the cold limit.

*The slope ≈ 2 is the exponent of the $n_{\max}(n_{\max}+1)$ degeneracy of the lowest shell of the posited generator **[MEASURED]**; an earlier reading of it as $\dim(\partial W) = 2$ (a two-dimensional entangling surface) is withdrawn, since for a CFT_3 the entangling surface of a hemispherical cap is a circle (Paper 50, 2026-09-28). This is structurally weaker than the QFT Bekenstein–Hawking area-law $S \sim A \cdot \Lambda^2/4$, which would scale as a power of the UV cutoff times the boundary dimension.*

Structural reading. *The operator-system truncation on which the finite-cutoff period closure holds (§ VIII.F, Sprint L1) does* not *preserve the continuum QFT vacuum entanglement that would*

give the Bekenstein–Hawking area-law. The modular-flow period closure (a posited 2π period, tagged M1) and the Bekenstein–Hawking area-law are structurally distinct features of the gravitational thermodynamics chain; at finite cutoff the GeoVac framework exhibits the former only in the automatic integer-spectrum sense, and not the latter. Consistent with the structural-skeleton-scope pattern (project development-frontier log, sprints H1, LS-8a, HF-3/4/5, multi-focal-composition wall, W3): the framework supplies an integer-spectrum candidate for the algebraic skeleton of modular flow ($K = 2J_z$ posited; its Bisognano–Wichmann identification is open, Paper 42's O0) but does not autonomously generate the calibration data (here: Newton's constant) that links wedge-entropy to physical area.

Cross-references: Paper 34 §V.B Bekenstein–Hawking probe row; Sprint TD Track 4 (Hawking temperature, machinery-witness, M1 closure); the May-16 BH scoping memo (`debug/bekenstein_hawking_sprint`; full Connes–Chamseddine cigar route) remains the longer-route alternative if a literal $S_{BH} = A/(4G)$ closure is pursued, but with the honest scope statement that Newton's constant is calibration input not derivation.

8.1 The unified $U(1) \times SU(2) \times SU(3)$ gauge content and its limits

The case-exhaustion theorem of Sec. 8 identifies M1 (the $k = 0$ volume slot) as the mechanism by which the Wilson plaquette-Haar projection injects π into spectral-triple observables. This subsection records a complementary structural fact about the family of GeoVac sub-triples that engage M1: they realise all three Standard-Model gauge groups as Wilson lattice constructions, on three structurally distinct sub-manifolds, with a single unified weak-coupling coefficient $1/(4\,N_c)$. The construction is honest about what it is and is not: a unified *catalogue* of three Wilson-Yang–Mills instances of the Marcolli–van Suijlekom gauge-network framework, not a unified spectral triple in the Connes–Chamseddine sense [5].

Three Wilson constructions, one kinetic coefficient

[PANEL-VERIFIED] Three independent sprints have produced gauge-invariant, finite, Haar-normalisable non-abelian Wilson lattice gauge theories on GeoVac sub-graphs:

- *Paper 25* [36] constructs the abelian $U(1)$ Wilson–Hodge structure on the Fock-projected S^3 Coulomb graph (Paper 7), with phases χ_v on nodes and A_e on edges covariant under the standard incidence operator.
- *Paper 30* [38] constructs the non-abelian SU(2) Wilson lattice gauge theory on the same $S^3 = \mathrm{SU}(2)$ graph, with link variables $U_e \in \mathrm{SU}(2)$, plaquettes from primitive non-backtracking walks, and Haar measure dU. Paper 30 §5.2 verifies the weak-coupling kinetic coefficient symbolically and numerically.
- *Sprint ST-SU3* (May 2026, modules `geovac/su3_wilson_s5.py`, `tests/test_su3_wilson_s5.py`, 47 pass and 2 skipped of 49 as of 2026-09-02) constructs the non-abelian SU(3) Wilson lattice gauge theory on the Bargmann–Segal S^5 graph of Paper 24 [35], with link variables $U_e \in \mathrm{SU}(3)$ in the fundamental representation (independent of the shell index N), plaquettes from primitive non-backtracking walks, and SU(3) Haar measure. Sprint ST-SU3 §4 verifies Theorem 2 (kinetic coefficient 1/12) symbolically and numerically at $N_{\max} = 2, 3$ (7 765 plaquettes), with maximal Cartan reduction to a coupled $U(1) \times U(1)$ theory verified at machine precision.

The three constructions are summarised in Table 5. The columns are not independent discoveries: each is a specialisation of the gauge-networks framework of Marcolli–van Suijlekom [6], with the Perez-Sanchez correction [11] that the continuum limit of such gauge networks (whose lattice action is of Wilson type) is the Yang–Mills action *without* a Higgs sector.

In each construction the edge Laplacian $L_1 = B^\top B$, where B is the signed node–edge incidence matrix, plays the role of weak-coupling kinetic operator on $\mathfrak{g}$-valued 1-cochains. Paper 25's "discrete Hodge–1 Laplacian" is therefore a single combinatorial object across all three constructions; only its target Lie algebra changes ($\mathfrak{u}(1)$, $\mathfrak{su}(2)$, $\mathfrak{su}(3)$). **[OBSERVATION]** This is the single mathematical identity unifying the three sectors at the kinetic level.

The two-step SU(3) story

The SU(3) row of Table 5 requires a specific clarification, because an earlier Sprint 5 Track S5 (April 2026, memo `debug/s5_gauge_structure_memo.md`) returned a clean negative when asking whether the S^5 Bargmann–Segal lattice carries a non-abelian gauge structure analogous to Paper 25's $U(1)$. The two findings are not in tension: they answer two different questions, and read together they give a two-step honest picture.

- *Step 1 (Sprint 5 Track S5, negative).* An adjacency-preserving SU(3) action on the $(N, 0)$ tower of the Bargmann graph fails Wilson's fixed-group-on-every-link requirement, because the transitions $(N, 0) \to (N + 1, 0)$ are Clebsch–Gordan intertwiners between distinct SU(3) irreducible representations of growing dimension $\dim(N, 0) = (N + 1)(N + 2)/2$, not group elements of a single SU(3). The natural shell-respecting gauge group of the Bargmann graph is $U(1)$, and this is recorded in Paper 25 §VII.A as a structural asymmetry between the S^3 and S^5 sectors.
- *Step 2 (Sprint ST-SU3, positive at the gauge level).* A *shell-independent* Wilson lattice gauge theory with link variables $U_e \in \mathrm{SU}(3)$ in the fundamental representation (not in any of the $(N, 0)$ irreps) is a structurally different object that bypasses the Step 1 obstruction. The links live as external gauge degrees of freedom on edges; they do not act on shell labels. Sprint ST-SU3 verified gauge-invariance of the Wilson action under node-local $U_e \mapsto g_{\mathrm{src}(e)} U_e g^\dagger_{\mathrm{tgt}(e)}$ to machine precision and computed the kinetic coefficient $1/12$ both symbolically and numerically.
- *Matter coupling (Sprint ST-SU3 Theorem 3, mixed).* When one tries to couple this Wilson SU(3) to the natural shell matter on the $(N, 0)$ tower, the original Step 1 obstruction is rediscovered: a fundamental-representation link cannot uniformly couple to representations of growing dimension, and the matter coupling reverts to the Clebsch–Gordan intertwiner structure. Wilson SU(3) on the Bargmann graph is therefore self-consistent as a *pure-gauge* theory but does not admit a natural matter coupling in the sense of Paper 30's spin-1/2 Dirac fermions on the Coulomb graph.

The two-step picture is consistent with Paper 24's Coulomb/HO asymmetry and adds to it (seven layers catalogued, six standing, Papers 24 and 31): to the prior standing asymmetries (calibration π; Wilson gauge-with-natural-matter; the spectrum-computing role of L_0 once listed here is withdrawn, 2026-09-25, since L_0 does not converge to Δ_{S^3}) this sharpens, rather than adds to, the Wilson gauge-with-natural-matter layer: on the S^3 Coulomb side the gauge group acts in the same representation as the matter, so matter–gauge coupling is structurally available (argued, not constructed). On the S^5 Bargmann side the combinatorial Wilson–Hodge vocabulary is universal but the gauge–matter coupling is not.

Marcolli–vS lineage of the unified content

Each construction in Table 5 is a specific gauge-networks instance in the sense of Marcolli and van Suijlekom [6], with the Perez-Sanchez correction [11] that the continuum limit of such a gauge network is the continuum Yang–Mills action (the lattice action being of Wilson type) *without* a Higgs. The vertex algebra is $\mathcal{A}_v = \mathbb{C}$ (or $C(V)$ globally) in all three rows; the vertex Hilbert space and the gauge group differ row-by-row, as listed in the table. In each row the gauge-network spectral action reduces to the standard Wilson action plus a quadratic kinetic term in the weak-coupling expansion, with coefficient $1/(4N_c)$.

The lineage placement is identical to that of Sec. 9, and each Wilson construction inherits the same status: each is a published-machinery instance with a specific manifold and gauge group; what is not duplicated in the literature is the *simultaneous* existence of all three SM gauge groups as siblings across GeoVac sub-manifolds.

What this is not: structural distance to a Standard Model

Table 5 is a catalogue, not a unified theory, and the project's rhetoric rule forbids treating it as the latter. The structural distance from this catalogue to Connes' SM construction lives in four named gaps.

G1 (Three generations of matter.) Each construction in Table 5 carries a single matter species: Paper 30 has one generation of Dirac fermions (Paper 14 Tier 2 [31]); Sprint ST-SU3 has one tower of $(N, 0)$ irreps. Sprint 4H Track SM-B (April 2026) tested the naive shell-to-generation map $\sum_f N_c Q_f^2 = 8 = |\lambda_3|$ and returned a clean negative: the per-generation contribution is charge-universal across the three SM generations whereas every per-shell S^3 invariant varies with n, so no shell→generation map is consistent with charge-universal per-generation structure. The $8 = 8$ equality was a numerical coincidence with no representation-theoretic content. There is no GeoVac mechanism currently selecting a generation tripling of the Hilbert space.

G2 (Higgs as inner fluctuation of D.) In Connes' SM, the Higgs field arises as the off-diagonal block of an inner fluctuation $D \mapsto D + \omega + \epsilon' J\omega J^{-1}$ ($\omega = \sum_i a_i[D, b_i]$) for $a_i, b_i \in \mathcal{A}_F = \mathbb{C} \oplus \mathbb{H} \oplus M_3(\mathbb{C})$ in an almost-commutative algebra. The Marcolli–vS gauge-network lattice action [6] does carry a finite vertex Dirac and hence vertex-Higgs terms; Perez-Sanchez 2025 [11] shows that the Higgs does not survive the *continuum limit*, which is Yang–Mills without a Higgs. To keep a Higgs in the limit, an almost-commutative finite factor with an off-diagonal $\mathbb{C} \leftrightarrow \mathbb{H}$ Yukawa block is needed. The natural extension for GeoVac is $\mathcal{A}_{\mathrm{GV}} \otimes \mathcal{A}_F$ with the electroweak slice $\mathcal{A}_F = \mathbb{C} \oplus \mathbb{H}$ (deferring color), and with a finite Dirac D_F on the four-dimensional electroweak fiber whose off-diagonal block encodes the Yukawa coupling matrix. Inner fluctuations on this extension formally split into a gauge-1-form sector (which must reproduce the unified $U(1) \times \mathrm{SU}(2)$ content of Papers 25–30) and a candidate Higgs scalar sector living in the off-diagonal $(\mathbb{C}, \mathbb{H})$ component. Whether the required D_F can be selected from GeoVac structure rather than imposed is the load-bearing open question; we report the result of this test (Sprint H1) below in Sec. 8.2. Note that the operator-system non-multiplicative-closure of Remark 1 (witness pair $M^2_{2,1,0}$, 14.9% residual against $\mathcal{O}_{n_{\max}=2}$) is a separate, finite-resolution feature of the truncated operator system itself; inner fluctuations are defined operationally as $\omega + J\omega J^{-1}$ with $\omega = \sum_i a_i[D, b_i]$, which uses ALGEBRA elements (the multiplier matrices, which are an algebra over $\mathbb{C}$ at the level of $M_{\dim H}(\mathbb{C})$ even though their span $\mathcal{O}_{n_{\max}}$ is not closed under multiplication), so the $a \cdot b \notin \mathcal{O}$ feature does not block

the fluctuation calculus. Sprint H1 (Sec. 8.2) closes this gap up to the Yukawa-undetermined verdict; Sprint G3 (Sec. 8.3) supplies the structural reason that Y is undetermined — γ_{GV} and γ_F are independent commuting $\mathbb{Z}_2$ gradings, the Yukawa lives on the γ_F flip, and no GeoVac-side data couples to that flip. G2 and G3 are therefore one structural fact, not two.

G3 (Hypercharge / electroweak unification.) The SM relation $Q = T_3 + Y_W/2$ requires co-located $U(1)_Y$ and $\mathrm{SU}(2)_L$ on a common Hilbert space with a left–right chiral asymmetry. Paper 25's $U(1)$ and Paper 30's SU(2) are co-located on the same S^3 graph (the co-location Sprint G3 tested and closed in the negative, §8.3; until 2026-09-04 this parenthesis still called it "the most concrete near-term unification target"), but the Camporesi–Higuchi Dirac (Sec. 3 Definition 3) is non-chiral by default. The chirality grading introduced in Sprint TS Track E2 (R3.5, full Dirac with both chiralities) is a $\mathbb{Z}_2$-grading of the spinor bundle, not the SM weak-isospin chirality, and it does not distinguish left- and right-handed sectors of a charged matter field. Sprint G3 (Sec. 8.3) closed this in the negative on S^3 alone: γ_{GV} and γ_F are independent commuting $\mathbb{Z}_2$ gradings on $\mathcal{H}_{\mathrm{GV}} \otimes \mathcal{H}_F$, with operator-norm residual exactly 2 at every $n_{\max} \in \{1, 2, 3\}$. Identification is structurally impossible by tensor-product factorisation, and the electroweak co-location route is therefore pushed to G4.

G4 (Cross-manifold obstruction, the dominant gap.) Connes' SM uses a *single* spectral triple $(\mathcal{A}_{\mathrm{SM}}, \mathcal{H}_{\mathrm{SM}}, D_{\mathrm{SM}})$ on a single base manifold, with all three SM gauge groups emerging from inner automorphisms of one almost-commutative algebra. The GeoVac construction has, at present, three separate spectral triples on two different graphs (the S^3 Coulomb graph for $U(1)$ and SU(2); the S^5 Bargmann graph for SU(3)). There is no GeoVac operator that geometrically unifies the S^3 and S^5 graphs into a single spectral triple containing both as commuting factors: at best the Bargmann and Coulomb graphs can be encoded as orthogonal factors of a tensor-product Hilbert space, but the resulting object is a direct sum of two Wilson-YM instances rather than a unification. This is the most consequential gap. It is also the most directly tied to the Coulomb/HO asymmetry of Sec. 7 (seven layers catalogued, six standing): the absence of cross-manifold unification is a structural symptom of the same asymmetry that forbids a calibration-π analog and a natural Wilson matter coupling on S^5. (An earlier version also named a spectrum-computing L_0; that layer is withdrawn, 2026-09-25, since L_0 is spectrally inert on S^3 too.) Sprint G4 scoping (Sec. 8.4) sharpens this gap into G4a (Connes SM on $\mathcal{T}_{S^3}$ alone with $\mathcal{A}_F = \mathbb{C} \oplus \mathbb{H} \oplus M_3(\mathbb{C})$, bounded sprint sequence, predicted positive-thin) and G4b (genuine cross-manifold $\mathcal{T}_{S^3} \otimes \mathcal{T}_{S^5}$, blocked at the NCG-framework level by the Riemannian-vs-Hardy-sector asymmetry). After G3 NEGATIVE, G4a is the only sprint-scale electroweak-unification path remaining.

A footnote on the GUT-embedding direction: Sprint 4H Track SM-C (April 2026) tested whether $1/40 = \Delta$ in Paper 2's $K = \pi(B + F - \Delta)$ formula appears naturally in SU(5), E_8, or Spin(10) as a Chern number, η-invariant, or Chern–Simons level, and returned a clean negative: the dual-Coxeter rewrite $1/40 = (3/8)/15 = \sin^2\theta_W^{\mathrm{GUT}}/\dim(\bar{\mathbf{5}} \oplus \mathbf{10})$ is tautological, recycling the integers 5 and 8 already present in Paper 2. The naive shell-to-generation embeddings $\mathrm{shell}_n \to \bar{\mathbf{5}}$ fail at the $\mathrm{SU}(3) \times \mathrm{SU}(2)$ branching level. GeoVac therefore has no current contact with the GUT-embedding lineage of the SM, beyond the tautological numerical coincidences.

Unification targets after G3: what remains

Of the four gaps, G4 (cross-manifold) is the most consequential and — noting that G3 is *not* among the live ones, Sprint G3 having closed it in the negative (§8.3: identification is structurally impossible

by tensor-product factorisation) — the only sprint-scale path among the three that remain is G4a (§8.3, matching the assessment there); G1 has no current selecting mechanism (§8.1). *Corrected 2026-09-04:* this sentence inserted the G3 hedge and left the predicate "G3 (electroweak unification) is the most reachable" standing, so it announced a proven negative as the nearest-term target in the same breath as recording it closed. Paper 25's $U(1)$ and Paper 30's SU(2) already live on the same underlying graph (the Fock-projected S^3 Coulomb graph) and share the same Camporesi–Higuchi Dirac operator, the same incidence structure B, and the same edge Laplacian $L_1 = B^\top B$. The two constructions differ only at the gauge-group level, and Paper 30 §5.1 (Result 1) establishes that Paper 25 is exactly the maximal-torus reduction of Paper 30. The natural unified construction is therefore an $\mathrm{SU}(2) \times U(1)$ Wilson lattice gauge theory on the same graph, with link variables $U_e \in \mathrm{SU}(2) \times U(1)$ and a chosen embedding of hypercharge as the $U(1)$ phase. Whether such a construction admits a Higgs-style inner fluctuation through an almost-commutative extension (closing G2 within the electroweak sector) is an open structural question. Whether the Camporesi–Higuchi Dirac admits a chirality grading compatible with weak-isospin is *not* open: Sprint G3 closed it in the negative on S^3 alone (§8.3), and the text here presented that proven negative as "the second open structural question" until 2026-09-03. *That open question* — the almost-commutative lift of Paper 30's SU(2) Wilson instance, *not* the chirality-grading question Sprint G3 closed — is a clean S^3-only target that does not require any progress on the S^5 side. It is not by itself an electroweak construction: what it would deliver is a non-chiral $\mathrm{SU}(2) \times \mathrm{U}(1)$ Wilson product on S^3; chiral electroweak co-location on S^3 is what Sprint G3 closed (§8.3), and the remaining electroweak-unification path is G4a (Connes SM on the S^3 triple alone, below). (Clarified 2026-09-04: an earlier version of this paragraph labelled the instance $\mathrm{SU}(2)_L$ — the weak-isospin identification G3 refuted — and promised "unified electroweak Wilson construction on S^3".)

We do not claim such a unification has been achieved, or even that its component pieces have been verified. The status of this subsection is: a recorded structural fact (the unified $1/(4N_c)$ coefficient and the lineage placement are real); a recorded honest gap analysis (G1–G4 are real); and a recorded open direction (the almost-commutative lift of Paper 30's SU(2) Wilson instance is the most concrete S^3-only target, and G4a the remaining electroweak-unification path; the chirality-grading route is closed in the negative, §8.3). Per the project's rhetoric rule, no claim is made that GeoVac contains the Standard Model; the claim is only that it contains the three SM gauge groups as separate Wilson lattice gauge theories on its own sub-manifolds, with a unified weak-coupling kinetic coefficient and a unified gauge-network lineage.

8.2 Sprint H1: Higgs as inner fluctuation on the GeoVac electroweak almost-commutative extension

The G2 gap of Sec. 8.1 (Higgs as inner fluctuation) admits a constructive sprint-scale test: build the minimal electroweak almost-commutative extension $\mathcal{T}_{\mathrm{AC}} = \mathcal{T}_{\mathrm{GV}} \otimes \mathcal{T}_F$ with $\mathcal{A}_F = \mathbb{C} \oplus \mathbb{H}$ on the truthful Camporesi–Higuchi GeoVac triple, compute the inner fluctuation $D \mapsto D + \omega + \epsilon' J\omega J^{-1}$ explicitly at finite $n_{\max}$, and ask whether the off-diagonal $\mathbb{C} \leftrightarrow \mathbb{H}$ Yukawa block of the finite Dirac D_F that turns Marcolli–vS-without-Higgs into a Higgs construction can be *selected from GeoVac structure* rather than imposed. This is the H1 sprint reported in `debug/h1_ac_extension_memo.md`; we summarize its content here.

Construction. Following Connes–Marcolli 2008 Ch. 1 §13 (the finite noncommutative geometry) [2], the finite electroweak Hilbert space is doubled into matter and antimatter sectors: $\mathcal{H}_F = \mathcal{H}_F^{\mathrm{matter}} \oplus \mathcal{H}_F^{\mathrm{antimatter}} = \mathbb{C}^4 \oplus \mathbb{C}^4$, with $\mathcal{A}_F = \mathbb{C} \oplus \mathbb{H}$ acting on the matter sector only (the opposite-algebra action on antimatter is realized via J). The 8 by 8 finite Dirac is $D_F =$

block_diag$(M, \bar{M})$ where the matter-sector M is the 4 by 4 Yukawa-Dirac with off-diagonal $L \leftrightarrow R$ block $Y = \mathrm{diag}(y_\nu, y_e)$. The combined Dirac is $D = D_{\mathrm{GV}} \otimes \mathbb{1}_F + s_{\mathrm{GV}} \otimes D_F$ with $s_{\mathrm{GV}} = \mathrm{sign}(D_{\mathrm{GV}})$ (Track 2 closure): this commutes with D_{GV} on the truthful CH Dirac, and is *not* the chirality-doubling grading $\gamma_{\mathrm{GV}} = \sigma_x \otimes \mathbb{1}$ of Proposition 5(b) used below. **[SYMBOLIC + MEASURED]** The combined real structure is $J = J_{\mathrm{GV}} \otimes J_F$ with J_F swapping matter and antimatter via complex conjugation; matter-antimatter doubling makes $J_F^2 = +\mathbb{1}$ and $J_F D_F = +D_F J_F$ (the KO-dim 6 signs of $\mathcal{A}_F$ in the Connes–Marcolli convention). The signs of the combined J follow factor by factor: $J^2 = J_{\mathrm{GV}}^2 \otimes J_F^2 = -\mathbb{1}$, and since $s_{\mathrm{GV}} = \mathrm{sign}(D_{\mathrm{GV}})$ is a function of D_{GV}, $JDJ^{-1} = D$, i.e. $JD = +DJ$. The combined sign pair $(\varepsilon, \varepsilon') = (-, +)$ is the sign pair of the GV factor, which J_F leaves unchanged; in Connes' sign table [1] (in the three-sign convention of [3]) it is shared by the KO-2, KO-3 and KO-4 columns, KO-3 being the odd continuum factor's own label. It is *not* the KO-dimension $3 + 6 \equiv 1 \pmod 8$ that the additive rule for products of real spectral triples would assign: KO-1 has $(\varepsilon, \varepsilon') = (+, -)$, and the additive rule is a theorem about the product whose Dirac operator carries the grading of the even factor, $D_{\mathrm{GV}} \otimes \gamma_F + \mathbb{1} \otimes D_F$ (the $\widetilde{D}$ of Dabrowski–Dossena [4], eq. (18), which their §4.2 prescribes when the first factor is odd; their eq. (10) is the classical Cartesian-product analogue); for that product their Table 5 (row 3_+) shows that Connes' KO-6 real structure on the finite factor (column 6_+) admits no consistent ε' at all, and the alternative real structure $J_F \gamma_F$ (column 6_-) gives KO-1 (entry 1_-) with $J^2 = +\mathbb{1}$, $JD = -DJ$. Our D is the ungraded sum above, so the additive rule does not apply to it, and we label it by its signs (derived factor by factor above and confirmed numerically). Both signs are verified to machine precision at $n_{\max} \leq 3$ (`tests/test_almost_commutative.py`, `test_J_combined_sign_pair`). **[PANEL-VERIFIED]** The pair $(-, +)$ alone does not fix a KO-dimension: the KO-2, KO-3 and KO-4 columns of Connes' table [1, 3] share it; KO-3 is the label of the odd continuum GV factor (S^3 carries no chirality). At finite cutoff the chirality-doubling grading $\gamma_{\mathrm{GV}} = \sigma_x \otimes \mathbb{1}$ (`geovac/chirality_grading.py`), extended as $\gamma_{\mathrm{GV}} \otimes \mathbb{1}_F$, squares to one, anticommutes with the combined D and commutes with J bit-exactly at $n_{\max} \leq 3$, so the finite combined triple built on the chirality-doubled model is *even* with $(\varepsilon, \varepsilon', \varepsilon'') = (-, +, +)$, the KO-4 column, i.e. the sign triple of that column (measured 2026-09-02; pinned in the same test). We therefore attach *no* KO-dimension label to the finite model truncation (the true window carries no grading, Proposition 5(c)) and record the measured sign triple $(-, +, +)$ instead (PI direction, 2026-09-02): the continuum factor is KO-3 and the finite grading reads as KO-4, so any single label at finite cutoff is a convention, not a result. (An earlier version of this paragraph carried the label KO-3; before 2026-09-02 the paper carried KO-1, from the additive rule that does not apply here.)

Inner fluctuations split cleanly into gauge and Higgs sectors. For $\omega = \sum_i a_i [D, b_i]$ with $a_i, b_i \in \mathcal{A}_{\mathrm{GV}} \otimes \mathcal{A}_F$, we have

$$\omega = \underbrace{\sum_i a_i^{\mathrm{GV}} [D_{\mathrm{GV}}, b_i^{\mathrm{GV}}] \otimes a_i^F b_i^F}_{\text{gauge piece}} + \underbrace{\sum_i a_i^{\mathrm{GV}} b_i^{\mathrm{GV}} s_{\mathrm{GV}} \otimes a_i^F [D_F, b_i^F]}_{\text{Higgs piece}} . \tag{4}$$

For the diagonal $\mathcal{A}_{\mathrm{GV}}$ the gauge piece vanishes identically ($[D_{\mathrm{GV}}, b^{\mathrm{GV}}] = 0$, Proposition 2); the module (`geovac/almost_commutative.py`) takes b^{GV} from the operator-system multipliers of $\mathcal{O}_{n_{\max}}$ (`FullDiracTruncatedOperatorSystem`), which do not commute with D_{GV}, and the nonzero $\|\omega_{\mathrm{gauge}}\|$ values below are on that representative. The gauge piece is read as reproducing Papers 25 ($U(1)$ from the $\mathbb{C}$-summand) and 30 (SU(2) from the $\mathbb{H}$-summand) **[OBSERVATION]**; an operator-level identification with those papers' link variables is not established here. The Higgs piece is non-zero exactly when $[D_F, b_i^F]$ has off-diagonal $\mathbb{C} \leftrightarrow \mathbb{H}$ matrix elements — which happens

iff the imposed Yukawa Y is non-zero. At $n_{\max} = 1, 2, 3$, with three test choices of Y:

D_F choice	$\|\Phi\|_{\max}$	$\|\omega_{\text{gauge}}\|_{\max}$	**Falsifier**
$Y = 0$	0.000	0.000–0.580	**HOLDS**
$y_e = 0.1$, $y_\nu = 0$	0.046–0.080	0.000–0.580	FAILS
$y_e = 0.3$, $y_\nu = 0.2$	0.171–0.271	0.000–0.580	FAILS

(50 random generators per cell; data `debug/data/h1_falsifier.json`).

Verdict: positive-thin. The Higgs construction is *well-defined* on the GeoVac AC extension at finite $n_{\max}$: the algebra, Hilbert space, Dirac, and real structure satisfy the relevant Connes axioms (order zero/one on the GV factor excepted, as in Table 1), and inner fluctuations, taken over the operator-system multipliers, decompose into a gauge sector read as recovering Papers 25/30 plus a Higgs sector Φ that is non-trivial whenever the imposed Yukawa Y is non-zero.

However, GeoVac structure does **not autonomously select** a non-trivial Y. Two candidate sources for D_F off-diagonal data were flagged in the H1 scoping memo (`debug/almost_commutative_scoping_memo.` candidate A used the offdiag CH chirality bridging variant of the GeoVac Dirac, but Track 2 (`debug/real_structure_finite_nmax_memo.md` §3.D) ruled this out by showing $JD = +DJ$ fails at residual ~ 2.0 on offdiag CH; candidate B was a Sturmian / DUCC inter-shell coupling whose physical interpretation as a Yukawa is itself speculative and was not pursued.

The strong falsifier of the scoping memo §5 ("every Hermitian D_F derivable from GeoVac structure forces the Higgs sector to vanish") therefore does not fire as a clean negative. The construction admits a Higgs given any Y; the question is which Y, and GeoVac data is silent. This is exactly the positive-thin reading of the scoping memo §0: GeoVac sits on the *Marcolli–vS-without-Higgs* side of the 2014/2024 distinction at the structural level — not because inner fluctuations cannot define a Higgs (they can), but because no GeoVac mechanism selects the Yukawa.

What this means for G2. Gap G2 of Sec. 8.1 is sharpened to the more precise statement: *the Higgs construction goes through, but the Yukawa is a free input*, not closed to either "GeoVac contains a Higgs" (which would require a structural selection of Y) or "GeoVac excludes a Higgs" (which would require a structural obstruction to non-zero Y). This places G2 in the same status as G1 (three generations are formally definable but not selected by GeoVac structure): definable but not derivable.

What this means for G3. Gap G3 (hypercharge / electroweak unification, requiring identification of the GV chirality grading γ_{GV} with the SM weak-isospin chirality) is *not* closed by H1. In our construction γ_{GV} is the chirality-doubling grading $\sigma_x \otimes \not{k}$ of the chirality-doubled model truncation (Proposition 5(b); not $\text{sign}(D_{\text{GV}})$, which commutes with D_{GV}), while the SM weak-isospin chirality is a separate $\mathbb{Z}_2$-grading on the matter sector of $\mathcal{H}_F$. The two are added independently; no GeoVac mechanism currently identifies them.

What this means for the operator-system non-multiplicative-closure caveat. Remark 1's observation that $\mathcal{O}_{n_{\max}}$ is not closed under multiplication (witness pair $M^2_{2,1,0}$, 14.9% residual at $n_{\max} = 2$) is a separate, finite-resolution feature of the operator system itself. Inner fluctuations are defined operationally as $\omega + J\omega J^{-1}$ with $\omega = \sum_i a_i[D, b_i]$, where the multiplier matrices a_i, b_i are individual elements of $M_{\dim H}(\mathbb{C})$ (which is an algebra over $\mathbb{C}$); the lack of closure of *their span* $\mathcal{O}_{n_{\max}}$ under multiplication does not block the fluctuation calculus. Order-zero $[a, JbJ^{-1}] = 0$ holds

automatically by the matter–antimatter doubling (a acts on matter, JbJ^{-1} on antimatter, sectors are disjoint).

Scope and limitations. H1 establishes the constructive existence of the AC extension at finite $n_{\max}$ on the truthful CH GeoVac triple. H1 does *not* establish a Higgs vev as a derived quantity (that would require a continuum spectral-action computation; deferred to Track 1 R2.5 L4 closure for the rigorous interpretation), does *not* include the SM color factor $M_3(\mathbb{C})$ (deferred per Sec. 8.1 G4), and does *not* identify the GV and weak-isospin chiralities (G3 stays open). Per the project's rhetoric rule, no claim is made that GeoVac contains the Higgs: the claim is that it admits a Higgs given an imposed Yukawa, and that the Yukawa is not derived from GeoVac structure.

Files. Implementation `geovac/almost_commutative.py`; tests `tests/test_almost_commutative.py` (56 passing as of 2026-09-03); data `debug/data/h1_falsifier.json`; memo `debug/h1_ac_extension_memo.md`.

Thermal extension of the H1 verdict (Sprint TD Track 3, May 2026). The same Yukawa-undetermined structural reading extends to the thermal Higgs effective potential $V_{\text{eff}}(\phi, T)$ on the combined triple $T_{\text{GV}} \otimes T_{S^1_\beta} \otimes T_F$. At one-loop, the thermal mass squared takes the form

$$m_T^2(T) \;=\; \frac{T^2}{12} \cdot \mathrm{Tr}_{\text{inner}}\big(Y^2_{\text{diag}}\big) \;+\; (\text{gauge contribution}),$$

where the outer Stefan–Boltzmann backbone $T^2/12$ comes from Sprint TD Track 1's $M_1 \times M_2$ factorisation (Paper 35, tensor-product verification section), and the inner $\mathrm{Tr}_{\text{inner}}(Y^2_{\text{diag}})$ is a $k = 2$ Dirichlet sum in the Yukawa eigenvalues (Paper 18 § IV.6; η-trivialization-theorem-style filtering applies). Any candidate T_F whose squared Yukawa diagonal reproduces the empirical SM spectrum reproduces the SM-data-fit value $m_T^2(v, T_C) \approx (160\ \text{GeV})^2$; GeoVac admits any compatible spectrum but selects none.

Empirical falsification of the strict $Y = 0$ reading. The strict Marcolli–vS-without-Higgs case ($Y = 0$) predicts $\mathrm{Tr}_{\text{inner}}(Y^2_{\text{diag}}) = 0$ and hence no high-T symmetry-restoring quadratic correction to $V_{\text{eff}}(\phi, T)$. This is empirically falsified by the electroweak phase transition / sphaleron crossover at $T_C \approx 160$ GeV inferred from baryogenesis phenomenology. The empirical universe forces $Y_F > 0$. This does not derive Y_F, but it closes the H1 verdict's residual ambiguity: at finite T, the framework plus observed sphaleron phenomenology rules out the strict $Y = 0$ slice. The "thin" part of the positive-thin verdict is now thinner — five thermal observables (Sprint TD Track 3 catalogue) are individual probes of distinct inner-factor structures (Yukawas, color algebra, generation count, charge assignment, Higgs sector), and the inner factor is required to deliver a specific value at each. **[CONDITIONAL]** Three further candidates are *forbidden* by the η-trivialization theorem (Paper 18 § IV.6) on any Krajewski-class even spectral triple — a positive structural prediction, conditional on the observable being carried by the η-trivialised inner factor: axial chemical potential thermodynamics, parity-odd transport, and finite-T EDMs vanish identically on the η-trivialised inner factor of any Connes–Chamseddine-compatible AC extension.

Modular propinquity reformulation of the H1 falsifier (Sub-sprint M-H1, May 2026). The dual modular propinquity construction [58] provides a natural test of whether the Higgs sub-bimodule structure imposes a constraint on the Yukawa Y beyond what the Connes axioms already enforce. The bimodule reformulation of the H1 verdict (Sub-sprint M-H1, 2026-05-23) closes *negatively* on this question: the dual modular duality requirement reduces to Hermiticity of D_F plus matter–antimatter doubling, both already imposed by Connes' axiom system; the Higgs cross-block

Φ lives entirely in the fiber sub-bimodule $\mathcal{A}_F = \mathbb{C} \oplus \mathbb{H}$, structurally decoupled from any GeoVac-side bimodule data; and the Morita-triviality constraint ("could $\Phi = u^* v$ for GeoVac-derived u, v?") does not survive matter–antimatter doubling. The H1 POSITIVE-THIN verdict survives unchanged. The "dual" in dual modular propinquity is, on our reading of Latrémolière's construction [58] (the abstract does not say), the bridge $\leftrightarrow$ tunnel construction duality, not Morita equivalence; the Morita route was structurally barred independently. *Net:* the framework gains substantive new content on the chemistry / bound-state QED side (Sub-sprints M-X, M-Y, M-Z) but not on the SM-unification side (M-H1). Source: `debug/sprint_modular_propinquity_mH1_higgs_memo.md`, `debug/sprint_modular_propinquity_synthesis_memo.md`.

Sprint M-Z partition tested on a second concrete case (W1c chemistry wall, Sprint β-neg 2026-05-23). The Sub-sprint M-Z cross-shift / endomorphism partition was tested on a second concrete case beyond the bound-state QED context where it was originally derived: the W1c-residual chemistry wall at NaH alkali hydrides (Paper 19 [60], Sec. `sec:w1c_residual`). The bridge sprint (`debug/sprint_w1c_mz_partition_analysis_memo.md`, 2026-05-23) proposed a *three-bucket refinement* — cross-shift (handled) / basis-closable cross-shift (handled at sufficient basis size) / endomorphism (external input) — and tested it via three prespecified falsifiable predictions for NaH max_n=3 (`debug/data/sprint_w1c_mz_partition_predictions.json`). The F1 max_n=3 sprint (`debug/sprint_f1_maxn3_predictions_test_memo.md`) returned **CLEAN NEGATIVE**: 1 of 3 predictions pass; P1 (internal PES minimum) fails on the exact falsification criterion, the proposed basis-closable cross-shift sub-class evaporates. *The two-bucket M-Z partition (cross-shift / endomorphism) stands as the canonical framing.*

The substantive structural finding the test produced is a sharper localization of chemistry-side endomorphisms: at max_n=3 under combined W1c + multi-zeta architecture, the framework DOES construct the right bonding orbital (dominant natural orbital $-0.698 \cdot \phi_{\text{Na, 3s}} - 0.687 \cdot \phi_{\text{H, 1s}}$ with 50/50 amplitude split), but the constructed bonding combination has higher energy than the separated configuration at every R. The framework can BUILD the right orbital but cannot energetically PREFER it. Sub-layer 3 of the W1c hierarchy reclassifies from "basis-closable cross-shift" to "module endomorphism (basis-irreducible at tested scales)" and joins sub-layer 2 in the endomorphism bucket.

Same-day same-arc follow-on sprints (F2 + F3) sharpen the structural reading further into an architectural-extension ladder. The F1 max_n=3 verdict "module endomorphism, basis-irreducible at tested scales" was further refined by two same-day same-arc follow-on sprints (F2 + F3) executing the diagnostic-before-engineering rule at the sprint-cycle level. **F2** (cross-V_{ne} kernel-shape substitution diagnostic, KERNEL-NOT-IT verdict): the multipole expansion at default $L_{\max} = 4$ is bit-faithful to converged 3D numerical quadrature within $\sim 2 \times 10^{-5}$ Ha at NaH $R_{\text{eq}} = 3.566$ bohr, three to four orders of magnitude below the bond-energy scale (maximum differential 2.7×10^{-4} of $D_e = 0.075$ Ha in the sprint memo's own table). Ruled out kernel-shape substitution as a closure mechanism. Substantive new content: the framework's h1 in `composed_qubit.build_composed_hamiltonian` is strictly block-diagonal — the cross-block off-diagonal h1 slot is *architecturally absent*, not just kernel-error-flawed. New sub-wall name **W1d** (cross-block h1 architectural extension), structurally distinct from a missing calibration input. **F3** (cross-block h1 architectural extension): implemented the missing matrix slot via new production module `geovac/cross_block_h1.py`; FCI naturals at NaH max_n=2 R=3.5 transform $[1.0000, 1.0000] \to [\mathbf{1.9991}, \mathbf{0.0007}]$ — a three-order-of-magnitude change ($1.0000 \to 0.0007$) in dominant natural-orbital occupation signature, with the bonding combination as a doubly-occupied FCI ground state. **W1d CLOSED at the FCI level.** But F3 mini-PES remains monotonically

descending with $D_e^{\mathrm{F3}} = +4.37$ Ha $= 58\times$ experimental: cross-block h1 enables bonding without supplying opposing Pauli repulsion against the [Ne] frozen core. **New sub-layer W1e (inner-region overattraction)**, F4 target via a bonding-orbital Phillips-Kleinman extension.

Net refined reading of the chemistry sub-walls under M-Z: chemistry sub-walls split into THREE structurally distinct sub-categories within the two-bucket M-Z partition: cross-shifts (handled by existing architecture), module endomorphisms (split chemistry-side mitigable vs QED-side strictly-external), and **architectural-absence** (missing matrix slots in the existing architecture, conditionally closable via architectural extension). The architectural-absence category is a structurally distinct refinement WITHIN the two-bucket partition, not a contradiction of it. The architectural-extension ladder (W1c-cross-screening → W1c-multi-zeta-basis → W1d-cross-block-h1 → W1e?) is the chemistry-side analog of refinements that the QED side does not natively support: QED endomorphisms (LS-8a Z_2-1, δm; Sprint H1 Yukawa) are strictly externally specified via UV-completion physics, while chemistry endomorphisms are intrinsically external but framework-internally computable (FrozenCore + atomic-physics fits) AND admit architectural extensions (cross-block h1, bonding-orbital PK). This explains why the W1c arc produced a five-sub-layer hierarchy (W1c, W1c-multi-zeta, W1d, W1e, plus the FALSIFIED three-bucket M-Z refinement candidate) while the QED-side LS-8a wall has remained at a single-counterterm-input characterization.

Sources: `debug/sprint_beta_neg_synthesis_memo.md` (Sprint β-neg synthesis, F1-maturity; superseded for framing by the full-arc memo); `debug/sprint_f2_cross_vne_kernel_memo.md` (F2 KERNEL-NOT-IT + architectural-absence finding, W1d named); `debug/sprint_f3_cross_block_h1_memo.md` (F3 W1d CLOSED at FCI level + W1e newly named); `debug/sprint_w1c_full_arc_synthesis_memo.md` (comprehensive full-arc synthesis at F3 maturity, supersedes Sprint β-neg synthesis at the framing level). Cross-references Paper 19 §`sec:w1c_residual` (W1c-residual wall + F1-F2-F3 arc extension with revised five-sub-layer hierarchy), Paper 17 [59] §6.10 (composed-architecture cross-reference, updated to F3 maturity), Paper 34 §`sec:conv_three_bucket_falsified` (running-catalogue entry, parent), and Paper 34 §`sec:conv_w1c_f2_f3` (running-catalogue entry, F2+F3 follow-on).

F4–F6 refinement of W1e characterization (β-update, 2026-05-23). Three diagnostic sprints in immediate succession (F4, F5, F6) further refined the W1e (inner-region overattraction) sub-layer characterization beyond F3's working hypothesis of "missing Pauli repulsion against the [Ne] frozen core," tested all three candidate closure mechanisms F3 §6 had named, and sharpened the chemistry-side sub-mechanism dissection of W1e in the M-Z partition. **F4** (bonding-orbital Phillips–Kleinman extension): RULED OUT at Step 1 diagnostic — rank-1 PK on the F3-constructed bonding orbital saturates at ∼43% closure ceiling in the FCI sensitivity test, far below the 96% required for the 2× experimental closure window. **W1e is therefore NOT a single-particle Pauli- orthogonality wall** (which would have been the cross-shift-class mechanism within M-Z); it is a multi-determinant FCI correlation effect. **F5** (explicit-core Hartree treatment of [Ne] electrons via cross-block $\sum_c (J-K)$): RULED OUT at Step 1 diagnostic — right SIGN (repulsive, +1.12 Ha) but only 25.7% of wall depth; predicted residual $D_e^{\mathrm{F5}} \approx +3.25$ Ha $=$ 43× experimental. **W1e is NOT addressable by mean-field Hartree-class core-bonding interaction** (which would have been closable by extended-bimodule machinery of standard-Hartree class operators within M-Z). **F6** (valence basis enlargement to NaH max_n=4 with full F3 stack): PARTIAL_IMPROVEMENT — 10.2% PES-level closure (the 2-point gate suggests 26.1% but is artifactual; the inner-region collapse continues from $R = 3.5$ down to $R = 2.0$ with another 0.696 Ha descent the 2-pt gate misses). **W1e is NOT addressable by basis enlargement alone at max_n=4** (which would have been a basis-truncation cross-shift candidate within M-Z).

Net structural reading of the chemistry-side sub-mechanism dissection of W1e under M-Z: the chemistry-side dissection of W1e refines the framework's understanding of where "correlation" sits in the M-Z partition. W1e correlation is neither single-particle Pauli (which is the cross-shift-class

mechanism within M-Z, ruled out by F4) nor mean-field Hartree-level (which would be closable by extended-bimodule machinery for standard-Hartree-class operators, ruled out by F5), but **genuine multi-determinant content that requires beyond-sprint-scale architectural lift** (Schmidt orthogonalization at the basis-construction level, DMRG-class beyond-FCI methods, or explicit correlation factors — all are beyond-sprint-scale architectural investments). W1e characterizes a **three-sub-sub-mechanism decomposition with structurally-independent closure ceilings**: basis-truncation (∼10% PES, F6 measured) + Hartree-pressure (∼25% 2-pt, F5 predicted) + deep-correlation-residual (∼65–90% of wall, untested by sprint-scale compute). The chemistry-side architectural-extension ladder (W1c-cross-screening → W1c-multi-zeta-basis → W1d-cross-block-h1) has plausibly hit a natural pause point at W1e: the next architectural extension candidates are beyond-sprint-scale commitments rather than sprint-scale diagnostic-then-implementation cycles, and the chemistry arc's forward direction shifts from "find the right closure mechanism" to "characterize the framework's structural-skeleton scope limits empirically for second-row hydride binding" — consistent with the broader multi-focal-composition wall taxonomy of the working-hypothesis register finding (the framework's structural-skeleton scope does not autonomously deliver binding energetics at this precision).

Methodological side finding for future M-Z sub-mechanism sprints. The F6 2-point-vs-PES discrepancy reveals that 2-point gates at fixed reference geometry ($R = 3.5$ bohr vs $R = 10$ bohr) overestimate PES-derived closure by ∼2.5× on monotonically-descending PES in the W1c-residual regime. Future sprint-cycle diagnostics in this regime should use PES well depth at $R_{\min}$ as the load-bearing closure metric rather than 2-point energy differentials. The F3-reported $D_e^{\mathrm{F3}} = +4.37$ Ha is itself a 2-pt value; the corresponding F3 PES well depth at $R_{\min} = 2.0$ is likely larger (untested but expected by analogy to F6). This is a methodological refinement of the diagnostic-before-engineering discipline (working-hypothesis register) at the sprint-cycle level: not just "run a diagnostic before production wiring," but also "choose the diagnostic geometry that captures the load-bearing physical content rather than a convenient reference."

Sources (F4–F6 refinement): `debug/sprint_f4_bonding_pk_memo.md` (F4 STOP-at-Step-1, rank-1 PK insufficient); `debug/sprint_f5_explicit_core_memo.md` (F5 STOP-at-Step-1, mean-field Hartree insufficient); `debug/sprint_f6_maxn4_nah_memo.md` (F6 PARTIAL_IMPROVEMENT, basis enlargement to max_n=4); `debug/sprint_beta_update_f4f6_memo.md` (β-update F4-F6-APPENDED-TO-F3-BASELINE synthesis). Cross-references Paper 19 §`sec:w1e_refinement_f4_f6` (refined six-sub-layer hierarchy with three W1e sub-sub-mechanisms), Paper 17 §6.10 (composed-architecture cross-reference, updated to β-update F4–F6 maturity), Paper 34 §`sec:conv_w1e_f4_f6_refinement` (running-catalogue entry, F4–F6 follow-on to F3 entry).

W1e sprint-scale candidate space closed at Schmidt + core correlation Day-1 diagnostics (2026-05-23). The two beyond-sprint-scale candidates F4–F6 named (Schmidt orthogonalization at basis-construction level; fully correlated [Ne] cores) were tested by Day-1 diagnostics. **Both ruled out.** Schmidt-orthogonalizing H 1s against the Na [Ne] core at $R = 3.5$ bohr produces $\sum_c |S_c|^2 = 1.00 \times 10^{-3}$ (only 0.1% projection mass; the H atom at $R = 3.5$ is structurally outside the [Ne] core's significant amplitude) and a total diagonal-element differential $\Delta_{\mathrm{total}} = +8.6$ mHa — $\sim 12\times$ below the prespecified GO threshold and $\sim 500\times$ below the wall depth. [Ne] core-correlation differential at NaH R_{eq} converges across three independent literature anchors (Iron-Oren-Martin 2003, Sylvetsky-Martin 2018, atomic Na correlation × active-fraction) to ∼ 0.001–0.003 Ha, $10^3\times$ below the wall depth. *Five independent sprint-scale W1e closure mechanisms are now ruled out* (F4, F5, F6, Schmidt, core correlation). The chemistry-side architectural-extension ladder (W1c-cross-screening → W1c-multi-zeta-basis → W1d-cross-block-h1) has reached the sprint-scale boundary; further closure candidates are substantial architectural commitments. The W1e wall is the **sixth structurally-independent instance of**

the multi-focal-composition wall pattern across the framework, joining five physics observables (Sprint H1 Yukawa, LS-8a $Z_2/\delta m$, HF-3 recoil, HF-4 Zemach, HF-5 multi-loop a_e). *The pattern's cross-domain transferability — five physics + one chemistry, two structurally distinct calibration-data classes hitting the same structural-skeleton-scope boundary — is the substantive new content of this closeout, extending the structural-skeleton scope statement from physics-only to physics+chemistry. Sources:* `debug/sprint_w1e_schmidt_diagnostic_memo.md` (Schmidt Day-1 STOP), `debug/sprint_w1e_core_correlation_estimate_memo.md` (core correlation DEFER). Cross-references Paper 19 §`sec:w1e_schmidt_core_correlation` and Paper 34 §`sec:conv_w1e_cross_domain_wall`

8.3 Sprint G3: Electroweak chirality co-location on S^3

Sprint H1 (Sec. 8.2) closed G2 up to the Yukawa-undetermined verdict and left G3 (electroweak unification, requiring identification of γ_{GV} with the SM weak-isospin chirality on $\mathcal{H}_F$) as the next physics question. The G3 sprint (May 2026) is a constructive test: build γ_{GV} on the GeoVac side and γ_F on the AC side as separate $\mathbb{Z}_2$ gradings each satisfying the standard NCG axioms, lift both to the tensor-product Hilbert space $\mathcal{H}_{\mathrm{GV}} \otimes \mathcal{H}_F$, and ask whether the residual

$$\Delta := \gamma_{\mathrm{GV}} \otimes \mathbb{1}_F - \mathbb{1}_{\mathrm{GV}} \otimes \gamma_F \tag{5}$$

vanishes (or vanishes on the physical sector under standard left/right convention swap). $\Delta = 0$ would close G3 in the positive; $\Delta \neq 0$ robustly would close G3 in the negative on S^3 alone and push electroweak unification onto G4 cross-manifold territory.

Constructions. The sprint executed three parallel tracks (memos `debug/g3a_chirality_memo.md`, `debug/g3b_chirality_F_audit_memo.md`, `debug/g3c_tensor_chirality_memo.md`). Track G3-A constructed $\gamma_{\mathrm{GV}} = \sigma_x \otimes \mathbb{1}_w$ on the R3.5 full Dirac operator system at $n_{\max} \in \{1, 2, 3\}$, with $w = \mathrm{spinor_dim}(n_{\max})$ and the basis ordering $[\chi = +1, \chi = -1]$ from `full_dirac_basis`. All three $\mathbb{Z}_2$-grading axioms are bit-exact at every tested $n_{\max}$: $\gamma_{\mathrm{GV}}^2 = \mathbb{1}$, $\{\gamma_{\mathrm{GV}}, D_{\mathrm{truthful}}\} = 0$, and $\gamma_{\mathrm{GV}} M \gamma_{\mathrm{GV}} = M$ as a matrix equality for every scalar multiplier $M \in \mathcal{O}_{n_{\max}}$. The natural σ_z candidate (the D-eigenvalue sign) commutes with D_{truthful} rather than anticommuting and is exposed only as a diagnostic. Module `geovac/chirality_grading.py` (50/50 tests passing).

Track G3-B added `ElectroweakFiniteTriple.chirality_F()` to `geovac/almost_commutative.py` with the standard Connes–Marcolli SM convention at KO-dim 6 of $\mathcal{T}_F$:

$$\gamma_F^{\mathrm{matter}} = \mathrm{diag}(+1, +1, -1, -1), \qquad \gamma_F^{\mathrm{anti}} = -\gamma_F^{\mathrm{matter}}. \tag{6}$$

The antimatter sign-flip is forced by $\{J_F, \gamma_F\} = 0$: the naive "identical on both copies" convention would put $\mathcal{T}_F$ at KO-dim 0 and silently break the H1 module's $\{J_F, D_F\}$ axiom suite. All grading axioms ($\gamma_F^2 = \mathbb{1}$, $\{\gamma_F, D_F\} = 0$ at any imposed Yukawa, $\gamma_F \pi_F(a) \gamma_F = \pi_F(a)$, $\gamma_F = \gamma_F^*$, $\{J_F, \gamma_F\} = 0$) hold to bit-exact zero residual; 15 new tests in `TestChiralityF`, all passing. The finite factor therefore carries the KO-dim 6 signs $(\varepsilon_F, \varepsilon'_F, \varepsilon''_F) = (+, +, -)$; **[SYMBOLIC + MEASURED]** the combined real structure on H1's combined Dirac carries the sign pair $(\varepsilon, \varepsilon') = (-, +)$, the GV factor's pair — with $\varepsilon'' = +1$ under the finite grading, and no finite-cutoff KO label attached (§8.2). The additive rule $3 + 6 \equiv 1 \pmod 8$ does not apply to H1's ungraded sum $D_{\mathrm{GV}} \otimes \mathbb{1}_F + s_{\mathrm{GV}} \otimes D_F$; it governs the graded product of Dabrowski–Dossena [4], which the paper does not use.

Tensor-product diagnostic. Track G3-C computed Δ at $n_{\max} \in \{1, 2, 3\}$ using the production γ_{GV} and γ_F from G3-A and G3-B. Sanity invariants are bit-exact: $\gamma_{\mathrm{GV}}^2 = \mathbb{1}$, $\gamma_F^2 = \mathbb{1}$, and $[\gamma_{\mathrm{GV}} \otimes \mathbb{1}_F, \mathbb{1}_{\mathrm{GV}} \otimes \gamma_F] = 0$ (the two operators commute because they act on disjoint tensor factors). The residual itself is the structurally cleanest possible non-zero object:

Convention swaps do not absorb the residual. Three standard convention swaps were tested at every $n_{\max}$: (a) global sign-flip $\gamma_F \to -\gamma_F$ (left/right convention), (b) restriction to the matter sector of $\mathcal{H}_F$, and (c) restriction to the Weyl chirality of $\mathcal{H}_{\mathrm{GV}}$. Swap (a) leaves the baseline invariant. Swaps (b) and (c) reduce $\|\Delta\|_F$ by dropping basis vectors, but $\|\Delta\|_{\mathrm{op}}$ remains 2 (or 1 under swap (c) with σ_x-γ_{GV}, where the Weyl restriction collapses $\gamma_{\mathrm{GV}}|_w$ to 0_w and $\Delta_c = -\mathbb{1}_w \otimes \gamma_F$, still non-zero). Combined swaps yield the same conclusion. No standard NCG sign convention absorbs Δ.

Robustness. A σ_x-vs-σ_z cross-check at $n_{\max} = 2$ confirms that the spectrum of Δ is identical under both γ_{GV} conventions (both produce $\{-2{:}32,\ 0{:}64,\ +2{:}32\}$, since σ_x and σ_z share eigenvalues ± 1 with identical multiplicities and both commute with γ_F). The structural verdict is therefore convention-independent.

Sanity check on $\gamma_5 := \gamma_{\mathrm{GV}} \otimes \gamma_F$. The product operator $\gamma_5 := \gamma_{\mathrm{GV}} \otimes \gamma_F$ is the canonical Connes–Chamseddine combined chirality on $\mathcal{T}_{\mathrm{AC}}$ and satisfies $\gamma_5^2 = \mathbb{1}$ to bit-exact precision at every $n_{\max}$, with spectrum $\{-1, +1\}$ in equal multiplicities. The product works as an involution; the *difference* does not. γ_5 is a well-defined $\mathbb{Z}_2$ operator, but it is a *grading of D* only for the production $\gamma_{\mathrm{GV}} = \sigma_x \otimes \mathbb{1}$ at vanishing Yukawa: with $s_{\mathrm{GV}} = \mathrm{sign}(D_{\mathrm{GV}})$ in its place one has $\|\{\gamma_5, D\}\|_F = 17, 52, 110$ at $n_{\max} = 1, 2, 3$, and with the production grading $4.7, 9.3, 14.8$ at $Y \neq 0$ (two distinct operators carried the name γ_{GV} in this section; disambiguated 2026-09-03, the sign operator renamed s_{GV} as in the Higgs section). γ_{GV} and γ_F do not identify across the tensor product.

Verdict: NEGATIVE on S^3 alone. The two chirality gradings γ_{GV} (the chirality-doubling grading $\sigma_x \otimes \mathbb{1}$ of Proposition 5(b)) and γ_F (weak-isospin grading on the AC factor) are *independent commuting* $\mathbb{Z}_2$ gradings on $\mathcal{H}_{\mathrm{GV}} \otimes \mathcal{H}_F$. They cannot be identified by any standard NCG sign convention because, by construction, γ_{GV} acts as identity on $\mathcal{H}_F$ and γ_F acts as identity on $\mathcal{H}_{\mathrm{GV}}$: they are different operators, acting on different tensor factors. Electroweak chirality co-location does not close on S^3 alone.

The G2–G3 corollary. Sprint H1 left G2 (Higgs as inner fluctuation) at the verdict "construction admits a Higgs given an imposed Yukawa Y, but no GeoVac mechanism selects Y." G3's NEGATIVE supplies the structural reason. The Yukawa sits in the off-diagonal $\mathbb{C} \leftrightarrow \mathbb{H}$ block of D_F that anticommutes with γ_F; the candidate GeoVac sources for that block (offdiag CH chirality bridging, Sturmian inter-shell coupling) have no kinematic relation to γ_F, because γ_{GV} is structurally independent of γ_F. Equivalently: no GeoVac-side data fixes Y because the GeoVac chirality lives on a different $\mathbb{Z}_2$ grading from the one the Yukawa flips. G2 and G3 are therefore one structural fact, not two. This is a *sharpening* of H1's G2 verdict (the Yukawa-undetermined gap is named structurally), not a new gap.

Files. Modules `geovac/chirality_grading.py` (G3-A, 50/50 tests) and the `chirality_F` additions to `geovac/almost_commutative.py` (G3-B); tests `tests/test_chirality_grading.py` and `tests/test_almost_commutative.py` (56 passing, `TestChiralityF` block); diagnostic `debug/g3c_tensor_chirali`

data `debug/data/g3a_chirality.json`, `debug/data/g3c_tensor_chirality.json`; memos `debug/g3a_chirality`
`debug/g3b_chirality_F_audit_memo.md`, `debug/g3c_tensor_chirality_memo.md`.

8.4 Sprint G4: Cross-manifold scoping (G4a reachable, G4b NCG-framework-blocked)

With G2 and G3 collapsed into one structural fact and electroweak unification on S^3 alone closed in the negative, G4 (cross-manifold) becomes *the* remaining route to a GeoVac-internal SM construction rather than the dominant gap among several. A scoping pass (May 2026, memo `debug/g4_cross_manifold_sco`) sharpens G4 into two sub-gaps with sharply different reachability.

G4a (reachable): Connes SM on $\mathcal{T}_{S^3}$ alone with $\mathcal{A}_F = \mathbb{C} \oplus \mathbb{H} \oplus M_3(\mathbb{C})$. The minimal extension of Sprint H1's electroweak slice $\mathcal{A}_F = \mathbb{C} \oplus \mathbb{H}$ to the full Connes SM finite algebra is reachable as a bounded sprint-scale extension extending `geovac/almost_commutative.py`. The $M_3(\mathbb{C})$ summand supplies SU(3) at the inner-fluctuation level on the existing GeoVac S^3 triple, without recourse to the S^5 Bargmann construction. Predicted outcome: *positive-thin*, in the exact sense of Sprints H1 and G3. Inner fluctuations decompose into a gauge-1-form sector (recovering $U(1) \times \mathrm{SU}(2) \times \mathrm{SU}(3)$ at the operator level, with the abelian and SU(2) parts already verified by H1) plus a Higgs sector that is non-trivial whenever the Yukawa-Dirac off-diagonal blocks are imposed. The G2–G3 corollary applies verbatim: G4a defines the construction, but no GeoVac mechanism selects the Yukawa, hypercharge assignments, or generations. G4a therefore does not close the SM-derivation question; it relocates the open data-selection items (Y, hypercharge, generation count) from "electroweak slice" (H1, G3) to "full Connes finite algebra," without changing their status. G4a remains worth pursuing because it makes Sprint ST-SU3's Bargmann SU(3) construction structurally redundant for SM-derivation purposes (one S^3 construction with $M_3(\mathbb{C})$ delivers the same gauge content) and because it is the most published-precedent-compatible target available.

G4b (NCG-framework-blocked): genuine cross-manifold $\mathcal{T}_{S^3} \otimes \mathcal{T}_{S^5}$. The structurally deeper version of G4 asks whether the GeoVac S^3 Coulomb triple and the Paper 24 S^5 Bargmann lattice can be combined into a single spectral triple with both as commuting factors, so that the SU(3) of Sprint ST-SU3 is honestly cross-manifold-located rather than recapitulated on S^3. This sub-gap is blocked at the *NCG-framework level*, not at the GeoVac level. Connes' tensor product of spectral triples [2] requires both factors to be Riemannian spectral triples in the same sense. $\mathcal{T}_{S^3}$ carries the Camporesi–Higuchi Riemannian spinor Dirac (KO-dim 3). Paper 24's S^5 Bargmann construction is built from the holomorphic Hardy sector $H^2(S^5)$ restricted to symmetric SU(3) irreps $(N, 0)$; its natural diagonal operator is the first-order complex Euler operator $\hat{N} = \sum_i z_i \partial_i$, not a Riemannian Dirac. Two candidate $\mathcal{T}_{S^5}$ exist, and neither preserves both standard NCG axioms and the GeoVac-specific content of Sprints 25–30 / ST-SU3:

- *Option A (Riemannian round-S^5).* Take D_{S^5} to be the standard Riemannian spinor Dirac on round S^5 (KO-dim 5). The tensor product is a textbook Connes object with combined KO-dim $3 + 5 = 8 \equiv 0 \pmod 8$, but it discards the $(N, 0)$-tower structure that made Paper 24's S^5 "natural," and the bit-exact π-free certificate of Paper 24 §III is irrelevant to round-S^5 (calibration π reappears via standard Weyl asymptotics).
- *Option B (Bargmann–Euler-based).* Promote the Euler operator $\hat{N}$ on the Bargmann graph to a Dirac-like object via chiral doubling. This preserves Sprint ST-SU3 SU(3) Wilson content and the π-free certificate, but the resulting object is *not* a Riemannian spectral triple in

the published sense: KO-dimension is not classically defined for a Hardy-sector first-order complex operator, and there is no published prescription in the NCG literature for tensoring a Riemannian spectral triple with a Hardy-sector / Berezin–Toeplitz first-order complex object. The closest analogs (Berezin–Toeplitz quantisation of the unit disk; Hekkelman 2022 [14] on the circle; Hekkelman–McDonald 2024 [15] on general spectrally truncated triples) stay within Berezin–Toeplitz on a single manifold or within Riemannian on a single manifold, never crossing.

The structural reading is that G4b is the Coulomb/HO asymmetry of Paper 24 §V resurfacing at the spectral-triple level. Paper 24 §5.3 records these layers of this asymmetry (seven catalogued, six standing, Paper 31), the standing ones all Coulomb-specific; the first four: (i) spectrum-computing role of L_0 (*withdrawn* 2026-09-25: L_0 is spectrally inert on both spheres); (ii) calibration π; (iii) Wilson gauge with natural matter coupling (argued), Sec. 8.1; and (iv) modular-Hamiltonian Pythagorean orthogonality on the hemispheric wedge, the structural mechanism behind the L2-E signature-independence finding (Paper 42 §11.7 [53]; Remark 47 above). G4b is a *sibling* of these layers, not a lone additional layer: of the four ingredients listed next, the Coulomb S^3 side carries a Riemannian spectral triple in the standard NCG sense and supports the finite-cutoff wedge period closure (for the chosen integer-spectrum generator; signature-blind, so the $(3,1)$ Krein carrier adds no Lorentzian content), while the Bargmann S^5 side has none of the four ingredients (no spinor bundle on the $(N,0)$ Hardy tower; no second-order-vs-first-order distinction with a separate Dirac; no half-integer wedge; no calibration π or natural Wilson matter coupling). The Connes tensor-product framework cannot bridge them as currently published. Closing G4b is therefore not a sprint-scale GeoVac task; it requires either an extension of NCG to handle Riemannian–Hardy mixed tensor products, or the identification of a unifying ambient manifold whose GeoVac substructure contains both S^3 Coulomb and S^5 Bargmann as natural sub-objects. Both directions are recorded as honest open structural questions, not as sprint targets.

Net. After Sprints H1 + G3 + G4 scoping, the four-gap analysis of Sec. 8.1 compresses to:

- G2 and G3 are one structural fact: the Yukawa is undetermined because γ_{GV} and γ_F are independent commuting $\mathbb{Z}_2$ gradings on the tensor product (G3 NEGATIVE).
- G4 splits cleanly: G4a (Connes SM on $\mathcal{T}_{S^3}$ alone with $M_3(\mathbb{C})$) is **closed at POSITIVE-THIN** (2026-05-31): $\mathrm{U}(1)\times\mathrm{SU}(2)\times\mathrm{SU}(3)$ recovered from inner fluctuations, the J-relations, self-adjointness and $\gamma_F^2 = I$ bit-exact (the combined order-zero/one residuals carry the GV factor's operator-system residual; see the scope correction below), Yukawa/hypercharge/generations remain imposed; G4b (genuine cross-manifold) is blocked at the NCG-framework level by the Coulomb/HO structural asymmetry.
- G1 (three generations) is unchanged from H1: formally definable, not selected by GeoVac structure.

GeoVac does not autonomously generate the Standard Model; the construction admits SM gauge content given imposed Yukawa, hypercharge, and generation choices, and the GeoVac data is silent on each of those choices. This is a clean structural verdict, in the sense of the project's rhetoric rule: the framework is consistent with a GeoVac SM construction, but the framework does not select one.

G4a closure (Sprint G4a, 2026-05-31). The full Connes SM finite algebra $\mathcal{A}_F = \mathbb{C}\oplus\mathbb{H}\oplus M_3(\mathbb{C})$ has been implemented and verified on $\mathcal{T}_{S^3}$ at $n_{\max} \in \{1,2,3\}$ with $\dim \mathcal{H}_F = 32$ (16 matter + 16 antimatter; $\dim \mathcal{H}_{\text{total}} = 1280$ at $n_{\max} = 3$). **[PANEL-VERIFIED]** The axioms $J^2 = -I$, $JD = +DJ$, $D = D^\dagger$ and $\gamma_F^2 = I$ hold at machine zero at every tested $n_{\max}$. *Scope correction (2026-09-03):* the order-zero and order-one residuals were also machine zero, but for a restricted reason — the module's finite algebra acted on the matter sector only while JbJ^{-1} acts on antimatter, so the two commute by disjoint support (and the module's quark action $\text{kron}(ew, m)$ was not a linear $*$-representation of the direct sum; corrected in `geovac/standard_model_triple.py`). With the corrected representation the finite-algebra order-zero and order-one residuals are exactly zero, while the combined-triple residuals (1.49 and 1.26 at $n_{\max} = 2$) are the GV factor's finite-resolution residual (the operator-system multipliers do not commute; cf. the order-zero/one rows of Table 2), factorising through the operator-system commutator with residual 0 (`tests/test_standard_model_triple.py`); the gauge census still returns $\text{U}(1) \times \text{SU}(2) \times \text{SU}(3)$ at $n_{\max} = 2$ (colour now read from the antiquark block); and the combined grading $s_{\text{GV}} \otimes \gamma_F$ built from $s_{\text{GV}} = \text{sign}(D_{\text{GV}})$ does *not* anticommute with D ($\|\{\gamma, D\}\|_F = 34, 104, 220$ at $n_{\max} = 1, 2, 3$), so the "no finite-resolution degradation" reading is withdrawn; with the production grading $\gamma_{\text{GV}} = \sigma_x \otimes \mathbb{1}$ the anticommutator vanishes. Inner fluctuations decompose into $\text{U}(1)\times\text{SU}(2)\times\text{SU}(3)$ gauge 1-forms plus a Higgs scalar that is non-trivial whenever the Yukawa-Dirac off-diagonal blocks are imposed and identically zero otherwise. Matter-antimatter and lepton-quark cross-sectors decouple exactly. **[PANEL-VERIFIED]** The predicted POSITIVE-THIN verdict is confirmed: the construction admits the full SM gauge content with imposed Yukawa, hypercharge, and generation choices, and the GeoVac data is silent on each of those choices. Module `geovac/standard_model_triple.py` ($\sim$ 530 lines), 47 tests passing.

H1 Higgs inner fluctuation (Sprint H1, May 2026). The full fluctuated Dirac $D_A = D + A + JAJ^{-1}$ has been constructed at $n_{\max} \in \{2, 3\}$ ($\dim \mathcal{H}_{\text{total}} = 512, 1280$). Inner fluctuations decompose cleanly into gauge ($\text{U}(1) \times \text{SU}(2) \times \text{SU}(3)$, matching G4a) plus Higgs (complex SU(2) doublet Φ in the L–R off-diagonal block, non-trivial iff Yukawa $Y \neq 0$). The spectral-action Mexican-hat sign structure $V(\Phi) = -\mu^2|\Phi|^2 + \lambda|\Phi|^4$ is the standard CCM result; the checks $\text{Tr}(D_F^2) > 0$ and $\text{Tr}(D_F^4) > 0$ hold for any nonzero Hermitian D_F **[SYMBOLIC]**, so they are consistency checks, not independent confirmation.

Theorem 6 (H1 Yukawa non-selection)**.** ***[PROOF-BY-ARGUMENT]*** *Within the almost-commutative spectral triple $(\mathcal{A}_{\text{GV}} \otimes \mathcal{A}_F, \mathcal{H}_{\text{total}}, D)$ with $\mathcal{A}_F = \mathbb{C} \oplus \mathbb{H} \oplus M_3(\mathbb{C})$, the space of admissible finite Dirac operators D_F satisfying all Connes axioms (Hermiticity, chirality grading, J-reality, order-one condition) carries a finite-dimensional real moduli space. (Under the degenerate algebra sample used before 2026-09-03 the matter-sector reductions read $512 \to 256 \to 128 \to 128$, with order-one appearing to impose no further constraint; that sample was an artefact, and with a linear $*$-representation of $\mathbb{C} \oplus \mathbb{H} \oplus M_3(\mathbb{C})$ the matter-sector projection of the full-axiom moduli space has rank 16 and the full-axiom dimension is 32 — see Theorem 7. The Yukawa non-selection conclusion is independent of this count.) For one generation, $Y = \text{diag}(y_\nu, y_e, y_u, y_d)$ accounts for 8 of these. Given the imposed $\mathcal{A}_F$, the gauge sector $\text{U}(1) \times \text{SU}(2) \times \text{SU}(3)$ follows from its inner automorphisms; the Yukawa values are* ***not selected*** *by any morphism in $\mathcal{A}_{\text{GV}}$.*

Proof sketch. The parameter count follows from direct enumeration on the chirality-graded J-real subspace of Hermitian operators on $\mathcal{H}_F$. The non-selection of the Yukawa is the G3 structural finding (Sprint G3, §8.3): γ_{GV} and γ_F are independent commuting $\mathbb{Z}_2$'s on $\mathcal{H}_{\text{GV}} \otimes \mathcal{H}_F$, so no GeoVac-side data couples to the γ_F flip that carries the Yukawa. The full fluctuated Dirac $D_A = D + A + JAJ^{-1}$ was constructed at $n_{\max} \in \{2, 3\}$ ($\dim \mathcal{H}_{\text{total}} = 512, 1280$) and the inner fluctuations decompose

cleanly into the gauge piece fixed by $\mathcal{A}_F$ (matching G4a) plus the Higgs piece (complex SU(2) doublet Φ in the L–R off-diagonal block, non-trivial iff $Y \neq 0$). The Mexican-hat sign structure is the standard CCM result (the trace positivity checks are automatic for nonzero Hermitian D_F). See `debug/h1_higgs_inner_fluctuation_memo.md`. □

The forced/free seam (forcing-catalogue Door 4, 2026-06-01). **[OBSERVATION]** The Yukawa non-selection theorem admits a sharper structural reading. The almost-commutative factorization $D^2 = D_{\mathrm{GV}}^2 \otimes 1 + 1 \otimes D_F^2$ (the cross term vanishes by chirality anticommutation *for the graded product* $D = D_{\mathrm{GV}} \otimes 1 + \gamma_{\mathrm{GV}} \otimes D_F$ with $\{\gamma_{\mathrm{GV}}, D_{\mathrm{GV}}\} = 0$; for the module's $s_{\mathrm{GV}} = \mathrm{sign}(D_{\mathrm{GV}})$ in its place the cross term is $2|D_{\mathrm{GV}}| \otimes D_F \neq 0$, norms $4.3, 13.2, 28.0$ at $n_{\max} = 1, 2, 3$; scoped 2026-09-03) factorizes the master Mellin engine into an *outer* ring — the M1/M2/M3 transcendentals of the GeoVac sector — tensored with an *inner* Yukawa Dirichlet ring $\mathbb{Q}[y_i^{-2s}]$ generated by the Yukawa eigenvalues. The two rings share no generator: the η-trivialization ($\{\gamma_F, D_F\} = 0$ annihilates the odd-k heat traces that would couple them), together with the G3 commuting-$\mathbb{Z}_2$ result, leaves the inner Yukawa ring sharing no generator with any forced outer ring. This is the precise content of the structural-skeleton scope (§1.5 of the project guidelines): the bone (the forced, integer/rational skeleton) and the calibration data (free) live in rings sharing no generator, and the Yukawa is free *because* it generates its own ring — a quantity cannot be derived from a ring it generates. The reading also explains, after the fact, why the H1, W3, and Koide attempts to select the Yukawa all returned clean negatives: each probed *inside* the almost-commutative category, where the seam already forbids selection.

The inner algebra structure is mostly forced (Door 4b, 2026-06-01). The seam places the Yukawa *values* on the free side; it does *not* place the inner *algebra structure* there. **[PANEL-VERIFIED]** A finite-algebra enumeration shows that among all finite semisimple real $*$-algebras with at most three summands and factor size at most three, exactly *two* reproduce the gauge group $\mathrm{U}(1) \times \mathrm{SU}(2) \times \mathrm{SU}(3)$ as their unitary/inner-automorphism content: the Standard Model finite algebra $\mathbb{C} \oplus \mathbb{H} \oplus M_3(\mathbb{C})$ and $\mathbb{C} \oplus M_2(\mathbb{C}) \oplus M_3(\mathbb{C})$. Under the Bertrand × complex-Hopf-tower reading at $n \leq 3$ (Door 4b; an organizing reading recorded in the forcing catalogue, not an argument given in Sec. 8.1, which is a catalogue of gauge constructions on distinct manifolds), one factor is assigned per Hopf rung $n = 1, 2, 3$, which fixes the factor count (3), the $n = 1$ factor ($\mathbb{C}$), and the $n = 3$ factor ($M_3(\mathbb{C})$); the residual non-uniqueness collapses to a single binary fork — $\mathbb{H}$ versus $M_2(\mathbb{C})$ at the $n = 2$ rung. This fork was conjectured (Door 4b) to have a GeoVac-native handle: the outer triple's $J_{\mathrm{GV}}^2 = -1$ (the continuum factor's KO-3 sign; no KO label is attached at finite cutoff) (the real-structure result of Sec. 3, the pseudoreal $\mathrm{Spin}(3) = \mathrm{SU}(2)$ two-spinor sign — the *same* $S^3 = \mathrm{SU}(2)$ fact that produces the gauge SU(2)) might select $\mathbb{H}$ over $M_2(\mathbb{C})$. **[SYMBOLIC PROOF]** A direct J sign-table audit (Door 4c, 2026-06-01) closed this conjecture *negative*: $J_{\mathrm{GV}}^2 = -1$ *admits* but does not *force* $\mathbb{H}$. The decisive fact is that over $\mathbb{C}$ the two candidates are the same complex algebra ($\mathbb{H} \otimes_{\mathbb{R}} \mathbb{C} \cong M_2(\mathbb{C})$); the $\mathbb{H}$-vs-$M_2(\mathbb{C})$ distinction is a choice of *real form*, fixed by the sign of an *internal* $\mathbb{C}^2$ antilinear involution ($J_{\mathrm{int}}^2 = -1 \Leftrightarrow \mathbb{H}$; $J_{\mathrm{int}}^2 = +1 \Leftrightarrow M_2(\mathbb{C})$), which lives on the algebra side and is invisible to the combined $J = J_{\mathrm{GV}} \otimes J_F$. The combined J^2 and the combined $(\varepsilon, \varepsilon')$ signs are *bit-identical* for both candidates at $n_{\max} \in \{1, 2, 3\}$, and the Chamseddine–Connes–Marcolli order-one-with-Yukawa selector is satisfied by both (matter/antimatter decoupling). The $\mathbb{H}$ choice is therefore a *literature import* (the CCM complex-fermion-representation / second-order condition on the finite factor), not a GeoVac forcing; the fork stays open. The honest ledger: *fixed under the Hopf-tower reading* are the factor count and the $\mathbb{C}$ and $M_3(\mathbb{C})$ summands; *admitted but not forced* is the $n = 2$ real form ($\mathbb{H}$ vs $M_2(\mathbb{C})$); *free* are the generation count (invisible to inner automorphisms),

the inner KO-dimension, and the Yukawa values. The net sharpening is real but bounded and conditional on the Hopf-tower reading: it fixes the factor count and two of the three matrix factors — more internal algebraic *shape* than the gauge group alone, which then follows from the inner automorphisms of the admitted $\mathcal{A}_F$ — while the third factor's real form, the generation multiplicity, and the Yukawa values remain unselected. Sources: `debug/door4_gauge_yukawa_boundary_memo.md`, `debug/door4b_inner_algebra_forcing_memo.md`, `debug/door4c_j_signtable_audit_memo.md`.

A geometric handle on the $n = 2$ fork (Door 4d, 2026-06-02). A second handle, structurally orthogonal to the combined-J sign table of Door 4c, is the *Division-Algebra-of-the-Sphere* (DAS) criterion: *at each Hopf rung n, the inner algebra is the natural associative real normed division algebra whose unit-norm sphere IS the rung's Hopf bundle total space S^{2n-1}.* The identifications are bit-exact at all three rungs: S^1 is the unit-norm sphere of $\mathbb{C}$ under complex multiplication (closure 1.1×10^{-16}); S^3 is the unit-norm sphere of $\mathbb{H}$ under quaternion multiplication and is bit-exactly isomorphic to $\mathrm{Sp}(1) = \mathrm{SU}(2)$ via the standard $\mathbb{H} \to M_2(\mathbb{C})$ embedding (closure 2.2×10^{-16}, SU(2)-isomorphism error 3.3×10^{-16}); and by Hurwitz's classification of finite-dimensional real normed division algebras (only $\mathbb{R}, \mathbb{C}, \mathbb{H}$ are associative, with unit spheres S^0, S^1, S^3 respectively), S^5 has no associative-division-algebra realization, so the inner algebra at $n = 3$ falls back to a matrix algebra. DAS *agrees* with Door 4b's elimination forcings at $n = 1$ ($\mathbb{C}$) and $n = 3$ ($M_3(\mathbb{C})$), and *closes* the $n = 2$ fork by selecting $\mathbb{H}$ — the direct realization with $\mathrm{U}(\mathbb{H}) = \mathrm{Sp}(1) = \mathrm{SU}(2)$ on the 3-dimensional S^3, in contrast to the $M_2(\mathbb{C})$ realization whose unit-norm sphere is the 4-dimensional U(2) with SU(2) reached only via the U(2)/U(1) unimodularity quotient. The DAS argument never invokes any J or sign-table data; it is a purely geometric criterion identifying the rung-n sphere with the unit-norm subset of the rung's associative division algebra. The verdict is *partial-door*: the existing Hopf-rung reading (Door 4b) extracts only the gauge group from the Hopf-rung sphere; adopting DAS as a forcing requires the additional structural theorem that the AC tensor product $\mathcal{H} = \mathcal{H}_{\mathrm{GV}} \otimes \mathcal{H}_F$ transfers the rung-n sphere's division-algebra structure to a constraint on $\mathcal{H}_F$'s inner algebra. This theorem is sprint-reachable and paper-level; if it closes, the inner-algebra status upgrades from "mostly forced with the $\mathbb{H}$-vs-$M_2(\mathbb{C})$ fork closed by literature import (CCM second-order condition)" to "mostly forced with the fork closed by a GeoVac-coherent geometric criterion," leaving only the generation count, the inner KO-dimension, and the Yukawa values outside the forcing. Source: `debug/door4d_division_algebra_sphere_memo.md`.

Falsifier A′ on DAS (Door 4e, 2026-06-02). The DAS handle introduced above is partial-door because the existing Hopf-rung reading (Door 4b) extracts only the gauge group from the Hopf-rung sphere. The natural test was whether any standard AC tensor-product compatibility condition fills the gap, i.e., distinguishes $\mathbb{H}$ from $M_2(\mathbb{C})$ as the $n = 2$ inner algebra. Three candidates were tested literally: (C1) SU(2)-equivariance under the Ad action $g \cdot m \cdot g^{-1}$; (C2) Hopf U(1)-equivariance under scalar phase $e^{i\theta} I$ acting from the right; (C3) principal-bundle compatibility with the Hopf U(1) fiber. All three returned *negative*: (C1) both $\mathbb{H}$ and $M_2(\mathbb{C})$ are SU(2)-Ad-invariant real $*$-subalgebras of $M_2(\mathbb{C})$ (the former is the unique minimal non-abelian such, the latter is the maximal; the distinction requires a minimality criterion that is itself extra input); (C2) every 2×2 complex matrix commutes with the scalar phase $e^{i\theta} I$ trivially, so the condition is vacuous; (C3) both candidate actions are principal-bundle compatible at machine precision by associativity of matrix multiplication and centrality of scalar phases. *No standard AC tensor-product compatibility condition forces $\mathbb{H}$ over $M_2(\mathbb{C})$.* The substantive content of the test is not the negative outcome (which was the expected one) but the *import comparison* it enables: the standard Chamseddine–Connes–Marcolli derivation of $\mathbb{H}$ at the $n = 2$ factor invokes *three* axioms beyond the

bare Connes spectral-triple data (second-order condition with nonzero finite Dirac, complex chiral fermion representation, and the $2N^2 = 32$ dimension count), whereas DAS invokes *one* (the rung-n sphere's division-algebra structure is inherited by the inner algebra). Both are imports relative to the bare axioms; DAS is leaner. The path to upgrading partial-door to door is therefore not a theorem about existing CCM machinery but a paper-level structural axiom: adopting "the inner algebra at rung n is the $*$-algebra whose unit group IS the rung-n sphere S^{2n-1} as a Lie group, when one exists; else fallback $M_n(\mathbb{C})$." This reads the Hopf-rung sphere fully (Lie-group structure plus topology) where the existing construction reads it partially (gauge group only), and gives $\mathbb{C}$ at $n = 1$ (via $S^1 = \mathrm{U}(1) = \mathrm{U}(\mathbb{C})$), $\mathbb{H}$ at $n = 2$ (via $S^3 = \mathrm{Sp}(1) = \mathrm{U}(\mathbb{H})$), and $M_3(\mathbb{C})$ at $n = 3$ (since S^5 is not a Lie group at all, by Adams 1958). Source: `debug/door4e_falsifier_A_prime_memo.md`.

Closure of the Door 4 series (Door 4f, 2026-06-02; net axiom savings audited). The natural next test was whether adopting Upgrade B introduces any new downstream constraint on existing GeoVac observables. Twelve sub-tests were run: inner KO-dimension (T1), chirality grading γ_F (T2), Yukawa eigenvalues (T3), generation count (T4), Connes order-zero (T5), order-one (T6), and second-order (T7) conditions, bosonic spectral-action gauge-sector coefficient (T8), Higgs vacuum manifold (T9), cross-rung Yukawa coupling (T10), extensibility to rung $n \geq 4$ via Adams 1958 (T11), and consistency with the Bertrand $\times$ Hopf-tower reading of Door 4b (T12). *All twelve* return either SILENT (Upgrade B unconstrained on the target), COMPATIBLE (the pre-existing constraint passes for $\mathbb{H}$, the mechanism unaffected), IDENTICAL to CCM (same observable from either $\mathbb{H}$ or $M_2(\mathbb{C})$ post-unimodularity), or a CLEAN extensibility/strengthening rule (T11, T12). Zero sub-tests introduce a new constraint; zero contradict any existing GeoVac result. Upgrade B is therefore a *clean reformulation* of the CCM $\mathbb{H}$-selection with no new predictive content.

Honest axiom-savings audit. The naive comparison in the preceding paragraph (CCM 3 axioms vs DAS 1 axiom) over-counts. Sub-test T7 (Connes second-order condition) shows that the second-order condition is *still* required in the Upgrade-B formulation, because it constrains the off-diagonal Yukawa structure of D_F (which $\mathbb{H}$-selection alone does not determine). Upgrade B therefore replaces *two* of CCM's three $\mathbb{H}$-selection axioms — the complex chiral fermion representation requirement and the $2N^2 = 32$ dimension count — with the single sphere-Lie-group axiom; the second-order condition persists in both formulations. Net axiom savings: *one*. The Door 4 series is structurally complete at **partial-door (final)**. The remaining open questions are paper-level (whether to adopt Upgrade B's axiom-minimality advantage as the working construction principle) and the deep wall (force N_{gen} and the inner KO-dim from a packing principle; no known handle; deferred). Source: `debug/door4f_falsifier_A_double_prime_memo.md`.

The deep wall is a relative necessity, not a gap (Direction 2 scoping, 2026-06-03). The "deep wall" named above — forcing N_{gen} and the inner KO-dimension from a packing principle — is sharper than an open gap: relative to the packing construction of Paper 0 as presently formulated, it is a *necessity*. Two established facts compose. First, the Door 4 seam: the inner Yukawa/real-structure data lives in a ring (and a set of $(\mathcal{H}_F, D_F, J_F, \gamma_F)$ data) sharing no generator with any forced outer ring (an Observation-level reading, §8.4 below; "provably disjoint" withdrawn 2026-09-03). Second, Paper 0 §VII: the packing construction is *kinematic* — it emits the label set (n, ℓ, m, s) and the graph topology, and emits no operator, no real structure, no coupling, and no Hilbert-space multiplicity. But N_{gen} is exactly a Hilbert-space multiplicity (the factor in $\mathcal{H}_F = \mathbb{C}^{N_{\mathrm{gen}}} \otimes \mathcal{H}_F^{(1)}$, invisible to every algebra- and automorphism-level mechanism), and the inner KO-dimension is exactly a (J_F, γ_F) datum — both of the type the packing construction does not produce. The packing axiom therefore cannot constrain either: the residue is packing-inaccessible by composition

of these two facts. The single logically-open escape is a *packing-native multiplicity* — a datum the construction emits on the outer labels that would map to a factor-N_{gen} tripling of $\mathcal{H}_F$. It is empty: Paper 0's entire multiplicity inventory is the $2\ell + 1$ angular slots (odd, and *growing* with ℓ, hence not a generation-universal multiplicity) together with a single $\mathbb{Z}_2$ already consumed by spin; no constant factor of three appears anywhere. This promotes the structural-skeleton scope from an observation to a *relative theorem of necessity* for the N_{gen}/inner-KO-dimension residue, and gives a single structural reason the H1, W3, and Koide selection attempts failed: all probed inside the almost-commutative category, on the side of the seam where the disjointness forbids selection and the packing axiom has no reach. The qualifier "relative" is load-bearing: the statement forecloses the residue for the *current* packing construction, not for every conceivable discrete one. Lifting it would require a *sibling axiom* that emits fiber multiplicities — a new primitive Paper 0 does not contain — recorded as the open (long-range) direction with no present handle. The inner *algebra shape* $\mathbb{C} \oplus \mathbb{H} \oplus M_3(\mathbb{C})$ is the one inner datum the construction does reach (Door 4b/4d, via the Hopf tower and Hurwitz's classification); the residue it cannot reach is precisely the multiplicity- and real-structure-valued data. Source: `debug/seam_packing_scoping_memo.md`.

The Forced-Count Theorem (synthesis, 2026-06-03). The eight preceding paragraphs (Sprint H1 through Door 4f and the Direction 2 scoping) compose into a single Bertrand-shaped statement that crystallizes the structural content already proved across the sub-sprints. **[SYMBOLIC + MEASURED]** Stated as a theorem it makes explicit what the distributed prose shows in pieces: the framework forces the *dimension* of the D_F moduli space, but does not (and at the current packing primitive cannot) force the points within it.

Theorem 7 (Forced D_F moduli dimension)**.** *Let $\mathcal{T} = (\mathcal{A}_{\text{GV}} \otimes \mathcal{A}_F,\ \mathcal{H}_{\text{GV}} \otimes \mathcal{H}_F,\ D_{\text{GV}} \otimes 1 + \gamma_{\text{GV}} \otimes D_F,\ J,\ \gamma)$ be the GeoVac almost-commutative spectral triple of Sec. 3 (the graded form is displayed for reference; the modules build the ungraded sum $D_{\text{GV}} \otimes 1 + 1 \otimes D_F$, §8.2; the count below concerns D_F alone and is the same for both), assuming:*

1. *The Bertrand-with-complex-Hopf-tower reading $n \leq 3$ (Door 4b; an organizing reading, not an argument of Sec. 8.1), with the Upgrade B sphere-Lie-group axiom of Door 4e (an adopted axiom), selecting $\mathcal{A}_F = \mathbb{C} \oplus \mathbb{H} \oplus M_3(\mathbb{C})$.*
2. *The canonical Chamseddine–Connes–Marcolli SM representation of $\mathcal{A}_F$ on $\mathcal{H}_F$, with generation multiplicity N_{gen} imported as an external input (Direction 2: not packing-derived).*
3. *The standard real-spectral-triple axioms: Hermiticity, KO-dim-6 chirality grading $\{\gamma_F, D_F\} = 0$, J-reality $[J_F, D_F] = 0$, and Connes order-one.*

Then the space of admissible finite Dirac operators D_F has real dimension forced by these inputs alone, with the full-axiom reduction chain at $N_{\text{gen}} = 1$ (on the full 32-dimensional finite geometry):

$$\dim_{\mathbb{R}} \mathcal{M}(D_F) :\ 2048 \xrightarrow{\text{Herm.}} 1024 \xrightarrow{\{\gamma_F, D_F\}=0} 512 \xrightarrow{[J_F, D_F]=0} 272 \xrightarrow{\textit{order-one}} 32. \tag{7}$$

[SYMBOLIC + MEASURED] *The endpoint 32 is computed with a linear $*$-representation of $\mathcal{A}_F$ on $\mathbb{C}^{32}$ (particles colour-blind, $M_3(\mathbb{C})$ acting on antiquarks; three independent linear-algebra routes agree, singular-value gap 12.4 vs 10^{-13}; `tests/test_trunk_qa_forced_count_moduli.py`). Its projection onto the 16-dimensional matter sector has rank 16 — the* full *lepton 2×2 and quark-flavour $2 \times 2 \otimes \mathbb{1}_3$ blocks, of which the four colour-diagonal Yukawas y_ν, y_e, y_u, y_d with their $L \leftrightarrow R$ placements are 4 complex, i.e.* half *(corrected 2026-09-04: the statement named only those*

four, which the proof below already calls false and which is 8 *real, not* 16*) and onto the matter–antimatter (Majorana) block rank* 16. Correction (2026-09-03): *the endpoint printed before,* 260 *(with a matter-sector chain* $512 \to 256 \to 128 \to 128$*), was an artefact of a degenerate algebra sample — eighteen of the twenty-four basis elements were identically zero and the remaining six had no quark block, because the module's quark action* $\mathrm{kron}(ew, m)$ *vanished at* $ew = 0$ *or* $m = 0$ *— so the order-one condition added almost no constraint; every "*128*" quoted earlier in this paper as a matter-sector moduli count refers to that degenerate sample. The physical-slice dimension under restriction to diagonal Yukawa* $Y = \mathrm{diag}(y_\nu, y_e, y_u, y_d)$ *is* 8 *per generation. The* N_{gen}*-generation count with full generation-mixing Yukawa matrices is not re-derived here (the* $128 \cdot N_{\mathrm{gen}}^2$ *stated before 2026-09-03 was the degenerate-sample figure).*

Proof sketch. The full-axiom chain in eq. (7) is the direct parameter-count of the constraint solution space (`tests/test_trunk_qa_forced_count_moduli.py`) on the full 32-dimensional finite geometry at $n_{\max} = 2$, $N_{\mathrm{gen}} = 1$: Hermiticity halves the free parameters; $\{\gamma_F, D_F\} = 0$ kills the chirality-diagonal blocks; J-reality identifies the matter and antimatter sectors ($512 \to 272$); the order-one condition, imposed with a linear $*$-representation of $\mathcal{A}_F$, removes 240 further parameters ($272 \to 32$): with $M_3(\mathbb{C})$ acting on the antiquark colour index, $[[D_F, a], JbJ^{-1}] = 0$ forces the quark Yukawa blocks to be colour-diagonal and colour-independent, and strongly constrains the matter→antimatter block P (it does *not* annihilate it — the rank-16 real Majorana slice survives, as the next paragraph details).

What survives, measured (corrected 2026-09-03). The sketch printed here previously said the condition "kills every lepton–quark and off-flavour entry" and described the rank-16 matter projection as the four colour-diagonal Yukawas with their $L \leftrightarrow R$ placements. Both descriptions are false, and the support pattern of the solution space is the theorem's actual content. The matter block (rank 16 real = 8 complex) is *full* on the lepton 2×2 — including $\nu_L \to e_R$ and $e_L \to \nu_R$ — and full on the quark flavour $2 \times 2 \otimes \mathbb{1}_3$, including $u_L^c \leftrightarrow d_R^c$; the four named Yukawas are 4 complex, i.e. *half* of it. The Majorana block (rank 16 real) carries 2 complex ν_R entries and *six* complex $\mathrm{lepton}_R \leftrightarrow$ antiquark-u_R entries — so twelve of its sixteen real moduli are precisely the lepton–quark operators the old sentence said were killed. The endpoint 32 and the two rank-16 projections are unchanged. Each constructed-triple axiom-consistency check passes at $n_{\max} \in \{2, 3\}$ in module `geovac/standard_model_triple.py` (backed by `tests/test_standard_model_triple.py`, 47 tests). *QA correction (2026-06-14):* those 47 tests verify the axiom consistency of one constructed D_F, *not* the moduli-dimension counts of the chain; the counts above are the direct enumeration of `tests/test_trunk_qa_forced_count_moduli.py`. *Statement relabelling (2026-09-01):* eq. (7) originally printed the mixed chain $1024 \to 512 \to 128 \to 128$, which matched neither the full-axiom nor the matter-sector convention; it now states the full-axiom count with the matter-sector restriction labelled explicitly, applying the relabelling this QA correction directed. (No N_{gen} scaling is claimed: as the statement says, the N_{gen}-generation count is not re-derived.) The status of the inputs: (1) rests on the Door 4b Hopf-tower reading (which fixes the factor count) plus the Upgrade B sphere-Lie-group axiom of Door 4e, which selects $\mathbb{H}$ at $n = 2$ as an *adopted axiom*, not a derivation (the Door 4 record is partial-door; Door 4c found $\mathbb{H}$ admitted but not forced); (3) is the standard real-spectral-triple axiom set; (2) is the one external input, recorded by the Direction 2 paragraph as packing-unreachable in the current Paper 0 construction. □

Corollary 5 (Bertrand analogy)**.** *Theorem 7 is the inner-factor analogue of Bertrand's theorem on closed-orbit central forces. Bertrand forces the* form *of the central potential (*$-k/r$ *or* kr^2*) without forcing the coupling* k*; Theorem 7 forces the* dimension *of the* D_F *moduli space without forcing the* D_F*-entries themselves. The two seams are structurally identical: the framework determines what the moduli space* is*, not where in it the realised system sits.*

Remark 49 (Paired with the Door 4 negative half)**.** *The positive content of Theorem 7 sits beside two negative companions already recorded above:*

- ***Door 4 seam reading (Yukawa-value wall):*** *the η-trivialization plus tensor factorization plus the G3 commuting-$\mathbb{Z}_2$ result place the D_F-entries inside $\mathcal{M}(D_F)$ in a Dirichlet ring sharing no generator with any forced outer ring, so the framework does not derive them from the forced rings (an Observation-level reading).* ***[PANEL-VERIFIED]*** *Sprint Yukawa-PSLQ (2026-06-03) added empirical confirmation: a 162-cell sweep of the nine measured Yukawa values against the forced inner-factor pure-Tate basis $\mathbb{Q}[\pi, \pi^{-1}]$ (basis sharpened by the η-trivialization audit ruling out M_3/Dirichlet L on the inner factor) returned zero hits at $M \leq 10^3$, at both MS-bar M_Z and MS-bar $\mu = 2 \times 10^{16}$ GeV, across three transforms $(y_f,\ y_f^2,\ \log y_f)$. Source:* `debug/sprint_yukawa_pslq_memo.md`.
- ***Direction 2 packing-reach theorem:*** *N_{gen} and the inner KO-dimension are packing-unreachable in the current Paper 0 construction (preceding paragraph, source* `debug/seam_packing_scoping_m`

[CONDITIONAL] *Theorem 7 thus crystallises the* full *forced/free seam at the level of D_F: the forced data is the moduli space and its dimension; the free data is (i) the point in the moduli space (Yukawa values, Door 4 wall), (ii) the multiplicity factor (N_{gen}, Direction 2), and (iii) the real-structure signature (inner KO-dimension, Direction 2). All three free data are calibration inputs in the structural-skeleton-scope sense (§1.5 of the project guidelines).*

Theorem 8 (N_{gen} non-selection under canonical CCM representation)**.** ***[PROOF-BY-ARGUMENT]*** *(Here and in the non-selection results below, "a morphism in $\mathcal{A}$" is informal: $\mathcal{A}$ is a list of axioms, and "no morphism selects X" means that the axioms are satisfied for every admissible value of X.) Let $\mathcal{A}$ denote the framework's current axiom set {Paper 0 packing axiom; standard real-spectral-triple axioms (Hermiticity, J-reality, KO-dim chirality, order-one); Hopf-tower truncation (Hurwitz); Bertrand's classical theorem; Upgrade B (sphere-Lie-group axiom)}, and fix the canonical Chamseddine–Connes SM representation of $\mathcal{A}_F = \mathbb{C} \oplus \mathbb{H} \oplus M_3(\mathbb{C})$ on the inner Hilbert space.*

For every integer $N \geq 1$, let $\mathcal{T}_N$ denote the GeoVac spectral triple with inner Hilbert-space multiplicity N (i.e., $\dim \mathcal{H}_F = 32N$ under the canonical rep). Then $\mathcal{T}_N$ satisfies every axiom in $\mathcal{A}$. Consequently, $\mathcal{A}$ contains no morphism that selects a specific value of N.

Proof sketch. Each axiom in $\mathcal{A}$ is independent of N:

- The Paper 0 packing axiom outputs (n, l, m, m_s) labels for a single S^3 outer graph; N is not in this output.
- The standard real-spectral-triple axioms constrain the relations (D, J, γ) on $\mathcal{H} = \mathcal{H}_{\text{GV}} \otimes \mathcal{H}_F$. $\dim \mathcal{H}_F$ scales linearly with N, but Hermiticity, J-reality, KO-dim chirality, and order-one are structural relations independent of multiplicity.
- Hopf-tower truncation (Hurwitz, Doors 4b/4d/4e) operates on $\mathcal{A}_F$; the algebra $\mathcal{A}_F = \mathbb{C} \oplus \mathbb{H} \oplus M_3(\mathbb{C})$ is fixed independent of N.
- Bertrand's classical theorem selects the outer manifold S^3; N-independent.
- Upgrade B constrains the inner-algebra structure via the sphere-Lie-group identification of Door 4e; N-independent.
- The canonical CCM SM representation distributes 32 fermion degrees of freedom per generation across the three algebra factors; the multiplicity N enters as an external Hilbert-space multiplicity rather than as a constraint emergent from the rep structure.

The Read 2 scoping (`debug/sprint_read2_n_gen_scoping_memo.md`, 2026-06-03) provides three independent obstructions to any candidate Hopf-tower-to-representation shortcut that would correlate N with the algebra factor count under canonical representation: the SM fermion DOF count per generation distributes across all three algebra factors; the Yukawa-Higgs cross-generation coupling structure required for CKM/PMNS phenomenology prevents algebra-factor reorganisation; and the alternative "3 factors = 3 generations" assignment gives 12 DOF per gen versus the 32 DOF of the SM. Each obstruction independently closes the candidate shortcut.

Therefore $\mathcal{T}_N$ is a valid GeoVac spectral triple for every $N \geq 1$ under canonical representation; no axiom in $\mathcal{A}$ has been exhibited that selects a specific value. □ □

Remark 50 (Scope and the unconditional question)**.** *Theorem 8 is conditional on the canonical CCM SM representation choice. The unconditional question — whether any SM-phenomenologically-consistent representation of $\mathcal{A}_F$ on $\mathcal{H}_F$ could correlate N with the algebra factor count under an alternative rep — remains the open long-range structural- research target named by the Read 2 sharpest falsifier: exhibit a representation in which the* 32 *fermion DOF per generation are distributed across the three algebra factors in a way that makes the three generations the three rep blocks of the algebra, and the cross-generation Yukawa-Higgs coupling (CKM/PMNS) is naturally encoded as the algebra cross-factor coupling. Absent such a construction, Theorem 8 stands as the conditional impossibility result corresponding to the Read 2 NO-GO scoping verdict.*

The Theorem identifies N_{gen} as Class 1 calibration data (external input on which the framework's outer-factor mechanism is structurally silent), formalising the Direction 2 packing-reach and Read 2 representation-level NO-GO scopings into a single structural statement. It is the multiplicity-side analogue of the H1 Yukawa non-selection theorem: both are non-selection results for inner-factor content, both conditional on the canonical rep, and together with the Forced-Count Theorem (7) they characterise the full free-side content of the inner-factor structural-skeleton boundary at the canonical-representation level.

[CONDITIONAL] *Paper 57 §6.1 (the Forced/Free Seam observations paper) records this theorem as the C3 closure of the empirical Direction 2 / Read 2 NO-GO scopings, and as one of three proof-by-argument non-selection results in the corpus (alongside H1 Yukawa non-selection and the Forced-Count Theorem).*

Theorem 9 (Inner KO-dim is external calibration data under canonical-rep convention)**.** ***[PROOF-BY-ARGUMENT]*** *(The combined triple's own sign data is the measured $(-,+,+)$ of §8.2; no additive KO rule applies to it, since for the graded product Dabrowski–Dossena's Table 5 admits no consistent ε' with a KO-6 finite factor. The statement below concerns only the* inner *factor's label.) Let $\mathcal{A}$ denote the framework's current axiom set (as defined in* Theorem 8*). The Paper 0 packing axiom is kinematic (Paper 0 § VII.B "What the construction does not provide"): it outputs the labels (n, l, m, m_s) and the S^3 graph topology, but neither a Dirac operator, nor a real structure, nor a chirality grading. The inner KO-dimensional signature is a property of the real-structure data (D_F, J_F, γ_F), none of which is in the packing axiom's output.*

The remaining axioms in $\mathcal{A}$ (standard real-spectral-triple axioms, Hopf-tower truncation, Bertrand's classical theorem, Upgrade B) constrain the consistency of (D_F, J_F, γ_F) with the outer GeoVac triple's signs (KO-dimension 3 *for the continuum factor; sign triple $(-,+,+)$ and no label at finite cutoff), but do not uniquely select an inner KO-dim signature. Different choices of (D_F, J_F, γ_F) that are consistent with these constraints yield different inner KO-dimensional signatures, each producing a valid GeoVac spectral triple. The canonical Chamseddine–Connes SM representation (inner KO-dim* 6*) is one such choice, supplied as external input to the framework.*

Proof sketch. The argument is a direct composition of two established facts:

Fact 1 (Paper 0 §VII.B, explicit scope statement). The packing construction is *kinematic*: it produces labels and graph topology, and explicitly does *not* produce operators, real structures, couplings, or fiber-Hilbert-space multiplicities.

Fact 2 (Door 4f T1, Door 4b Q3). The inner KO-dimensional signature is defined as a property of the real-structure data (D_F, J_F, γ_F). The chirality-grading axiom $\{\gamma_F, D_F\} = 0$ and the J_F-reality conditions $J_F^2 = \pm\mathbf{1}$, $J_F D_F = \pm D_F J_F$, $J_F \gamma_F = \pm \gamma_F J_F$ specify the residue classes mod 8 that are consistent, but do not uniquely select one in the absence of additional constraints (such as the phenomenological constraints imposed by SM fermion doubling and Majorana neutrino-mass structure).

Composition. $\mathcal{A}$ produces no (D_F, J_F, γ_F) data autonomously (Fact 1), and the inner KO-dim is a function of that data (Fact 2). Therefore $\mathcal{A}$ does not autonomously select an inner KO-dim signature. The canonical CCM SM rep supplies a specific (D_F, J_F, γ_F) triple (KO-dim 6); alternative specifications consistent with the outer signs and the standard real-spectral-triple axioms produce alternative KO-dim signatures, each consistent with $\mathcal{A}$. □ □

Remark 51 (The full inner-factor structural-skeleton boundary)**.** *Theorems 7, the H1 Yukawa non-selection (Theorem 6), 8, and 9 together characterise the full free-side content of the inner-factor structural-skeleton boundary at the canonical-representation level:*

- ***Forced data:*** *the moduli space $\mathcal{M}(D_F)$ and its dimension (full-axiom:* 32 *at $N_{\text{gen}} = 1$, matter-sector projection rank* 16*,* 8 *per generation on the diagonal-Yukawa slice; the $128\, N_{\text{gen}}^2$ matter count quoted before 2026-09-03 was a degenerate-sample artefact — see Theorem 7).*
- ***Free data:*** *(i) the point in $\mathcal{M}(D_F)$ (Yukawa values — H1 non-selection), (ii) the Hilbert-space multiplicity (N_{gen} — Theorem 8), (iii) the real-structure signature (inner KO-dim — Theorem 9).*

These four theorems together formalise the structural-skeleton scope pattern at the inner-factor level for the GeoVac framework's current axiom set. Each is conditional on the canonical CCM SM representation. The unconditional versions — whether alternative phenomenologically-consistent representations could correlate any of the three free data with framework-derivable structure — remain long-range structural-research targets; Read 2 sharpest falsifier names the candidate construction for N_{gen}, and analogous targets exist for KO-dim (a non-canonical rep that admits a different KO-dim while retaining SM phenomenology) and for Yukawa values (a packing-derivable structure on $\mathcal{A}_F$ that selects within the diagonal slice).

Paper 57 §6.3 (Forced/Free Seam observations paper) records the trio plus Forced-Count Theorem as the C3 closure of the inner-factor non-selection arc.

Definition 4 (Multi-cutoff spectral action structure)**.** *A multi-cutoff spectral action on a real spectral triple $\mathcal{T} = (\mathcal{A}, \mathcal{H}, D)$ is a functional of the form*

$$S^{\text{multi}}(D, \{\Lambda_i\}_{i=1}^k, \{f_i\}_{i=1}^k) \;=\; \sum_{i=1}^{k} c_i \,\operatorname{Tr} f_i(D/\Lambda_i)$$

or more generally any functional of D depending on $k \geq 2$ independent energy scales $\Lambda_1, \ldots, \Lambda_k$ in a manner not reducible to a function of a single scale by reparametrization.

Theorem 10 (Single-cutoff spectral action)**.** ***[PROOF-BY-ARGUMENT]*** *The framework's current axiom set $\mathcal{A}$ (as defined in* Theorem 8*) specifies the Chamseddine–Connes spectral action $S(D, \Lambda) = \operatorname{Tr} f(D/\Lambda)$ as a single-cutoff functional in the sense of* Definition 4*. No morphism derivable from $\mathcal{A}$ produces a multi-cutoff spectral action structure.*

Proof sketch. The argument is a structural inspection of each axiom in $\mathcal{A}$:

1. **Spectral action axiom (CCM).** The standard CC spectral action $S(D, \Lambda) = \mathrm{Tr}\, f(D/\Lambda)$ specifies a single-cutoff functional by stipulation. The argument D/Λ is a dimensionless ratio against a single energy scale.
2. **Paper 0 packing axiom + standard real-spectral-triple axioms + Hopf-tower truncation + Bertrand + Upgrade B.** These axioms are algebraic, kinematic, and topological: they specify $(\mathcal{A}_F, \mathcal{H}_F, D_F, J_F, \gamma_F)$ and the outer manifold geometry, but introduce no energy-scale data.
3. **Inner fluctuations $D \to D + A$ for $A = a[D, b]$.** These preserve the single-Dirac structure of the underlying spectral triple. The fluctuated operator $D' = D + A$ enters the spectral action as $S(D', \Lambda)$, still a single-cutoff functional.
4. **Tensor products $\mathcal{T}_1 \otimes \mathcal{T}_2$ (Paper 39).** These produce a new spectral triple with combined Dirac $D_1 \otimes 1 + 1 \otimes D_2$, on which the spectral action $S(D_1 \otimes 1 + 1 \otimes D_2, \Lambda)$ remains single-cutoff with the new combined D. No second cutoff is introduced by the tensor-product construction.

No axiom in $\mathcal{A}$ introduces a second independent energy scale, and no morphism derivable from $\mathcal{A}$ produces a functional of the form $S^{\mathrm{multi}}(D, \Lambda_1, \Lambda_2, \ldots, \Lambda_k, \ldots)$ with $k \geq 2$ independent scales. Therefore the framework's spectral action machinery is single-cutoff. □ □

Corollary 6 (Multi-loop renormalization counterterms are external calibration data)**.** *Multi-loop QED renormalization counterterms (wavefunction renormalization Z_2, mass counterterm δm, vertex counterterm Z_1, etc.) require a multi-cutoff structure (Λ_{UV}, μ_R) where Λ_{UV} is the ultraviolet cutoff for the bare integrand and μ_R is the renormalization scale at which finite subtraction is performed. By* Theorem 10*, $\mathcal{A}$ contains no morphism producing such a structure. Therefore the multi-loop QED counterterms are external calibration data, supplied to the framework rather than derived from it.*

Consequently the following catalogue entries from Paper 57 [56] are classified as multi-cutoff renormalization calibration data:

- ***E7:*** *two-loop self-energy counterterms $Z_2, \delta m$.*
- ***E8:*** *multi-loop QED corrections beyond the one-loop LS-7 closure.*
- ***G3:*** *HF-5 multi-loop QED on hyperfine.*
- ***G4:*** *LS-8a two-loop SE renormalization (identical to E7 under the cross-domain unity of Paper 57 §4).*

The LS-8a sprint (May 2026) empirically confirmed this structural verdict: the framework's iterated Chamseddine–Connes spectral action reproduces the two-loop UV-divergent integrand at the right prefactor, sign, and divergence rate $\sim N^{3.79}$ (Paper 36 LS-8a), but cannot autonomously generate the $Z_2, \delta m$ counterterms required for finite multi-loop closure. Theorem 10 formalises the structural reason for the LS-8a wall: single-cutoff machinery cannot produce multi-cutoff observables.

Remark 52 (Honest scope)**.** *Theorem 10 is a proof-by-argument result within the framework's current axiom set $\mathcal{A}$. Two scope clarifications:*

(i) The theorem does not claim "no multi-cutoff structure exists." *It claims that no morphism derivable from $\mathcal{A}$* as currently stated *produces a multi-cutoff structure. A sibling axiom*

*adding renormalization-group flow machinery would extend the axiom set and could in principle produce multi-cutoff observables. Such a sibling axiom is the analog of the N_{gen} sibling-axiom direction (*Theorem 8 *unconditional version): a logically-possible escape, named here, that requires new primitive content beyond $\mathcal{A}$.*

(ii) The theorem does not address the spatial-composition sub-sector of the multi-focal-composition wall. *The catalogue entries G1 (HF-3 recoil cross-register $V_{eN}(\hat{r}_e, \hat{R}_N)$), G2 (HF-4 Zemach magnetization density), and G5 (W1e chemistry inner-region correlation) lie outside this theorem's scope: they are spatial × spatial composition walls, structurally distinct from the multi-cutoff renormalization sub-sector. **[OPEN]** A general multi-focal-composition wall theorem unifying both sub-sectors remains the named open structural-research target.*

Paper 57 §4 (cross-domain unity) and Paper 57 §6.3 (Sprint E7/E8 closure) record this theorem as the renormalization sub-sector closure of the multi-focal-composition wall.

Theorem 11 (Tensor-product radial coupling does not autonomously match physical two-body profiles)***. [PROOF-BY-ARGUMENT, with an empirical leg: the Pearson check of step (2.4) (***0.846**,** 0.796 ***at $n_{\max} = 2, 3$) is a measurement, so the argument covers only the structural part (2.1)–(2.3)]*** *Let $\mathcal{T}_a = (\mathcal{A}_a, \mathcal{H}_a, D_a^{\text{CH}})$ and $\mathcal{T}_b = (\mathcal{A}_b, \mathcal{H}_b, D_b^{\text{CH}})$ be two GeoVac spectral triples on the same outer manifold S^3 at distinct focal lengths $\lambda_a \neq \lambda_b$, equipped with the Camporesi–Higuchi spinor Dirac operators. The framework's natural tensor-product spectral triple $\mathcal{T}_a \otimes \mathcal{T}_b$ [43] produces composed two-body operators whose:*

1. ***Angular structure** is forced: Gaunt selection rules, m-conservation, and monopole structure are all autonomously reproduced by the composed algebra (Paper 54 [55] Thm 3).*
2. ***Radial coupling profile** is determined autonomously by the combined Dirac structure (the Clifford-doubled graded product of Paper 39, since the naive $D_a^{\text{CH}} \otimes 1 + \gamma_a \otimes D_b^{\text{CH}}$ is unrealisable with $\gamma_a = 0$ on S^3), but does not autonomously coincide with the physical 1/r Coulomb form, the dipole-magnetization form, or other specific two-body radial profiles required by empirical observables.*

The framework's axiom set $\mathcal{A}$ contains no morphism enforcing that the composed radial coupling profile matches a specific physical two-body observable; matching a given observable requires external specification.

Proof sketch. The two assertions of the theorem stand on independent structural content:

(1) Angular structure forcing. Paper 54 Thm 3 [55] establishes that the composed algebra structure on $\mathcal{A}_a \otimes \mathcal{A}_b$ acts on the two-body Hilbert space through Wigner 3j coefficients in a way that exactly reproduces the angular selection rules (Gaunt, m-conservation, monopole hierarchy) of physical two-body Coulomb-style interactions. This is a direct consequence of the SU(2) representation-theoretic structure of the angular momentum operators on S^3, which the framework inherits via the Fock projection (Paper 7) and the Hopf-tower truncation.

(2) Radial coupling non-reproduction. The structural reason is the non-commutativity of the Fock-projection conformal factor with the tensor-product construction:

1. Paper 7's chordal-distance identity maps continuum $-Z/r$ on $\mathbb{R}^3$ to the Laplace–Beltrami operator on the unit S^3 (whose harmonics the graph's nodes label; the graph Laplacian is not its discretisation and does not converge to it, Paper 7) via a focal-length- dependent stereographic conformal factor $c(\lambda)$. Two different focal lengths λ_a, λ_b give two different conformal factors c_a, c_b.

2. The tensor-product spectral triple $\mathcal{T}_a \otimes \mathcal{T}_b$ has continuum limit on the product manifold $S^3_a \times S^3_b$. Under Fock projection, this corresponds to a six-dimensional space $\mathbb{R}^3_a \times \mathbb{R}^3_b$ parametrised by two independent position vectors, with combined conformal factor $c_{ab} \neq c_a \cdot c_b$ in general.

3. Physical two-body observables ($V_{eN}(\hat{r}_e, \hat{R}_N)$, Zemach magnetization, chemistry inner-region $V_{ee}(r_{12})$) are functions of the *relative* coordinate $\vec{r}_a - \vec{r}_b$ on $\mathbb{R}^3$ (three dimensions, not six). The reduction from the six-dimensional product manifold to the three-dimensional relative manifold is not in the framework's axiom set.

4. Empirical verification: Sprint resolvent_two_body (2026-06-01, `debug/sprint_resolvent_two_body_memo.md`) tested four weightings of the resolvent $(D_a^{\mathrm{CH}})^{-2}$ against the physical 1/r Coulomb profile and found Pearson correlations of at most 0.846 ($n_{\max} = 2$) and 0.796 ($n_{\max} = 3$) for the best weighting (values after the 2026-08-29 ERI-evaluator fix), decreasing with $n_{\max}$. Paper 54 verified the angular selection rules but reported only partial radial matching (connected fraction 77% at $n_{\max} = 2$, 32% at $n_{\max} = 3$). Together these indicate that the framework's natural composition mechanisms (tensor product, resolvent, gauged spectral action) do not autonomously match physical two-body radial profiles.

No axiom in $\mathcal{A}$ supplies the relative-coordinate reduction from the six-dimensional product manifold to the three-dimensional relative manifold, nor enforces a specific composed radial profile on the tensor-product spectral triple. Therefore the framework specifies the angular structure of two-body operators autonomously (via the SU(2) representation theory of S^3) but does not autonomously match the radial structure of any specific physical two-body observable. □ □

Corollary 7 (Spatial-composition catalogue entries are calibration data)**.** *The following catalogue entries from Paper 57 [56] are classified as spatial-composition calibration data:*

- ***G1:*** *HF-3 recoil cross-register $V_{eN}(\hat{r}_e, \hat{R}_N)$ (Track NI cross-register, Sprint HF 2026-05-07);*
- ***G2:*** *HF-4 Zemach magnetization density (Sprint HF 2026-05-07);*
- ***G5:*** *W1e chemistry inner-region overattraction (Sprint W1c arc 2026-05-23).*

Each requires a specific physical radial coupling profile that Theorem 11 establishes is not autonomously produced by the framework's tensor-product composition mechanism. The angular content is forced; the radial content is external calibration.

Remark 53 (Multi-focal-composition wall bound by argument)**.** *Together with Theorem 10 + Corollary 6 (E7, E8, G3, G4), Theorem 11 + Corollary 7 (G1, G2, G5) close seven catalogue entries of the multi-focal-composition wall family (Paper 57 §4 cross-domain unity) at proof-by-argument level (Theorem 11 with an empirical leg) under two structurally distinct theorems:*

- ***Renormalization sub-sector*** *(E7, E8, G3, G4): closed by Theorem 10 — spectral action is single-cutoff; no multi-cutoff structure in $\mathcal{A}$.*
- ***Spatial-composition sub-sector*** *(G1, G2, G5): closed by Theorem 11 — tensor-product composition reproduces angular structure but not radial structure; no relative-coordinate-reduction morphism in $\mathcal{A}$.*

An eighth catalogue entry (G6) is the H1 Yukawa non-selection, closed by Theorem 8's companion result (Theorem 6). The entries are finer than the instances: all six independent multi-focal-composition wall instances of Paper 57 §4 (H1, LS-8a, HF-3/4/5, W1e; G6 = F1 is not independent) are now characterised at proof-by-argument level by one of the three theorems.

The multi-focal-composition wall is bound by argument at the canonical-representation level. Together with the inner-factor results (the Forced-Count Theorem 7, a dimension count, and the non-selection theorems 8, 9 and H1), the corpus now carries six proof-by-argument results covering the inner-factor structural-skeleton boundary, the multi-loop QED renormalization sub-sector, and the spatial- composition sub-sector.

Paper 57 §6.3 (Sprint G1/G2/G5 closure) records this completion.

Observation 3 (No single-mechanism generation of the combination rule $K = \pi(B + F - \Delta)$ has been found)**.** ***[OBSERVATION]*** *Twelve candidate mechanisms for single-principle generation of the empirical combination rule $K = \pi(B+F-\Delta)$ for α^{-1} (Paper 2) have been empirically eliminated (Paper 2 §IV.G; Phases 4B–4I, April 2026, nine mechanisms; Sprint A α-SP, 2026-04-18, Connes–Chamseddine direct spectral-action derivation; Sprint K-CC, 2026-05-03, three further mechanisms via PSLQ at 100 dps against a 244-element basis, a natural Λ search from S^3 spectral data, and a unified CC heat-kernel route blocked by the algebraic obstruction Theorem T9 of Paper 28). The three ingredients have separate spectral homes ($B = 42$ a finite Casimir trace, $F = \pi^2/6$ a Fock-degeneracy Dirichlet value, $\Delta = 1/40 = (g_3^{\mathrm{Dirac}})^{-1}$ an inverse Dirac mode degeneracy; Observation 1), and no single mechanism is currently exhibited that produces their combination with the weights $(+1, +1, -1)$ and the external π prefactor.*

Remark 54 (E6 scope: why this is an Observation, not a theorem)**.** *An earlier version of this section stated the above as a theorem with a two-fact structural proof.The proof does not stand. Its sub-ring partition assigned the rationals $B = 42$ and $\Delta = 1/40$ to different Mellin sub-rings as though the sub-rings were exclusive; but $\mathbb{Q}$ lies in both the M_1 ring $\mathbb{Q}[\pi, \pi^{-1}]$ and the M_2 ring $\bigoplus_k \pi^{2k}\mathbb{Q}$, so $B+F-\Delta$ lies wholly in the M_2 ring and the claimed mixing of sub-rings does not occur. Its first fact invoked the π-source case-exhaustion (Theorem 4) for* every *transcendental, whereas that theorem is [CONDITIONAL] and covers only the first fifteen of the twenty-eight Paper 34 projections; it also sits in tension with the remark that the spectral action engages M_1 and M_2 jointly (Sec. 8). What survives is the empirical record of twelve eliminations, recorded above as an Observation. It does not derive K, and it does not show that K cannot be generated by a single mechanism from some extended axiom set.*

Per the project's standing hard prohibition, the Paper 2 combination rule is labeled an Observation *(not a conjecture or derivation); nothing in this section promotes or demotes that label. Paper 2's status as Observations-folder paper (post 2026-05-02 curve-fit-audit migration) is unchanged. Catalogue entry E6 (the combination rule for α^{-1}) is classified as calibration data with no single mechanism found: the framework provides three structural homes for the ingredients but no principle for their combination.*

Theorem 12 (Cutoff function f and its Mellin moments are external calibration data)**.** ***[PROOF-BY-ARGUMENT]*** *The framework's axiom set $\mathcal{A}$ specifies the Chamseddine–Connes spectral action functional $S(D, \Lambda, f) = \mathrm{Tr}\, f(D/\Lambda)$ on a real spectral triple. The cutoff function f enters as a Schwartz-class test function on $[0, \infty)$ (Paper 51) but is not constrained by any axiom in $\mathcal{A}$. Consequently the Mellin moments*

$$\varphi(s) \;=\; \int_0^\infty x^{s-1} f(x)\, dx$$

are external calibration data: no morphism in $\mathcal{A}$ selects the moment values $\varphi(0), \varphi(1), \varphi(2), \ldots$.

The Wald two-term-exactness ⇒ pure Einstein ⇒ action-G ≡ entropy-G argument (Paper 51 §G7) forces relations *among the moments via the sector-wise Mellin moment map (tip ↔ $\varphi(0)$, Einstein–Hilbert ↔ $\varphi(1)$, Λ_{cc} ↔ $\varphi(2)$); these relations follow by argument within $\mathcal{A}$ (the Wald step itself is a theorem). The moment* values *are not constrained by $\mathcal{A}$.*

Corollary 8 (CC fine-tuning is external moment selection)**.** *The observed cosmological-constant problem in the framework's sector-wise Mellin moment formulation (Paper 51 §G4-5) requires*

$$\Lambda_{cc} \cdot G_{\text{eff}} \;\approx\; 36\pi \cdot \frac{\varphi(2)}{\varphi(1)^2} \;\approx\; 10^{-122}$$

(dimensionless), which is consistent with the framework's structural relations between G_N, Λ_{cc}, and S_{BH} (Wald, theorem-grade) provided $\varphi(2)/\varphi(1)^2 \approx 10^{-124}$ with $\varphi(0), \varphi(1) = O(1)$. No natural cutoff function f produces this fine-tuned moment ratio, and no axiom in $\mathcal{A}$ constrains the moment values to it.

Catalogue entries D5 (cutoff function moments $\varphi(0), \varphi(1), \varphi(2)$) and D6 (cosmological constant value Λ_{cc}) are external calibration data inheriting from the cutoff function selection.

Remark 55 (D5/D6 scope and the sibling-axiom direction)**.** *Theorem 12 formalises the precise location of the cosmological-constant problem in the GeoVac framework: the gravitational* relations *are forced (Wald, Paper 51 §G7); the cutoff-function* moments *are external. The CC problem in this framing is a moment-selection problem on the cutoff function f, structurally similar to the Higgs–Yukawa selection problem (*Theorem *H1) but on a different calibration object.*

A sibling axiom that constrains the cutoff function or its moments (e.g., a renormalisation-group flow axiom, a regulator-independence principle, an asymptotic-safety constraint) would extend $\mathcal{A}$ and could in principle select the moment values. None is currently exhibited; the long-range structural-research direction is named here as the natural analog of the inner-factor sibling-axiom directions.

Paper 57 §6.3 records this theorem as the gravity sector closure of the C-arc.

Remark 56 (Chemistry-side η-trivialization analog: open structural-research target)**.** *The SM-side η-trivialization theorem (Paper 18 `thm:eta_trivialization`) forces the inner-factor period ring to $M_1 \cup M_2$ on any Krajewski-class even finite spectral triple satisfying the chirality grading axiom $\{\gamma_F, D_F\} = 0$: for all odd k, $\mathrm{Tr}(D_F^k \cdot e^{-tD_F^2}) \equiv 0$, so M_3 (vertex-parity Hurwitz / Catalan G / Dirichlet L) vanishes on the inner factor. This is the structural reason F1 (Yukawa values) and F4–F7 (Higgs VEV, CKM, PMNS, neutrino masses) live outside the master Mellin engine's structural skeleton: the inner-factor Yukawa values occupy the M1/M2 slots, but their generators are tied to the couplings y_i and are not among those of the rings the framework's outer-factor mechanism generates (an Observation-level reading; the rings are not disjoint, since both contain $\mathbb{Q}$).*

The chemistry-side analog would be a structural property of the FrozenCore $Z_{\text{eff}}(r)$ pipeline that mirrors $\{\gamma_F, D_F\} = 0$ on the chemistry inner-factor side: a chirality-style grading on the Clementi–Raimondi $Z_{\text{eff}}(r)$ structure that forces a similar period-ring restriction on chemistry inner-factor data. Such a property is not currently exhibited. The most concrete candidate direction would identify a discrete grading on the FrozenCore radial profile pipeline (e.g., parity of multi-zeta exponents, partition of core orbital basis into chirality-graded blocks) that produces an η-trivialization-style theorem for chemistry inner-factor data.

Sprint-scale path is unclear: unlike the SM-side η-trivialization, which follows directly from a single axiom ($\{\gamma_F, D_F\} = 0$, established in Krajewski 1998 and canonically adopted in CCM 2010), the chemistry-side analog would need to identify the analog grading axiom from scratch. This is a substantial NCG-research target.

Catalogue entries H6 (Clementi–Raimondi Z_{eff} exponents) and H7 (multi-zeta coefficients in second-row valence basis) remain empirical-only pending this structural-research target's completion. The empirical-grade content (W1e period-class diagnostic, `debug/sprint_w1e_period_class_memo.md`,

2026-06-04) establishes that H6, H7 are not in $M_1 \cup M_2 \cup M_3$ at audit ceiling 100; the open question is whether they sit in a structurally distinct chemistry-side ring forced by a hypothetical analog axiom.

Paper 57 §6.3 records this remark as the named substantial structural-research target for the chemistry sector; the C-arc's sprint-scale run closes at v3.103.0 with this target named but not attempted.

Remark 57 (The C-arc terminal state)**.** *After Sprints C2 + C3 + F3 + E7/E8 + G1/G2/G5 + E6 + D5/D6 + Chemistry-Analog (all 2026-06-08, v3.97.0–v3.103.0), the corpus carries* seven *proof-by-argument results (the Forced-Count dimension count and six non-selection results) and one empirical non-selection Observation across the following sectors (the E6 item was demoted from theorem to Observation on 2026-09-28, Remark 54):*

***Inner-factor sector** (closed at canonical-rep level):*

1. *Forced-Count Theorem (7).*
2. *H1 Yukawa non-selection (Theorem 6).*
3. *N_{gen} non-selection (8).*
4. *Inner KO-dimension non-selection (9).*

***Multi-focal-composition wall** (bound by argument under two sub-sector theorems, one with an empirical leg):*

5. *Single-cutoff spectral action (10).*
6. *Spatial-composition radial wall (11).*

*α **combination rule sector** (single-mechanism generation not found; empirical, not closed):*

7. *No single-mechanism generation of $K = \pi(B + F - \Delta)$ found (Observation 3; twelve mechanisms eliminated).*

***Gravity sector** (cutoff function external-selection closed):*

8. *Cutoff function f and its moments are external (12).*

The chemistry-side η-trivialization analog (Remark 56) is the named substantial structural-research target; catalogue entries H6, H7 remain empirical-only pending its completion.

The remaining empirical-only catalogue entries (F4–F7 Higgs VEV / CKM / PMNS / neutrino mass values, structurally inheriting from F1 and F4; I-entries foundational calibration) are subsidiary residuals inheriting from theorems above (Higgs admission B7 + H1 Yukawa non-selection) or sitting outside the structural-skeleton scope (Born rule via Gleason).

The full forced/free seam (Paper 57) is therefore in a stable terminal state for the path-#1 vein C arc: seven proof-by-argument results and one empirical Observation across four sectors, one named substantial follow-on (chemistry η-trivialization analog), all subsidiary residuals accounted for.

Files. Scoping memo `debug/g4_cross_manifold_scoping_memo.md` ($\sim 5\,100$ words). G4a sprint memo `debug/sprint_g4a_connes_sm_memo.md`. H1 sprint memo `debug/h1_higgs_inner_fluctuation_memo.md`. Sprint C3 N_{gen} non-selection theorem memo `debug/sprint_c3_n_gen_non_selection_memo.md` (2026-06-08). G4b is recorded as an open structural question.

8.5 Frontier-of-field framing: GeoVac-internal walls vs. spectral-action open problems

[**OBSERVATION**] The five walls catalogued in the multi-focal-composition audit (`debug/multifocal_phase_a_s` May 2026) are not all GeoVac-specific. Reading them through the spectral-triple construction of Sec. 3 and the Marcolli–van Suijlekom lineage of Sec. 9, three are GeoVac-internal architectural gaps that are tooling-addressable, and two sit at the broader frontier of the spectral-action / NCG programs. We name them here for honesty and to fix the scope of the open questions of Sec. 10.

GeoVac-internal walls (W1a, W1b, W1c). [OPEN]

- **W1a (cross-register coordinate operator):** no two-body spatial operator $V(\hat{\mathbf{r}}_e, \hat{\mathbf{R}}_n)$ has been implemented across the nuclear and electronic registers of the composed nuclear–electronic Hamiltonian (Paper 23 [34] §VI; `geovac/nuclear/nuclear_electronic.py`). The framework couples the registers through spin-spin hyperfine and a classical finite-size shift but evaluates spatial operators at the classical proton position. This is an architectural gap, not a no-go theorem; the calibrated atomic-physics target is the Pachucki–Patkoš–Yerokhin recoil Hamiltonian (PRL 130, 023004, 2023), which is exact in the mass ratio at $(Z\alpha)^6$.
- **W1b (magnetization-distribution operator):** no operator on the proton register represents the Zemach radius r_Z as a Layer-2 magnetization focal length distinct from the charge radius r_p (Paper 34 [40] § III; Sprint HF-4, May 2026). Calibrated against Eides et al. *Phys. Lett. B* (2024). Architectural gap; QCD-internal magnetization density is input data, not GeoVac-internal.
- **W1c (cross-center frozen-core screening):** the `FrozenCore` $Z_{\text{eff}}(r)$ provides same-center screening only; cross-center screening of bare V_{ne} from the opposite nucleus is not implemented (Sprint 7 NaH/MgH_2 balanced FCI; project development-frontier log). Sprint 7b made partial progress for the spin-orbit splitting; the PES extension is mechanical once cross-center screening is wired through the multipole expansion. Architectural gap.

[**OBSERVATION**] These three walls are tooling-addressable inside the present framework: the missing operators exist in principle, are calibrated against published atomic-physics targets, and require no new mathematics in the broader NCG literature. Each is addressable with existing tooling, though each is a substantial program rather than a bounded sprint. Their existence does not undermine the spectral-triple framing of Sec. 3; they catalogue what has not yet been built rather than what cannot be.

Frontier-of-field walls (W2a, W2b). [OPEN]

- **W2a (multi-loop UV/IR composition):** the bare iterated Connes–Chamseddine spectral action on the Camporesi–Higuchi spectrum reproduces UV-divergent integrands of multi-loop QED (Sprint LS-8a, May 2026; Paper 36 [42] §VII) but cannot autonomously generate Z_2, δm, Z_3 counterterms for finite extraction at two loops. This is not GeoVac-specific. The Fathizadeh–Marcolli periods-and-motives analysis of Robertson–Walker spectral actions [7] partially addresses spectral-action coefficient structure on homogeneous compact backgrounds, and the subsequent Hekkelman–McDonald 2024 noncommutative integral approximation [15] provides a single-cutoff Szegő-style asymptotic theory in the Connes–van Suijlekom paradigm; neither closes the multi-cutoff matching question that renormalization counterterm generation requires. The published QFT archetype is the matched effective-field-theory chain (e.g. HQET → NRQED → pNRQED → SCET; Fontes and Szafron, JHEP

05 (2025) 171), where each matching step is at a single physical scale and imports $\overline{MS}$ counterterms by hand. No published spectral-action framework reproduces this autonomously. W2a is therefore an open question of the spectral-action program, of which GeoVac is one specific instance; it is not a defect of the present construction relative to the lineage.

- **W2b (cross-manifold spectral-triple composition):** the tensor product of two infinite metric spectral triples $(\mathcal{T}_{S^3}, d_{D_{S^3}}) \otimes (\mathcal{T}_{S^5}, d_{D_{S^5}})$, needed to compose the GeoVac Coulomb sector (Sec. 3) with the Bargmann–Segal HO sector of Paper 24 [35], has no published Latrémolière-propinquity convergence theorem. The almost-commutative case (one infinite Riemannian factor + one finite factor) is addressed by Latrémolière 2026 [18] (*Spectral continuity of almost commutative manifolds for the C^1 topology on Riemannian metrics*), in the sense of metric continuity of almost-commutative models $C(M) \otimes \mathfrak{B}$ as the metric varies; the two-infinite case is not handled in that paper (implicitly, by its construction), nor in the closely related Farsi–Latrémolière–Packer 2024 inductive-sequence work (Adv. Math. 437, 109442). This is the structural content of the G4b cross-manifold blocker (Sec. 8.4): not a GeoVac-specific limitation but a genuinely open NCG question. Closing it would itself be a substantial mathematical contribution, parallel to but separate from Paper 38's SU(2) state-space GH convergence theorem.

Forecast (without claiming closure). We do not claim W1a/W1c can be closed without surprises; the multi-λ Shibuya–Wulfman algebraic check that gates the W1a attack (multipole termination across mismatched exponents) is not in the published Avery-school literature and could fail to preserve Gaunt-style closure (Track 3 literature memo, May 2026, §1). W1b is bounded by external QCD input. W2a and W2b are open in the broader literature, not just in the GeoVac record. The honest forecast is that closing W1a/b/c is internal-effort plausible as a substantial program per wall; closing W2a or W2b would be a contribution to the spectral-action / NCG programs rather than to GeoVac alone, and we do not commit to it as a deliverable here.

Implication for the Marcolli–vS lineage placement. Sec. 9 read the present construction as a graph spectral triple in the lineage of [6, 10, 11]. The frontier-of-field framing above sharpens this: the lineage placement is correct, and the W2a/W2b walls are the lineage's own open problems, not GeoVac's separate concerns. This is consistent with Observation 4 (each structural ingredient has published precedent; what is not duplicated is the α observation: K matches α^{-1} to 4.8×10^{-7}, and the root of the cubic $1/\alpha = K - \alpha^2$ matches to 8.8×10^{-8}). The W1 walls remain GeoVac-side because they are about the architectural extension of the present triple to multi-particle real-space coupling, which the SM-style almost-commutative construction does not address.

8.6 Lorentzian transfer audit and the master Mellin engine under signature extension (Sprint L0, May 2026)

The case-exhaustion theorem (Theorem 4, **[CONDITIONAL]**, first fifteen Paper 34 projections) classifies every π in such a chain as engaging one of three sub-mechanisms of a master Mellin engine $\mathcal{M}[\mathrm{Tr}(D^k\, e^{-tD^2})]$ with $k \in \{0, 1, 2\}$: M1 ($k = 0$ volume slot), M2 (Seeley–DeWitt heat-kernel coefficients), M3 (vertex-parity Hurwitz / Dirichlet-L content). The GH-convergence theorem (Theorem 5) closes WH1's commutative-factor leg on the Riemannian side (the almost-commutative structure's convergence is open); the Bisognano–Wichmann reading (Remark 43, with Unruh pendant) is a structural correspondence alongside it, whose operator-level falsifier is open. We close

out Sec. 8 with a brief structural appendix that records what Sprint L0 (May 2026) found when the audit was extended to the Riemannian $(3,0) \to$ Lorentzian $(3,1)$ extension of the framework.

Sprint L0 verdict on the master Mellin engine. Sprint L0 (`debug/lorentzian_l0_audit_memo.md`, May 2026; machine-readable table at `debug/data/lorentzian_l0_audit.json`) classifies all twenty-eight entries of Paper 34's projection dictionary [40] by their transfer behavior under signature extension; the full per-projection table sits at Paper 34 § V.E (`sec:lorentzian_transfer_audit`), the structural framing of the Sig/Op partition is in Paper 31 § 8 (`sec:signature_partition`) [39]. Two sharpenings of the master Mellin engine fall out of the L0 audit.

(i) M1 sub-mechanism is the load-bearing free-transfer engine. The M1 sub-mechanism ($k = 0$) on a temporal S^1_β or proper-time Rindler η_E-circle inherits a Lorentzian reading via the continuum Hawking / Sewell / Bisognano–Wichmann / Unruh correspondence, which is published prior art (the framework adds no finite-cutoff realization of it; Remark 43). Four entries of Paper 34's dictionary sit in the WICK-MAP FREE bucket via this mechanism: observation / temporal-window (§ III.15), vector-photon promotion (§ III.11) at the $1/(4\pi)$ per-loop level, gauge choice (§ III.26) at the Coulomb-gauge per-loop level, and the Wick-rotation projection itself (§ III.27). None requires Krein-space machinery for its metric-functional-level reading. The 2π of the Euclidean circle and the 2π of the continuum modular-flow / boost period coincide by the published continuum theorems; at finite cutoff the framework's 2π period is posited with the chosen generator, not derived, and its $\mathrm{Vol}(S^1)$ (M1) tag is a classification of that value (an earlier version said "the same generator on both sides", withdrawn 2026-09-28).

(ii) M3 trivialization at $(3,1)$ — a prediction since falsified or tautological (Sprint L2-D, § 8.9). The L0 audit derived the following prediction from the Bizi–Brouder–Besnard [50] signature classification; Sprint L2-D found it **FALSIFIED** under the n_{Fock}-parity convention and only *tautologically* confirmed under the chirality-pairing convention, so it is not a live structural prediction and is recorded here as history: at signature $(s,t) = (3,1)$, i.e. BBB $(m,n) = (4,6)$, the BBB sign table flips the $\{J, \gamma_5\}$ anticommutation relation that drives M3's vertex-parity content on the Riemannian side (Paper 28's $\zeta(s, 3/4)$ and $\zeta(s, 5/4)$ Hurwitz reductions producing Catalan's G and $\beta(4)$). On chirality-symmetric spectrum at $(3,1)$, the vertex-parity asymmetry vanishes, M3's half-integer-Hurwitz content becomes structurally empty, and the case-exhaustion theorem (Theorem 4) trivializes to a *two-mechanism partition* M1 $\cup$ M2 on the Lorentzian side. The Riemannian theorem is stated at signature $(3,0)$, so neither the prediction nor its failure bears on it. The L2-D outcome is that the BBB sign flip is a spectral-triple-axiom statement, not a vertex-parity-sum statement: M3's mechanism is sectional in n_{Fock}-parity, not in spacetime signature (§ 8.9).

What this does and does not change. The Riemannian case-exhaustion theorem stands; the commutative-factor convergence theorem of WH1 (Theorem 5) stands. The L0 audit is a documentation-level extension that classifies the framework's transfer behavior under signature change without re-deriving any of the Riemannian results. The falsifier it names would deepen the Wick-rotation reading from structural correspondence to an operator-level identification by *deriving* the modular Hamiltonian K on $\mathcal{T}_{n_{\max}}$ for a Rindler-wedge-restricted Camporesi–Higuchi state, verifying $\sigma_{i\cdot 2\pi} =$ identity at finite $n_{\max}$, and taking the GH limit using Theorem 5's lemmas; that would bear on all four Wick-rotation faces simultaneously via the shared KMS-Tomita-Takesaki algebra. Sprint L1 did not do this: it *chose* an integer-spectrum $K = 2J_z$, for which the period closure is automatic for any integer-spectrum generator (Remark 43; regression-pinned in `tests/test_modular_period_closure_automatic.py`), so the falsifier remains open.

The Nieuviarts 2025 [52] twisted-spectral-triple morphism is the named trigger that motivated reopening the May 2026 NO-GO scoping; a parallel L2-Nieuviarts-scoping pass tests whether the construction applies cleanly to $S^3 = \mathrm{SU}(2)$ (odd-dimension caveat) and would shorten blocker B1 (Lorentzian propinquity, long-range) substantially if applicable. The current audit does *not* assume the morphism applies.

8.7 Krein space construction at signature $(3,1)$ (Sprint L2-B, 2026-05-16)

The first concrete deliverable of the Sprint L2 arc lifts the Hilbert space and the fundamental symmetry of the truncated triple from signature $(3,0)$ to signature $(3,1)$, following the van den Dungen 2016 prescription [51]. The construction is at the *Hilbert-space level only*; the Lorentzian Dirac and the Connes axiom audit are deferred to L2-C and L2-D below.

Krein space. For $n_{\max} \geq 1$ and $N_t \geq 1$ define

$$\mathcal{K}_{n_{\max},N_t} \;:=\; \mathcal{H}_{\mathrm{GV}}^{n_{\max}} \otimes L^2(\mathbb{R}_t)_{\mathrm{cutoff},N_t}, \tag{8}$$

where $\mathcal{H}_{\mathrm{GV}}^{n_{\max}}$ is the Riemannian Hilbert space of Definition 2 and $L^2(\mathbb{R}_t)_{\mathrm{cutoff},N_t}$ is the truncated temporal slot: a uniform real grid $t_k = -T_{\max} + k \cdot 2T_{\max}/(N_t - 1)$ for $N_t \geq 2$, or the singleton $\{0\}$ for $N_t = 1$. The temporal slot is NOT compactified to S^1_β (which would create closed timelike curves per Geroch's theorem and is structurally incompatible with the bounded-wedge reading of Paper 42).

Conventions. We use the West-coast metric $\eta = \mathrm{diag}(+1,-1,-1,-1)$ and the Peskin–Schroeder chiral (Weyl) basis for the four $\mathrm{Cl}(3,1)$ gamma matrices. In this basis, $\gamma^5 = \mathrm{diag}(-I_2, +I_2)$ is diagonal and `FullDiracLabel.chirality` is identified with the γ^5 eigenvalue, so the Riemannian limit identification with $\mathcal{H}_{\mathrm{GV}}$ is a transparent basis-label match (no similarity transform required). The Dirac basis alternative (where γ^0 is diagonal instead) is unitarily equivalent at the operator level but would have required a unitary similarity transform for the Riemannian-limit check; the chiral basis was chosen on Riemannian-limit-clarity grounds.

Fundamental symmetry J. The fundamental symmetry is

$$J \;=\; J_{\mathrm{spatial}} \otimes I_{N_t}, \tag{9}$$

where J_{spatial} is the lift of γ^0 to the chirality-doubled $\mathcal{H}_{\mathrm{GV}}$: on $|n_D, \kappa, m_j, \chi\rangle$ the action is the chirality swap $|n_D, \kappa, m_j, -\chi\rangle$. In the chiral basis, $\gamma^0 = \left(\begin{smallmatrix} 0 & I_2 \\ I_2 & 0 \end{smallmatrix}\right)$ is the off-diagonal block-swap, so J_{spatial} is a real $\{0,1\}$-valued permutation matrix. The Krein inner product is $\langle \psi, \phi \rangle_{\mathcal{K}} := \langle \psi, J\phi \rangle$.

Krein axiom verification. At every panel cell $(n_{\max}, N_t) \in \{1,2,3\} \times \{1, 11, 21\}$, the construction satisfies

1. $J^2 = +I$ (bit-exact Frobenius residual $= 0.0$),
2. $J^* J = I$ (J Hilbert-unitary, bit-exact),
3. $J^* = J$ (J Hilbert-Hermitian, bit-exact),
4. $\mathcal{K} = \mathcal{K}^+ \oplus \mathcal{K}^-$ with $\dim \mathcal{K}^\pm = \dim \mathcal{K}/2$ and $J|_{\mathcal{K}^\pm} = \pm I$ on the eigenspaces (rank $\dim \mathcal{K}/2$ each, bit-exact).

The bit-exact-zero residuals follow from the structural fact that $J = J_{\text{spatial}} \otimes I_{N_t}$ is the Kronecker product of a $\{0,1\}$-valued integer permutation matrix and an identity matrix, both involving only operations exact in IEEE 754 arithmetic on real integers.

Load-bearing Riemannian-limit check. At $N_t = 1$ the temporal slot collapses to the singleton $\{0\}$ and $\mathcal{K}_{n_{\max},N_t=1}$ recovers $\mathcal{H}_{\text{GV}}^{n_{\max}}$ *bit-identically* at every $n_{\max} \in \{1,2,3\}$: the dimension formula $\dim \mathcal{K}_{n_{\max},1} = N_{\text{Dirac}}(n_{\max}) = \frac{2}{3} n_{\max}(n_{\max}+1)(n_{\max}+2) = 4, 16, 40$ matches the Paper 32 § III formula exactly, the basis labels are bit-identical under `FullDiracLabel.__eq__`, and the J matrix at $N_t = 1$ equals J_{spatial} as numpy arrays (Frobenius difference = 0.0). This confirms that the Camporesi–Higuchi spatial spinor bundle is structurally compatible with the $\text{Cl}(3,1)$ gamma-matrix embedding in the West-coast chiral basis; the commutative-factor convergence theorem of WH1 (Theorem 5) is not re-opened by this result. Source: `debug/l2_b_krein_construction_memo.md`, raw data `debug/data/l2_b_krein_construction.json`, implementation `geovac/krein_space_construction.py` regression tests `tests/test_krein_space_construction.py` (109 cases, all pass; zero regression in 132 upstream tests).

Honest scope. The Krein-space construction lifts the Hilbert space and the fundamental symmetry only. The Lorentzian Dirac, the algebra $\mathcal{A}_L$ on $\mathcal{K}$, the operator-algebra Krein-adjoint structure, and the modular Hamiltonian on the Krein space are deferred to L2-C, L2-D, and L2-E respectively.

8.8 Lorentzian Dirac operator D_L on $S^3 \times \mathbb{R}$ (Sprint L2-C, 2026-05-16)

Sprint L2-C constructs the Lorentzian Dirac operator on the Krein space (8) following the i^t Krein-triple pattern of van den Dungen Proposition 4.1 [51] (whose operator is i^t times the *pseudo*-Riemannian Dirac of (M,g), with the Riemannian metric g_r entering only through completeness and the positive-definite inner product): Wick-rotate the Lorentzian metric g on $M = S^3 \times \mathbb{R}$ to its Riemannian counterpart g_r, construct the standard Riemannian Dirac $\not{D}_{g_r}$ on (M, g_r), and define $D_L = i\ \not{D}_{g_r}$. The Krein self-adjointness of this D_L is verified directly ($\{\gamma^0, D_{\text{GV}}\} = 0$, ∂_t anti-Hermitian; `geovac/lorentzian_dirac.py`), not inherited from the proposition; the identification of $i\not{D}_{g_r}$ with van den Dungen's $i\not{D}_g$ on the static product is not established here (citation corrected 2026-09-03).

Construction. The Wick-rotated Riemannian Dirac decomposes via spatial-temporal splitting:

$$\not{D}_{g_r} \;=\; \gamma^0 \otimes \partial_t + D_{\text{GV}} \otimes I_{N_t}, \tag{10}$$

with γ^0 from the Sprint L2-B chiral basis, D_{GV} the Paper 32 § III Camporesi–Higuchi truthful full-Dirac matrix on $\mathcal{H}_{\text{GV}}^{n_{\max}}$, and ∂_t the temporal-derivative matrix on the $L^2(\mathbb{R}_t)_{\text{cutoff},N_t}$ grid. The Lorentzian Dirac is then

$$D_L \;=\; i \cdot \left[\gamma^0 \otimes \partial_t + D_{\text{GV}} \otimes I_{N_t}\right] \;\in\; \mathbb{C}^{\dim\mathcal{K}\times\dim\mathcal{K}}. \tag{11}$$

Temporal derivative. The temporal-derivative ∂_t uses centered finite differences with Dirichlet (zero) boundary conditions:

$$(\partial_t)_{k,l} \;=\; \frac{1}{2\Delta_t}\big(\delta_{l,k+1} - \delta_{l,k-1}\big), \qquad \Delta_t = \frac{2T_{\max}}{N_t - 1}. \tag{12}$$

This choice is constrained but not unique: what the vdD recipe requires for Krein-self-adjointness $D_L^{\times} = D_L$ is an anti-Hermitian ∂_t ($\partial_t^{\dagger} = -\partial_t$), and centered FD with Dirichlet-zero boundary

conditions is the simplest (lowest-order) discretization with that property; any antisymmetric stencil (e.g. fourth-order centered FD) is also anti-Hermitian and would serve. Forward, backward, and upwind schemes give $\partial_t^\dagger \neq -\partial_t$ (breaking Krein-self-adjointness); periodic boundary conditions would re-introduce closed timelike curves (Sprint L2-A audit § 3.8).

Sign of i^t. With $t = 1$, the prescription gives $i^t = i$ (not $-i$). The sign is derived from the structural requirement $D_L^\times = D_L$ with $D_L^\times := \gamma^0 D_L^\dagger \gamma^0$, using $\{\gamma^0, D_{\mathrm{GV}}\} = 0$ (the chirality swap γ^0 anticommutes with the chirality-diagonal D_{GV}) and $\partial_t^\dagger = -\partial_t$. Explicit symbolic calculation (sympy verification) gives $D_L^\times = +D_L$ for $i^t = +i$ and $D_L^\times = -D_L$ for $i^t = -i$.

Verification at finite cutoff. The construction verifies at every panel cell $(n_{\max}, N_t) \in \{1, 2, 3\} \times \{1, 11, 21\}$:

1. *Krein-self-adjointness.* $\|D_L^\times - D_L\|_F = 0$ bit-exact at every cell.
2. *Real spectrum on Krein-positive cone.* Restricting D_L to $\mathcal{K}^+$ (the $+1$ eigenspace of J) gives real eigenvalues to machine precision ($\max|\mathrm{Im}(\lambda)| \leq 2.32 \times 10^{-17}$).
3. *Riemannian-limit recovery (LOAD-BEARING).* At $N_t = 1$, $D_L|_{N_t=1} = i \cdot D_{\mathrm{GV}}$ bit-identically. The absolute-value spectrum $\{|\lambda_n| = n_{\mathrm{Fock}} + 1/2\}$ with multiplicities $g_n^{\mathrm{Dirac}} = 2n(n+1)$ matches D_{GV} bit-exactly at $n_{\max} = 1, 2, 3$.

Chirality finding (non-zero $\{\gamma^5, D_L\}$). A non-trivial structural finding emerges: the BBB universal relation $\chi D = -D\chi$ (Sec 5(v) of [50]) does *not* hold on truthful D_{GV}. The anticommutator $\{\gamma^5, D_L\}$ has Frobenius norm $2\|\gamma^5 D_{\mathrm{GV}}\|_F = 2\|D_{\mathrm{GV}}\|_F$ at $N_t = 1$ (values $6.00, 18.33, 38.88$ at $n_{\max} = 1, 2, 3$), because GeoVac's truthful D_{GV} is chirality-diagonal in the Peskin–Schroeder chiral basis and therefore commutes with γ^5. This finding is structural, not a bug: it is a consequence of two locked Riemannian-side conventions (chirality-as-γ^5 from L2-B and chirality-diagonal D_{GV} from Paper 32 § III). The full diagnostic and the three resolutions are documented in Sec. 8.9 below.

Sources. Implementation: `geovac/lorentzian_dirac.py`. Regression tests: `tests/test_lorentzian_dirac.p` (108 cases, all pass; zero regression in 205 upstream tests). Sprint memo: `debug/l2_c_lorentzian_dirac_memo.md` raw data `debug/data/l2_c_lorentzian_dirac.json`.

8.9 Connes axiom audit at $(m, n) = (4, 6)$ (Sprint L2-D, 2026-05-16)

Sprint L2-D performs the Connes axiom audit on the pair (D_L, γ^5, J_L) built in Sprints L2-B/C, against the Bizi–Brouder–Besnard 2018 [50] classification at $(m, n) = (4, 6)$.

BBB sign determination. Reading the BBB Table 1 entries at $(m, n) = (4, 6)$ directly from arXiv:1611.07062 v2: $\varepsilon = +1$, $\varepsilon'' = -1$, $\kappa = -1$, $\kappa'' = +1$. These four primitive signs, combined with the four BBB defining relations $J^2 = \varepsilon$, $J\chi = \varepsilon'' \chi J$, $J\eta = \varepsilon\kappa\, \eta J$, $\eta\chi = \varepsilon''\kappa''\, \chi\eta$, predict at $(4, 6)$: $J^2 = +I$, $\{J, \chi\} = 0$, $\{J, \eta\} = 0$, $\{\eta, \chi\} = 0$. **[SYMBOLIC + MEASURED]** At the bare Cl$(3, 1)$ 4-spinor level with charge conjugation $U_4 = i\gamma^2$, all four relations hold bit-exact in the chiral basis (Frobenius residual $= 0.0$). The translation $(s, t) = (3, 1) \leftrightarrow (m, n) = (4, 6)$ uses BBB Table 3: $m \equiv t + s$, $n \equiv t - s \pmod 8$, giving $m = 4$, $n = -2 \equiv 6 \pmod 8$.

Lift to the Krein space. The 4-spinor charge conjugation $i\gamma^2 = (i\sigma_y)_{\text{chir}} \otimes (i\sigma^2)_{\text{spin}}$ decomposes into a chirality-swap-with-sign factor times the standard spin-1/2 charge conjugation. The lift to $\mathcal{K}_{n_{\max},N_t}$ is

$$J_L \;=\; J_{L,\text{spatial}} \otimes K_t, \tag{13}$$

where $J_{L,\text{spatial}}$ acts on $|n,l,m_j,\chi\rangle$ as $\sigma_{\text{chir}}(\chi)\cdot\sigma_{\text{spin}}(l,-m_j)\cdot|n,l,-m_j,-\chi\rangle$ with $\sigma_{\text{chir}}(\pm1) = \mp1$ and $\sigma_{\text{spin}}(l,m_j) = (-1)^{(2l+1-2m_j)/2}$, and K_t is complex conjugation on $\mathbb{C}^{N_t}$. The matrix $U_L := J_L \cdot K^{-1}$ is a real $\{0,\pm1\}$-valued unitary; the antilinear $J_L = U_L K$ satisfies $J_L^2 = +I$ bit-exact via $U_L\overline{U_L} = U_L^2$ in IEEE 754.

Six-axiom panel. At every panel cell $(n_{\max}, N_t) \in \{1,2,3\} \times \{1,11,21\}$ on truthful D_{GV}:

Axiom	Predicted at $(4,6)$	Residual
(i) $J_L^2 = +I$ (BBB $\varepsilon = +1$)	$+I$	0.0 bit-exact
(ii) $\{J_L, \chi\} = 0$ (BBB $\varepsilon'' = -1$)	0	0.0 bit-exact
(iii) $\{J_L, \eta\} = 0$ (BBB $\varepsilon\kappa = -1$)	0	0.0 bit-exact
(iv) $J_L D_L = +D_L J_L$ (BBB universal Sec 5(v))	$JD = +DJ$	0.0 bit-exact
(v) $\{\chi, D_L\} = 0$ (BBB universal Sec 5(v))	$\chi D = -D\chi$	STRUCTURAL FINDING (non-zero)
(vi) order-zero $[a, J_L b J_L^{-1}] = 0$	0	≤ 0.0675 (max entry, sample-of-3)
(vii) order-one $[[D_L, a], J_L b J_L^{-1}] = 0$	0	≤ 0.1014 (max entry, sample-of-3)

Axioms (i)–(iv) hold bit-exactly at every cell. Axioms (vi)–(vii) exhibit the same nonzero residuals as the Riemannian-side audit (Table 1, order zero and order one on $\mathcal{O}_{n_{\max}}$); over the full 55-element multiplier basis at $n_{\max} = 3$ the maxima are 0.0785 and 0.2026 (measured 2026-09-02), i.e. the sample-of-3 figures understate the order-one residual by a factor of two. The residuals are on the operator-system multiplier basis (on the diagonal $\mathcal{A}_{\text{GV}}$ both conditions hold trivially), and on the Riemannian side they grow with $n_{\max}$ (item 5 of the finite-$n_{\max}$ audit; **[MEASURED]**, mechanism open). Axiom (v) is the structural finding documented in Table 2 and discussed below.

Structural finding on $\{\chi, D_L\}$. The BBB universal axiom $\chi D = -D\chi$ fails on truthful D_{GV}: $\|\{\gamma_K^5, D_L\}\|_F = 2\|D_{\text{GV}}\|_F$ at $N_t = 1$, scaling as $\|D_{\text{GV}}\|_F\sqrt{N_t}\cdot(2+\text{small})$ at $N_t > 1$. The mechanism is the chirality-diagonal structure of D_{GV} in the chiral basis (§ 8.8). This is the load-bearing scope finding of Sprint L2-D and is the same structural fact captured in Table 2. The three resolutions (R1 accept, R2 use offdiag CH, R3 redefine γ^5) are documented in the discussion immediately following Table 2, with R1 + R2 in parallel as the recommended path for Sprint L2-E.

M3 trivialization is convention-dependent. Sprint L0 (§ 8.6 above) predicted that the M3 sub-mechanism of the master Mellin engine would *trivialize at* $(3,1)$ *on chirality-symmetric spectrum*, via the BBB sign-flip $\{J, \gamma^5\} = 0$ removing the asymmetry that drives M3's vertex-parity content. Sprint L2-D tests this prediction and finds: **the prediction is convention-dependent**.

Under the n_{Fock}*-parity convention* (the Paper 28 §QED-vertex convention that accesses Catalan G and Dirichlet $\beta(4)$ via quarter-integer Hurwitz shifts), the vertex-parity sum $D_{\text{even}}(4) - D_{\text{odd}}(4)$ on the chirality-symmetric $(3,1)$ truncation is bit-identical to the Riemannian value. Each full-Dirac level n_{Fock} contributes its full degeneracy $g_n = 2n(n+1)$ summed over both chirality blocks; the parity is read from the chirality-independent n_{Fock} label. The L0 prediction is therefore **FALSIFIED** under this reading at every $n_{\max} \in \{1,\ldots,5\}$ (bit-identical difference).

Under the *chirality-pairing convention* (pair "even" with $\chi = +1$ Weyl and "odd" with $\chi = -1$ anti-Weyl), the chirality-symmetric truncation gives $D_+ = D_-$ identically by chirality symmetry, so the difference vanishes bit-exact. The L0 prediction is **trivially confirmed** under this reading, but tautologically (any chirality-symmetric truncation gives the result by symmetry alone, independent of signature).

The structural refinement is therefore: **the BBB sign-flip $\{J, \gamma^5\} = 0$ at $(m, n) = (4, 6)$ is a spectral-triple-axiom statement, not a vertex-parity-sum statement.** It does not directly imply M3 trivialization at any natural parity convention. M3's mechanism is sectional in n_{Fock}-parity, NOT in spacetime signature. Paper 18 § III.7 records the corresponding master-Mellin-engine sharpening.

Sources. Implementation: `geovac/connes_axiom_audit_31.py` (700 lines incl. docstrings). Regression tests: `tests/test_connes_axiom_audit_31.py` (87 cases, all pass). Sprint memo: `debug/l2_d_connes_ax` raw data `debug/data/l2_d_connes_axiom_audit_31.json` (BBB-Table-1 signs, 4-spinor verification, full axiom panel, M3 trivialization panel).

Sprint L2-E handoff. The four BBB-predicted-sign axioms hold bit-exactly, so the operator-algebra structure on the Krein space is fully BBB-compatible at the J-relation level. Sprint L2-E (the Lorentzian-level Paper 42 redo) should construct the Krein-side modular Hamiltonian $K_L = -\log \Delta_L$ on the Krein-positive cone $\mathcal{K}^+$ and verify the analog of Paper 42's $\sigma_{2\pi}(O) = O$ at the operator-system level on the Krein space. The resolution choice R1 (truthful D_{GV}, preserves Riemannian-limit recovery) versus R2 (offdiag CH, satisfies BBB axiom (v)) is recommended to be tested in parallel. The commutative-factor convergence theorem of WH1 (Theorem 5) is not re-opened by these results.

8.10 Krein-level compact-period closure (signature-blind) at finite cutoff for a chosen generator (Sprint L2-E, 2026-05-17)

[PANEL-VERIFIED] computation; [SYMBOLIC (trivial)] content Sprint L2-E (`geovac/modular_hamilt` $\sim$ 860 lines including docstrings; tests in `tests/test_modular_hamiltonian_lorentzian.py`, 74 fast plus 3 slow, all passing) builds the operator-system-level Krein-side modular Hamiltonian on the hemispheric wedge of $S^3 \times \mathbb{R}$ at signature $(3, 1)$ and verifies the four LOAD-BEARING falsifiers of the Sprint L2-F catalogue (`debug/sprint_l2_falsifiers.md` § 3) plus the Riemannian-limit recovery LOAD-BEARING falsifier plus the headline H_{local} verdict and the six-witness collapse, all bit-exact at every $(n_{\max}, N_t) \in \{1, 2, 3\} \times \{1, 11, 21\}$. The closure mirrors Paper 42 §§ 4–8 [53] verbatim with the temporal slot added via the Sprint L2-B Krein space.

Lorentzian wedge, KMS state, and BW choice. The hemispheric wedge on the Krein space is $W_L = P_W^{\text{spatial}} \otimes P_{t \geq 0}$, where $P_W^{\text{spatial}} = (1/2)(I + R_{\text{polar}})$ is the Paper 42 hemispheric wedge on $\mathcal{H}_{\text{GV}}$ (the m_j-reflection involution) and $P_{t \geq 0}$ is the $t \geq 0$ half-line projector on $\mathbb{C}^{N_t}$. The wedge KMS state under the BW choice $H_{\text{local}} := K_L^{\alpha, W}/\beta$ at $\beta = 2\pi$ is

$$\rho_W^L = \frac{e^{-K_L^{\alpha, W}}}{Z},$$

β-independent at the algebra-action level, with $K_L^{\alpha, W}$ the wedge-restricted BW-α generator inheriting integer eigenvalues two_m_j from Paper 42 Definition 5.1.

The Krein-level period closure at finite cutoff. The Krein-side BW-α (chosen integer-spectrum generator) and BW-γ (Tomita–Takesaki on the Krein-GNS Hilbert–Schmidt space $M_{\dim W_L}(\mathbb{C})$) constructions both verify $\sigma^L_{2\pi}(O) = O$ bit-exactly at every panel cell:

$(n_{\max}, N_t)$	$\dim K$	$\dim W_L$	$r_W^{L,\alpha}$	$r_W^{L,\mathrm{TT}}$
$(1,1)$	4	2	0.0	$\leq 10^{-16}$
$(1,11)$	44	12	$\leq 10^{-16}$	$\leq 10^{-16}$
$(1,21)$	84	22	$\leq 2\times 10^{-16}$	$\leq 2\times 10^{-16}$
$(2,1)$	16	8	$\leq 10^{-16}$	$\leq 10^{-16}$
$(2,11)$	176	48	$\leq 3\times 10^{-16}$	$\leq 3\times 10^{-16}$
$(2,21)$	336	88	$\leq 4\times 10^{-16}$	$\leq 4\times 10^{-16}$
$(3,1)$	40	20	$\leq 5\times 10^{-16}$	$\leq 5\times 10^{-16}$
$(3,11)$	440	120	$\leq 4\times 10^{-15}$	$\leq 4\times 10^{-15}$
$(3,21)$	840	220	$\leq 7\times 10^{-15}$	$\leq 7\times 10^{-15}$

Verdict at every cell: COMPACT_PERIOD_CLOSURE (signature-blind; the closure cannot distinguish the $(3,1)$ carrier from the Euclidean one — see the retraction note below). Residuals scale as $O(\sqrt{\dim}\cdot\varepsilon_{\mathrm{machine}})$ — pure round-off, no structural drift. Flow conjugacy $\sigma_t^{L,\mathrm{TT}}(O) = \sigma_{-t}^{L,\alpha}(O)$ at general $t \in \{1, \pi, 2\pi\}$ holds bit-exactly with the same residual bound, and the six-witness cross-consistency residual is exactly 0.0 at every cell.

Statement (Krein-level compact-period closure (signature-blind) at finite cutoff). *On the Krein space $\mathcal{K}_{n_{\max},N_t} = \mathcal{H}_{\mathrm{GV}}^{n_{\max}} \otimes \mathbb{C}^{N_t}$, with the wedge $W_L = P_W^{\mathrm{spatial}} \otimes P_{t\geq 0}$, the chosen integer-spectrum generator $K_L^{\alpha,W}$, the state $\rho_W^L = e^{-K_L^{\alpha,W}}/Z$, and the unit normalisation $\kappa_g = 1$, the following hold. They hold for* any *integer-spectrum generator, so they identify no Bisognano–Wichmann modular flow (regression-pinned in* `tests/test_modular_period_closure_automatic.py`*); deriving K from a wedge-restricted Camporesi–Higuchi state is the open falsifier (Remark 43):*

- *BW-α period closure:* $\sigma^{L,\alpha}_{2\pi}(O) = O$ bit-exactly for all $O \in B(\mathcal{K}_{W_L})$.
- *BW-γ Tomita period closure:* $\sigma^{L,\mathrm{TT}}_{2\pi}(a) = a$ bit-exactly for all $a \in B(\mathcal{K}_{W_L})$ on the Krein-GNS space.
- *Flow conjugacy:* $\sigma_t^{L,\mathrm{TT}}(a) = \sigma_{-t}^{L,\alpha}(a)$ bit-exactly at general t.
- *Six-witness collapse:* all six instantiations $\{\mathrm{BW}, \mathrm{HH}_{M=1}, \mathrm{HH}_{M=2}, \mathrm{Sew}_{M=1}, \mathrm{Unruh}_{a=1}, \mathrm{Unruh}_{a=2}\}$ give bit-identical Δ_L and K_L^{TT}.

All four statements verified bit-exact at every tested $(n_{\max}, N_t) \in \{1,2,3\} \times \{1,11,21\}$. Proofs mirror Paper 42 Theorems 5.4, 6.3, 7.1, and Cor. 8.1 [53].

The 2π of the period is the period chosen with the integer-spectrum generator — posited, not derived — on the Riemannian side and on the Krein carrier alike; its tag as $\mathrm{Vol}(S^1)$ (master Mellin engine $k = 0$, M1) classifies that posited value and is not an output of the case-exhaustion theorem. The closure holds at every Krein cutoff because the chosen generator has integer spectrum, and it is *signature-blind* (withdrawn 2026-09-03, per the 2026-07-04 retirement recorded in the C16 registry): the finite generator $K = 2J_z$ is *posited*, not derived as a truncated boost, and its flow is compact (integer spectrum, $e^{2\pi i K} = \mathbb{1}$); the period closure is the compact $\beta = 2\pi$ circle, and the $(3,1)$ label is a carrier choice that the closure cannot distinguish from the Euclidean one (Papers 45/47; WH7). **[ARGUED]** The Euclidean-only reading of the closure; a Wick/KMS reading is conditional on

Paper 42's open question O0 [53]. An earlier version of this paragraph read the closure as an operator-level identification of the four-witness statement at finite cutoff; that reading is retired.

Riemannian-limit recovery (load-bearing falsifier). At $N_t = 1$, $P_{W_L} = P_W^{\text{spatial}}$, $K_L^{\alpha,W} = K_\alpha^W$, $\rho_W^L = \rho_W$, $\Delta_L = \Delta$, and $K_L^{\text{TT}} = K_{\text{TT}}$ bit-identically (Frobenius residual exactly 0.0, not merely machine-precision) at every $n_{\max} \in \{1, 2, 3\}$. This is the load-bearing falsifier that ensures Sprint L2-E is the N_t-carrier *extension* of Paper 42 § 6 (the Tomita BW-γ construction) [53], not a parallel-but-different construction; it does not make the extension Lorentzian in any metrically visible sense (withdrawn reading, 2026-09-03).

H_{local} at the Riemannian limit (Sprint L2-E). Paper 42 § 7.2 / open question O3 names a load-bearing scope finding: the framework's intrinsic Dirac D_W is not the local Hamiltonian whose $\beta = 2\pi$ Gibbs state is $\rho_W = e^{-K_\alpha^W}/Z$; $H_{\text{local}} := K_\alpha^W/\beta$ is, with $K_\alpha^W = 2J_z$ posited (whether ρ_W is a BW vacuum is open, Paper 42's O0). Sprint L2-E tests whether this finding holds at signature $(3, 1)$. **Result:** at the Riemannian limit ($N_t = 1$) on the Lorentzian Krein space, $\|H_{\text{local}}|_{(3,1),N_t=1} - D_L^W|_{N_t=1}\|_F$ EQUALS the Riemannian-side residual bit-exact at every tested $n_{\max} \in \{1, 2, 3\}$ (values 2.1332, 6.5275, 13.854). At $N_t > 1$ the Lorentzian-side residual is *refined upward* by the temporal-derivative content of $D_L = i(\gamma^0 \otimes \partial_t + D_{\text{GV}} \otimes I)$. At $N_t = 1$ the Lorentzian Krein space reduces to the Riemannian one, so the equality there holds by construction: it is a consistency check, not evidence of signature-independence, and the non-trivial content is the $N_t > 1$ refinement. Whether the spectral-action–vs–modular-Hamiltonian generator distinction is a deeper structural feature of the framework, rather than of the round S^3 truncation, is not settled by this test; Paper 42 § 10 O3 stays open [53].

Honest scope. Closure is at finite $n_{\max}$ and finite N_t, NOT in the continuum (GH-limit) limit. No published Lorentzian quantum-metric convergence construction exists as of May 2026 (Latrémolière 2017/2026 and Hekkelman–McDonald 2024 are strictly Riemannian), and the framework's own attempt was DESCOPED on 2026-06-09 by the K^+ annihilation theorem (Paper 45): the finite generator $K = 2J_z$ is posited and its flow is compact, so no Lorentzian metric statement is available at finite cutoff (the Euclidean-only reading is **[ARGUED]**; a Wick/KMS reading is conditional on Paper 42's O0) (updated 2026-09-03; an earlier version named this as the Sprint L3 target). The Sprint L2-D structural finding (BBB universal $\chi D = -D\chi$ fails on truthful D_{GV}) is preserved in L2-E: the wedge construction is $K_L^{\alpha,W}$-driven and does not invoke this axiom. Cross-manifold W2b ($\mathcal{T}_{S^3} \otimes \mathcal{T}_{\text{Hardy}(S^5)}$) remains **open** as a separate frontier (see § VIII.D "frontier-of-field framing"), structurally distinct from the signature-blind $(3, 1)$ Krein-carrier sign audit within $\mathcal{T}_{S^3}$ recorded here (not a Lorentzian extension; the programme is descoped, Paper 45).

Sources. Memo: `debug/l2_e_modular_hamiltonian_lorentzian_memo.md`. Computational driver: `debug/l2_e_modular_hamiltonian_lorentzian_compute.py`. Raw data: `debug/data/l2_e_modular_hamilton` Implementation: `geovac/modular_hamiltonian_lorentzian.py`. Tests: `tests/test_modular_hamiltonian_lor` (74 fast + 3 slow, all pass; zero regression in 355 upstream tests). The math.OA-style standalone writeup of the (since descoped) Lorentzian-extension programme was Paper 43, archived 2026-09-28 after both of its headline results were withdrawn (the finite four-witness closure is automatic for the posited generator, and the Pythagorean $1/\pi^2$ is the posited normalisation squared); its surviving content — the Krein carrier and its axiom audit, the automatic Krein-carrier period closure, and the Pythagorean orthogonality as a chirality-parity identity — is now Paper 42 §11, which makes no metrically visible Lorentzian claim (the programme is descoped, Paper 45).

8.11 Modular Hamiltonian at finite $n_{\max}$: Sprint L1 period closure on the Riemannian side (2026-05-16)

[SYMBOLIC (trivial)] This subsection records the Sprint L1 computation. It does *not* close the falsifier named in Remark 43 and § 8.6. The generator K on the truncated Camporesi–Higuchi spectral triple $\mathcal{T}_{n_{\max}}$ is *chosen* as $K = 2J_z$ (`geovac/modular_hamiltonian.py`), not derived from a wedge-restricted state; it has integer spectrum, so the period closure $\sigma_{2\pi}(O) = O$ holds bit-exactly for any O of integer spectrum (regression-pinned in `tests/test_modular_period_closure_automatic.py`), and the four witness labels (Bisognano–Wichmann, Hartle–Hawking, Sewell, Unruh) are the same K with κ_g relabelled, so they agree by construction. The four-witness statement therefore remains a structural correspondence (Remark 43); it is not made framework-internal at finite cutoff by this computation.

Architecture. Sprint L1 implements the locked architectural decisions of L1-A (`debug/l1_modular_hamiltonian`
L1-B (`debug/l1_infrastructure_audit_memo.md`), and L1-C (`debug/l1_witness_spec_memo.md`):

- **Wedge (W1):** hemispheric P_W on S^3 aligned with the Hopf-base axis, defined as $P_W = (1/2)(I + R_{\text{polar}})$ where $R_{\text{polar}} : m_j \mapsto -m_j$ is the equatorial reflection involution on the spinor basis; $P_W^2 = P_W$ and $P_W^* = P_W$ are bit-exact (no smoothing needed; the reflection is an exact involution on the discrete basis).
- **Unit normalisation (U-1):** natural rapidity units $\kappa_g = 1$ for the BW witness; the four witnesses parameterize as Hartle–Hawking $\kappa_g = 1/(4M)$, Unruh $\kappa_g = a$, Sewell $\kappa_g = 1/(4M)$.
- **Construction:** the BW-α "geometric realization" is the *choice* $K = J_{\text{polar}} = 2J_z$ (rotation generator around the wedge axis), with integer eigenvalues two_m$_j$ on the full-Dirac basis (odd integers in $\{-(2l+1), \ldots, +(2l+1)\}$). The Tomita BW-$\gamma$ realization is also instantiated as a sanity check on the trace cyclicity of the KMS state at expectation level; both realizations close consistently, the Tomita one by construction, since the state is built from the same K.
- **Dirac:** truthful Camporesi–Higuchi (`camporesi_higuchi_full_dirac_matrix`), the only Dirac choice satisfying $J_{GV} D = +D J_{GV}$ on the truncated operator system (see § 4). R3.2's n-degeneracy of the Connes-distance SDP — a property of the doubled Fock/Weyl model, absent on the true Camporesi–Higuchi window (Paper 38) — would not bite for modular flow in any case, which requires only cyclic-separatingness of the cyclic vector.

Result. The maximum periodicity residual $\max_O \|\sigma_{2\pi}(O) - O\|_F$, taken over the wedge-restricted versions of the first five non-identity multipliers in the `FullDiracTruncatedOperatorSystem` at cutoff $n_{\max}$, is at machine precision $O(\sqrt{\dim \mathcal{H}_{n_{\max}}} \cdot \varepsilon_{\text{machine}})$ for every tested $n_{\max}$ and every witness:

$n_{\max}$	$\dim_H$	wedge dim	$\max \|\sigma_{2\pi}(O) - O\|_F$	verdict
2	16	8	1.80×10^{-16}	STRONG_IDENTIFICATION
3	40	20	4.76×10^{-16}	STRONG_IDENTIFICATION
4	80	40	9.33×10^{-16}	STRONG_IDENTIFICATION
5	140	70	1.59×10^{-15}	STRONG_IDENTIFICATION

The STRONG_IDENTIFICATION verdict (a code label, not a claim of identification) labels a bit-exact $\sigma_{2\pi} = \text{id}$ compact-period closure — a compact-flow feature of the integer modular spectrum (signature-blind, not a metric strengthening; cf. the retraction note in §8.10).

The $n_{\max} = 3$ cutoff is the one at which the cumulative node count $\dim_H = 4 + 12 + 24 = 40$ numerically coincides with Paper 2's $g_3^{\text{Dirac}} = \Delta^{-1} = 40$, the single-level degeneracy at $k = 3$ (different objects: $N_{\text{Dirac}}(n_{\max}) = n_{\max}\, g^{\text{D}}_{n_{\max}}/3$, so they agree only at $n_{\max} = 3$). The L1 closure at this cutoff carries no additional content: the closure holds identically at every cutoff.

Cross-witness collapse. The cross-witness consistency residual $\big|\|\sigma_{2\pi}^{(\text{witness})}(O) - O\| - \|\sigma_{2\pi}^{\text{BW}}(O) - O\|\big|$ is exactly 0 at every tested $n_{\max}$ between BW ($\kappa_g = 1$), HH at $M = 1$ ($\kappa_g = 1/4$), HH at $M = 2$ ($\kappa_g = 1/8$), Sewell at $M = 1$ ($\kappa_g = 1/4$), Unruh at $a = 1$ ($\kappa_g = 1$), and Unruh at $a = 2$ ($\kappa_g = 2$). The period closure is independent of κ_g because the posited generator $K_\alpha^W = 2J_z$ has integer spectrum: $\mathrm{e}^{\mathrm{i}\cdot 2\pi\cdot n} = 1$ for any integer n. The four labels collapse to a single test at the unit normalisation U-1, by construction.

Structural reading. The period closure $\sigma_{2\pi}(O) = O$ is finite-cutoff exact, not a limit-statement. It does not require the Gromov–Hausdorff machinery of Theorem 5 (although it is consistent with it via the state-space GH-rate cross-check: residual / $\gamma_{n_{\max}}$ ratio is $O(\varepsilon_{\text{machine}})$, several orders below the rate prediction). The reason is elementary: the chosen $K = 2J_z$ has integer spectrum on the full-Dirac basis (odd integers two_m$_j$, since m_j is half-integer). This is *not* a framework-internal version of Bisognano–Wichmann's continuum statement, which is about the modular flow of a given state; that version requires the open falsifier.

The π-source case-exhaustion theorem (Theorem 4) is unaffected by this result. The 2π in the modular period is the posited period of the chosen generator, tagged Vol(S^1) (M1) as a classification, not derived. The $\sigma_{2\pi}$ period closure holds at the operator level, but it is *signature-blind* (a compact-flow feature of the integer modular spectrum, not a metric strengthening): the Lorentzian boost-orbit reading of that period continues to run through the published Wick-rotation chain, the closure being unable to distinguish the $(3, 1)$ carrier from the Euclidean one.

Honest scope. The L1 closure is at signature $(3, 0)$ (Riemannian). Three distinctions remain open:

- A Krein-carrier sign audit at signature $(3, 1)$ via the Bizi–Brouder–Besnard 2018 [50] classification was carried out as Sprint L2 (§8.9, §8.10); it is signature-blind, the Sprint L0 prediction that M3 trivializes at $(3, 1)$ was falsified under the n_{Fock}-parity convention (§8.9), and the Lorentzian programme is descoped (Paper 45).
- Non-tracial states would require explicit Tomita polar decomposition of the modular S. The tracial Gibbs state on $\mathcal{T}_{n_{\max}}$ gives the trivial reduction $K_{\text{mod}} = \beta H_{\text{local}} + \log Z$ (BW-γ); non-tracial extension is an open structural question.
- Cross-manifold modular structure (the Paper 24 § V cross-manifold $\mathcal{T}_{S^3} \otimes \mathcal{T}_{\text{Hardy}(S^5)}$ blocker) remains unaddressed and is not the subject of Sprint L1.

The implementation is at `geovac/modular_hamiltonian.py` (production module, fresh) with `tests/test_modular_hamiltonian.py` (41 tests, 39 fast + 2 slow, all pass). Computational driver: `debug/l1_modular_hamiltonian_compute.py`. Machine-readable results JSON: `debug/data/l1_modular_hamil` Closure memo: `debug/l1_modular_hamiltonian_results_memo.md`.

Sprint L1-tighten (same day): Tomita–Takesaki consistency check. The L1 computation above uses the chosen generator $K = J_{\text{polar}}$ as the load-bearing object; the Tomita BW-γ realization was instantiated only as an expectation-level KMS sanity check. Sprint L1-tighten (same day, 2026-05-16 evening) rebuilds the BW-γ construction at the full Tomita–Takesaki level on the GNS Hilbert–Schmidt space $H_{\text{GNS}} = M_{\dim_W}(\mathbb{C}) \cong \mathbb{C}^{\dim_W^2}$, computes the modular operator Δ as a $(\dim_W^2 \times \dim_W^2)$ matrix via the polar decomposition $S = J_{\text{TT}}\Delta^{1/2}$, extracts $K_{\text{TT}} = -\log\Delta$, and verifies $\sigma_{2\pi}^{\text{TT}}(O) = O$ at the operator-action level for the same five generators tested by BW-α.

The construction uses the natural local Hamiltonian $H_{\text{local}} := K_\alpha^W/\beta$ on the wedge sub-algebra (so $\beta H_{\text{local}} = K_\alpha^W = 2J_z$ is a *posited* integer-spectrum generator, spectrum two_m$_j$, chosen by analogy with the Wightman BW vacuum $\propto e^{-2\pi K_{\text{boost}}}$ on the Rindler wedge; its identification as a BW modular Hamiltonian is open, Paper 42's O0 [53]). The wedge state $\rho_W = e^{-K_\alpha^W}/Z$ is the unit-β Gibbs state, not BW-normalised: since $K_\alpha^W = 2J_z$, the BW-normalised state would be $e^{-\pi K_\alpha^W}/Z$ (Paper 50 [63]).

Result (UNIFIED_STRONG — a bit-exact $\sigma_{2\pi} = \mathrm{id}$ compact-period closure, signature-blind, not a metric strengthening): at every tested $n_{\max} \in \{2,3,4,5\}$ for all six witness instantiations (BW, HH$_{M=1}$, HH$_{M=2}$, Sewell$_{M=1}$, Unruh$_{a=1}$, Unruh$_{a=2}$):

$n_{\max}$	$\dim_W$	$\dim_{\text{GNS}}$	$\sigma_{2\pi}^{\alpha}$ residual	$\sigma_{2\pi}^{\text{TT}}$ residual	$\sigma_t^{\text{TT}} - \sigma_{-t}^{\alpha}$ at $t=1$	$J_{\text{TT}}^2 - I$
2	8	64	1.10×10^{-16}	1.09×10^{-16}	4.18×10^{-17}	0
3	20	400	2.22×10^{-16}	1.07×10^{-15}	9.79×10^{-17}	5.00×10^{-17}
4	40	1600	3.51×10^{-16}	3.03×10^{-15}	3.18×10^{-16}	7.22×10^{-17}
5	70	4900	4.97×10^{-16}	4.79×10^{-15}	4.03×10^{-16}	2.18×10^{-17}

All residuals at machine precision. The four-witness cross-collapse is preserved at the Tomita–Takesaki level (in fact, the density matrix $\rho_W = e^{-K_\alpha^W}/Z$ is itself β-independent, so Δ and K_{TT} are identical across the six witnesses); all six instantiations give bit-identical $\sigma_{2\pi}^{\text{TT}}$ residuals at every $n_{\max}$.

Structural relation between BW-α and BW-γ. The two constructions generate the same modular automorphism on the algebra, up to the sign of t:

$$\sigma_t^{\text{TT}}(a) \;=\; \rho^{it}\, a\, \rho^{-it} \;=\; e^{-itK_\alpha^W}\, a\, e^{+itK_\alpha^W} \;=\; \sigma_{-t}^{\alpha}(a). \tag{14}$$

The cross-flow difference $\|\sigma_1^{\text{TT}}(O) - \sigma_1^{\alpha}(O)\|_F \sim 0.16$ at non-period $t = 1$ is non-zero in general (the conjugate flows are not equal at arbitrary t); but $\|\sigma_1^{\text{TT}}(O) - \sigma_{-1}^{\alpha}(O)\|_F < 10^{-16}$ verifies equation (14) bit-exactly. At $t = 2\pi$ the sign is invisible because $e^{\pm i2\pi n} = 1$ for any integer n, so both flows close to the identity bit-exactly.

$J_{\text{TT}}^2 = +I$ at the FULL Tomita level. The polar-decomposition antilinear conjugation $J_{\text{TT}}(X) = \rho^{1/2}X^*\rho^{-1/2}$ satisfies $J_{\text{TT}}^2 = +I$ to machine precision at every $n_{\max}$ (residuals in the table above). This is structurally distinct from the L1 verification, which only checked the canonical sanity case $U = I$; L1-tighten verifies the $\rho^{1/2}$-weighted Tomita conjugation itself, which is the genuine state-dependent antilinear operator defined by the polar decomposition. This separates J_{TT} from the real structure J_{GV} (which has $J_{\text{GV}}^2 = -I$); the two antilinear operators coexist on $H_{n_{\max}}$ without conflation, as required by L1-A § 10e.

Verdict. The two constructions agree at the operator-action level (up to sign of t; identical at the period), but the agreement is circular: ρ_W is built as $e^{-K_\alpha^W}/Z$, so its modular generator is K_α^W by construction, and Eq. (14) restates that. The Tomita machinery is verified correct; no Bisognano–Wichmann flow is identified, and the four-witness statement is not closed at the operator-system level by either reading (the falsifier of Remark 43 remains open).

Honest scope. The unified closure rests on the choice $H_{\text{local}} = K_\alpha^W/\beta$, which makes the wedge KMS state the unit-β Gibbs state $\rho_W = e^{-K_\alpha^W}/Z$ of a posited generator chosen by analogy with the Wightman BW vacuum $\propto e^{-2\pi K_{\text{boost}}}$ (not BW-normalised, which would be $e^{-\pi K_\alpha^W}/Z$; the BW identification is Paper 42's open O0). An alternative choice $H_{\text{local}} = D_W$ (wedge-restricted Camporesi–Higuchi Dirac) has half-integer spectrum, so its operator-level period lift is $e^{i2\pi D_W} = -I$ (the spinor double cover) rather than the $+I$ of the odd-integer K_α^W; the conjugation flow still closes for both (the global $-I$ cancels), but only K_α^W lifts the period to the identity operator, so the framework's intrinsic Dirac is not the canonical local Hamiltonian for the wedge KMS state at $\beta = 2\pi$. The L1-tighten closure is for the wedge state built from the chosen K_α^W; whether that state is the restriction of a Camporesi–Higuchi vacuum is the open falsifier.

The L1-tighten implementation extends `geovac/modular_hamiltonian.py` with the `TomitaModularStructure` class (load-bearing BW-γ) and accessor methods (`tomita_structure`, `k_alpha_geometric`, `k_tomita`, `compare_alpha_vs_tomita`) on the `ModularHamiltonian` class. Test coverage: `tests/test_modular_hamiltonia` extended with 26 fast + 2 slow new tests at the time (the file now holds 70 tests as of 2026-09-02: 66 pass, 4 slow-marked and skipped by default; 0 regressions of the 41 L1 tests). Computational driver: `debug/l1_tighten_tomita_compute.py`. Machine-readable results: `debug/data/l1_tighten_tomita_results.j` Closure memo: `debug/l1_tighten_tomita_results_memo.md`.

9 Marcolli–van Suijlekom Placement: Instance + GeoVac-Original Extension

The construction of Sec. 3 is an *instance* of the gauge networks framework of Marcolli and van Suijlekom [6] (with the Perez-Sanchez correction [11] that the continuum limit is Yang–Mills without Higgs) at the data level, together with three GeoVac-original structural extensions: the forced-graph claim (Bertrand's theorem × Fock projection forces S^3), the master Mellin engine $\mathcal{M}[\mathrm{Tr}(D^k e^{-tD^2})]$ at $k \in \{0, 1, 2\}$ classifying the framework's transcendentals, and the cosmic-Galois layer U^* constructed in Paper 56 [65]. MvS 2014 does not predict a Tannakian dual; the U^* sits on top of the gauge-network scaffolding as additional structure, not as an MvS corollary. The key parallels at the data level:

- *Vertex algebra.* Marcolli–vS take $\mathcal{A}_V = C(V)$ on a vertex set V; we take $\mathcal{A}_{\text{GV}} = \mathbb{C}^{V_{\text{Fock}}}$ on the Fock-projected S^3 vertex set (Sec. 3).
- *Edge connection.* Marcolli–vS take a connection on edges valued in a compact unitary group (edge maps in $U(N)$); Paper 25 takes $G = U(1)$ and Paper 30 takes $G = \mathrm{SU}(2)$.
- *Spectral action.* Marcolli–vS show that the spectral action of the resulting almost-commutative spectral triple equals a Wilson lattice gauge action on the connection; Paper 30 verifies this explicitly for SU(2) at $n_{\max} = 3, 4$.
- *Continuum limit.* Per Perez-Sanchez, the continuum limit of the gauge-network spectral action is Yang–Mills without a Higgs sector; Paper 28's heat-kernel Seeley–DeWitt expansion is the spectral-action expansion in this limit.

Observation 4 (Lineage placement)**.** *Each structural ingredient of the GeoVac spectral triple has published precedent: finite almost-commutative spectral triples ([19, 20, 21]), graph spectral triples ([6, 10]), spectral truncations ([12, 14, 15]), and the Standard Model spectral action ([5]). What is not duplicated in the literature is the α observation (K matches α^{-1} to 4.8×10^{-7}; the root of the cubic $1/\alpha = K - \alpha^2$ matches to 8.8×10^{-8}) from a discrete spectral construction; that remains the original empirical Observation of Paper 2.*

This is part of what the working hypothesis WH1 in the project register rests on (working-hypothesis register; its commutative-factor convergence leg proved as of 2026-06-10, the almost-commutative structure's convergence open): the structural framing is in the published lineage; the empirical claim is not duplicated.

10 Open Questions

The construction of Sec. 3 and the audit of Sec. 4 leave the following structural questions genuinely open.

Q1. Order one beyond rank 1. **[OPEN]** The order-one condition holds vacuously for the diagonal $\mathcal{A}_{\mathrm{GV}}$ (whose commutators with D_{GV} vanish, Proposition 2), *fails* at finite $n_{\max} \geq 2$ on the Fock/Weyl model of the operator-system representative (Table 1) and, from $n_{\max} = 1$, on the true window (audit item 6), and non-trivially constrains the rank-1 non-abelian extension $\mathcal{A}_{\mathrm{GV}} \otimes M_2(\mathbb{C})$ of Paper 30 in the gauge-network setting. At rank ≥ 2 (e.g., $M_3(\mathbb{C})$ or sums of matrix algebras as in the Standard Model triple), order-one becomes a strong constraint and typically requires modification of the Dirac operator (*first-order condition* $\Rightarrow$ Higgs sector in [5]). Whether GeoVac admits a natural rank-≥ 2 extension—and what its spectral action would predict—is an open question. The Bargmann–Segal S^5 HO triple of Paper 24 was probed for an SU(3) extension in Sprint 5 (project development-frontier log) with negative outcome; that is one negative data point, not the full picture. The electroweak slice $\mathcal{A}_F = \mathbb{C} \oplus \mathbb{H}$ (rank up to 2 via $\mathbb{H} \cong M_2(\mathbb{C})$ over $\mathbb{C}$, a real $*$-subalgebra) is the most reachable test, and Sprint H1 addresses it; see Sec. 8.2 below for the verdict.

Q2. Real structure at finite $n_{\max}$ (resolved for the J-relations; order zero/one open). **[PANEL-VERIFIED]** Proposition 4 establishes a real structure on the continuum spinor bundle that restricts to the truncation. The axiom audit of Sec. 4 pins the finite-$n_{\max}$ behaviour down to machine precision: the restricted $J_{\mathrm{GV}}^{(n_{\max})}$ satisfies $J^2 = -\mathbb{1}$ (*not* $+\mathbb{1}$), $JD = +DJ$ and $J\mathcal{O}J^{-1} = \mathcal{O}$ bit-exactly at every $n_{\max}$ examined in the audit (`test_real_structure`), so for these three relations the finite-versus-limit question is closed in favour of finite-$n_{\max}$ exactness. No KO-dimension label is attached at finite cutoff: the chirality-doubled model truncation is even (Proposition 5(b)) with sign triple $(-,+,+)$, the true window that converges is odd like the continuum (measured, $n_{\max} \leq 3$; Proposition 5(c)) with only the pair $(\varepsilon, \varepsilon') = (-,+)$ defined, and the KO-3 label belongs to the continuum factor. **[MEASURED]** **[OPEN]** The order-zero and order-one conditions on the Fock/Weyl model of the operator-system representative fail at finite $n_{\max}$ with residuals that grow with the cutoff (order zero over $n_{\max} = 2, 3, 4$, order one over $n_{\max} = 2, 3$). **[MEASURED]** On the true Camporesi–Higuchi window of the convergence theorem the three J-relations hold exactly and order zero and order one fail from $n_{\max} = 1$, with residuals growing toward the generators' sup-norm scale, identified as a Toeplitz boundary term confined to the top Dirac levels (audit item 6, `tests/test_true_window_real_structure.py`). Open: the mechanism on the model, and the sense (if any) in which the continuum identities are recovered — on the true window the norm residual does not decay, while its normalised Hilbert–Schmidt weight falls over $n_{\max} = 2$–5 (measured in the audit driver, not test-pinned).

Q3. KO-dimension and the four-way coincidence. **[OPEN]** Working hypothesis WH4 asks: is the four-way S^3 coincidence (Sec. 1) forced by the KO-dimension 3 structure of the present triple's continuum factor, or is it independent geometric data? The relevant test is whether any other rank-1

spectral triple of KO-dimension 3 fails to produce all four roles. S^3 is unique among compact rank-1 non-abelian simply-connected groups, so the question reduces to: within the Marcolli–vS gauge-network setting, does KO-dimension 3 + non-abelian fiber + non-trivial Hopf bundle uniquely select S^3? The lineage literature does not appear to address this directly.

Q4. Spectral-action selection of Λ. **[OPEN]** Paper 28's spectral-action computation produced an exact two-term form $\mathrm{Tr}\, e^{-D^2/\Lambda^2} = (\sqrt{\pi}/2)\Lambda^3 - (\sqrt{\pi}/4)\Lambda + O(e^{-\pi^2/\Lambda^2})$ but did not autonomously select Λ. Setting the trace equal to K/π gave a closed-form Cardano solution $\Lambda_\infty = 3.7102\ldots$ (50 dps), which was tautological in the sense that it was determined by demanding the equality. Whether there is a structural principle that selects Λ is open; if so, K/π would become a derived quantity rather than a coincidence. This is the most direct route by which Paper 2 could be promoted from observation to derivation, and it is also the one most resistant to incremental progress.

Q5. Higher-form structure (Breit / rank-2 connections). **[OPEN]** Paper 25 §VII.B asked whether the Breit interaction (rank-2 in electron-electron coupling) lifts the $U(1)$ 1-form picture to a higher-form gauge theory. In spectral-triple language this is the question whether the GeoVac triple admits a natural extension to a rank-2 fiber that produces the Breit retarded integrals. Paper 30's SU(2) result is rank 1 (vector); rank 2 (tensor) is a distinct extension; the answer is unknown.

Q6. Convergence beyond the commutative factor. **[OPEN]** Theorem 5 proves state-space convergence of the commutative factor on the true window, under the translation seminorm. Two convergence statements are not proved: the same convergence in the Dirac-commutator seminorm, and convergence of the almost-commutative (gauge-network, $\mathcal{A} \otimes M_n(\mathbb{C})$) structure that WH1 names (Remark 46).

11 Conclusion

[INTERNAL THEOREM] This paper has constructed an explicit finite spectral triple $(\mathcal{A}_{\mathrm{GV}}, \mathcal{H}_{\mathrm{GV}}, D_{\mathrm{GV}})$ on the Fock-projected S^3 graph and audited it against the standard Connes axioms (Table 1, Theorem 2). The diagonal triple is a spectral, not a metric, object ($[D_{\mathrm{GV}}, \pi(a)] = 0$); the metric role and the continuum limit belong to the operator-system representative on the true Camporesi–Higuchi window (Paper 38), metrized by the left-translation seminorm: convergence of that commutative factor is proved (explicit asymptotic rate **[INTERNAL THEOREM + MEASURED]**), while Dirac-seminorm convergence and convergence of the almost-commutative structure are **[OPEN]** (Q6). At finite cutoff, on the Fock/Weyl model objects, the three J-relations hold exactly, the model truncation is even (sign triple $(-,+,+)$, no KO label), and order zero and order one fail with residuals that grow with the cutoff (**[MEASURED]**, mechanism open); on the true window the three J-relations also hold exactly and order zero and order one also fail, by a Toeplitz boundary term of the compression (**[MEASURED]**). **[OBSERVATION]** Papers 25, 28, 30, 31 are read as projections of the resulting structure (Table 3), Papers 25 and 30 through the gauge-network Dirac rather than D_{GV}. The construction sits inside the published Marcolli–van Suijlekom gauge-network framework with the Perez-Sanchez correction. The combination rule $K = \pi(B + F - \Delta)$ is now phrased as a sum across three structurally distinct sectors of the same triple (Observation 1), without changing its status as an observation. **[OBSERVATION]** The Coulomb/HO asymmetry of Paper 31 is the statement that the GeoVac triple's continuum limit is one specific real odd KO-dimension-3 triple, not a generic class (Observation 2); its true-window truncation is

odd like the limit (measured, $n_{\max} \leq 3$), and only the chirality-doubled model truncation is even (Proposition 5).

Four things this paper did not do. It did not derive α; Paper 2 stays in Observations. It did not claim the Standard Model; GeoVac is rank 1, the SM is higher rank. It did not prove that the four-way S^3 coincidence is forced by KO-dimension 3; that is open question Q3. It did not identify a Bisognano–Wichmann modular flow at finite cutoff: the finite-cutoff period closure is automatic for the chosen integer-spectrum generator, and deriving the generator from a wedge-restricted state is open (Remark 43).

Three things this paper did do. It made the spectral-triple framing explicit and auditable. It placed the framework in a published mathematical lineage. It made the open questions about α, the fiber rank, and the four-way coincidence into precise spectral-triple questions instead of free-floating conjectures.

References

[1] A. Connes, "Noncommutative geometry and reality," *J. Math. Phys.* **36**, 6194–6231 (1995).

[2] A. Connes and M. Marcolli, *Noncommutative Geometry, Quantum Fields and Motives*, AMS Colloquium Publications **55** (2008).

[3] W. D. van Suijlekom, *Noncommutative Geometry and Particle Physics*, Mathematical Physics Studies, Springer (2015).

[4] L. Dabrowski and G. Dossena, "Product of real spectral triples," *Int. J. Geom. Methods Mod. Phys.* **8**, 1833–1848 (2011), arXiv:1011.4456.

[5] A. H. Chamseddine and A. Connes, "Spectral action principle," *Commun. Math. Phys.* **186**, 731–750 (1997); "Why the Standard Model," *J. Geom. Phys.* **58**, 38–47 (2008).

[6] M. Marcolli and W. D. van Suijlekom, "Gauge networks in noncommutative geometry," *J. Geom. Phys.* **75**, 71–91 (2014), arXiv:1301.3480.

[7] F. Fathizadeh and M. Marcolli, "Periods and motives in the spectral action of Robertson–Walker spacetimes," *Comm. Math. Phys.* **356**, 641–671 (2017), arXiv:1611.01815.

[8] P. Deligne, "Le groupe fondamental unipotent motivique de $\mathbb{G}_m - \mu_N$, pour $N = 2, 3, 4, 6$ ou 8," *Publ. Math. IHES* **112**, 101–141 (2010).

[9] C. Glanois, "Motivic unipotent fundamental groupoid of $\mathbb{G}_m \setminus \mu_N$ for $N = 2, 3, 4, 6, 8$ and Galois descents," arXiv:1411.4947 (v2, 2 Sep 2015); published version *J. Number Theory* **160**, 334–384 (2016).

[10] C. I. Perez-Sanchez, "Bratteli networks and the Spectral Action on quivers," arXiv:2401.03705 (2024).

[11] C. I. Perez-Sanchez, "Comment on 'Gauge networks in noncommutative geometry',", arXiv:2508.17338 (2025).

[12] A. Connes and W. D. van Suijlekom, "Spectral truncations in noncommutative geometry and operator systems," *Commun. Math. Phys.* **383**, 2021–2067 (2021).

[13] W. D. van Suijlekom, "Gromov–Hausdorff convergence of state spaces for spectral truncations," *J. Geom. Phys.* **162**, 104075 (2021), arXiv:2005.08544.

[14] E.-M. Hekkelman, "Truncated geometry on the circle," *Lett. Math. Phys.* **112**, 20 (2022), arXiv:2111.13865.

[15] E.-M. Hekkelman and E. A. McDonald, "A noncommutative integral on spectrally truncated spectral triples, and a link with quantum ergodicity," *J. Funct. Anal.* (2025, published online, ScienceDirect PII S0022123625003362); arXiv:2412.00628.

[16] W. D. van Suijlekom, "A generalization of K-theory to operator systems," arXiv:2409.02773 (2024).

[17] Y. Gaudillot-Estrada and W. D. van Suijlekom, "Convergence of spectral truncations for compact metric groups," *Int. Math. Res. Not. (IMRN)* **2025** (2025), rnaf197; arXiv:2310.14733.

[18] F. Latrémolière, "Spectral continuity of almost commutative manifolds for the C^1 topology on Riemannian metrics," arXiv:2603.19128 (2026).

[19] T. Krajewski, "Classification of finite spectral triples," *J. Geom. Phys.* **28**, 1–30 (1998).

[20] M. Paschke and A. Sitarz, "Discrete spectral triples and their symmetries," *J. Math. Phys.* **39**, 6191–6205 (1998).

[21] B. Ćaćić, "Real structures on almost-commutative spectral triples," *Lett. Math. Phys.* **103**, 793–816 (2013), arXiv:1209.4832.

[22] R. Camporesi and A. Higuchi, "On the eigenfunctions of the Dirac operator on spheres and real hyperbolic spaces," *J. Geom. Phys.* **20**, 1–18 (1996).

[23] P. B. Gilkey, *Invariance Theory, the Heat Equation, and the Atiyah–Singer Index Theorem*, 2nd ed., CRC Press, Boca Raton, 1995.

[24] D. V. Vassilevich, "Heat kernel expansion: user's manual," *Phys. Rep.* **388**, 279–360 (2003), arXiv:hep-th/0306138.

[25] J. Karamata, *Theory and Applications of Stieltjes Integrals*, SANU Posebna izdanja CLIV, Belgrade, 1949.

[26] J. Korevaar, *Tauberian Theory: A Century of Developments*, Grundlehren der mathematischen Wissenschaften **329**, Springer, Berlin, 2004.

[27] G. Tenenbaum, *Introduction to Analytic and Probabilistic Number Theory*, 3rd ed., Graduate Studies in Mathematics **163**, American Mathematical Society, Providence, RI, 2015.

[28] GeoVac Project, "Angular Momentum as Information Geometry: Isotropic Packing and the Recurring Angular Cross-Section," Paper 0 (2026).

[29] GeoVac Project, "The Fine Structure Constant as a Spectral Coincidence on the Hopf Fibration," Paper 2 (2026; observation status, May 2026).

[30] GeoVac Project, "The Dimensionless Vacuum: Recovering the Schrödinger Equation on a Scale-Invariant Label Lattice," Paper 7 (2026).

[31] GeoVac Project, "Structurally Sparse Qubit Hamiltonians from Spectral Graph Theory," Paper 14 (2026).

[32] GeoVac Project, "Spectral-Geometric Exchange Constants: Projecting Discrete Spectra onto Continuous Manifolds," Paper 18 (2026).

[33] GeoVac Project, "Potential-Independent Angular Sparsity for Qubit Hamiltonians in Spherical Fermion Systems," Paper 22 (2026).

[34] GeoVac Project, "Nuclear Shell Model Hamiltonians on the Hyperspherical Lattice," Paper 23 (2026).

[35] GeoVac Project, "The Bargmann–Segal Lattice: π-Free Discretization of the Harmonic Oscillator on S^5," Paper 24 (2026).

[36] GeoVac Project, "The Hopf Graph as a Lattice Gauge Structure: A Framework Observation," Paper 25 (2026).

[37] GeoVac Project, "QED on S^3: Spectral Geometry of Perturbative Transcendentals," Paper 28 (2026).

[38] GeoVac Project, "SU(2) Wilson Lattice Gauge Structure on the GeoVac Hopf Graph: The Non-Abelian Sibling of Paper 25," Paper 30 (2026).

[39] GeoVac Project, "The Universal/Coulomb Partition of the GeoVac Spectral Triple," Paper 31 (2026).

[40] GeoVac Project, "GeoVac as a Two-Layer Framework: Projection Taxonomy and the Empirical Matches Catalog," Paper 34 (2026).

[41] GeoVac Project, "Time as Projection: The Klein-Gordon Spectrum on $S^3 \times \mathbb{R}$ and the Algebraic / Observation Split," Paper 35 (2026).

[42] GeoVac Project, "Bound-State QED on S^3: One-Loop Closure of the Hydrogen Lamb Shift," Paper 36 (2026; observation status).

[43] J. Loutey, "State-space Gromov–Hausdorff convergence of truncated tensor-product Camporesi–Higuchi spectral triples on $S^3 \times S^3$," Paper 39 (2026; companion to Paper 38), DOI via Zenodo.

[44] J. J. Bisognano and E. H. Wichmann, "On the duality condition for a Hermitian scalar field," *J. Math. Phys.* **16**, 985–1007 (1975).

[45] J. J. Bisognano and E. H. Wichmann, "On the duality condition for quantum fields," *J. Math. Phys.* **17**, 303–321 (1976).

[46] G. L. Sewell, "Quantum fields on manifolds: PCT and gravitationally induced thermal states," *Annals of Physics* **141**, 201–224 (1982).

[47] J. B. Hartle and S. W. Hawking, "Path-integral derivation of black-hole radiance," *Phys. Rev. D* **13**, 2188–2203 (1976).

[48] W. G. Unruh, "Notes on black-hole evaporation," *Phys. Rev. D* **14**, 870–892 (1976).

[49] E. Witten, "APS Medal for Exceptional Achievement in Research: Invited article on entanglement properties of quantum field theory," *Rev. Mod. Phys.* **90**, 045003 (2018), arXiv:1803.04993.

[50] N. Bizi, C. Brouder, and F. Besnard, "Space and time dimensions of algebras with applications to Lorentzian noncommutative geometry and quantum electrodynamics," *J. Math. Phys.* **59**, 062303 (2018), arXiv:1611.07062.

[51] K. van den Dungen, "Krein spectral triples and the fermionic action," *Math. Phys. Anal. Geom.* **19**, 4 (2016), arXiv:1505.01939.

[52] G. Nieuviarts, "Emergence of Lorentz symmetry from an almost-commutative twisted spectral triple," arXiv:2502.18105 (2025).

[53] J. Loutey, *Tomita–Takesaki modular structure on truncated $SU(2)$ spectral triples: the four-witness 2π period at finite cutoff* (Paper 42 of the GeoVac series, Zenodo, 2026).

[54] J. Loutey, *Gravity from the GeoVac spectral action: continuum results and the discrete-substrate program* (Paper 51 of the GeoVac series, Zenodo, 2026).

[55] J. Loutey, *Two-body interactions from the tensor-product spectral triple: Selection rules from the algebra, coupling strengths from the metric* (Paper 54 of the GeoVac series, Zenodo, 2026).

[56] J. Loutey, *Forced and Free Content in the GeoVac Framework: An Observations Paper* (Paper 57 of the GeoVac series, Zenodo, 2026).

[57] J. Loutey, *The Generalized-Sturmian Secular Equation Block-Encodes Without a Metric* (Paper 60 of the GeoVac series, Zenodo, 2026).

[58] F. Latrémolière, "The Dual Modular Gromov–Hausdorff Propinquity and Completeness," *J. Noncommut. Geom.* **15**, 347 (2021). arXiv:1811.04534.

[59] J. Loutey, "Composed Natural Geometries for Core-Valence Diatomics: LiH and BeH^+ from Discrete Graph Topology," GeoVac Paper 17 (2026).

[60] J. Loutey, "Toward PK-Free Molecular Hamiltonians: Two-Center Sturmian Integrals and Coupled Composition," GeoVac Paper 19 (2026).

[61] J. Loutey, "State-space Gromov–Hausdorff convergence of truncated Camporesi–Higuchi spectral triples on S^3," GeoVac Paper 38 (2026).

[62] J. Loutey, "State-space Gromov–Hausdorff convergence of spectral truncations of compact semisimple Lie groups with bi-invariant metric," GeoVac Paper 40 (2026).

[63] J. Loutey, "Boundary CFT_3-on-S^3 observables and the GeoVac spectral triple: partition function and wedge KMS entropy," GeoVac Paper 50 (2026).

[64] J. Loutey, "Periods of GeoVac: a cyclotomic mixed-Tate classification of the master Mellin engine," GeoVac Paper 55 (2026).

[65] J. Loutey, "The Tannakian substrate of GeoVac," GeoVac Paper 56 (2026).

[66] J. Loutey, "The Three-Center Electron-Repulsion Integral Is an Elliptic Bessel Moment: Genus One at the Third Center in Momentum-Space Slater Theory," GeoVac Paper 59 (2026).

[67] P. R. Halmos, "Two subspaces," *Trans. Amer. Math. Soc.* **144**, 381 (1969).

[68] A. Böttcher and I. M. Spitkovsky, "A gentle guide to the basics of two projections theory," *Linear Algebra Appl.* **432**, 1412 (2010).

Axiom	Status at finite $n_{\max}$	Status in continuum limit
Faithful $*$-representation	Holds rigorously	Holds rigorously
Bounded commutators	Holds ($[D, \pi(a)] = 0$ on $\mathcal{A}_{\mathrm{GV}}$)	Holds rigorously
Compact resolvent	Auto (finite dim)	Holds (CH summability)
Finite summability	Auto	3^+-summable
Order one, $\mathcal{A}_{\mathrm{GV}}$ (diagonal)	Holds vacuously	—
Real structure J	Holds (audited)	Holds
$J^2 = -\mathbb{1}$	Holds exactly $\forall n_{\max}$	Holds
$JD = +DJ$ (truthful CH)	Holds exactly $\forall n_{\max}$	Holds
$J\mathcal{O}J^{-1} = \mathcal{O}$	Holds exactly $\forall n_{\max}$	Holds
Order zero, $\mathcal{O}_{n_{\max}}$ (model)	Fails, $n_{\max} \geq 2$ (grows)	Holds
Order one, $\mathcal{O}_{n_{\max}}$ (model)	Fails, $n_{\max} \geq 2$	Holds
J^2, JD, $J\mathcal{O}J^{-1}$ (true window)	Hold exactly, $n_{\max} \leq 4$ (pinned)	Holds
Order zero (true window)	Fails, $n_{\max} \geq 1$ (grows; boundary term)	Holds
Order one (true window)	Fails, $n_{\max} \geq 1$ (grows; boundary term)	Holds
$\mathbb{Z}_2$-grading (model)	Even ($\sigma_x \otimes \mathbb{1}$)	Absent (odd)
$\mathbb{Z}_2$-grading (true window)	None (commutant of $\mathcal{O}$ = scalars, $n_{\max} \leq 3$)	None (odd)

Table 1: Axiom audit for the GeoVac triple at signature $(3, 0)$. Rows marked $\mathcal{A}_{\mathrm{GV}}$ refer to the diagonal representative, rows marked $\mathcal{O}_{n_{\max}}$ to the Fock/Weyl *model* of the operator-system representative (Remark 1), not the true Camporesi–Higuchi window that converges; rows marked *true window* are measured on the true Camporesi–Higuchi window (audit item 6), all other finite-cutoff entries on model objects. The continuum column lists properties of the *target* triple, the round S^3 with the Camporesi–Higuchi Dirac (real, odd, KO-dimension 3, classical facts); the convergence theorem identifies the limiting state space with that target's, up to isometry, and does not by itself deliver these properties. At finite $n_{\max}$ the three J-relations hold exactly (the rows marked $\forall\, n_{\max}$ additionally follow from the construction of Proposition 4); **[MEASURED]** order zero and order one on the model fail and the failure grows (order zero in residual and failing fraction over $n_{\max} = 2, 3, 4$; order one in failing fraction, $22\% \to 44\%$ over $n_{\max} = 2, 3$; normalised by the generator norms its max residual is nearly flat, a measurement not frozen in the tests; mechanism open), and on the true window they fail from $n_{\max} = 1$ and grow, as a boundary term of the compression; the chirality-doubled model truncation is even (Proposition 5(b)), with sign triple $(-, +, +)$ and no KO label at finite cutoff, while the true window is odd (measured, $n_{\max} \leq 3$; Proposition 5(c)). A finite-cutoff Krein-carrier sign audit at BBB $(m, n) = (4, 6)$ (signature-blind; the Lorentzian programme is descoped, Paper 45) is recorded in Table 2 below.

Axiom	$(3,0)$, finite cutoff	Krein carrier, BBB $(m,n) = (4$
J^2	$-\mathbb{1}$ (bit-exact, $\forall n_{\max}$)	$+\mathbb{1}$ (bit-exact, BBB $\varepsilon = +1$)
JD	$+DJ$ (bit-exact, $\forall n_{\max}$)	$+DJ$ (bit-exact, BBB universal
$J\chi$	$J\gamma_{\mathrm{GV}} = +\gamma_{\mathrm{GV}}J$ (doubling grading)	$\{J,\chi\} = 0$ (bit-exact, BBB $\varepsilon'' =$
$J\eta$	n/a (no η in Riemannian)	$\{J,\eta\} = 0$ (bit-exact, BBB $\varepsilon\kappa =$
$\eta\chi$	n/a	$\{\eta,\chi\} = 0$ (bit-exact, BBB $\varepsilon''\kappa'$
χD	$\{\gamma_{\mathrm{GV}}, D\} = 0$ (Prop. 5(b))	$-D\chi$ (BBB universal); FAILS f
Order zero $[a, JbJ^{-1}]$	7.9×10^{-2} max entry ($n_{\max} = 3$; grows with $n_{\max}$)	same (full basis, bit-identical to
Order one $[[D,a], JbJ^{-1}]$	2.03×10^{-1} max entry ($n_{\max} = 3$)	same (full basis, bit-identical; s

Table 2: Finite-cutoff Krein-carrier sign audit at BBB $(m,n) = (4,6)$ via Sprint L2-B/C/D, 2026-05-16 (signature-blind; it does not make the extension Lorentzian in any metrically visible sense, and the Lorentzian programme is descoped, Paper 45) (`debug/l2_{b,c,d}_*_memo.md`, `debug/data/l2_d_connes_axiom_audit_31.json`). **[PANEL-VERIFIED]** Four BBB-predicted-sign axioms hold bit-exactly across the $(n_{\max}, N_t) \in \{1,2,3\} \times \{1,11,21\}$ panel. The BBB universal $\chi D = -D\chi$ axiom fails structurally on the truthful Camporesi–Higuchi D_{GV}: this is a structural feature of GeoVac, not a bug, and is the load-bearing scope finding of Sprint L2-D. Resolution path R1+R2 (truthful for axioms requiring $JD = +DJ$, offdiag CH for axioms requiring $\{\chi, D\} = 0$) mirrors the Riemannian-side R3.5/Paper 42 trade. The commutative-factor convergence theorem of WH1 (Theorem 5) is NOT re-opened by these results; the sign audit neither extends the framework to Lorentzian signature nor re-tests its Riemannian foundation.

Paper	Object in this paper's language	Sector
Paper 7	Fock projection / label set	Manifold sector[a]
Paper 22	Selection rules from $\mathcal{A}_{\mathrm{GV}}$	$\mathcal{A}$ sector (universal)
Paper 23	D-replacement: nuclear shell model	Different D, same $\mathcal{A}$
Paper 24	Different $\mathcal{A}$: S^5 holomorphic	Distinct triple (HO)
Paper 25	$U(1)$ gauge-network fluctuation[b]	Cartan torus
Paper 28	Tr $f(D_{\mathrm{GV}}^2/\Lambda^2)$	Spectral action (one loop)
Paper 30	SU(2) gauge network, M_2 fiber[b]	Non-abelian extension
Paper 31	$\mathcal{A}/D$ split (spectral data)	Structural partition

Table 3: **[OBSERVATION]** Sub-sector identification of the major framework papers as projections of the GeoVac spectral triple. [a]The continuum limit is carried by the operator-system representative on the true Camporesi–Higuchi window, $\mathcal{O}^{\mathrm{CH}}_{n_{\max}}$ (Theorem 5), not by $\mathcal{A}_{\mathrm{GV}}$. [b]Via the gauge-network (edge-hopping) Dirac of Papers 25/30, not the diagonal D_{GV}, which has no inner fluctuations.

Layer	Coulomb (this paper)	Harmonic oscillator (Paper 24)
Manifold algebra $\mathcal{A}$	Functions on S^3 Fock graph	Functions on S^5 holomorphic Hardy graph
Projection order	Second-order (scalar Laplace–Beltrami)*	First-order (complex Euler)
Spectrum	Quadratic n^2-1 (scalar LB)*	Linear $E_N = N + 3/2$
Calibration constant	π (volume slot $\mathrm{Vol}(S^2)/4$)	None (π-free in exact arithmetic)
Wilson + natural matter	SU(2), matter rep = link rep (argued)	Pure gauge only; matter needs CG
L_0 spectrum (withdrawn)	No ($L_0 \not\to \Delta_{S^3}$)	No (spectrum diagonal)

Table 4: **[OBSERVATION]** The Coulomb/HO asymmetry from Paper 31, in spectral-triple language. The two systems are distinct spectral triples sharing only the algebra type (functions on a graph) and the trace structure of the spectral action. *The rows describe the Fock projection's scalar sector; this paper's Dirac operator D_{GV} is first order with spectrum $\pm(n+3/2)$, and the order contrast of the row is between the two *projections* (conformal vs. holomorphic).

Construction	G	**Host manifold**	**Vertex Hilbert space $\mathcal{H}_v$**	$\frac{1}{4N_c}$
Paper 25	$U(1)$	S^3 (Hopf total space)	$\mathbb{C}$ (scalar)	$\frac{1}{4}$
Paper 30	SU(2)	$S^3 = \mathrm{SU}(2)$	$\mathbb{C}^2$ (Dirac)	$\frac{1}{8}$
ST-SU3	SU(3)	S^5 Bargmann–Segal	$\bigoplus_N \mathbb{C}^{\dim(N,0)}$	$\frac{1}{12}$

Table 5: The three Wilson lattice gauge constructions as instances of the Marcolli–van Suijlekom gauge-network framework on GeoVac sub-manifolds. The kinetic coefficient $\frac{1}{4N_c}$ in the rightmost column is the weak-coupling expansion coefficient of the Wilson action $S_W = \beta \sum_P \big(1 - \frac{1}{N_c}\mathrm{Re}\,\mathrm{tr}\,U_P\big)$ around the trivial vacuum, in the standard normalisation $\mathrm{tr}(T^a T^b) = \frac{1}{2}\delta^{ab}$. **[SYMBOLIC + MEASURED]** The three rows verify this coefficient independently: Paper 30 §5.2 and Sprint ST-SU3 §4 to machine precision; the abelian Paper 25 case is the trivial $N_c = 1$ limit of either non-abelian construction.

$n_{\max}$	$\dim \mathcal{H}_{\mathrm{GV}}$	$\dim \mathcal{H}_{\mathrm{total}}$	$\|\Delta\|_{\mathrm{op}}$	$\|\Delta\|_F$	spec(Δ)
1	4	32	2.000	8.000	$\{-2{:}8,\ 0{:}16,\ +2{:}8\}$
2	16	128	2.000	16.000	$\{-2{:}32,\ 0{:}64,\ +2{:}32\}$
3	40	320	2.000	25.298	$\{-2{:}80,\ 0{:}160,\ +2{:}80\}$

Table 6: Tensor-product chirality residual $\Delta = \gamma_{\mathrm{GV}} \otimes \not{K}_F - \not{K}_{\mathrm{GV}} \otimes \gamma_F$ at $n_{\max} \in \{1,2,3\}$, computed in `debug/g3c_tensor_chirality.py` from the production γ_{GV} (G3-A, $\sigma_x \otimes \not{K}_w$) and γ_F (G3-B, KO-dim 6). Operator norm is exactly 2 at every $n_{\max}$; eigenvalue spectrum is $\{-2, 0, +2\}$ with multiplicities in ratio 1:2:1, $n_{\max}$-independent. Frobenius norm scales as $\sqrt{2 \cdot \dim \mathcal{H}_{\mathrm{total}}}$ exactly. Data `debug/data/g3c_tensor_chirality.json`.