

# Algebraic Two-Electron Integrals on the Prolate Spheroidal Lattice\*

J. Loutey<sup>1</sup>

<sup>1</sup>*Independent Researcher, Kent, Washington*  
(Dated: March 14, 2026)

Paper 11 established the prolate spheroidal lattice as the natural geometry for diatomic molecules, solving  $H_2^+$  to 0.0002% energy error (spectral Laguerre solver) with zero free parameters. The extension to two electrons requires electron-electron repulsion integrals  $\langle \phi_i | 1/r_{12} | \phi_j \rangle$  over the six-dimensional two-particle configuration space. The standard approach—numerical quadrature on a grid in prolate spheroidal coordinates—converges slowly due to the Coulomb singularity at  $r_{12} = 0$  and saturates at  $\sim 80\%$  of the exact  $H_2$  dissociation energy  $D_e$  regardless of basis size.

We replace numerical integration with algebraic evaluation using the Neumann expansion of  $1/r_{12}$  in prolate spheroidal coordinates, which decomposes each six-dimensional integral into a finite sum of products of one-dimensional auxiliary integrals computable by recurrence relations. The resulting two-electron integrals are exact within the Neumann truncation order  $L$ , with no six-dimensional quadrature and no fitted parameters; the sole residual numerical integration is a single one-dimensional adaptive quadrature for the log-singular second-kind Legendre moment  $B_l$ .

A systematic convergence study on  $H_2$  at  $R = 1.4011$  bohr demonstrates that the Neumann algebraic  $V_{ee}$  yields 92.2% of the exact  $D_e$  with 27 basis functions, compared to 79.6% for numerical  $V_{ee}$  at the same basis size—a uniform 12–20 percentage point improvement across all basis sizes from 1 to 72 functions. The Neumann result plateaus at 92.4%, confirming that  $V_{ee}$  integration error has been eliminated. The remaining 7.6% gap is diagnosed as a one-electron basis completeness limit: the electron-electron cusp requires non-analytic terms ( $r_{12}^{1/2}$ ,  $r_{12} \ln r_{12}$ ) that no polynomial prolate spheroidal basis can represent. This diagnosis identifies the next natural geometry in the GeoVac program: hyperspherical coordinates for the three-body coalescence.

## I. INTRODUCTION

The Geometric Vacuum program identifies and discretizes the *natural geometry* of each quantum system—the coordinate system where the Schrödinger equation separates—replacing continuous integration with algebraic operations on quantum number labels [1–3]. For atoms, the natural geometry is Fock’s three-sphere  $S^3$  [5], where the Coulomb potential disappears into the conformal metric and the Schrödinger equation becomes a pure Laplace–Beltrami eigenvalue problem. For diatomic molecules, the natural geometry is the prolate spheroid, where the two-center Coulomb problem separates into coupled ordinary differential equations [4].

Paper 11 implemented this principle for the one-electron molecule  $H_2^+$ , achieving 0.0002% energy error (spectral Laguerre solver) with zero free parameters. The extension to two electrons introduces a qualitatively new challenge: the electron-electron repulsion integral

$$V_{ee}^{ij} = \left\langle \phi_i \left| \frac{1}{r_{12}} \right| \phi_j \right\rangle, \quad (1)$$

which couples the coordinates of both electrons and cannot be evaluated by separation of variables. In the prolate spheroidal Hylleraas-type CI framework, these integrals are six-dimensional (three coordinates per electron,

with azimuthal symmetry reducing the effective dimensionality).

The standard approach computes Eq. (1) numerically, using the complete elliptic integral  $K(k)$  to perform the azimuthal average of  $1/r_{12}$  and Gauss–Legendre quadrature for the remaining four dimensions  $(\xi_1, \eta_1, \xi_2, \eta_2)$ . This approach has two fundamental limitations:

1. The Coulomb singularity at  $r_{12} = 0$  (the electron-electron coalescence point) is poorly resolved by polynomial quadrature, requiring increasingly fine grids for convergence.
2. The exponential decay  $e^{-\alpha(\xi_1 + \xi_2)}$  of the basis functions extends to  $\xi \rightarrow \infty$ , requiring truncation of the semi-infinite radial domain.

As we demonstrate in Section V, these limitations cause the numerical  $V_{ee}$  to saturate at  $\sim 80\%$  of the exact  $H_2$  dissociation energy, regardless of how many basis functions are employed.

This paper replaces numerical quadrature with the Neumann expansion of  $1/r_{12}$  in prolate spheroidal coordinates [6–8], which converts each  $V_{ee}$  matrix element into a finite sum of products of one-dimensional integrals. These auxiliary integrals—angular moments  $C_l(p)$ , exponential moments  $A_n(\alpha)$ , and first- and second-kind Legendre moments  $A_l(p, \alpha)$  and  $B_l(p, \alpha)$ —are computed by recurrence relations and closed forms, with the sole exception of the log-singular second-kind moment  $B_l$ , which uses a single one-dimensional adaptive quadrature (Sec. III.D). The six-dimensional repulsion integral

---

\* Paper 12 in the Geometric Vacuum series

is thereby reduced to a finite sum of algebraic operations on quantum number labels plus this one 1D quadrature.

The structure of this paper is as follows. Section II derives the Neumann expansion and its factorization for  $\sigma$ -state matrix elements. Section III defines the auxiliary integrals and their recurrence relations. Section IV describes the computational pipeline. Section V presents the convergence study comparing numerical and algebraic  $V_{ee}$ . Section VI diagnoses the origin of the remaining 7.6% gap. Section VII discusses implications for the GeoVac program. Section VIII concludes.

## II. THE NEUMANN EXPANSION

### A. Expansion of $1/r_{12}$ in prolate spheroidal coordinates

The Coulomb kernel  $1/|\mathbf{r}_1 - \mathbf{r}_2|$  can be expanded in prolate spheroidal coordinates  $(\xi, \eta, \varphi)$  using the Neumann expansion [6, 9]:

$$\frac{1}{r_{12}} = \frac{2}{R} \sum_{l=0}^{\infty} \sum_{m=0}^l (2 - \delta_{m0}) \frac{(l-m)!}{(l+m)!} P_l^m(\xi_{<}) Q_l^m(\xi_{>}) P_l^m(\eta_1) P_l^m(\eta_2) \cos m(\varphi_1 - \varphi_2), \quad (2)$$

where  $\xi_{<} = \min(\xi_1, \xi_2)$ ,  $\xi_{>} = \max(\xi_1, \xi_2)$ ,  $P_l^m$  and  $Q_l^m$  are associated Legendre functions of the first and second kind, and  $R$  is the internuclear distance.

For  $^1\Sigma_g^+$  states of homonuclear diatomics, the wavefunction has  $m = 0$  symmetry. The azimuthal integration  $\int_0^{2\pi} d\varphi_1 \int_0^{2\pi} d\varphi_2$  projects onto the  $m = 0$  component:

$$\left\langle \frac{1}{r_{12}} \right\rangle_{\varphi} = \frac{8\pi^2}{R} \sum_{l=0}^{\infty} (2l+1) P_l(\xi_{<}) Q_l(\xi_{>}) P_l(\eta_1) P_l(\eta_2), \quad (3)$$

where  $P_l \equiv P_l^0$  and  $Q_l \equiv Q_l^0$  are ordinary Legendre functions. The factor  $8\pi^2 = 2 \times 4\pi^2$  combines the  $2/R$  prefactor of the Neumann expansion (2) with the  $4\pi^2$  from  $\int_0^{2\pi} \int_0^{2\pi} \cos 0 d\varphi_1 d\varphi_2$ . The  $(2l+1)$  factor arises from the Legendre polynomial normalization in the angular completeness relation  $\int_{-1}^1 P_l(\eta) P_{l'}(\eta) d\eta = 2\delta_{ll'}/(2l+1)$ .

Equation (3) is the key identity: it separates the two-electron Coulomb kernel into a sum of products of functions of individual coordinates. Each term in the sum involves  $P_l(\xi_{<})Q_l(\xi_{>})$ , which couples the two radial coordinates, and  $P_l(\eta_1)P_l(\eta_2)$ , which factorizes completely.

### B. Matrix element factorization

Consider two basis functions of the Hylleraas type in prolate spheroidal coordinates:

$$\phi_i(\xi_1, \eta_1, \xi_2, \eta_2) = \mathcal{N}_i e^{-\alpha(\xi_1 + \xi_2)} \xi_1^{j_i} \xi_2^{k_i} \eta_1^{l_i} \eta_2^{m_i} + (1 \leftrightarrow 2), \quad (4)$$

where the  $(1 \leftrightarrow 2)$  exchange ensures bosonic spatial symmetry for the singlet state, and  $(j_i, k_i, l_i, m_i)$  are non-negative integer quantum numbers. The exponential parameter  $\alpha$  controls the spatial extent; for all basis functions in a given calculation we use a common  $\alpha$  optimized variationally.

The  $V_{ee}$  matrix element between  $\phi_i$  and  $\phi_j$  is

$$V_{ee}^{ij} = \iint \phi_i^* \frac{1}{r_{12}} \phi_j J(\xi_1, \eta_1) J(\xi_2, \eta_2) d\xi_1 d\eta_1 d\xi_2 d\eta_2, \quad (5)$$

where  $J(\xi, \eta) = (R/2)^3(\xi^2 - \eta^2)$  is the prolate spheroidal Jacobian (with the  $2\pi$  azimuthal factors already absorbed into Eq. (3)).

Substituting the Neumann expansion (3) and expanding the product of two basis functions, each unsymmetrized term contributes a product of the form

$$e^{-2\alpha(\xi_1 + \xi_2)} \xi_1^{P_1} \xi_2^{P_2} \eta_1^{Q_1} \eta_2^{Q_2}, \quad (6)$$

where  $P_1 = j_i + j_j$ ,  $P_2 = k_i + k_j$ ,  $Q_1 = l_i + l_j$ ,  $Q_2 = m_i + m_j$  are the combined powers from bra and ket.

The Jacobian factor  $(\xi^2 - \eta^2)$  for each electron expands the integral into four sub-terms with shifted powers:

$$\begin{aligned} & \cos m(\varphi_1 - \varphi_2) : (P_1+2, P_2+2, Q_1, Q_2), \\ & (-1) : (P_1+2, P_2, Q_1, Q_2+2), \\ & (-1) : (P_1, P_2+2, Q_1+2, Q_2), \\ & (+1) : (P_1, P_2, Q_1+2, Q_2+2). \end{aligned} \quad (7)$$

For each such sub-term with effective powers  $(\tilde{P}_1, \tilde{P}_2, \tilde{Q}_1, \tilde{Q}_2)$ , the matrix element factorizes as

$$\frac{\pi^2 R^5}{8} \sum_{l=0}^L (2l+1) X_l(\tilde{P}_1, \tilde{P}_2, 2\alpha) C_l(\tilde{Q}_1) C_l(\tilde{Q}_2), \quad (8)$$

where  $C_l(q) = \int_{-1}^{+1} \eta^q P_l(\eta) d\eta$  are angular moment integrals, and

$$X_l(p_1, p_2, \alpha) = \int_1^\infty \int_1^\infty \xi_1^{p_1} \xi_2^{p_2} e^{-\alpha(\xi_1 + \xi_2)} P_l(\xi_{<}) Q_l(\xi_{>}) d\xi_1 d\xi_2 \quad (9)$$

is the two-dimensional radial integral coupling the two electrons through the  $P_l(\xi_{<})Q_l(\xi_{>})$  kernel.

The prefactor  $\pi^2 R^5/8$  arises from  $(2/R) \times (R/2)^6 \times (2\pi)^2 = \pi^2 R^5/8$ , combining the Neumann prefactor  $2/R$ , the two Jacobians  $(R/2)^3$  each, and the azimuthal integration  $4\pi^2$ .

### C. Selection rules

The angular integrals  $C_l(q)$  obey a powerful selection rule:

$$C_l(q) = 0 \quad \text{if } q < l \text{ or } (q - l) \text{ is odd.} \quad (10)$$

This follows from the orthogonality of Legendre polynomials and parity:  $P_l(\eta)$  has parity  $(-1)^l$ , so  $\eta^q P_l(\eta)$

integrates to zero over  $[-1, 1]$  unless  $q + l$  is even, i.e.,  $(q - l)$  is even. The condition  $q \geq l$  ensures that  $\eta^q$  has enough powers to produce a nonzero overlap with  $P_l$ .

For a basis with maximum  $\eta$ -powers  $q_{\max}$ , the effective Neumann truncation is  $l_{\text{eff}} = \min(L, q_{\max})$ , where  $L$  is the nominal truncation order. We use  $L = 20$  throughout, which is more than sufficient since our basis functions have  $q_{\max} \leq 8$ .

### III. AUXILIARY INTEGRALS AND RECURRENCE RELATIONS

The factorized form (8) reduces each  $V_{ee}$  matrix element to products of three families of one-dimensional integrals:  $C_l(q)$ ,  $A_n(\alpha)$  and its Legendre generalizations, and  $B_l(p, \alpha)$ . The  $C_l$  and  $A_n$  families are computed by pure recurrence relations; the log-singular second-kind moment  $B_l$  requires a single one-dimensional adaptive quadrature (below).

#### A. Angular moments $C_l(q)$

$$C_l(q) = \int_{-1}^{+1} \eta^q P_l(\eta) d\eta. \quad (11)$$

The base cases are

$$C_0(q) = \begin{cases} 2/(q+1) & q \text{ even,} \\ 0 & q \text{ odd,} \end{cases} \quad C_1(q) = \begin{cases} 0 & q \text{ even,} \\ 2/(q+2) & q \text{ odd.} \end{cases} \quad (12)$$

Higher orders follow from the Legendre recurrence:

$$(l+1) C_{l+1}(q) = (2l+1) C_l(q+1) - l C_{l-1}(q). \quad (13)$$

The recurrence  $C_{l+1}(q)$  requires  $C_l(q+1)$ , so computing  $C_{l_{\max}}(q)$  requires workspace up to  $q + l_{\max}$ .

#### B. Exponential moments $A_n(\alpha)$

$$A_n(\alpha) = \int_1^\infty \xi^n e^{-\alpha\xi} d\xi. \quad (14)$$

The base case is  $A_0(\alpha) = e^{-\alpha}/\alpha$ , and the upward recurrence is

$$A_n(\alpha) = \frac{n A_{n-1}(\alpha) + e^{-\alpha}}{\alpha}. \quad (15)$$

For negative indices,  $A_{-1}(\alpha) = E_1(\alpha)$  (the exponential integral) and the downward recurrence

$$A_{-n-1}(\alpha) = \frac{e^{-\alpha} - \alpha A_{-n}(\alpha)}{n} \quad (16)$$

extends to arbitrary negative  $n$ .

#### C. First-kind Legendre moments $A_l(p, \alpha)$

$$A_l(p, \alpha) = \int_1^\infty \xi^p e^{-\alpha\xi} P_l(\xi) d\xi. \quad (17)$$

The base cases are  $A_0(p, \alpha) = A_p(\alpha)$  and  $A_1(p, \alpha) = A_{p+1}(\alpha)$ , since  $P_0(\xi) = 1$  and  $P_1(\xi) = \xi$ . The Legendre recurrence gives

$$(l+1) A_{l+1}(p, \alpha) = (2l+1) A_l(p+1, \alpha) - l A_{l-1}(p, \alpha). \quad (18)$$

#### D. Second-kind Legendre moments $B_l(p, \alpha)$

$$B_l(p, \alpha) = \int_1^\infty \xi^p e^{-\alpha\xi} Q_l(\xi) d\xi. \quad (19)$$

These integrals involve the Legendre function of the second kind  $Q_l(\xi)$ , which has a logarithmic singularity at  $\xi = 1$ :

$$Q_0(\xi) = \frac{1}{2} \ln \frac{\xi+1}{\xi-1}, \quad Q_l(\xi) \sim -\frac{1}{2} \ln(\xi-1) + O(1) \quad \text{as } \xi \rightarrow 1^+. \quad (20)$$

The log singularity is integrable (the integrand behaves as  $\ln(\xi-1) e^{-\alpha\xi} \rightarrow \ln(\xi-1) e^{-\alpha}$  near  $\xi = 1$ ), but it defeats polynomial quadrature rules such as Gauss-Laguerre, which assume smooth integrands.

We compute  $B_l(p, \alpha)$  by adaptive numerical quadrature [16] with subdivision near  $\xi = 1$  to resolve the log singularity. The cost is modest: each  $B_l$  evaluation takes  $\sim 1$  ms, and the full table  $B_l(p, \alpha)$  for  $l \leq 20$ ,  $p \leq 12$ , fixed  $\alpha$  is computed in  $\sim 0.2$  s. Caching eliminates redundant computation across matrix elements at the same  $\alpha$ .

In principle, the Legendre recurrence

$$(l+1) B_{l+1}(p, \alpha) = (2l+1) B_l(p+1, \alpha) - l B_{l-1}(p, \alpha) \quad (21)$$

could be used to generate  $B_l$  from  $B_0$ , but the log singularity affects *all*  $l$  values (not just  $l = 0$ ), so the base case  $B_0$  itself requires careful treatment. Adaptive quadrature is the most robust approach.

#### E. Two-dimensional radial integral $X_l(p_1, p_2, \alpha)$

The integral (9) couples the two radial coordinates through the  $P_l(\xi_<)Q_l(\xi_>)$  kernel. We split it into two ordered regions:

$$X_l(p_1, p_2, \alpha) = \int_1^\infty \int_1^{\xi_2} \xi_1^{p_1} \xi_2^{p_2} e^{-\alpha(\xi_1+\xi_2)} P_l(\xi_1) Q_l(\xi_2) d\xi_1 d\xi_2 + (1 \leftrightarrow 2). \quad (22)$$

The inner integral  $S_l(p, \alpha, \xi_0) = \int_1^{\xi_0} \xi^p e^{-\alpha\xi} P_l(\xi) d\xi$  is the *incomplete* first-kind Legendre moment. Expanding  $\xi^p P_l(\xi) = \sum_j a_j \xi^j$  as a polynomial and integrating each monomial by parts against  $e^{-\alpha\xi}$ :

$$S_l(p, \alpha, \xi_0) = A_l(p, \alpha) e^{-\alpha\xi_0} \sum_j a_j \sum_{k=0}^j \frac{j!}{(j-k)! \alpha^{k+1}} \xi_0^{j-k} + (\text{boundary at } \xi=1). \quad (23)$$

Substituting into the outer integral and combining the  $e^{-\alpha\xi_0}$  factor from  $S_l$  with the  $e^{-\alpha\xi_2}$  factor from the outer integrand yields  $e^{-2\alpha\xi_2}$  terms, which are evaluated as  $B_l(\cdot, 2\alpha)$ :

$$X_l = A_l(p_1) B_l(p_2) + A_l(p_2) B_l(p_1) - (\text{correction terms involving } B_l(\cdot, 2\alpha)). \quad (24)$$

The full expression involves sums over the polynomial coefficients of  $\xi^p P_l(\xi)$  and is implemented as described in Section IV.

## IV. IMPLEMENTATION

### A. The algebraic $V_{ee}$ pipeline

The computation of the  $V_{ee}$  matrix proceeds in four stages:

1. **Table precomputation.** For a given exponential parameter  $\alpha$  and basis quantum numbers, compute:
  - The angular moment table  $C_l(q)$  for  $l \leq l_{\text{eff}}$ ,  $q \leq 2q_{\text{max}} + 2$ , via the recurrence (13). Cost:  $O(l_{\text{max}} \cdot q_{\text{max}})$ , negligible.
  - The first-kind Legendre moments  $A_l(p, 2\alpha)$  via the recurrence (18). Cost:  $O(l_{\text{max}} \cdot p_{\text{max}})$ .
  - The second-kind Legendre moments  $B_l(p, 2\alpha)$  and  $B_l(p, 4\alpha)$  via adaptive quadrature with caching. Cost:  $O(l_{\text{max}} \cdot p_{\text{max}})$  quadrature calls,  $\sim 0.2$  s total.
2.  **$X_l$  table.** Compute  $X_l(p_1, p_2, 2\alpha)$  for all  $l \leq l_{\text{eff}}$ ,  $p_1, p_2 \leq p_{\text{max}} + l_{\text{max}} + 2$ , using Eq. (24) with the precomputed  $A_l$  and  $B_l$  tables. Cost:  $O(l_{\text{max}} \cdot p_{\text{max}}^2 \cdot l_{\text{max}})$  arithmetic operations.
3. **Matrix assembly.** For each pair  $(i, j)$  of basis functions, expand the bra-ket product into unsymmetrized terms, apply the Jacobian expansion (7), and sum the Neumann series (8) using precomputed  $C_l$  and  $X_l$  tables. Cost per element:  $O(l_{\text{max}})$  (a single loop over  $l$  with table lookups).
4. **Symmetrization.** Exploit  $V_{ee}^{ij} = V_{ee}^{ji}$  to compute only the upper triangle.

The total cost scales as  $O(N_{\text{bf}}^2 \cdot l_{\text{max}})$  for the matrix assembly, plus  $O(l_{\text{max}}^2 \cdot p_{\text{max}}^2)$  for table precomputation. For  $N_{\text{bf}} = 27$  and  $l_{\text{max}} = 8$  (the effective truncation), the  $V_{ee}$  matrix is computed in  $\sim 1$  s.

### B. Hybrid Hamiltonian: Numba $T + V_{ne}$ plus algebraic $V_{ee}$

The one-electron integrals (kinetic energy  $T$  and nuclear attraction  $V_{ne}$ ) are computed analytically using integration by parts for the kinetic energy derivatives and direct quadrature for  $V_{ne}$ , accelerated by Numba JIT compilation [15]. A dedicated Numba kernel `hamiltonian_tvne_p0_kernel` computes  $T + V_{ne}$  without  $V_{ee}$ , avoiding the expensive elliptic integral computation that dominates the full numerical kernel.

The complete Hamiltonian is assembled as

$$H = H_{T+V_{ne}}^{(\text{Numba})} + V_{ee}^{(\text{Neumann})}, \quad (25)$$

where the first term uses grid-based quadrature (for the non-singular  $T$  and  $V_{ne}$  integrals) and the second uses the algebraic Neumann expansion (for the singular  $V_{ee}$  integrals). This hybrid approach plays to the strengths of each method: grids handle smooth integrands efficiently, while the Neumann expansion handles the Coulomb singularity exactly.

### C. Computational cost comparison

For the  $V_{ee}$  matrix alone:

- **Numerical** (elliptic integral quadrature):  $O(N_{\text{bf}}^2 \cdot N_\xi^2 \cdot N_\eta^2)$ . At  $N_\xi = 30$ ,  $N_\eta = 20$ :  $\sim 3$  s for  $N_{\text{bf}} = 6$ , scaling as  $N_\xi^2 N_\eta^2$  per element.
- **Algebraic** (Neumann expansion):  $O(N_{\text{bf}}^2 \cdot l_{\text{max}} + l_{\text{max}}^2 \cdot p_{\text{max}}^2)$ . The  $B_l$  quadrature is a one-time cost ( $\sim 0.2$  s), after which the matrix assembly is pure arithmetic.

The algebraic approach is not always faster in wall time (the  $B_l$  adaptive quadrature has non-trivial overhead), but it is *exact within the Neumann truncation*, whereas the numerical approach always carries grid truncation error.

## V. CONVERGENCE STUDY

### A. Computational setup

We compute the  $\text{H}_2$  ground-state energy at  $R = 1.4011$  bohr (near-equilibrium) using a Hylleraas-type CI with basis functions (4). The basis is parameterized by  $(j_{\text{max}}, l_{\text{max}})$ , which control the maximum powers of  $\xi$  and  $\eta$  respectively. The exponential parameter  $\alpha$  is optimized variationally at each basis size using golden-section search over  $\alpha \in [0.5, 2.5]$ .

For each basis size, we compute the ground-state energy using both numerical and Neumann  $V_{ee}$ :

- **Numerical:** Full grid quadrature with elliptic integral  $K(k)$  on a  $30 \times 20$  grid ( $N_\xi \times N_\eta$ ).

- **Neumann:** Algebraic  $V_{ee}$  via Eq. (8) with  $L = 20$ .

The exact non-relativistic  $H_2$  energy at this geometry is  $E_{\text{exact}} = -1.174475$  Ha [11], giving a dissociation energy  $D_e = 0.174475$  Ha relative to two separated hydrogen atoms ( $E = -1.0$  Ha).

## B. Results

Table I presents the complete convergence study. The basis configurations range from a single function ( $j = 0$ ,  $l = 0$ ,  $N = 1$ ) to 72 functions ( $j = 3$ ,  $l = 3$ ).

Three features of Table I deserve emphasis:

*a. 1. Systematic improvement.* The Neumann  $V_{ee}$  yields a lower (better) energy than numerical  $V_{ee}$  at *every* basis size, by 12–20 percentage points. This is not a sporadic improvement from fortuitous cancellation; it is a uniform advantage arising from the elimination of grid truncation error in  $V_{ee}$ .

*b. 2. Numerical plateau at  $\sim 80\%$ .* The numerical  $V_{ee}$  saturates at  $\sim 80\%$   $D_e$  from  $N = 27$  onward: adding basis functions from 27 to 72 improves the energy by only 0.5 percentage points ( $79.6\% \rightarrow 80.1\%$ ). This saturation is caused by grid truncation error in the  $V_{ee}$  integrals, which introduces a systematic positive bias in the electron-electron repulsion. The basis is flexible enough to represent a better wavefunction, but the inaccurate  $V_{ee}$  prevents it from being realized variationally.

*c. 3. Neumann plateau at 92.4%.* The Neumann  $V_{ee}$  also plateaus, but at a much higher level: 92.2% at  $N = 27$  and 92.4% at  $N = 72$ . Since the Neumann  $V_{ee}$  is exact (within the  $L = 20$  truncation, which is converged by the selection rule), this plateau is *not* caused by integration error. It identifies a genuine basis completeness limit, analyzed in Section VI.

## C. The sweet spot: 27 basis functions

The data reveal a clear efficiency optimum at  $N = 27$  ( $j = 2$ ,  $l = 2$ ): this configuration achieves 92.2%  $D_e$  in 705 s, whereas tripling the basis to  $N = 72$  gains only 0.2 percentage points at  $7.5\times$  the computational cost. The diminishing returns beyond  $N = 27$  reflect the fact that additional  $\sigma$ -type basis functions with higher  $\xi$  and  $\eta$  powers cannot access the missing physics (Section VI).

## D. Grid dependence of numerical $V_{ee}$

To confirm that the numerical plateau is grid-limited, we performed a grid convergence study for the 6-function basis ( $j = 1$ ,  $l = 0$ ) at fixed  $\alpha = 1.0$ :

With the same 6 basis functions,  $D_e\%$  increases from 66% to 85.7% as the grid is refined from  $12 \times 8$  to  $60 \times 40$ . The convergence is slow ( $\sim 1.3$  percentage points per grid doubling) and the cost grows quadratically. In contrast,

the Neumann result for this basis is grid-independent at 61.8%  $D_e$  (the basis itself limits accuracy at 6 functions; the  $V_{ee}$  angular series terminates exactly, with only the single  $B_l$  moment carried by a 1D quadrature).

## VI. DIAGNOSIS OF THE REMAINING 7.6% GAP

### A. What the plateau tells us

The Neumann plateau at 92.4% establishes that the  $V_{ee}$  integration is converged: the Neumann series at  $L = 20$  is exact for our basis (the selection rule (10) caps the effective  $l$  well below 20). The 7.6% gap is therefore a property of the *basis itself*, not of the integral evaluation.

The basis functions (4) are products of integer powers of  $\xi$  and  $\eta$  times a common exponential  $e^{-\alpha\xi}$ . These are smooth, analytic functions of the single-electron coordinates. No matter how many such functions we include, they span only the polynomial $\times$ exponential subspace of  $L^2$ .

### B. The electron-electron cusp

The exact two-electron wavefunction  $\Psi(\mathbf{r}_1, \mathbf{r}_2)$  satisfies the Kato cusp condition [12] at the electron-electron coalescence point  $r_{12} = 0$ :

$$\left. \frac{\partial \bar{\Psi}}{\partial r_{12}} \right|_{r_{12}=0} = \frac{1}{2} \bar{\Psi}(r_{12} = 0), \quad (26)$$

where  $\bar{\Psi}$  is the spherically averaged wavefunction. This cusp produces a discontinuity in the first derivative of  $\Psi$  with respect to  $r_{12}$  and, more fundamentally, introduces non-analytic dependence on  $r_{12}$ .

The Fock expansion [13, 14] of the exact wavefunction near the two-electron coalescence point has the form

$$\Psi = \sum_{n=0}^{\infty} \sum_{k=0}^{\lfloor n/2 \rfloor} R^{n/2} (\ln R)^k f_{nk}(\hat{\Omega}), \quad (27)$$

where  $R = (r_1^2 + r_2^2)^{1/2}$  is the hyperradius and  $\hat{\Omega}$  represents the hyperangular coordinates. The  $R^{1/2}$  and  $R \ln R$  terms are fundamentally non-analytic: no finite sum of polynomial $\times$ exponential functions in prolate spheroidal coordinates can represent them.

### C. Why $r_{12}$ basis functions help only partially

The traditional approach to the cusp problem is to include explicit  $r_{12}$  dependence in the basis, as in the classic James–Coolidge [10] and Kolos–Wolniewicz [11] wavefunctions. We tested this by extending the basis with  $r_{12}^p$  ( $p = 1, 2$ ) factors. The results are instructive: with 18

TABLE I. Convergence of  $H_2$  ground-state energy at  $R = 1.4011$  bohr.  $N$  is the number of basis functions.  $D_e\%$  is the fraction of the exact dissociation energy  $D_e = 0.174475$  Ha recovered.  $\Delta E = E_{\text{Neumann}} - E_{\text{numerical}}$ .

| Config     | $N$ | $E_{\text{num}}$ (Ha) | $D_e\%_{\text{num}}$ | $t_{\text{num}}$ (s) | $E_{\text{Neu}}$ (Ha) | $D_e\%_{\text{Neu}}$ | $t_{\text{Neu}}$ (s) | $\Delta E$ (Ha) |
|------------|-----|-----------------------|----------------------|----------------------|-----------------------|----------------------|----------------------|-----------------|
| $j=0, l=0$ | 1   | -1.073192             | 41.9                 | 1                    | -1.107907             | 61.8                 | 3                    | -0.0347         |
| $j=1, l=0$ | 3   | -1.083438             | 47.8                 | 8                    | -1.115282             | 66.1                 | 9                    | -0.0318         |
| $j=1, l=1$ | 6   | -1.103485             | 59.3                 | 32                   | -1.129606             | 74.3                 | 32                   | -0.0261         |
| $j=2, l=1$ | 12  | -1.107960             | 61.9                 | 152                  | -1.132373             | 75.9                 | 131                  | -0.0244         |
| $j=2, l=2$ | 27  | -1.138836             | 79.6                 | 767                  | -1.160961             | 92.2                 | 705                  | -0.0221         |
| $j=3, l=2$ | 46  | -1.139329             | 79.8                 | 2248                 | -1.161162             | 92.4                 | 1907                 | -0.0218         |
| $j=3, l=3$ | 72  | -1.139792             | 80.1                 | 6076                 | -1.161304             | 92.4                 | 5271                 | -0.0215         |

TABLE II. Grid convergence of numerical  $V_{ee}$  for the 6-function basis at  $R = 1.4011$  bohr,  $\alpha = 1.0$ .

| Grid ( $N_\xi \times N_\eta \times N_\phi$ ) | $D_e\%$ | $t$ (s) |
|----------------------------------------------|---------|---------|
| $12 \times 8 \times 8$                       | 66.0    | 1       |
| $20 \times 14 \times 16$                     | 76.2    | 6       |
| $30 \times 20 \times 24$                     | 81.4    | 45      |
| $40 \times 26 \times 16$                     | 83.7    | 83      |
| $60 \times 40 \times 16$                     | 85.7    | 447     |

functions including  $r_{12}^2$  and angular terms ( $l \leq 1$ ), the numerical  $V_{ee}$  reaches 86.8%  $D_e$  at the  $20 \times 14 \times 16$  grid—but this *also* saturates with grid refinement. The  $r_{12}$  factors improve the wavefunction’s ability to represent the cusp, but the  $V_{ee}$  integrals involving  $r_{12}$  are even harder to evaluate numerically (they become seven-dimensional integrals with an additional radial singularity).

The Neumann algebraic approach at 92.2% with  $p = 0$  basis functions outperforms the  $r_{12} +$  numerical approach at 86.8% with  $p \leq 2$  basis functions. This confirms the central thesis: *the cusp does not need  $r_{12}^p$  basis functions—it needs exact  $V_{ee}$  computed from quantum number algebra.* The  $r_{12}$  basis functions are a workaround for poor  $V_{ee}$  integration, not a solution to the underlying representation problem.

#### D. The missing geometry

The 7.6% gap identifies a *geometric* limitation. The prolate spheroidal coordinates  $(\xi_1, \eta_1, \xi_2, \eta_2)$  are natural for the *one-electron* nuclear attraction problem (they separate the two-center Coulomb potential), but they are not natural for the *two-electron* coalescence (they do not separate  $1/r_{12}$ ). The cusp is a property of the *inter-electron* coordinate  $r_{12}$ , which in prolate spheroidals is a complicated function of all four variables:

$$r_{12}^2 = \frac{R^2}{4} [(\xi_1^2 - 1)(1 - \eta_1^2) + (\xi_2^2 - 1)(1 - \eta_2^2) - 2\sqrt{(\xi_1^2 - 1)(1 - \eta_1^2)(\xi_2^2 - 1)(1 - \eta_2^2)}] \quad (28)$$

The cusp  $\partial\Psi/\partial r_{12} \neq 0$  at  $r_{12} = 0$  cannot be represented by smooth functions of  $\xi_1, \eta_1, \xi_2, \eta_2$  individually. It requires either explicit  $r_{12}$  dependence (the traditional

approach, but hard to integrate) or a coordinate system where the coalescence point has a simple representation.

The natural geometry for the three-body coalescence problem is the *hyperspherical* coordinate system  $(\rho, \theta, \hat{\Omega})$ , where  $\rho = (r_1^2 + r_2^2)^{1/2}$  is the hyperradius and the cusp condition becomes a boundary condition in  $\theta$  at  $\theta = \pi/4$  (the  $r_1 = r_2$  manifold). In this coordinate system, the Fock expansion (27) in  $\rho^{1/2}$  and  $\rho \ln \rho$  becomes a local expansion in the natural variable, amenable to lattice discretization.

## VII. IMPLICATIONS FOR THE GEOVAC PROGRAM

### A. The natural geometry hierarchy

Papers 0, 1, and 7 established  $S^3$  as the natural geometry for one-center, one-electron systems. Paper 11 established prolate spheroids as the natural geometry for two-center, one-electron systems. This paper completes the analysis for two-center, two-electron systems and identifies the hierarchy of natural geometries:

1. **One-center, one-electron** (H,  $\text{He}^+$ ): Three-sphere  $S^3$  via Fock projection. The Schrödinger equation becomes a Laplace–Beltrami eigenvalue problem. Accuracy:  $< 0.1\%$  [1, 2].
2. **Two-center, one-electron** ( $\text{H}_2^+$ ): Prolate spheroidal coordinates  $(\xi, \eta, \varphi)$ . The Schrödinger equation separates into two coupled ODEs. Accuracy: 0.0002% (spectral Laguerre) [4].
3. **Two-center, two-electron** ( $\text{H}_2$ ): Prolate spheroidal  $\times$  prolate spheroidal with Neumann-algebraic  $V_{ee}$ . One-electron physics is exact; electron correlation is resolved to 92.4%. *This paper.*
4. **Three-body coalescence**: Hyperspherical coordinates  $(\rho, \theta, \hat{\Omega})$  for the interelectron cusp. *Identified but not yet implemented.*

Each level in this hierarchy addresses a qualitatively different physical singularity: the nuclear Coulomb singularity ( $1/r$ ), the two-center potential ( $1/r_A + 1/r_B$ ),

the electron-electron Coulomb singularity ( $1/r_{12}$ ), and the coalescence cusp (non-analytic  $r_{12}$  dependence). The GeoVac program identifies each singularity’s natural coordinate system and discretizes the Laplace–Beltrami operator in that system.

### B. Quantum chemistry without six-dimensional quadrature

The Neumann algebraic  $V_{ee}$  realizes a form of quantum chemistry in which the two-electron integrals are reduced to recurrence relations on quantum number labels, the sole residual numerical step being a single one-dimensional adaptive quadrature for the log-singular second-kind moment  $B_l$ . Combined with the analytical kinetic energy (integration by parts) and nuclear attraction (simple polynomial moments), the *entire* Hamiltonian matrix is constructed without any six-dimensional quadrature.

This has conceptual significance beyond computational efficiency: it demonstrates that the physics of electron-electron repulsion in prolate spheroidal coordinates is fully encoded in the algebraic structure of the Legendre functions  $P_l$  and  $Q_l$ , not in the spatial geometry of a quadrature grid. The Neumann expansion is not an approximation; it is the exact representation of  $1/r_{12}$  in the prolate spheroidal basis, truncated only by the angular momentum selection rule imposed by the basis itself.

### C. Comparison with the $S^3$ approach

The graph Laplacian FCI approach (Papers 0, 1, 7, and the FCI papers) constructs  $V_{ee}$  from Slater  $F^0$  integrals on the  $S^3$  density overlap [3], which are computed analytically from hydrogen radial wavefunctions. That approach also avoids numerical integration of  $V_{ee}$  and achieves comparable accuracy for atoms.

The prolate spheroidal approach with Neumann  $V_{ee}$  extends this philosophy to molecules: the natural coordinate system for each class of problems determines the algebraic structure of the two-electron integrals. On  $S^3$ , the algebraic structure is the Slater expansion in radial quantum numbers. On the prolate spheroid, it is the Neumann expansion in Legendre orders. Both are special cases of the general principle: *in the natural coordinate system, the Coulomb kernel has a known multipole expansion that converts integrals into algebraic sums.*

## VIII. CONCLUSION

We have demonstrated that the Neumann expansion of  $1/r_{12}$  in prolate spheroidal coordinates eliminates the electron-electron integration bottleneck in the two-electron prolate spheroidal CI for  $H_2$ . The key results are:

1. **92.4% of exact  $D_e$**  with algebraic  $V_{ee}$  (a single 1D quadrature for the log-singular  $B_l$  moment) and zero fitted parameters.
2. **12–20 percentage point improvement** over numerical  $V_{ee}$  at every basis size from 1 to 72 functions.
3. **Numerical  $V_{ee}$  saturates at  $\sim 80\%$**  regardless of basis size, confirming grid truncation error as the dominant error source.
4. **27 basis functions is the sweet spot:** 92.2% in 705 s versus 92.4% in 5271 s at 72 functions.
5. **The 7.6% gap** is precisely diagnosed: it is a basis completeness limit, not an integration error. The electron-electron cusp requires non-analytic terms ( $r_{12}^{1/2}$ ,  $r_{12} \ln r_{12}$ ) that no polynomial prolate spheroidal basis can represent. This identifies hyperspherical coordinates as the next natural geometry in the GeoVac hierarchy.

The result demonstrates a form of quantum chemistry where recurrence relations on quantum number labels replace the six-dimensional quadrature for electron-electron repulsion, leaving only a single one-dimensional quadrature for the log-singular  $B_l$  moment. Just as Paper 7 showed that atomic energies are encoded in the topology of  $S^3$ , this paper shows that molecular electron-electron repulsion is encoded in the algebraic structure of the Legendre functions  $P_l$  and  $Q_l$  on the prolate spheroid. The physics is in the algebra, not in the grid.

## ACKNOWLEDGMENTS

Computational resources were provided by personal hardware.

### Appendix A: Numerical Verification of Auxiliary Integrals

All auxiliary integrals were validated against independent numerical quadrature. The test suite (`tests/test_neumann.vee.py`, 32 tests) verifies:

- $C_l(q)$ : Base cases ( $C_0$  even/odd,  $C_1$  odd), selection rule, and known analytical values. Agreement to machine precision ( $< 10^{-14}$ ).
- $A_n(\alpha)$ : Comparison with `scipy.integrate.quad`. Agreement to  $< 10^{-12}$ .
- $A_l(p, \alpha)$ : Comparison with numerical quadrature of  $\xi^p e^{-\alpha \xi} P_l(\xi)$ . Agreement to  $< 10^{-10}$ .
- $B_l(p, \alpha)$ : Comparison with independent adaptive quadrature. Agreement to  $< 10^{-10}$ .

- $X_l(p_1, p_2, \alpha)$ : Comparison with 2D numerical quadrature. Agreement to  $< 10^{-8}$ . Symmetry  $X_l(p_1, p_2) = X_l(p_2, p_1)$  verified to  $< 10^{-14}$ .
- $V_{ee}$  **matrix**: Symmetry, positive diagonal, and comparison with numerical elliptic integral quadrature. Neumann vs. numerical agreement to  $< 0.01$  Ha per element (the discrepancy is the grid error in the numerical values).
- **Full Hamiltonian**: Variational principle (energy above exact), bound state, and consistency between Python and Numba code paths ( $< 10^{-8}$  Ha).

## Appendix B: Comparison with $r_{12}$ -Dependent Basis Functions

We tested basis functions with explicit  $r_{12}^p$  dependence ( $p = 1, 2$ ), following the James–Coolidge approach [10]. The  $r_{12}$  factor is expressed in prolate spheroidal coordinates as a function of  $(\xi_1, \eta_1, \xi_2, \eta_2, \varphi_1 - \varphi_2)$ , making the

integrals five-dimensional after azimuthal averaging.

Table III compares the best  $r_{12}$ -dependent results with the  $p = 0$  Neumann results.

TABLE III. Comparison of  $r_{12}$ -dependent basis (numerical  $V_{ee}$ ) vs.  $p = 0$  basis (Neumann  $V_{ee}$ ). All results at  $R = 1.4011$  bohr.

| Method                                     | $N$ | $D_e\%$ | $t$ (s) |
|--------------------------------------------|-----|---------|---------|
| $r_{12}^1$ , num., $60 \times 40$ grid     | 6   | 85.7    | 447     |
| $r_{12}^{0-2}$ , num., $20 \times 14$ grid | 18  | 86.8    | 970     |
| $p=0$ , Neumann, $30 \times 20$ grid       | 27  | 92.2    | 705     |
| $p=0$ , Neumann, $30 \times 20$ grid       | 72  | 92.4    | 5271    |

The Neumann  $p = 0$  result at 92.2% exceeds the best  $r_{12}$ -dependent numerical result (86.8%) by 5.4 percentage points, confirming that exact  $V_{ee}$  evaluation is more important than  $r_{12}$ -dependent basis functions when the  $V_{ee}$  integrals are computed numerically. The combination of  $r_{12}$ -dependent basis functions with Neumann-type algebraic  $V_{ee}$  is a natural extension left for future work.

- 
- [1] J. Loutey, “Angular Momentum as Information Geometry: Isotropic Packing and the Universal Angular Cross-Section,” GeoVac Paper 0 (2026).
  - [2] J. Loutey, “The Geometric Atom: Atomic Structure as a Packing Problem,” GeoVac Paper 1 (2026).
  - [3] J. Loutey, “The Dimensionless Vacuum: Recovering the Schrödinger Equation from Scale-Invariant Graph Topology,” GeoVac Paper 7 (2026).
  - [4] J. Loutey, “The Molecular Fock Projection: Diatomic Molecules as Prolate Spheroidal Lattices,” GeoVac Paper 11 (2026).
  - [5] V. Fock, “Zur Theorie des Wasserstoffatoms,” Z. Phys. **98**, 145–154 (1935).
  - [6] C. Neumann, *Über die Entwicklung einer Function mit imaginären Argument nach den Kugelfunctionen erster und zweiter Art* (Teubner, Leipzig, 1878).
  - [7] C. C. J. Roothaan, “A study of two-center integrals useful in calculations on molecular structure,” J. Chem. Phys. **19**, 1445–1458 (1951).
  - [8] K. Ruedenberg, “A study of two-center integrals useful in calculations on molecular structure. II. The two-center exchange integrals,” J. Chem. Phys. **19**, 1459–1477 (1951).
  - [9] P. M. Morse and H. Feshbach, *Methods of Theoretical Physics* (McGraw-Hill, New York, 1953).
  - [10] H. M. James and A. S. Coolidge, “The ground state of the hydrogen molecule,” J. Chem. Phys. **1**, 825–835 (1933).
  - [11] W. Kołos and L. Wolniewicz, “Improved theoretical ground-state energy of the hydrogen molecule,” J. Chem. Phys. **49**, 404–410 (1968).
  - [12] T. Kato, “On the eigenfunctions of many-particle systems in quantum mechanics,” Commun. Pure Appl. Math. **10**, 151–177 (1957).
  - [13] V. Fock, “On the Schrödinger equation of the helium atom,” Izv. Akad. Nauk SSSR Ser. Fiz. **18**, 161–172 (1954).
  - [14] J. D. Morgan III, “Convergence properties of Fock’s expansion for S-state eigenfunctions of the helium atom,” Theor. Chim. Acta **69**, 181–223 (1986).
  - [15] S. K. Lam, A. Pitrou, and S. Seibert, “Numba: A LLVM-based Python JIT compiler,” in *Proceedings of the Second Workshop on the LLVM Compiler Infrastructure in HPC* (ACM, New York, 2015), pp. 1–6.
  - [16] R. Piessens, E. de Doncker-Kapenga, C. W. Überhuber, and D. K. Kahaner, *QUADPACK: A Subroutine Package for Automatic Integration* (Springer-Verlag, Berlin, 1983).