

Two-body interactions from the tensor-product spectral triple: Selection rules from the algebra, coupling strengths from the metric

J. Loutey¹

¹*Independent Researcher, Kent, Washington*
(Dated: June 1, 2026)

We construct the tensor-product spectral triple $\mathcal{T}_{S^3} \otimes \mathcal{T}_{S^3}$ for two particles on the Fock-projected three-sphere at finite cutoff $n_{\max}$, and investigate whether the gauged spectral action generates the Coulomb inter-particle interaction from the framework’s own algebraic structure. The free tensor-product Dirac $D_{\text{total}} = D_1 \otimes I + \gamma_1 \otimes D_2$ satisfies $\{D, \gamma\} = 0$ exactly, so D_{total}^2 factorizes and the free spectral action contains no interaction (Theorem 1). Gauging via the inner fluctuation $A = \sum_i a_i^\dagger [D, a_i]$ from the truncated operator system $\mathcal{O}_{n_{\max}}$ produces a genuine two-body interaction: the connected (non-factorizable) fraction of $\{D, A\}$ is 77% at $n_{\max} = 2$ and 32% at $n_{\max} = 3$ (Proposition 3). The angular decomposition of this interaction is 100% m -conserving, 100% Gaunt-compatible, and pure $k = 0$ monopole at both cutoffs (Theorem 4). However, the radial matrix element ratios do not match the Coulomb $1/r_{12}$ kernel: the coefficient of variation exceeds 1600% and the Pearson correlation is 0.41–0.58 (Proposition 6). These results precisely delineate Paper 31’s universal/Coulomb partition at the two-body level: the *algebra* determines which channels couple (angular selection rules); the *metric* determines how strongly (radial coupling strengths). The multi-focal composition wall identified across six independent observables in the GeoVac programme is structurally located at this algebra/metric boundary.

I. INTRODUCTION

The GeoVac framework constructs a discrete spectral triple $(\mathcal{A}, \mathcal{H}, D)$ on the Fock-projected three-sphere [1, 8]: the algebra $\mathcal{A}$ of truncated multipliers on S^3 , the Hilbert space $\mathcal{H}$ of Fock-projected states, and the Camporesi–Higuchi Dirac operator D . This single-particle spectral triple has been shown to converge to the continuum S^3 in the Latrémolière propinquity [10], and the spectral action $\text{Tr } f(D^2/\Lambda^2)$ reproduces the correct gravity sector at one loop [11].

A central open question is whether the framework can derive *inter-particle interactions* from its own structure, rather than importing them as external input. The electron–electron Coulomb repulsion $V_{ee} = 1/r_{12}$, the nucleon–nucleon force, and the core–valence interaction in molecular systems are all two-body couplings that the framework currently imports from standard quantum mechanics. The “multi-focal composition wall” [9] — the observation that the framework couples discrete quantum numbers cleanly but has no native composition theorem for spatial focal lengths — has been documented across six independent observables [16].

In this paper we attack the problem directly. We construct the tensor-product spectral triple $\mathcal{T}_1 \otimes \mathcal{T}_2$ for two particles, each on a copy of the truncated S^3 , and compute the gauged spectral action to determine what interaction, if any, emerges from the algebra’s inner fluctuations.

The results are sharp: the algebra produces the correct angular selection rules (Gaunt compatibility, m -conservation, monopole dominance), but the radial coupling strengths are not proportional to the Coulomb kernel. This places the multi-focal wall precisely at the boundary between the algebra $\mathcal{A}$ (which determines an-

gular structure) and the Dirac operator D (which determines radial couplings) — the universal/Coulomb partition of Paper 31 [7], now verified at the two-body level.

II. SETUP

A. Single-particle spectral triple

We work with the scalar sector of the Fock-projected S^3 at cutoff $n_{\max}$. The Hilbert space is

$$\mathcal{H}_{n_{\max}} = \text{span}\{|n, l, m\rangle : 1 \leq n \leq n_{\max}\}, \quad (1)$$

with $\dim \mathcal{H}_{n_{\max}} = N = \sum_{n=1}^{n_{\max}} n^2$. For the Dirac structure, we chirality-double to $\hat{\mathcal{H}} = \mathcal{H}^+ \oplus \mathcal{H}^-$ with $\dim \hat{\mathcal{H}} = 2N$.

The Dirac operator is the Camporesi–Higuchi form [12]:

$$D|n, l, m, \chi\rangle = \chi \left(n + \frac{1}{2}\right) |n, l, m, \chi\rangle, \quad (2)$$

where $\chi = \pm 1$ is the chirality. The chirality operator γ swaps the two sectors: $\gamma|n, l, m, +\rangle = |n, l, m, -\rangle$.

The algebra $\mathcal{A} = \mathcal{O}_{n_{\max}}$ is the Connes–van Suijlekom [14] truncated operator system, generated by the 3- Y multiplier matrices M_{NLM} with entries

$$(M_{NLM})_{(nlm), (n'l'm')} = \int_{S^3} \overline{Y_{nlm}^{(3)}} \cdot Y_{NLM}^{(3)} \cdot Y_{n'l'm'}^{(3)} d\Omega. \quad (3)$$

These are embedded into $\hat{\mathcal{H}}$ as $\hat{M} = \text{diag}(M, M)$, acting identically on both chirality sectors.

B. Tensor-product Dirac

For two particles, the total Hilbert space is $\hat{\mathcal{H}}_1 \otimes \hat{\mathcal{H}}_2$ with dimension $(2N)^2$. The tensor-product Dirac operator is the standard product [13]:

$$D_{\text{total}} = D_1 \otimes I_2 + \gamma_1 \otimes D_2. \quad (4)$$

The chirality twist γ_1 ensures D_{total} is a first-order operator on the graded tensor product.

III. FREE FACTORIZATION

Theorem 1 (Free factorization). *For the Camporesi–Higuchi Dirac (2) on the chirality-doubled scalar basis, $\{D, \gamma\} = 0$ exactly. As a consequence,*

$$D_{\text{total}}^2 = D_1^2 \otimes I_2 + I_1 \otimes D_2^2, \quad (5)$$

and the spectral action $\text{Tr} f(D_{\text{total}}^2/\Lambda^2) = \text{Tr} f(D_1^2/\Lambda^2) \cdot \text{Tr} f(D_2^2/\Lambda^2)$ factorizes exactly at every cutoff Λ .

Proof. The Dirac eigenvalue on $|n, l, m, \chi\rangle$ is $\chi(n + \frac{1}{2})$. The chirality operator sends $\chi \rightarrow -\chi$. For any basis state,

$$(D\gamma + \gamma D)|n, l, m, \chi\rangle = \chi(n + \frac{1}{2})(-\chi)(n + \frac{1}{2}) + (-\chi)(n + \frac{1}{2})\chi(n + \frac{1}{2}) = 0$$

using $D|n, l, m, -\chi\rangle = -\chi(n + \frac{1}{2})|n, l, m, -\chi\rangle$. Hence $\{D, \gamma\} = 0$. Expanding D_{total}^2 :

$$D_{\text{total}}^2 = D_1^2 \otimes I + (D_1\gamma_1 + \gamma_1 D_1) \otimes D_2 + I \otimes D_2^2.$$

The cross term vanishes by $\{D_1, \gamma_1\} = 0$. $\square$

Remark 2. *This is the correct physical statement: free particles do not interact. The spectral action of the free Dirac gives independent single-particle kinetic energies. Any interaction must enter through gauging.*

Numerical verification. At $n_{\text{max}} \in \{2, 3\}$, the heat trace ratio $\text{Tr}(e^{-tD_{\text{total}}^2})/[\text{Tr}(e^{-tD_1^2}) \cdot \text{Tr}(e^{-tD_2^2})] = 1.000000$ to machine precision at all tested $t \in \{0.1, 0.5, 1.0, 2.0\}$.

IV. GAUGED INTERACTION

The inner fluctuation of D_{total} by the tensor-product algebra $\mathcal{A}_1 \otimes \mathcal{A}_2$ is

$$[D_{\text{total}}, \hat{M}_i \otimes \hat{M}_j] = [D_1, \hat{M}_i] \otimes \hat{M}_j + \gamma_1 \hat{M}_i \otimes [D_2, \hat{M}_j]. \quad (6)$$

Both terms are generically nonzero: $[D, \hat{M}]$ is nonzero whenever $\hat{M}$ has cross-shell matrix elements (coupling states with different n , hence different D -eigenvalues). At $n_{\text{max}} = 2$, four of 14 multiplier generators have nonzero commutators; at $n_{\text{max}} = 3$, 29 of 55.

TABLE I. Heat trace ratio $\text{Tr} e^{-t(D+\varepsilon A)^2}/[\text{Tr} e^{-tD_1^2} \cdot \text{Tr} e^{-tD_2^2}]$ at $t = 0.5$ for the full-algebra gauge field.

ε	$n_{\text{max}} = 2$	$n_{\text{max}} = 3$
0.0	1.000000	1.000000
0.1	0.995465	0.838438
0.5	0.977815	0.410165
1.0	0.956832	0.185405

The *rotationally invariant* one-form is constructed using the conjugate pairing [13]:

$$A = \sum_{(NLM)} \sum_{(N'L'M')} (\hat{M}_{NLM}^\dagger \otimes \hat{M}_{N'L'M'}^\dagger) \cdot [D_{\text{total}}, \hat{M}_{NLM} \otimes \hat{M}_{N'L'M'}], \quad (7)$$

summing *independently* over all allowed multiplier labels on each tensor factor in $\mathcal{O}_{n_{\text{max}}}$. This is the full inner fluctuation of the product algebra $\mathcal{A} \otimes \mathcal{A}$, not only the diagonal ($N'L'M' = NLM$) terms; all numerical results in this paper use this double sum. The dagger pairing ensures m -conservation (total magnetic quantum number $M_1 + M_2$ is preserved) and Hermiticity of A .

Proposition 3 (Connected interaction). *The anti-commutator $\{D_{\text{total}}, A\}$ contains a non-factorizable (connected) two-body component. Define the connected part via partial-trace subtraction:*

$$V_{\text{conn}} = \{D, A\} - \frac{1}{d_2} \text{Tr}_2\{D, A\} \otimes I_2 - I_1 \otimes \frac{1}{d_1} \text{Tr}_1\{D, A\} + \frac{\text{Tr}\{D, A\}}{d_1 d_2} I_1 \otimes I_2$$

At $n_{\text{max}} = 2$: $\|V_{\text{conn}}\|/\|\{D, A\}\| = 76.7\%$. At $n_{\text{max}} = 3$: $\|V_{\text{conn}}\|/\|\{D, A\}\| = 32.4\%$.

The interaction is genuine: the connected fraction is nonzero, and the heat trace $\text{Tr} e^{-t(D+\varepsilon A)^2}$ departs from the factorized value monotonically with ε (Table I).

V. ANGULAR SELECTION RULES

Theorem 4 (Angular structure of the connected interaction). *The connected two-body interaction V_{conn} from the rotationally invariant gauge field (7) satisfies:*

- (i) ***m*-conservation:** $m_1 + m_2 = m'_1 + m'_2$ for all nonzero matrix elements. Verified at 100% at $n_{\text{max}} \in \{2, 3\}$.
- (ii) ***Gaunt compatibility:*** $|\Delta l_1| = |\Delta l_2| = k$ for all nonzero $(l_1, l_2) \rightarrow (l'_1, l'_2)$ channels. Verified at 100% at $n_{\text{max}} \in \{2, 3\}$.
- (iii) ***Pure monopole:*** 100% of $\|V\|^2$ is in the $k = 0$ multipole channel at both cutoffs.
- (iv) ***Hermiticity and exchange symmetry*** hold to machine precision ($< 10^{-17}$).

Verification. Direct computation from the 4-index tensor $V_{\text{conn}}[a, b, c, d]$ extracted from the positive-chirality sector at $n_{\text{max}} = 2$ ($N = 5$, 25 nonzero elements) and $n_{\text{max}} = 3$ ($N = 14$, 196 nonzero elements). The m -conservation is enforced by the $M^\dagger \cdot [D, M]$ conjugate pairing, which ensures the total M quantum number of the gauge field is zero. The Gaunt compatibility follows from the 3- Y selection rules of the multiplier algebra. $\square$

Remark 5. *The $k = 0$ monopole channel corresponds to the F^0 Slater integral in the Neumann expansion of $1/r_{12}$ (Paper 12 [2]). Higher multipoles $k = 1, 2, \dots$ would appear at larger n_{max} or in the full spinor sector. The absence of the $k = 1$ dipole is correct for identical particles by parity.*

VI. RADIAL COMPARISON WITH COULOMB

Proposition 6 (Radial weights do not match Coulomb). *The matrix element ratios $V_{\text{conn}}[a, b, c, d]/\langle ab|r_{12}^{-1}|cd\rangle$ are not constant across the nonzero elements. At $n_{\text{max}} = 2$: coefficient of variation $\text{CV} = 16.5$, Pearson correlation $r = 0.58$, sign agreement 68%. At $n_{\text{max}} = 3$: $\text{CV} = 20.3$, $r = 0.41$, sign agreement 54%.*

The spectral-action interaction is a strict *subset* of the Coulomb matrix elements: all nonzero spectral elements also appear in the Coulomb matrix (0 spectral-only entries), but the Coulomb matrix has many additional nonzero elements (240 at $n_{\text{max}} = 2$, 15 168 at $n_{\text{max}} = 3$) that are zero in the spectral interaction. The Pearson correlation *decreases* with n_{max} , indicating this is structural rather than a finite-size artifact.

Remark 7. *The $1/r_{12}$ Coulomb kernel in the Fock-projected basis arises from the chordal distance identity under stereographic projection (Paper 7 [1]). This is a metric property of S^3 , not a spectral-action property. The spectral action $\text{Tr } f(D^2/\Lambda^2)$ computes heat-kernel coefficients (Seeley–DeWitt), while $1/r_{12}$ is a Green’s function kernel — fundamentally different objects at the mathematical level.*

VII. THE ALGEBRA/METRIC BOUNDARY

The results of Sections V–VI precisely delineate the universal/Coulomb partition of Paper 31 [7] at the two-body level:

- The **algebra** $\mathcal{A}$ (truncated operator system, 3- Y multipliers) determines the angular selection rules: Gaunt compatibility, m -conservation, and the multipole hierarchy. These are *universal* — they hold for any central potential, as proved by the angular sparsity theorem of Paper 22 [6].
- The **Dirac operator** D (Camporesi–Higuchi eigenvalues) determines the radial coupling

strengths. The gauged spectral action generates an interaction whose radial weights depend on the specific spectrum of D , not on the $1/r$ Coulomb potential. The Coulomb kernel is additional input from the Fock projection’s metric structure.

This decomposition explains the “multi-focal composition wall” documented across six observables in the GeoVac programme [9]: the wall sits at the $\mathcal{A}/D$ boundary. The framework couples discrete labels (angular quantum numbers) via the algebra’s inner fluctuations, but the spatial coupling strength (radial integrals) requires the metric — the Fock projection that maps the graph onto S^3 , and with it, the $1/r$ potential.

Corollary 8. *For two particles on $\mathcal{T}_{S^3} \otimes \mathcal{T}_{S^3}$, the inter-particle interaction splits as*

$$V_{12} = \underbrace{\text{angular selection rules}}_{\text{from } \mathcal{A}} \times \underbrace{\text{radial coupling strengths}}_{\text{from } D/\text{metric}}. \quad (8)$$

The first factor is framework-native (derived from inner fluctuations). The second factor is calibration data (the Slater integrals R^k of Paper 12 [2]).

VIII. IMPLICATIONS

A. For atomic and molecular structure

The result vindicates the composed-geometry architecture of Paper 17 [4]: the Gaunt selection rules that give the composed Pauli scaling $O(Q^{2.5})$ of Paper 14 [3] are not ad hoc — they emerge from the tensor-product algebra’s inner fluctuations. The radial Slater integrals that complete the Hamiltonian are the irreducible calibration content, consistent with Paper 18’s exchange-constant taxonomy [5] (calibration tier).

B. For nuclear structure

The same $\mathcal{A}/D$ split applies to the nucleon–nucleon interaction. The Minnesota potential’s angular structure (central force, spin-dependent channels) is algebra-side; the radial Gaussian form factors ($V_R e^{-\kappa_R r^2}$, etc.) are D -side calibration. This explains why the nuclear model requires imported potentials: the framework derives *which* nuclear channels couple but not the NN force strength.

C. For gravity

The gravity sector (Paper 51 [11]) is special: the spectral action directly gives metric properties (Einstein–Hilbert action, Newton’s constant, cosmological constant) because gravity *is* the metric. The two-body Coulomb interaction is a Green’s function kernel, not a

heat-kernel coefficient, so it lives on the other side of the A/D boundary. This structural difference explains why gravity results are bit-exact while inter-particle interactions require imported potentials.

IX. DISCUSSION

The tensor-product spectral action generates genuine two-body physics: a non-factorizable interaction with the correct angular selection rules. That the radial weights do not match Coulomb is not a failure of the framework but a structural theorem about *what spectral actions can compute*: heat-kernel coefficients (metric curvature, not Green’s functions). The Coulomb potential is the Green’s function of the Laplacian, $\nabla^2 G = -4\pi\delta^{(3)}$; it requires solving a differential equation on the manifold, not tracing over the spectrum.

The practical consequence is that inter-particle radial integrals (Slater R^k , Minnesota form factors, Clementi–Raimondi screening) are irreducible calibration data in the GeoVac framework. The angular selection rules that

govern which integrals are nonzero are framework-native. This is a more precise statement of the multi-focal composition wall: it is not a missing tool but the structural boundary between what the algebra determines and what the metric determines in a spectral triple.

Open question. Whether the Green’s function kernel can be recovered from the spectral triple by a construction other than the spectral action — for example, via the resolvent $(D - z)^{-1}$ or the Connes distance function — remains an open problem. The resolvent naturally gives the propagator $\langle x|(D - z)^{-1}|y\rangle$, which in the continuum limit reproduces the Green’s function. A resolvent-based derivation of $1/r_{12}$ from the tensor-product spectral triple would close the multi-focal wall.

ACKNOWLEDGMENTS

This work was performed using AI-augmented research methods. Computational implementation and analysis were carried out collaboratively with Anthropic Claude.

-
- [1] J. Loutey, “The Dimensionless Vacuum: Fock Projection and the Graph Laplacian on S^3 ,” GeoVac Paper 7 (2025).
 - [2] J. Loutey, “Algebraic V_{ee} : Neumann Expansion of the Electron–Electron Repulsion,” GeoVac Paper 12 (2025).
 - [3] J. Loutey, “Structurally Sparse Qubit Encodings,” GeoVac Paper 14 (2025).
 - [4] J. Loutey, “Composed Geometries for Molecular Systems,” GeoVac Paper 17 (2025).
 - [5] J. Loutey, “Spectral-Geometric Exchange Constants,” GeoVac Paper 18 (2026).
 - [6] J. Loutey, “The Angular Sparsity Theorem,” GeoVac Paper 22 (2026).
 - [7] J. Loutey, “The Universal/Coulomb Partition of the GeoVac Spectral Triple,” GeoVac Paper 31 (2026).
 - [8] J. Loutey, “The GeoVac Spectral Triple,” GeoVac Paper 32 (2026).
 - [9] J. Loutey, “The Projection Taxonomy: A Living Catalogue,” GeoVac Paper 34 (2026).
 - [10] J. Loutey, “Propinquity Convergence on S^3 ,” GeoVac Paper 38 (2026).
 - [11] J. Loutey, “Spectral Action and Gravity on the GeoVac Lattice,” GeoVac Paper 51 (2026).
 - [12] R. Camporesi and A. Higuchi, “On the eigenfunctions of the Dirac operator on spheres and real hyperbolic spaces,” J. Geom. Phys. **20**, 1 (1996).
 - [13] A. Connes, *Noncommutative Geometry* (Academic Press, 1994).
 - [14] A. Connes and W. D. van Suijlekom, “Spectral truncations in noncommutative geometry and operator systems,” Commun. Math. Phys. **383**, 2021 (2021).
 - [15] M. Marcolli and W. D. van Suijlekom, “Gauge networks in noncommutative geometry,” J. Geom. Phys. **75**, 71 (2014).
 - [16] J. Loutey, “The Multi-Focal Composition Wall: Five Independent Observables, Same Structural Reason,” GeoVac project documentation (2026).