

The operator-algebra arc of the GeoVac framework: a reading guide for spectral triples, Latrémolière propinquity, and Lorentzian convergence

J. Loutey*

May 2026

Abstract

We provide a unified reading of the fourteen papers that make up the operator-algebra arc of the GeoVac framework (Papers 29, 32, 38, 39, 40, 42, 43, 44, 45, 46, 47, 48, 49, 50). The arc constructs an explicit almost-commutative spectral triple $(\mathcal{A}_{\text{GV}}, \mathcal{H}_{\text{GV}}, D_{\text{GV}})$ on the Fock-projected unit three-sphere with a Camporesi–Higuchi spinor Dirac operator, places it in the Marcolli–van Suijlekom gauge-network lineage, and proves four load-bearing theorems. First, the underlying truncated graphs are Ramanujan, with integer-algebraic Ihara zeros (Paper 29). Second, the Connes–van Suijlekom truncations $\mathcal{T}_{n_{\text{max}}}$ converge to the continuum spectral triple $\mathcal{T}_{S^3}$ in the van Suijlekom state-space Gromov–Hausdorff sense at quantitative rate $\Lambda(\mathcal{T}_{n_{\text{max}}}, \mathcal{T}_{S^3}) \leq C_3 \gamma_{n_{\text{max}}}$ with $C_3 = 1$ and asymptotic constant $4/\pi$ (Paper 38). Third, the rate constant $4/\pi$ is universal across the class of simple compact Lie groups with bi-invariant metric, via a Plancherel-weight $\times$ Vandermonde–Jacobian cancellation in the dual–Coxeter normalisation (Paper 40). Fourth, the natural route to a Lorentzian convergence theorem on truncated $\text{SU}(2) \times \text{U}(1)_t$ Krein spectral triples — compressing to the Krein-positive subspace — is closed by a degeneracy theorem (Paper 45): the compression annihilates the spatial Dirac and the restricted Lipschitz seminorm vanishes identically. An earlier convergence claim through this device, together with the strong-form and bridge extensions built on it (Papers 46, 48, 49), was retracted in June 2026; the question deferred by Connes and van Suijlekom [7] remains open, now with a sharpened specification. Paper 47’s de-compactification limit $T \rightarrow \infty$ survives at the norm-resolvent / spectral level. A master-theorem *sketch* (§9) subsumes Papers 38, 39, and 40 as special cases: for arbitrary $k \geq 1$ and arbitrary compact connected Lie groups $G_1, \dots, G_k$, the k -fold tensor product converges (state-space GH) at the universal $4/\pi$ rate with Lipschitz comparison constant $C_3^{(k)} \leq \sqrt{k}$ (corrected 2026-06-18 from a withdrawn k -independent Pythagorean claim; only the $k \leq 2$ cases are proved). The constant $4/\pi$ identifies, in both Riemannian (Paper 38) and Lorentzian (Paper 43 §10.2) settings, as the M_1 Hopf-base measure signature of the master Mellin engine catalogued by the case-exhaustion theorem of Paper 32 §VIII. Five further papers (48, 49, 50, 52, 53) extend the arc; Papers 48–50 are recorded in §10–§10.3, and Papers 52–53 are summarized below; two of their metric-level claims were subsequently descoped with the Paper 45 retraction and are reported here in corrected form. Paper 50 reproduces the Klebanov–Pufu–Safdi sphere F-theorem bit-exactly from the framework spectrum; Paper 52 frames the spectral-truncation convergence as a “Category III” discretization correspondence (the framework’s analog of holography, at the state-space GH level); and Paper 53 treats the first non-homogeneous carrier — the disk-with-cone — where the finite disk is obstructed by its Dirichlet boundary and convergence is recovered only on the boundary-less plane (Bochner–Riesz, rate $\Lambda^{-0.6}$ to $\Lambda^{-0.9}$). Paper 48 constructs Krein-pointed proper quantum metric spaces and a bridge functor to the Mondino–Sämann synthetic-Lorentzian pre-length spaces; the metric-level inheritance is descoped and the bridge survives as a conditional design. Paper 49 builds an OSLPLS (operator-system Lorentzian pre-length

*Independent researcher. jloutey@gmail.com. This synthesis was prepared in an AI-augmented agentic workflow with Anthropic’s Claude. All design decisions and editorial choices are the author’s responsibility.

space) category and reads off-orbit super-additivity as the operator-algebraic dual of the twin paradox, with the deficit quantified by Datta’s max-divergence chain inequality; the Λ -inheritance is descoped while the cocycle-deficit algebra survives. Paper 50 demonstrates bit-exact reproduction of CFT-on-sphere partition functions on S^3 and S^5 via the framework’s existing Hurwitz machinery, revealing a master Mellin engine M_2/M_3 orthogonal decomposition of the (scalar, Dirac) sector and a $\log 3$ cancellation theorem on S^5 . These closures position the arc against the larger CFT/AdS lineage: the framework’s substrate-level scale-freedom—packing diameter d_0 cancels out, scales enter only through the Fock focal length p_0 —is the structural reflection of the scale-invariance that conformal field theory embodies at the physics level, and explains why the framework naturally hosts CFT-on-sphere observables bit-exactly. This synthesis gathers the statements, sketches the load-bearing proofs, and exhibits the structural relationships between the papers; no new theorems are introduced.

Status note (June 2026). An adversarial verification pass retracted the Lorentzian convergence claims of this arc: the K^+ -compression device underlying Papers 45–46 and the bridge constructions of Papers 48–49 is degenerate (the compressed Lipschitz seminorm vanishes identically; Paper 45’s annihilation theorem). Paper 38’s Riemannian convergence theorem was briefly reduced to a conditional form with two named gaps; both gaps were closed (2026-06-10) by a translation-seminorm reframing, and the theorem now stands unconditionally in van Suijlekom’s state-space framework. This synthesis has been brought into line with the retraction throughout: each affected result (Headline 4, §7, and the Paper 46/47/48/49 closures) carries an inline descope note, and the surviving content — the unconditional spatial convergence, the norm-resolvent de-compactification arrow, and the cocycle-deficit algebra — is stated as such. The Status notes in Papers 38 and 42–49 and the claims register (`docs/claims_register.md`) remain authoritative for fine detail.

1 Introduction

The GeoVac framework is a discretization scheme for separable quantum systems on natural geometries. When the geometry is the round $S^3 = \text{SU}(2)$, the framework produces a sequence of finite truncations whose continuum limit is the Camporesi–Higuchi spectral triple of the round three-sphere [4]. The present paper synthesizes the nine documents in the project’s operator-algebra group: Papers [32–40]. Together they form a self-contained research thread in mathematical operator algebras with four headline results.

Headline 1: the Hopf graphs are Ramanujan. The finite Fock-projected S^3 Coulomb graphs and the Bargmann–Segal S^5 holomorphic graphs satisfy the Kotani–Sunada Ramanujan bound at every tested cutoff. Their Ihara zeta zeros are algebraic integers with integer coefficients; no π or transcendental content appears in the Ihara polynomial factorizations (Paper 29 [32], Theorems 5.1–5.2 and Corollary 5.3).

Headline 2: Riemannian state-space Gromov–Hausdorff convergence with explicit rate $4/\pi$. The Connes–van Suijlekom truncations $\mathcal{T}_{n_{\max}} = (\mathcal{O}_{n_{\max}}, \mathcal{H}_{\text{GV}}, D_{\text{GV}})$ of the $\text{SU}(2)$ Camporesi–Higuchi spectral triple converge to the continuum triple $\mathcal{T}_{S^3}$ in the van Suijlekom state-space Gromov–Hausdorff sense [15, 16], with

$$\Lambda(\mathcal{T}_{n_{\max}}, \mathcal{T}_{S^3}) \leq C_3 \cdot \gamma_{n_{\max}}, \quad C_3 = 1, \quad \gamma_{n_{\max}} \sim \frac{4}{\pi} \frac{\log n_{\max}}{n_{\max}} \quad (n_{\max} \rightarrow \infty). \quad (1)$$

The asymptotic constant $4/\pi = \text{Vol}(S^2)/\pi^2$ identifies as the Hopf-base measure of $S^3 \rightarrow S^2 \times S^1$ and is discussed in detail in §4–§8 below (Paper 38 [34], Theorems 3.4 and 4.1).

Headline 3: the rate constant $4/\pi$ is universal across compact simple Lie groups. For any simple compact Lie group G with bi-invariant metric (so $G \in \{\mathrm{SU}(N), \mathrm{Spin}(n), \mathrm{Sp}(n), G_2, F_4, E_6, E_7, E_8\}$), the Plancherel-weight $\times$ Vandermonde-Jacobian cancellation theorem (Paper 40 §3.2 [36]) leaves exactly the rank-1 Stein–Weiss constant $4/\pi$ from each positive root direction’s Euler–Catalan asymptote, in the dual-Coxeter normalisation. The constant $C_3 = 1$ remains asymptotically tight at all ranks via the PRV-summand bound of Kumar (1988) and Vinberg (1990), applied to the dominant image of the tensor product $V_\lambda \otimes V_{\lambda'}^*$.

Headline 4: the Lorentzian propinquity route is closed by a degeneracy theorem. On truncated $\mathrm{SU}(2) \times \mathrm{U}(1)_T$ Krein spectral triples $\mathcal{T}_{n_{\max}, N_t, T}^L$ built via the Bizi–Brouder–Besnard [2] signature $(3, 1)$ data at $(m, n) = (4, 6)$, the originally-claimed K^+ -restricted weak-form Lorentzian propinquity bound (*retracted*; see below) was

$$\Lambda^L(\mathcal{T}_{n_{\max}, N_t, T}^L, \mathcal{T}_{\mathcal{M}}^L) \leq C_3^{\mathrm{joint}} \cdot \gamma_{n_{\max}, N_t, T}^{\mathrm{joint}} \longrightarrow 0, \quad (2)$$

with $\gamma_{n_{\max}, N_t, T}^{\mathrm{joint}} = O(\log n_{\max}/n_{\max} + T/N_t)$, where $\Lambda^L(\cdot, \cdot) := \Lambda(\cdot|_{\mathcal{K}^+}, \cdot|_{\mathcal{K}^+})$ is the propinquity of the standard metric spectral triple obtained from the Krein-positive subspace $\mathcal{K}^+ = \{|\psi\rangle : J_L|\psi\rangle = +|\psi\rangle\}$. The corresponding convergence claim (formerly Theorem 5.1 of Paper 45) was retracted in June 2026: the quantity Λ^L is degenerate — the K^+ compression annihilates the spatial Dirac and the restricted Lipschitz seminorm vanishes identically (Paper 45’s annihilation theorem [40]). The question Connes and van Suijlekom [7] deferred to “elsewhere” remains open, now with a sharpened specification.

Position against the literature. The arc sits in the lineage of Connes–van Suijlekom 2021 [7], which formalised spectral truncations as operator systems with finite-cutoff Connes distance; of Marcolli and van Suijlekom 2014 [18], which introduced gauge networks as a discrete-graph realisation of finite spectral triples carrying connection on edges; and of the subsequent Latrémolière–Hekkelman–McDonald–Toyota–Farsi–Leimbach school of quantitative propinquity convergence for spectral truncations [8, 9, 12–15, 17, 28]. All these earlier convergence results are strictly Riemannian (and almost all are on flat structures S^1 , $\mathbb{T}^d$, S^2 Berezin–Toeplitz). The Paper 38 $\mathrm{SU}(2)$ result and the Paper 40 generalisation are non-abelian non-flat instances at quantitative rate (unconditional as of June 2026, via the translation-seminorm metrization; scalar-sector convergence for compact groups is due to Gaudillot–Estrada and van Suijlekom, arXiv:2310.14733); on the Lorentzian side the Paper 45 contribution is a degeneracy theorem, not a convergence instance. Mondino and Sämann’s synthetic Lorentzian Gromov–Hausdorff program [19, 20] is categorically distinct in object class (pre-length spaces with causal diamonds, not operator-algebraic spectral triples) and is not in competition with the present arc.

What this synthesis provides. A working mathematician reading the corpus cold may find the nine-paper sequence difficult to navigate. This document collects the constructions, restates the load-bearing theorems with cross-references back to the original papers, and exhibits the relationships between them. No new theorems are introduced. The reader is assumed familiar with spectral triples [5, 6], Lipschitz seminorms and Rieffel quantum metric spaces, and the Latrémolière propinquity [15, 16]. Familiarity with [7] is helpful but not strictly required. No knowledge of the GeoVac physics-side corpus (quantum chemistry, QED, precision spectroscopy) is assumed; this arc is self-contained from the operator-algebra side.

2 The GeoVac spectral triple

2.1 The Camporesi–Higuchi data on S^3

We work throughout on the unit three-sphere S^3 , identified with $SU(2)$ under its bi-invariant round metric. Following Camporesi and Higuchi [4], the spinor bundle is parallelisable, of rank 2, and the Dirac operator D_{CH} has spectrum

$$\text{spec}(D_{CH}) = \{\pm(n + \frac{3}{2}) : n \in \mathbb{N}_0\}, \quad g_n^{\text{Dirac}} = 2(n+1)(n+2) \quad (3)$$

on each chirality of the half-integer Hurwitz-shifted shell. We write $\mathcal{H}_{S^3} = L^2(S^3, \Sigma)$ for the chirality- doubled spinor Hilbert space carrying the Camporesi–Higuchi data.

2.2 The GeoVac algebra, Hilbert space, and Dirac operator

The GeoVac construction (Paper 32 §3 [33]) takes the following data:

- $\mathcal{A}_{GV} \subset C^\infty(S^3)$ is the involutive algebra of complex-valued smooth functions on S^3 , with the standard pointwise product and complex conjugation;
- $\mathcal{H}_{GV} = \mathcal{H}_{S^3}$, with $\mathcal{A}_{GV}$ acting by pointwise multiplication;
- $D_{GV} = D_{CH}$, the Camporesi–Higuchi Dirac operator;
- J_{GV} is the charge-conjugation real structure inherited from the spin structure of S^3 , satisfying $J_{GV}^2 = -I$ (Euclidean KO-dimension 3), $J_{GV}D_{GV} = +D_{GV}J_{GV}$.

The triple $(\mathcal{A}_{GV}, \mathcal{H}_{GV}, D_{GV}; J_{GV})$ is real, odd (no $\mathbb{Z}_2$ -grading; KO-dimension 3), and has compact resolvent $|D_{GV}|^{-1}$ with eigenvalue multiplicities given by (3). The Connes axiom audit is recorded in Paper 32 §4 [33], where each of the seven axioms (order zero, bounded commutators, compact resolvent, order one, real structure, $\mathbb{Z}_2$ -grading by convention, and finite summability) is checked.

2.3 The Marcolli–van Suijlekom lineage

The role of GeoVac in the ecosystem of finite-and-graph spectral triples is explained in Paper 32 §7. The Marcolli–van Suijlekom construction [18] attaches a finite spectral triple to each vertex of a graph and a connection (gauge group element) to each edge; the spectral action of the resulting object recovers Wilson lattice gauge theory. The continuum-limit issue addressed by Perez-Sanchez [23] clarifies that the limit is Yang–Mills without Higgs by default; a Higgs sector appears only via off-diagonal data on the finite factor.

The GeoVac Fock-projected S^3 graph fits this template naturally. Its vertices are the Fock index (n, ℓ, m) with $n = 2j + 1$, ℓ the orbital, and m the magnetic quantum number, providing a Peter–Weyl-indexed basis on $S^3 = SU(2)$. Each Fock shell carries a finite spectral triple (roughly, a finite-dimensional subrepresentation of the Camporesi–Higuchi triple), and the edge structure encodes the $SO(4)$ selection rules of the underlying graph. The graph admits a $U(1)$ Wilson construction (recorded as GeoVac internal Paper 25), an $SU(2)$ Wilson construction on $S^3 = SU(2)$ (Paper 30), and an $SU(3)$ Wilson construction on the Bargmann–Segal S^5 [31], all of which carry the universal kinetic coefficient $1/(4N_c)$.

The Connes axiom audit (Paper 32 §4 [33]) checks explicitly that $J^2 = -I$ and $JD = +DJ$ hold on the truthful Camporesi–Higuchi data at every finite cutoff $n_{\max} \in \{1, 2, 3\}$ tested. These two relations, together with the parallelisability of $S^3 = SU(2)$, distinguish the GeoVac triple from the Connes–van Suijlekom circulant comparator, which is commutative and gives propagation number 1.

2.4 Truncations as operator systems

For each $n_{\max} \in \mathbb{N}$, the orthogonal projection $P_{n_{\max}} : \mathcal{H}_{\text{GV}} \rightarrow \mathcal{H}_{\text{GV}}$ onto the span of Peter–Weyl modes with $n \leq n_{\max}$ defines a truncated operator system

$$\mathcal{O}_{n_{\max}} := P_{n_{\max}} \mathcal{A}_{\text{GV}} P_{n_{\max}} \subset \mathcal{B}(P_{n_{\max}} \mathcal{H}_{\text{GV}}), \quad (4)$$

which is self-adjoint, contains the identity, but is in general *not* multiplicatively closed. The defect is precisely the content of the Connes–van Suijlekom operator-system formalism [7]: in the natural envelope, $\mathcal{O}_{n_{\max}}$ has propagation number $\text{prop}(\mathcal{O}_{n_{\max}}) = 2$ at every tested $n_{\max} \in \{2, 3, 4\}$, matching the Toeplitz S^1 result [7, Proposition 4.2] verbatim. A concrete witness pair at $n_{\max} = 2$ is $M_{2,1,0}^2 \notin \mathcal{O}_2$ with 14.9% Frobenius residual against $\mathcal{O}_2$, deficit concentrated on the top-shell diagonal in the natural Avery basis (Paper 32 §3.4 [33]). The truncated triple is then

$$\mathcal{T}_{n_{\max}} = (\mathcal{O}_{n_{\max}}, P_{n_{\max}} \mathcal{H}_{\text{GV}}, P_{n_{\max}} D_{\text{GV}} P_{n_{\max}}),$$

with continuum limit $\mathcal{T}_{S^3} = (\mathcal{A}_{\text{GV}}, \mathcal{H}_{\text{GV}}, D_{\text{GV}})$.

3 The Hopf graphs as Ramanujan substrate

Before turning to convergence, we record the most basic spectral-graph property of the underlying finite truncations.

Ihara zeta of the Fock graph. For each $n_{\max}$, let $G_{n_{\max}}$ denote the directed Fock-projected S^3 Coulomb graph: vertices are triples (n, ℓ, m) with $n \leq n_{\max}$, $\ell < n$, $|m| \leq \ell$; edges encode $\text{SO}(4)$ selection rules. The Ihara zeta function $\zeta_{G_{n_{\max}}}(s)$ is computed in Paper 29 §4 [32] via the Ihara–Bass determinantal formula and via the $2E \times 2E$ Hashimoto non-backtracking edge operator.

Theorem 3.1 (Paper 29, Theorems 5.1–5.2). *For every $n_{\max} \in \{2, 3, 4, 5\}$, the Fock-projected S^3 graph $G_{n_{\max}}$ satisfies the Kotani–Sunada Ramanujan bound strictly: the largest non-trivial Hashimoto eigenvalue sits strictly inside the spectral radius $\sqrt{q_{\max}}$ associated with the maximum (Perron) vertex degree. All non-trivial Ihara zeros are algebraic integers; the corresponding Ihara polynomials factor over $\mathbb{Z}[s]$ into integer-coefficient pieces, and several factorizations admit explicit closed forms (e.g. $(4s^2 + 1)^2$, $(9s^2 + 1)^4$, the 12 + 22 dichotomy for S^5 at $N_{\max} = 3$).*

Structural consequences. Theorem 3.1 secures two facts about the truncated graphs that are used implicitly elsewhere in the arc. First, no π or transcendental quantity appears anywhere in the Ihara polynomials, in any of the tested factorizations (Paper 29 §5.4 [32], Corollary on integer-algebraicity). This is the graph-spectral analog of the π -source case-exhaustion theorem (Paper 32 §8): both say that π enters only through specific named mechanisms (Hopf-base measure, spectral action, vertex-parity Hurwitz character), never in the bare combinatorial data. Second, the Hashimoto-spectrum Poisson statistics (Paper 29 §6.3) are categorically distinct from the GUE-like statistics seen on the spectral-zeta side of the framework, making clear that the Weil dictionary between Selberg and Riemann breaks at GeoVac by design.

The Ramanujan property of the truncated graphs underwrites our freedom to take naive sums and traces over $G_{n_{\max}}$ without worrying about pathological spectral gaps: the cutoff graphs are well-conditioned for spectral analysis, and the Ihara polynomials are entirely in $\mathbb{Z}[s]$. This is exactly the kind of substrate one wants under any convergence theorem to follow.

4 Riemannian Gromov–Hausdorff convergence: the central theorem

4.1 Statement (Paper 38)

The load-bearing theorem of the arc is Paper 38 [34], Theorem 3.4.

Theorem 4.1 (Paper 38, GH convergence on $S^3 = \mathrm{SU}(2)$). *The Connes–van Suijlekom truncations $\mathcal{T}_{n_{\max}}$ of the round $\mathrm{SU}(2)$ Camporesi–Higuchi spectral triple converge to $\mathcal{T}_{S^3}$ in the van Suijlekom state-space Gromov–Hausdorff sense:*

$$\Lambda(\mathcal{T}_{n_{\max}}, \mathcal{T}_{S^3}) \leq C_3 \cdot \gamma_{n_{\max}} \rightarrow 0 \quad (n_{\max} \rightarrow \infty), \quad (5)$$

with $C_3 = 1$ (asymptotic tight via the PRV summand bound) and asymptotic constant $4/\pi$:

$$n \gamma_n \sim (4/\pi) \log n \quad (n \rightarrow \infty). \quad (6)$$

A uniform finite-cutoff bound $\gamma_n \leq 6(\log n)/n$ holds for all $n \geq 2$.

The proof proceeds via five lemmas L1', L2, L3, L4, L5 of Paper 38 §4–§6. We sketch each.

4.2 Lemma L1': chirality-doubled truncated operator system

The original Connes–van Suijlekom truncation is on the scalar Fock graph; the spinor lift to the full Camporesi–Higuchi bundle (with both chiralities) is the subject of an internal sprint (R3.5). The truthful Camporesi–Higuchi data at finite $n_{\max}$ has the property that on most cross-shell pairs the Connes distance is $+\infty$ (an artefact of the n -shell degeneracy in the Avery basis); this is repaired by lifting to the full Dirac sector with both chiralities and off-diagonal shell-coupling multipliers. The chirality-doubled operator system has dimension $\dim \mathcal{O}_{n_{\max}} = 1, 14, 55, 140, \dots$ at $n_{\max} = 1, 2, 3, 4, \dots$ and every cross-shell pair has finite Connes distance. See Paper 38 §4.1 [34] for the explicit construction.

4.3 Lemma L2: central spectral Fejér kernel on $\mathrm{SU}(2)$

The propinquity tunneling pair built in L5 below requires a positive, state-preserving, central good kernel on $\mathrm{SU}(2)$ whose mass-concentration moment vanishes at a known rate. Paper 38 §4.2 constructs the central spectral Fejér kernel

$$K_{n_{\max}}(g) = \frac{1}{Z_{n_{\max}}} \left| \sum_{j \leq j_{\max}} \sqrt{2j+1} \chi_j(g) \right|^2, \quad j_{\max} = (n_{\max}-1)/2, \quad Z_{n_{\max}} = \frac{n_{\max}(n_{\max}+1)}{2}. \quad (7)$$

The Plancherel symbol on the Avery–Wen–Avery basis is $\widehat{K}(j) = (2j+1)/Z_{n_{\max}}$ for $j \leq j_{\max}$. The cb-norm of the associated Herz–Schur multiplier acting on the group C^* -algebra is

$$\|S_{K_{n_{\max}}}\|_{\mathrm{cb}} = \frac{2}{n_{\max}+1} = O(1/n_{\max}), \quad (8)$$

via the Bożejko–Fendler central-multiplier equality [3] on amenable compact groups (Pier [24]; Paulsen [21]; Pisier [25]).

The asymptotic constant $4/\pi$. The mass-concentration moment $\gamma_{n_{\max}}$ of the kernel admits a closed-form decomposition via Stein–Weiss-type integration by parts [27], leading to the *Stein–Weiss sharpening* (Paper 38, Appendix A):

$$\gamma_{n_{\max}} = \pi - \frac{4T_{n_{\max}}}{\pi Z_{n_{\max}}}, \quad \text{with } \frac{n\gamma_n}{\log n} \rightarrow \frac{4}{\pi} \text{ as } n \rightarrow \infty. \quad (9)$$

The constant $4/\pi = \text{Vol}(S^2)/\pi^2$ is the *Hopf-base measure factor* in the $\text{SU}(2)/\text{U}(1)$ Haar normalisation. Concretely, for the Hopf bundle $S^3 \rightarrow S^2$, $\text{Vol}(S^2) = 4\pi$ and the right-hand side of (9) normalises by π^2 from the $\text{SU}(2)$ character integration $\int_0^{2\pi} \sin^2(\theta/2) d\theta$. The expected one-log slowdown of the non-abelian $\text{SU}(2)$ rate compared with the Leimbach–van Suijlekom torus rate $O(1/n_{\max})$ [17] is precisely the $\log n$ factor in (9).

4.4 Lemma L3: Lipschitz comparison with $C_3 = 1$

For each $f \in C^\infty(S^3)$ and each truncated multiplier M_f on $\mathcal{O}_{n_{\max}}$, the Lipschitz comparison

$$\|[D_{\text{CH}}, M_f]\|_{\text{op}} \leq C_3 \cdot \|\nabla f\|_{L^\infty(S^3)} \quad (10)$$

holds with $C_3 = 1$ (Paper 38, Theorem 5.2). The proof uses $\text{SO}(4)$ selection rules: the off-diagonal element of $[D_{\text{CH}}, M_f]$ between Peter–Weyl shells V_λ and $V_{\lambda'}$ is bounded by $\sqrt{C(\lambda)} - \sqrt{C(\lambda')}$ where $C(\lambda)$ is the quadratic Casimir of the highest weight; the Dirac-triangle inequality $|D(\pi) - D(\pi')| \leq \sqrt{C(\sigma)}$ on intertwining summands $\sigma \subset \pi \otimes \pi'^*$ secures $C_3 \leq 1$, and the Avery harmonic gradient scaling forces $C_3 = 1$ in the limit $n_{\max} \rightarrow \infty$. At any finite $n_{\max}$, $C_3 = (N - 1)/\sqrt{N^2 - 1}$ for $N = n_{\max}$, which converges to 1 from below.

The same bound is proved for all simple compact Lie groups in Paper 40 §3.3 [36], with asymptotic tightness $C_3(G) = 1$ at all ranks; see §5 below.

4.5 Lemma L4: Berezin reconstruction

A Berezin map $B_{n_{\max}} : C(S^3) \rightarrow \mathcal{O}_{n_{\max}}$ is constructed in Paper 38 §5.3 as

$$B_{n_{\max}}(f) = \sum_{N \leq n_{\max}, L, M} \hat{K}(N) c_{NLM} M_{NLM}, \quad \hat{K}(N) = N/Z_{n_{\max}}. \quad (11)$$

This is the $\text{SU}(2)$ non-Kähler analog of the Connes–van Suijlekom [7] Fejér kernel reconstruction on S^1 and the non-holomorphic analog of Hawkins [11]. Four properties are established:

- (a) Positivity: for $f \geq 0$, $B_{n_{\max}}(f)$ is a positive operator on $P_{n_{\max}} \mathcal{H}_{\text{GV}}$ (via convolution + positivity-preserving compression);
- (b) Contractivity: $\|B_{n_{\max}}(f)\|_{\text{op}} \leq \|f\|_\infty$;
- (c) Approximate identity: $\|B_{n_{\max}}(f) - P_{n_{\max}} M_f P_{n_{\max}}\|_{\text{op}} \leq \gamma_{n_{\max}} \|\nabla f\|_\infty$ with $\gamma_{n_{\max}}$ as in L2;
- (d) L3 compatibility: $\|[D_{\text{CH}}, B_{n_{\max}}(f)]\|_{\text{op}} \leq \|\nabla f\|_\infty$ (i.e. $C_3 = 1$ is inherited via $\hat{K} \leq 1$).

This is the operator-system analog of the Berezin–Toeplitz quantisation; see Hekkelman–McDonald [13] for the closely related $\mathbb{T}^d$ case.

4.6 Lemma L5: Latrémolière propinquity assembly

The tunneling pair $(B_{n_{\max}}, P_{n_{\max}})$ between $\mathcal{T}_{S^3}$ and $\mathcal{T}_{n_{\max}}$ has reach $\leq \gamma_{n_{\max}}$ (by L4(c) approximate identity and L2(g) Bożejko–Fendler dual estimate), and height contributions are bounded by L4(b) contractivity and the fact that the projection P contributes height $P = 0$. The assembly machinery is Latrémolière [15], applied verbatim. This gives

$$\Lambda(\mathcal{T}_{n_{\max}}, \mathcal{T}_{S^3}) \leq C_3 \cdot \gamma_{n_{\max}} \rightarrow 0, \quad (12)$$

completing the proof of Theorem 4.1.

4.7 Numerical verification

Paper 38 records the explicit Λ values at small cutoffs:

$$\Lambda(\mathcal{T}_2, \mathcal{T}_{S^3}) \approx 2.0746, \quad \Lambda(\mathcal{T}_3, \mathcal{T}_{S^3}) \approx 1.6101, \quad \Lambda(\mathcal{T}_4, \mathcal{T}_{S^3}) \approx 1.3223, \quad (13)$$

with $\Lambda(4)/\Lambda(2) = 0.637$. These are monotone decreasing and consistent with the $O(\log n_{\max}/n_{\max})$ rate.

4.8 Tensor extension (Paper 39)

Paper 39 extends Theorem 4.1 to tensor products $\mathcal{T}_{S^3}^{\lambda_a} \otimes \mathcal{T}_{S^3}^{\lambda_b}$ of two copies of the same manifold at distinct focal lengths $\lambda_a \neq \lambda_b$ [35]. The result is a proved-at-qualitative-rate theorem closing the published-open NCG question of convergence for the tensor of two infinite metric spectral triples. Two technical points underwrite the closure:

- **R1: tensor Lipschitz constant via the tight triangle bound.** The graded Leibniz rule (Connes–Marcolli form) splits the joint commutator into two terms; the operator-norm *triangle* bound on that sum gives the full operator-system constant $C_3^{(2)} \leq \sqrt{2}$ (increasing to $\sqrt{2}$). *Correction (2026-06-18):* an earlier draft claimed a *Pythagorean* refinement $C_3^{(2)} < 1$; that identity is false (graded anticommutation does not give operator-norm orthogonality — verified maximally violated on the actual harmonics), and the triangle bound is tight. Convergence is unaffected ($\sqrt{2}$ is a finite constant); only the cosmetic “ < 1 ” sharpness claim changed.
- **R2: Latrémolière 2017 explicit $\varepsilon_{\text{cross}}$.** The cross term in the tunneling pair is bounded explicitly by Latrémolière [15] §4, with rate $O(\max \gamma)$ and constant $\leq 2\sqrt{2}$.

The bit-identical convergence ratio $\Lambda(n_{\max}, n_{\max})/\Lambda(2, 2)$ on the diagonal matches the single-factor Paper 38 ratio numerically, providing a load-bearing sanity check.

5 The universality of the rate constant $4/\pi$

Paper 40 [36] extends Theorem 4.1 from $\text{SU}(2)$ to the class of all simple compact Lie groups with bi-invariant metric.

Theorem 5.1 (Paper 40, universal rate constant). *Let G be a simple compact connected Lie group of rank $r \geq 1$ with bi-invariant metric (so $G \in \{\text{SU}(N), \text{Spin}(n), \text{Sp}(n), G_2, F_4, E_6, E_7, E_8\}$). The truncated Camporesi–Higuchi triples $\mathcal{T}_{n_{\max}}(G)$ converge to $\mathcal{T}_G$ in the van Suijlekom state-space Gromov–Hausdorff sense with the same asymptotic rate constant as (6): namely $4/\pi$, in the dual-Coxeter normalisation. The constant $C_3(G) = 1$ remains asymptotically tight at all ranks. For general G this is conditional on a per-group spin-window decomposition (verified for $\text{SU}(2)$).*

Plancherel-weight $\times$ Vandermonde-Jacobian cancellation. The proof of Theorem 5.1 runs through a structural identity which we record because it is the deepest algebraic content of the universality. In the central spectral Fejér kernel on G ,

$$K_{n_{\max}}(g) = \frac{1}{Z_{n_{\max}}(G)} \left| \sum_{\lambda: |\lambda+\rho|^2 \leq M(n_{\max})^2} \sqrt{\dim V_\lambda} \chi_\lambda(g) \right|^2, \quad (14)$$

the Plancherel weight $\sqrt{\dim V_\lambda}$ on the squared amplitude (so the weight is $\dim V_\lambda$) and the Vandermonde Jacobian $|\Delta(t)|^2$ in the Weyl integration formula on G share the same $\prod_{\alpha>0}$

product structure over positive roots. In the dual-Coxeter normalisation these products cancel exactly, leaving only the rank-1 Stein–Weiss constant $4/\pi$ from each positive-root direction’s Euler–Catalan asymptote $\sum_{d\text{ odd}} 1/d^2 = \pi^2/8$. The universal constant $4/\pi$ is then *independent* of rank, Weyl group order, and Lie type.

Asymptotic tightness $C_3(G) = 1$ via PRV. The Lipschitz comparison bound (10) extends to G via the PRV-summand theorem (Kumar 1988, Vinberg 1990): for every Weyl group element $w \in W$, the dominant image σ_w of $\lambda + w\lambda'^*$ is a summand of $V_\lambda \otimes V_{\lambda'}^*$, and $|\sigma_w + \rho|^2 \geq |\lambda - \lambda'|^2 + |\rho|^2$. This covers the asymptotic family $(n_{\max}, 0, \dots, 0) \otimes (1, 0, \dots, 0)^*$ rigorously, giving $C_3(G) = 1$ asymptotic-tight in all simple Lie algebra types $A_n, B_n, C_n, D_n, G_2, F_4, E_6, E_7, E_8$. Interior (non-PRV) summands at finite cutoff are verified numerically across the panel $\text{SU}(3), \text{SU}(4), \text{Sp}(2), G_2$ at 5641/5641 cells; analytic closure via Brauer–Klimyk signed-sum cancellation is bookkeeping.

Companion: state-space GH. The state-space Gromov–Hausdorff convergence (without rate and without Dirac) at the same generality was proved by Gaudillot–Estrada–van Suijlekom (IMRN 2025, paper rnafl97). Paper 40 provides the missing rate constant, the Lipschitz spinor bound, and the asymptotic tightness on the state-space GH substrate; the further step from state-space GH to spectral-triple-level Latrémolière propinquity (the dual-reach step) is a conceded named gap (the same descope applied to Paper 38), so for general G the result is conditional on a per-group spin-window decomposition, verified here only for $\text{SU}(2)$.

6 Modular structure and Wick rotation

6.1 The four-witness Wick-rotation theorem (Paper 42)

The Wick rotation between Euclidean and Lorentzian field theory is classically formulated through four equivalent presentations on a flat spacetime:

- Hartle–Hawking 1976 [10]: the temperature of the Euclidean cigar’s τ -circumference identifies as the Hawking temperature;
- Sewell 1982 [26]: the KMS state at inverse-temperature $\beta = 2\pi/\kappa_g$ is the wedge vacuum;
- Bisognano–Wichmann 1976 [1]: the modular automorphism group of the wedge von Neumann algebra acts as boosts around the wedge edge;
- Unruh 1976 [29]: an accelerated observer with proper acceleration a sees a $T_U = a/(2\pi)$ thermal bath in the vacuum.

On a flat background, the four are equivalent under standard Wick-rotation arguments. On a curved background like S^3 they are not obviously equivalent, and at finite cutoff (operator-system level) they require separate construction.

Paper 42 [37] establishes the four-witness theorem at the operator-system level on the truncated Camporesi–Higuchi spectral triple $\mathcal{T}_{n_{\max}}$ via two structurally independent but operator-action conjugate constructions:

Theorem 6.1 (Paper 42, BW- α geometric realisation). *For each $n_{\max} \in \{2, 3, 4, 5\}$, there exists a wedge projection $P_W = \frac{1}{2}(I + R_{\text{polar}})$ on $P_{n_{\max}}\mathcal{H}_{\text{GV}}$ (bit-exact idempotent and Hermitian) and a geometric modular Hamiltonian*

$$K_\alpha^W = J_{\text{polar}} \quad (\text{rotation generator preserving } W), \quad (15)$$

with integer spectrum $\{2\omega m_j\}$ on the full-Dirac basis. The modular flow satisfies $\sigma_{2\pi}(O) = O$ bit-exact (residual $O(\sqrt{\dim \mathcal{H}} \cdot \varepsilon_{\text{machine}})$) for the four witnesses BW, Hartle–Hawking, Sewell, Unruh, with cross-witness collapse residual identically zero.

Theorem 6.2 (Paper 42, BW- γ Tomita–Takesaki realisation). *On the Krein-GNS Hilbert–Schmidt space $\mathcal{H}_{\text{GNS}} = M_{\dim W}(\mathbb{C})$ with KMS state $\rho_W = e^{-K_\alpha^W}/Z$ and the BW choice $H_{\text{local}} := K_\alpha^W/\beta$, the polar decomposition $S = J_{\text{TT}} \cdot \Delta^{1/2}$ gives a Tomita–Takesaki modular Hamiltonian $K_{\text{TT}} = -\log \Delta$ with the same integer spectrum, and the operator-action conjugacy*

$$\sigma_t^{\text{TT}}(a) = \sigma_{-t}^\alpha(a) \quad (16)$$

holds bit-exact at every tested $t \in \{1, \pi, 2\pi\}$. The modular operator Δ satisfies $J_{\text{TT}}^2 = +I$, categorically distinct from the Connes J_{GV} ($KO\text{-dim } 3$, $J^2 = -I$).

Load-bearing scope finding (Paper 42 §7.2). A key derived structural finding is that the local Hamiltonian required by Tomita–Takesaki at $\beta = 2\pi$ *must* be $H_{\text{local}} = K_\alpha^W/\beta$ rather than the intrinsic Camporesi–Higuchi Dirac operator D_{CH} . The intrinsic D_{CH} has half-integer spectrum, so $\beta \cdot D_{\text{CH}}$ would not have integer eigenvalues and the bit-exact closure $e^{i \cdot 2\pi \cdot n} = 1$ would fail. Spectral-action (Chamseddine–Connes [5]) and thermal-time (Connes–Rovelli) paradigms point to different generators on the wedge KMS state. This is the open question O3 of Paper 42 §10, sharpened in Paper 43 to the signature-independence statement below.

6.2 Lorentzian extension at finite cutoff (Paper 43)

Paper 43 [38] extends the four-witness theorem to the Lorentzian Krein space at signature $(3, 1)$, built via the Bizi–Brouder–Besnard prescription [2] at $(m, n) = (4, 6)$ in the Peskin–Schroeder chiral basis, with Lorentzian Dirac

$$D_L = i(\gamma^0 \otimes \partial_t + D_{\text{GV}} \otimes I_{N_t}) \quad (17)$$

per van den Dungen 2016 Proposition 4.1 [30].

Theorem 6.3 (Paper 43 §5–§7, Lorentzian four-witness at finite cutoff). *At every tested $(n_{\text{max}}, N_t) \in \{1, 2, 3\} \times \{1, 11, 21\}$, the Krein-level analogues of (15) and (16) hold bit-exact:*

- (i) *The geometric Lorentzian modular Hamiltonian $K_L^{\alpha, W} = J_{\text{polar}}|_{W_L}$ has integer spectrum on the Krein hemispheric wedge.*
- (ii) *The Krein–Tomita modular operator Δ_L admits a polar decomposition $S = J_{L, \text{TT}} \cdot \Delta_L^{1/2}$; the Tomita modular Hamiltonian K_L^{TT} has integer spectrum.*
- (iii) *$\sigma_{2\pi}^{L, \alpha}(O) = O$ and $\sigma_{2\pi}^{L, \text{TT}}(a) = a$ bit-exact, with cross-witness collapse residual zero.*
- (iv) *At $N_t = 1$, the Lorentzian construction reduces to the Riemannian one of Theorems 6.1–6.2 bit-identically.*

Signature-independence of the load-bearing scope. The deepest structural finding of Paper 43 is that the “ $H_{\text{local}} \neq D_W^L$ ” statement is *signature-independent* at the Riemannian reduction. At $N_t = 1$ the residual norm $\|H_{\text{local}} - D_W^L\|_F$ on the Lorentzian Krein wedge equals the Riemannian-side value at every n_{max} (Paper 43 §7.1):

$$\|H_{\text{local}}|_{(3,1), N_t=1} - D_W^L|_{N_t=1}\|_F = \|H_{\text{local}}^{\text{Riem}} - D_W\|_F \quad (\text{at every } n_{\text{max}}).$$

At $N_t > 1$ the Lorentzian residual is refined upward by the temporal-derivative content of D_L : the spectral-action object D_L has more content than D_{GV} , so the gap between H_{local} (modular-Hamiltonian generator) and D_W^L (spectral-action object) widens with temporal cutoff. This refines the Paper 42 §10 open question O3 from “open about signature-dependence” to a deeper structural feature.

6.3 Pythagorean HS-orthogonality and the $1/\pi^2$ signature

Paper 43 §10.2 records a Pythagorean refinement of the modular vs spectral-action gap. Let $\langle \cdot, \cdot \rangle_{\text{HS}}$ denote the Hilbert–Schmidt inner product on the wedge operator algebra.

Corollary 6.4 (Paper 43 §10.2, HS-orthogonality on the Lorentzian wedge). *For every $\kappa_g > 0$ and every panel cell $n_{\max} \in \{1, \dots, 6\}$, $N_t \in \{1, 11\}$, and witness $w \in \{\text{BW}, \text{HH}_1, \text{HH}_2, \text{Sew}, \text{Unruh}_1, \text{Unruh}_2\}$:*

$$\langle H_{\text{local}}, D_W^L \rangle_{\text{HS}} = 0 \quad (\text{bit-exact}), \quad (18)$$

and the residual norm has the closed form

$$\|H_{\text{local}} - D_W^L\|_F^2 = \frac{\kappa_g^2}{4\pi^2} S(n) + D(n), \quad S(n) = \frac{n(n+1)(n+2)(2n^2+4n-1)}{15}, \quad D(n) = \frac{n(n+1)(n+2)(2n-1)}{20} \quad (19)$$

The closed form (19) is PSLQ-verified at 100 digits with coefficient ceiling 10^6 at 18 panel cells. The $1/\pi^2$ prefactor on $\|H_{\text{local}}\|^2$ is the same M_1 Hopf-base measure signature (9) that appears in the Paper 38 GH convergence rate constant $4/\pi$. This crosswiring between sections 4 and 6 is taken up systematically in §8.

Subspace decomposition. The mechanism is the orthogonal decomposition of the operator algebra $B(\mathcal{K}_W)$ into a diagonal subspace (where $H_{\text{local}} = K_\alpha^W/\beta$ lives, since K_α^W has eigenvalues two m_j on the full-Dirac basis) and an off-diagonal subspace with $\Delta n = \pm 1$ (where D_W^L lives, since the Camporesi–Higuchi Dirac intertwines $\pm m_j$ chirality partners at adjacent shells). Formal operator-theoretic proof of this decomposition is named as Paper 43 open question O4; the PSLQ-verified Pythagorean closed form (19) is empirically sharp at 100 digits and 18 panel cells.

7 Lorentzian propinquity

7.1 Operator-system substrate (Paper 44)

The first step in any Lorentzian propinquity proof is an operator-system substrate. Paper 44 [39] provides it on the chirality-doubled Camporesi–Higuchi spinor space $\mathcal{K}_{n_{\max}, N_t} = \mathcal{H}_{\text{GV}}^{n_{\max}} \otimes \mathbb{C}^{N_t}$ at BBB signature $(m, n) = (4, 6)$.

Theorem 7.1 (Paper 44, propagation number and envelope dependence). *The truncated Lorentzian operator system $\mathcal{O}_{n_{\max}, N_t}^L$ satisfies*

$$\text{prop}(\mathcal{O}_{n_{\max}, N_t}^L) = 2 \quad \text{under the Weyl-doubled achievable envelope}, \quad \text{prop}(\mathcal{O}_{n_{\max}, N_t}^L) = \infty \quad \text{under the full (diagonal)} \quad (20)$$

generalising the Riemannian $\text{prop} = 2$ result for $\mathcal{O}_{n_{\max}}$ (Paper 32 §3.4) to the Lorentzian Krein setting. The witness pair $(M^{2,1,0,0}, M^{2,1,0,0})$ at $n_{\max} = 2$ exhibits $\sim 38\%$ Frobenius residual against $\mathcal{O}_{2,3}^L$. Riemannian-limit recovery at $N_t = 1$ is bit-exact (Frobenius residual = 0.0 in float64) across $n_{\max} \in \{1, 2, 3\}$.*

K^+ -positivity is trivial at the operator-multiplier level. A key structural observation of Paper 44 is that all chirality-doubled scalar multipliers $M \oplus M$ commute with $J_L = \text{chirality-swap}$, and therefore preserve $\mathcal{K}^+$ automatically: $\mathcal{O}^{L,+} = \mathcal{O}^L$ exactly. Consequently the non-trivial Krein-positivity program shifts from operator level to *state* level (Wasserstein–Kantorovich on K^+ states), which is the natural setting for the Paper 45 propinquity proof below.

7.2 K^+ -restricted weak-form Lorentzian propinquity: the degeneracy theorem (Paper 45)

Paper 45 [40] closes off the Lorentzian-propinquity route via a degeneracy theorem (the natural K^+ -weak-form seminorm vanishes identically) on the substrate of Paper 44.

Definition 7.2 (Paper 45 §1.3, K^+ -weak-form Lorentzian propinquity). For two truncated Lorentzian spectral triples $\mathcal{T}_i^L$ ($i = 1, 2$), define

$$\Lambda^L(\mathcal{T}_1^L, \mathcal{T}_2^L) := \Lambda(\mathcal{T}_1^L|_{\mathcal{K}^+}, \mathcal{T}_2^L|_{\mathcal{K}^+}), \quad (21)$$

where Λ is the standard Latrémolière propinquity [15, 16] and the restriction to $\mathcal{K}^+ = \{|\psi\rangle : J_L|\psi\rangle = +|\psi\rangle\}$ is the standard metric spectral triple obtained from the Krein-positive subspace, on which the indefinite Krein product reduces to a positive-definite Hilbert inner product.

Theorem 7.3 (Paper 45, annihilation theorem). *The device of Definition 7.2 is degenerate. For D_L as in (17), Krein-self-adjointness of the anti-Hermitian spatial block forces $\{J_L, D_{\text{GV}} \otimes I\} = 0$, hence $P^+(D_{\text{GV}} \otimes I)P^+ = 0$: the compression annihilates the spatial Dirac. The surviving temporal block is momentum-diagonal and commutes with every momentum-diagonal temporal multiplier, so the restricted Lipschitz seminorm vanishes identically on the operator system and Λ^L of (21) is not a quantum Gromov–Hausdorff-type distance. (Bit-exact: restricted-seminorm kernel 42/42 and 275/275 multipliers at $(n_{\text{max}}, N_t) = (2, 3), (3, 5)$; frozen falsifier in the repository test suite.)*

An earlier revision of the arc claimed a convergence theorem through this device (formerly Theorem 5.1 of Paper 45); the claim was retracted in June 2026. The rate expression

$$C_3^{\text{joint}} \cdot \gamma_{n_{\text{max}}, N_t, T}^{\text{joint}} = O(\log n_{\text{max}}/n_{\text{max}} + T/N_t) \quad (22)$$

survives as a well-defined closed form whose meaning is an unconditional *spatial* statement (Paper 38), the temporal factor being a metrically invisible spectator.

7.3 Three structural simplifications

The original transfer of the Paper 38 five-lemma chain to the joint $\text{SU}(2) \times \text{U}(1)$ setting recorded three structural “simplifications.” Under the degeneracy analysis each is reinterpreted: they are *symptoms* of the temporal factor’s metric invisibility, not conveniences:

PURE_TENSOR Berezin map. The joint Berezin map $B^{\text{joint}} = B^{\text{SU}(2)} \otimes B^{\text{U}(1)}$ factorises as a pure tensor. Each of the four required properties (positivity, contractivity, approximate identity, L3 compatibility) inherits factor-by-factor; no cross-factor verification is needed.

Vanishing time-chirality cross-term. The joint Lichnerowicz commutator simplifies via the structural identity

$$[D_L, a_s \otimes a_t] = i[D_{\text{GV}}, a_s] \otimes a_t \quad (23)$$

because $\{\gamma^0, D_{\text{GV}}\} = 0$ on the chirality-doubled basis. (γ^0 anticommutes with the spatial Camporesi–Higuchi Dirac because D_{GV} is chirality-diagonal in the Peskin–Schroeder chiral basis.) The temporal direction contributes nothing to the joint commutator, and $C_3^{\text{joint}} = C_3^{\text{SU}(2)}$ verbatim. Read correctly, (23) is the *diagnosis*: the momentum-diagonal temporal algebra is invisible to the Dirac commutator.

Corrected cb-norm convention. The smoothing map of the joint central spectral Fejér kernel is unital completely positive, with cb-norm exactly 1; the quantity $2/(n_{\max} + 1)$ formerly quoted as the “joint cb-norm” is the maximum of the $SU(2)$ Plancherel *mass distribution*, a different invariant (the Bożejko–Fendler appeal is withdrawn; the correct statement is elementary). The N_t -dependence of the rate (22) lives entirely in the joint mass-concentration moment (the T/N_t Fejér-on-the-circle rate).

Riemannian-limit recovery at $N_t = 1$ bit-exact. At $N_t = 1$, the joint Berezin reduces bit-identically to $B^{SU(2)} \otimes B_{1,T}^{U(1)} = B^{SU(2)}$, since the $N_t = 1$ temporal Berezin map is the identity on the single Fourier mode. Hence $\Lambda^L(\mathcal{T}_{n_{\max},1,T}^L, \mathcal{T}_{\mathcal{M}}^L)$ agrees with the Riemannian Paper 38 bound bit-exactly. Under the degeneracy analysis this recovery is an automatic formula identity (the panel quantity only ever contained the spatial rate formula), not an operator-level falsifier; the falsifier with teeth is the restricted-seminorm check of Theorem 7.3.

7.4 Honest scope

Theorem 7.3 closes the natural device; the Lorentzian convergence question itself remains open *at the continuum / non-compact level*. The *finite-cutoff* metric question is now understood structurally (Paper 45 § open Q1, 2026-06): the truncated Bisognano–Wichmann boost is *compact* (the modular generator has integer spectrum, $e^{2\pi i K} = I$ at every finite cutoff $\Rightarrow$ the modular flow is the 2π -periodic KMS circle, not a non-compact boost). A compact circle carries only a Euclidean (forward-triangle) metric — which is why the forward chain inequality is the only universal direction and every Lipschitz / boost seminorm is signature-blind — so the finite-cutoff quantum metric is Euclidean and the Lorentzian signature is the Wick rotation $t \rightarrow it$ of that circle, a convention rather than a finite-cutoff metric. The genuine reverse-triangle Lorentzian signature requires the non-compact continuum ($n_{\max} \rightarrow \infty$, type-III) limit, which is the sense in which the question remains open. This is the metric-level dual of the WH7 reading (compactness $\Rightarrow$ discreteness $\Rightarrow \pi/\beta$). Several aspects of scope require explicit comment.

- (G1) **Strong-form Lorentzian propinquity (Paper 45 §1.4 G1).** The K^+ -restriction reduces the indefinite Krein product to a Hilbert inner product on $\mathcal{K}^+$ — but annihilates the spatial Dirac (Theorem 7.3), so no quantum-metric machinery applies on the compression. Strong-form Lorentzian propinquity (Latrémollière-style metric on Krein spectral triples in their own right, without K^+ -restriction) requires an indefinite-inner-product analog of the operator-norm Lipschitz seminorm with appropriate behaviour under Krein-self-adjoint D_L . This remains an open NCG-math problem; multi-month scale.
- (G2) **De-compactification $T \rightarrow \infty$.** The temporal factor is compactified to $U(1)_T$ with finite radius T . De-compactification to non-compact $\mathbb{R}_t$ is a separate program (internal Sprint L3c) requiring renormalisation of the cb-norm bound and reach moment.
- (G3) **Cross-manifold $\mathcal{T}_{S^3} \otimes \mathcal{T}_{\text{Hardy}(S^5)}$.** Blocked at the NCG-framework level by the Riemannian-vs-Hardy-sector asymmetry recorded in Paper 24 §V of the GeoVac corpus (a four-layer Coulomb/HO asymmetry); this is the Coulomb-side claim only.
- (G4) **Inner-factor calibration data.** Higgs / Yukawa selection question of the Connes–Marcolli SM construction; orthogonal to the present convergence theorem, addressed by Paper 32 §8.C verdict POSITIVE-THIN (construction admits Higgs structurally, but no autonomous Yukawa selection from the present data).

8 The master Mellin engine and the cross-arc connection

The most striking structural observation that emerges from reading the arc is that the same transcendental signatures appear in *different* settings. The constant $4/\pi$ of Paper 38 and the

constant $1/\pi^2$ of Paper 43 §10.2 are not numerical coincidences: they are two appearances of the same Hopf-base measure mechanism, the M_1 sub-case of the master Mellin engine catalogued in Paper 32 §8.

8.1 The π -source case-exhaustion theorem (Paper 32 §8)

Theorem 8.1 (Paper 32 §8, π -source case-exhaustion). *Let $\mathcal{C}$ be any finite composition of GeoVac projections applied to a Layer-1 datum on $(\mathcal{A}_{\text{GV}}, \mathcal{H}_{\text{GV}}, D_{\text{GV}})$ or its Bargmann–Segal sibling triple. The following are equivalent:*

- (i) $\mathcal{C}(\text{datum})$ contains a transcendental of the form π^k for $k \in \mathbb{Q}$;
- (ii) $\mathcal{C}$ contains at least one projection whose evaluation includes a continuous integration over a parameter promoted from the discrete graph spectrum;
- (iii) $\mathcal{C}$ engages at least one of three spectral-triple mechanisms,

- M_1 (Hopf-base measure): $\pi = \text{Vol}(S^2)/4$ via the Hopf bundle $S^3 \rightarrow S^2 \times S^1$, equivalently $4/\pi = \text{Vol}(S^2)/\pi^2$;
- M_2 (Seeley–DeWitt): $\sqrt{\pi}$ via heat-kernel coefficients on the Camporesi–Higuchi Dirac spectrum on S^3 (Paper 28 Theorem T9: $\zeta_{D^2}(s)$ is polynomial in π^2 with rational coefficients at every integer s);
- M_3 (vertex-topology Dirichlet L): Catalan G and $\beta(s)$ via quarter-integer Hurwitz shifts in two-loop sunset vertices.

The TS-E3 sharpening of Paper 32 further shows that M_1, M_2, M_3 are sub-cases of a single master Mellin-engine mechanism

$$\pi\text{-source} = \mathcal{M}[\text{Tr}(D^k e^{-tD^2})] \quad \text{with } k \in \{0, 1, 2\}, \quad (24)$$

where $k = 0$ selects M_1 , $k = 1$ selects M_3 , and $k = 2$ selects M_2 .

8.2 M_1 in the Riemannian GH rate

The rate constant $4/\pi$ in (6) is the M_1 Hopf-base measure signature. This is not a derived observation; it is forced by the Plancherel-weight $\times$ Vandermonde–Jacobian cancellation theorem (Paper 40 §3.2) which leaves only the rank-1 Stein–Weiss constant after the $\prod_{\alpha>0}$ products cancel. Equivalently, the single-positive-root Euler–Catalan sum

$$\sum_{d \text{ odd}} \frac{1}{d^2} = \frac{\pi^2}{8}$$

produces the π^2 in the denominator, and the Hopf bundle $S^3 \rightarrow S^2$ produces $\text{Vol}(S^2) = 4\pi$ in the numerator.

8.3 M_1 in the Lorentzian HS-orthogonality

The $1/\pi^2$ prefactor on $\|H_{\text{local}}\|^2$ in the Pythagorean closed form (19) is the same M_1 signature at the operator-system level: the Krein wedge BW choice $H_{\text{local}} = K_{\alpha}^W/(2\pi)$ introduces a factor of $1/(2\pi)$ from the modular-circle circumference, which squared gives $1/(4\pi^2)$ in $\|H_{\text{local}}\|^2$. The Hopf-base measure factor $4/\pi$ of Paper 38 and the $1/\pi^2$ of Paper 43 are two manifestations of the same $\text{Vol}(S^2) = 4\pi$ identity in the master Mellin engine.

Why this matters. The case-exhaustion theorem (Theorem 8.1) is a *spectral-triple-level* statement about which transcendentals can appear in evaluations of GeoVac projection chains. The constant $4/\pi$ of Paper 38 is a *metric-geometry-level* statement about the propinquity rate. The fact that these two abstractly-different levels agree on the same signature is non-trivial: it identifies the metric geometry of the GeoVac spectral triple as carrying the same M_1 signature that the abstract Mellin engine catalogues. Abstract taxonomy meets concrete propinquity rates. Paper 43 §10.2 takes this further: the Lorentzian Pythagorean decomposition at the operator-system level carries the same M_1 signature again, this time on the operator algebra of the Krein wedge.

9 Closures since drafting (May 23, 2026)

The synthesis above (§1–§8) was drafted around the arc Papers 29, 32, 38, 39, 40, 42, 43, 44, 45. On 2026-05-23, four substantive closures landed in a single session, adding Papers 46 and 47 to the arc and promoting two named open follow-ons within Papers 39 and 43. This section records the four closures in summary form.

9.1 Strong-form Lorentzian propinquity (Paper 46)

Status (June 2026): the strong-form convergence claim of this subsection is descoped pending repair with the Paper 45 retraction. On the natural chirality-doubled substrate the operator-norm Lipschitz seminorm L_{op} has kernel far exceeding the scalars (every temporal multiplier $\mathbf{1} \otimes M^{\text{temp}}$ is L_{op} -invisible), so neither side of (25) is a quantum-metric distance between quantum compact metric spaces; the bit-exact panel match below re-evaluates the same spatial rate formula and is not an independent convergence. The enlarged-substrate algebra of Appendix B survives and is the named starting point for the repair. See Paper 46’s Status note and docs/claims_register.md.

Paper 45 §1.4 G1 named the strong-form Lorentzian propinquity (a Latrémolière-style metric on Krein spectral triples without K^+ -restriction) as the principal open NCG-math problem of the Lorentzian arc. Paper 46 [41] *claimed to close* this question on the natural chirality-doubled scalar-multiplier substrate of Paper 44 [39] via the operator-norm Lipschitz seminorm $L_{\text{op}}(a) := \|[D_L, a]\|_{\text{op}}$:

$$\Lambda^{\text{strong}}(\mathcal{T}_{n_{\text{max}}, N_t, T}^L, \mathcal{T}_{\mathcal{M}}^L) \leq C_3^{\text{op}}(n_{\text{max}}) \cdot \gamma_{n_{\text{max}}, N_t, T}^{\text{joint}} \rightarrow 0, \quad (25)$$

with closed-form $C_3^{\text{op}}(n_{\text{max}}) = \sqrt{1 - 1/n_{\text{max}}} \rightarrow 1^-$ and rate $\gamma^{\text{joint}} = O(\log n_{\text{max}}/n_{\text{max}} + T/N_t)$ surviving verbatim from Paper 45. The numerical panel matches Paper 45’s K^+ -weak-form panel *bit-exactly* at every tested cell — the “free-upgrade” reading.

Paper 46’s Appendix B (added 2026-05-22) extends the closure to the *enlarged* operator system with chirality-flipping generators satisfying $\{J_L, M^{\text{flip}}\} = 0$. The substantive new content is the *gradient-norm absorption mechanism*: the strict-strong-form separation between enlarged and K^+ -weak-form substrates manifests at the Lipschitz seminorm level ($L_{\text{op}}(a^{\text{flip}}) > 0$ on enlarged generators while $L_{\text{P45}}^+(a^{\text{flip}}) = 0$ identically), but the separation is paid *entirely in a J -graded gradient-norm extension* $G^{\text{enlarged}} = G^{\text{natural}} + 2\|D_t\|_{\text{op}}\|f^{\text{flip}}\|_{\text{op}}$, leaving the Latrémolière propinquity bound bit-exactly preserved: $\Lambda^{\text{enlarged}}(n_{\text{max}}, N_t) = \Lambda^{\text{P46}}(n_{\text{max}}, N_t)$ at every panel cell.

9.2 De-compactification limit, norm-resolvent / spectral closure (Paper 47)

Status (June 2026): the outer norm-resolvent arrow of (26) and the three-carrier identification survive the Paper 45 retraction unchanged; the inner propinquity arrow (Paper 45/46 main theorem applied along the scaling) is descoped, so the composite is established at the norm-resolvent / spectral level only. See Paper 47’s Status note and docs/claims_register.md.

Paper 45 §1.4 G2 named the de-compactification limit $T \rightarrow \infty$ on the temporal carrier S_T^1 as a separate open program. Paper 47 [42] closes it at the *norm-resolvent / spectral* level via a *two-rate hybrid composite*:

$$\mathcal{T}_{n_{\max}, N_t, T(N_t)}^L \xrightarrow{\Lambda^L\text{-rate } O(\log n_{\max}/n_{\max} + T(N_t)/N_t)} \mathcal{T}_{S^3 \times S_{T(N_t)}^1}^L \xrightarrow{\text{nr-rate } O(e^{-|\Im z|T(N_t)/2})} \mathcal{T}_{S^3 \times \mathbb{R}_t}^L \quad (26)$$

along any *admissible scaling* $T(N_t) \rightarrow \infty$ with $T(N_t)/N_t \rightarrow 0$. The inner arrow is propinquity convergence at coupled cells (Paper 45/46 main theorem applied along the scaling); the outer arrow is norm-resolvent convergence of the Lorentzian Dirac on the compactified spacetime to the Lorentzian Dirac on the genuine spacetime, with explicit exponential tail bound $O(e^{-|\Im z|T/2})$ on compact-support subspaces via Reed–Simon [43] § XIII.16 / Bloch-decomposition machinery and the Krein-isometric embedding $\tilde{J}_T : L^2(S_T^1) \hookrightarrow L^2(\mathbb{R})$.

A strategic clarification appears in Paper 47 §1.1: the natural *inductive-limit propinquity* framework of Latrémolière [16] § 5 for AF C*-algebras is structurally wrong for the present setting because AF C*-algebras have totally-disconnected Gelfand spectrum while $\mathbb{R}$ is connected, so $C_0(\mathbb{R})$ cannot be an AF inductive limit. Norm-resolvent convergence is the right framework at the spectral level.

Paper 47 §6 also establishes a *three-carrier identification*: the periodic compact $S^3 \times S_T^1$ (Paper 45 substrate), the bounded Dirichlet $S^3 \times [-T/2, T/2]$ (Paper 43 / Sprint L2-B/E substrate), and the non-compact $S^3 \times \mathbb{R}_t$ (Paper 47 limit) all converge in norm-resolvent sense to the same physical Lorentzian Dirac as $T \rightarrow \infty$. This resolves the apparent tension between Paper 43’s bounded-interval architecture (chosen to preserve causal structure / no CTCs via Geroch’s theorem) and Paper 45’s periodic-compact architecture (chosen to enable Latrémolière propinquity machinery on a compact carrier). Both are valid compact approximations to the same physical Lorentzian Dirac.

A numerical-verification panel (Sprint L3c- α .2, 2026-05-23) at 24 cells $(n_{\max}, N_t) \in \{2, 3\} \times \{3, 5, 7, 11\}$ along three scalings ($T_0 = 2\pi$ canonical, $T_1 = N_t / \log N_t$ sub-linear, $T_2 = \sqrt{N_t}$ square-root) confirms Λ^L bit-exact T -and- N_t -independence at fixed $n_{\max}$: $\Lambda(n_{\max} = 2) = 2.074551$ and $\Lambda(n_{\max} = 3) = 1.610060$ to 6+ significant figures across all three scalings, matching Paper 45 Table 1 verbatim. Structural reason: $\gamma^{U(1)}(N_t, T)$ is subordinate to $\gamma^{\text{SU}(2)}(n_{\max})$ at every admissible-scaling cell, so the joint rate $\gamma^{\text{joint}} = \max(\gamma^{\text{SU}(2)}, \gamma^{U(1)}) = \gamma^{\text{SU}(2)}$ exactly. The empirical confirmation upgrades the L3c- α theorem from “analytical corollary of Paper 45” to “analytically AND empirically established.”

9.3 Pythagorean orthogonality formal proof (Paper 43 §10.2 promotion)

Paper 43 §11 O4 named a formal proof of the diagonal/off-diagonal subspace-decomposition mechanism behind the L2-F.1 (2026-05-17) empirical observation $\langle H_{\text{local}}, D_L^W \rangle_{\text{HS}} = 0$ as a follow-on. The formal proof landed on 2026-05-23 via a $\mathbb{Z}_2$ wedge-chirality parity argument and is now recorded in Paper 43 §10.2 as a formal theorem.

The argument has three structural inputs from Papers 42/43/44 [37–39]: (I1) the wedge chirality grading Π_W (BBB Krein-chirality γ^5 restricted to the wedge) is a unitary self-adjoint involution; (I2) $[\Pi_W, H_{\text{local}}] = 0$ (even parity of the geometric BW- α generator $K_\alpha^W = J_{\text{polar}}^W$ under wedge chirality); (I3) $\{\Pi_W, D_L^W\} = 0$ (odd parity of the wedge-restricted Lorentzian Dirac, via $\{\gamma^0, \gamma^5\} = 0$ on the temporal factor and the unfolded-wedge basis change on the spatial factor).

The orthogonality follows from a parity / cyclicity-of-trace lemma: for any unitary involution Π on a Hilbert space, $[\Pi, A] = 0$, $\{\Pi, B\} = 0$ imply $\text{Tr}(A^\dagger B) = 0$ via $\text{Tr}(X) = \text{Tr}(\Pi X \Pi^{-1})$ giving $\text{Tr}(A^\dagger B) = \text{Tr}(A^\dagger \cdot (-B)) = -\text{Tr}(A^\dagger B)$, forcing 0. The bit-exact L2-F.1 numerical panel at 18 cells is empirical corroboration of the now-algebraic identity at machine precision.

9.4 Master theorem of the math.OA arc

Paper 39 §6 (iii) and (v) named two open follow-ons: (iii) the k -fold extension of the Paper 39 two-factor tensor-product propinquity to $\mathcal{T}_{S_3^{\otimes k}}$, and (v) the higher-rank tensor product $\mathcal{T}_{G_1} \otimes \mathcal{T}_{G_2}$ at arbitrary compact connected Lie group factors. Both closed on 2026-05-23.

(iii) **k -fold extension — $\sqrt{k}$ constant (sketch).** The k -fold extension to $SU(2)$ factors uses the graded Leibniz rule and the operator-norm *triangle* bound, giving $C_3^{(k)} \leq \sqrt{k} \sqrt{(N_* - 1)/(N_* + 1)} \nearrow \sqrt{k}$ where $N_* := \max_j n_j$. *Correction (2026-06-18):* an earlier draft claimed a **k -independent** constant $\sqrt{(N_* - 1)/(N_* + 1)} \rightarrow 1^-$ via a “Pythagorean Leibniz” sum-of-squares identity $\|\sum_j X_j\|_{\text{op}}^2 = \sum_j \|X_j\|_{\text{op}}^2$. That identity is **false** (pairwise anticommutation of the $D^{(k)}$ summands does not make the commutator terms operator-norm orthogonal — verified maximally violated at $k = 2$ on the actual harmonics). The genuine $\sqrt{k}$ factor is real, not eliminated; for each fixed k convergence still holds (finite constant), but the constant grows with the number of factors. Only the $k = 2$ case is proved (Paper 39); $k > 2$ is a sketch.

(v) **Master theorem — universal compact-Lie-group factors (sketch).** Combining the k -fold extension with the universal $4/\pi$ rate of Paper 40 [36] gives the expected multi-factor form

$$\Lambda^{(k)}(\mathcal{T}_{n_1, \dots, n_k}^{(k)}, \mathcal{T}_{G_1 \times \dots \times G_k}^{(k)}) \leq C_3^{(k)} \cdot \max_j \gamma_{n_j}^{G_j}, \quad C_3^{(k)} \leq \sqrt{k} \sup_j C_3^{G_j}(n_j) \nearrow \sqrt{k}, \quad (27)$$

for arbitrary $k \geq 1$ and arbitrary compact connected Lie groups $G_1, \dots, G_k$ with bi-invariant metrics. *Correction (2026-06-18):* this was boxed as a closed “master theorem” with a **k -independent** constant $\sup_j C_3^{G_j}(n_j)$; with the corrected triangle bound the constant carries a genuine $\sqrt{k}$ and is not k -independent. The joint rate still inherits the universal $4/\pi$ asymptotic; but this multi-factor / general- G form is a **sketch** (only the $k = 2$, $SU(2)$ and $k = 1$ cases are proved).

The master theorem subsumes Papers [34] ($k = 1$, $G = SU(2)$), [36] ($k = 1$, general G), and the two-factor case treated above (§4.8, $k = 2$, $G_1 = G_2 = SU(2)$) as special cases. Paper 45’s compact-temporal Lorentzian extension at $S^3 \times S_T^1$ is the $k = 2$, $G_1 = SU(2)$, $G_2 = U(1)$ special case, providing independent justification of Paper 45’s main theorem from the master construction.

10 Further closures: Krein–Mondino–Sämman bridge, OSLPLS, and bit-exact CFT-on-sphere

The arc admits three further closures landed subsequent to the May 22, 2026 drafting of this synthesis. These add three further papers (48, 49, 50) and extend the math.OA standalone count from eleven to fourteen.

10.1 Krein-pointed proper QMS and the Mondino–Sämman bridge (Paper 48)

Status (June 2026): the metric-level content of this subsection is descoped with the Paper 45 retraction; the bridge design survives as a conditional proposal. See the Status notes in Papers 48–49 and docs/claims_register.md.

Paper 48 [44] constructs Krein-pointed proper quantum metric spaces **KreinMetaMet**_{pp} as the natural Krein-lift of Latrémolière’s recent pointed proper quantum metric space framework [53], and builds a *bridge functor* $W : \mathbf{KreinMetaMet}_{\text{pp}} \rightarrow \mathbf{LorPLG}_{\text{cov}}$ to the Mondino–Sämman synthetic Lorentzian pre-length space category [19, 20]. The bridge is identified via the Connes–Rovelli thermal-time hypothesis [52]: the modular distance K on the Krein side measures

thermal time (additive, forward triangle), while the proper-time distance ℓ^L on the Mondino–Sämman side measures geometric proper time (super-additive, reverse triangle); Connes–Rovelli identifies these as two readings of the same wedge KMS state, related by Wick rotation. The bridge is functorial, not isometric.

Conditional on the bridge construction — whose K^+ -weak-form input is descoped with the Paper 45 retraction — Bridge Theorem 6.4 with all four properties (B1)+(B2)+(B3)+(B4) would yield seven newly accessible theorems on the GeoVac hemispheric wedge, including: (T3) bit-exact panel inheritance — the $\Lambda(2, 3) = 2.0746$, $\Lambda(3, 5) = 1.6101$, $\Lambda(4, 7) = 1.3223$ panel of Paper 45 would transport verbatim; (T5) synthetic Pythagorean orthogonality with the $1/\pi^2$ master Mellin engine M_1 signature as the inverse-squared bridge cover slab-period; and (T6) a proposed G2 metric-level *route* for the GeoVac wedge specifically, via the bridge $\rightarrow$ MS pLGH workaround. This last item is the most substantive design content of Paper 48: it offers a conditional synthetic-side workaround for the Paper 45 §1.4 G2 metric-level problem (contingent on the descoped bridge, hence a proposed route rather than a closure), sidestepping the absence of a published non-compact Latrémolière propinquity by working through the Mondino–Sämman category instead.

Paper 48 is the tenth math.OA standalone in the arc, sibling to Papers 38, 42, 43, 44, 45, 46, 47, 29, 32, 40.

10.2 OSLPLS and the strong-form bridge: twin paradox as quantum information (Paper 49)

Status (June 2026): the Λ -inheritance and K^+ -framed claims of this subsection are descoped with the Paper 45 retraction; the cocycle-deficit algebra and the OSLPLS category design survive. See Paper 49’s Status note and `docs/claims_register.md`.

Paper 49 [45] closes Paper 48 §8.1’s open question Q1’ by constructing an *OSLPLS* (Operator-System Lorentzian Pre-Length Space) category and a strong-form bridge functor $W^{\text{flip}} : \mathbf{KreinMetaMet}_{\text{pp}}^{\text{flip}} \rightarrow \mathbf{OSLPLS}_{\text{cov}}$ on the Paper 46 Appendix B enlarged substrate (chirality-flipping generators $M^{\text{flip}} = \text{diag}(W, -W)$ where $\{J, M\} = 0$). The closure rests on a genuinely new operator-algebraic construction: the Connes–Rovelli thermal-time stack across *distinct* KMS states, extending Connes–Rovelli [52] from the single-KMS-state setting to multi-state stack consistency. Properties (B1’)+(B2’)+(B3’) hold theorem-grade; (B4’), the metric-level Λ -inheritance, is *descoped* (the Krein-side seminorm is degenerate, Paper 45/46).

The headline structural finding is the *twin-paradox-as-quantum-information* reading: strict super-additivity of the OSLPLS reverse triangle on off-orbit triples is the operator-algebraic dual of the special-relativistic twin paradox, with the deficit quantified by Datta’s max-divergence chain inequality [54]. The detoured worldline pays *two* max-divergence cocycle costs $\Delta_{\text{max}} S^{\sigma_1 \rightarrow \sigma_2} + \Delta_{\text{max}} S^{\sigma_2 \rightarrow \sigma_3}$; the direct worldline pays *one* cost $\Delta_{\text{max}} S^{\sigma_1 \rightarrow \sigma_3}$; the strict inequality follows from Datta 2009’s max-divergence chain inequality (the Umegaki relative entropy itself does not satisfy a chain inequality in general — see Paper 49 §5.4 and Q3’ for the honest scope). Numerical verification at $(n_{\text{max}}, N_t) \in \{(2, 3), (3, 5), (4, 7)\}$ confirms: Λ bit-exact at all three panel cells (rate-formula consistency, not a metric convergence), *illustrative* Umegaki relative-entropy deficits 66.998 / 68.720 / 81.256 nats positive and monotone-increasing in n_{max} (the load-bearing D_{max} chain is verified 96/96), Riemannian-limit recovery at $N_t = 1$ bit-exact.

The Bousso–Chandrasekaran–Rath–Shahbazi–Moghaddam 2020 “kink transform” / Connes cocycle flow gravity dual [51] appears here as a natural bulk reading of the OSLPLS construction (Paper 49 §11): the strict super-additivity of the OSLPLS reverse triangle on off-orbit triples translates, under the kink-transform dictionary, to a positive relative entropy on the gravitational dual — recovering the BCFM positivity statement from the operator-algebraic side.

Paper 49 is the eleventh math.OA standalone in the arc. Closed-form expression for the cocycle entropy production deficit as a function of OSLPLS state-space coordinates remains the named open question (Paper 49 §10, Q2’).

10.3 Bit-exact CFT-on-sphere partition functions and the master Mellin engine M_2/M_3 orthogonal decomposition (Paper 50)

Paper 50 [46] demonstrates that the framework’s existing Hurwitz-zeta machinery (used throughout Papers 38–47 for spectral sums on the Camporesi–Higuchi tower) reproduces continuum CFT-on-sphere partition functions *bit-exactly*, via a closed-form combination of $\zeta'(-2)$, $\zeta'(-4)$, $\zeta(3)$, $\zeta(5)$, $\log 2$, and π identified by PSLQ at 80–100 digit precision.

Bit-exact match on S^3 . For the conformally coupled scalar and Camporesi–Higuchi Dirac on round S^3 ,

$$\begin{aligned} F_{\text{scalar}}^{S^3} &= \frac{\log 2}{8} - \frac{3\zeta(3)}{16\pi^2} \approx 0.0638 && \text{(PSLQ integer relation [8, 2, -3])}, \\ F_{\text{Dirac}}^{S^3} &= \frac{\log 2}{4} + \frac{3\zeta(3)}{8\pi^2} \approx 0.21896 && \text{(symbolic Hurwitz expansion)}. \end{aligned}$$

These reproduce the Klebanov–Pufu–Safdi values [48] bit-exactly.

Orthogonal decomposition and log 3 cancellation theorem on S^5 . On the (scalar, Dirac) sector, the framework reveals an orthogonal decomposition:

$$F_D + 2F_s = \frac{\log 2}{2} \quad (M_2 \text{ sector}), \quad F_D - 2F_s = \frac{3\zeta(3)}{4\pi^2} \quad (M_3 \text{ sector}).$$

On S^5 , the pattern extends: $F_{\text{scalar}}^{S^5} = -\log(2)/32 - \zeta(3)/(32\pi^2) + 15\zeta(5)/(64\pi^4)$ (PSLQ relation [32, -2, -2, 15]), $F_{\text{Dirac}}^{S^5} = -(3\log 2)/128 - 5\zeta(3)/(128\pi^2) - 15\zeta(5)/(256\pi^4)$. The log 3 contributions that arise from zero-mode boundary structure in the minimal-scalar variant cancel algebraically against M_2 -sector identities (Paper 50 Theorem 7.2), restricting the substrate-natural ring to $\log 2 \cdot \mathbb{Q} + \zeta(3)/\pi^2 \cdot \mathbb{Q} + \zeta(5)/\pi^4 \cdot \mathbb{Q}$ on S^5 .

S^7 negative. On S^7 , no PSLQ relation in any tested natural ring closes the scalar value $\approx -0.001594\dots$, including bases extended with $\log 3$, $\log 5$, $\log 7$, Catalan G , polygamma at $1/2$, and zeta-zeta products of total weight up to 7 (Paper 50 §8). Cross-check against direct numerical computation of the Hurwitz expansion verifies that the framework machinery is faithful and the negative is structural, not a computational artifact. This identifies an open target: which spectral object on S^7 would close the substrate-natural ring at this multiplicity?

Cataloguing the master Mellin engine in physics. The S^3 and S^5 closures are recorded in Paper 50 §9 as the first two entries of a running catalogue of master-Mellin-engine appearances on the physics side. Paper 50 also flags structural correspondences to other M_2 -sector results in the QFT literature, including Henningson–Skenderis holographic Weyl anomaly [49] (M_2 Seeley–DeWitt content), TT-bar / cutoff AdS [50], and the Chamseddine–Connes spectral action [5] (also M_2). These correspondences are flagged as structural, not as new derivations.

Structural reading: scale-freedom on substrate $\leftrightarrow$ scale-invariance in physics. The bit-exact CFT closures on S^3 and S^5 are structurally explained by the framework’s substrate-level scale-freedom: the GeoVac packing axiom outputs a graph whose Laplacian eigenvalues are pure rationals $n^2 - 1$, the multiplicities are integers, the Wigner-symbol couplings are exact rationals, and the packing diameter d_0 *cancels out* of every spectral quantity. Physical scales enter only through the Fock projection focal length p_0 , which is a stereographic conformal map. This scale-freedom on the substrate is the geometric reflection of the same principle that conformal field theory [48] embodies on the physics side: a CFT has no preferred length scale at the level of its action and observables; the GeoVac graph has no preferred length scale at the

level of its eigenvalues and matrix elements. The framework’s ability to host continuum CFT-on-sphere observables bit-exactly reflects this structural alignment. This is *not* a claim that GeoVac *is* a CFT; it is a structural reflection that explains *why* the framework reproduces what it does. (The substrate-side account of this principle is developed in the Group 3 foundations synthesis [47].)

Paper 50 is the twelfth math.OA standalone in the arc.

11 Open questions and outlook

The arc closes the convergence side of the operator-algebra story on the Riemannian / spatial side (Paper 38, unconditional) and the modular-Hamiltonian side at the operator-system level on Riemannian (Paper 42) and Lorentzian (Paper 43) backgrounds; the Lorentzian *convergence* question is left open by Paper 45’s annihilation theorem. Several questions remain open.

Strong-form Lorentzian propinquity (Q1) — OPEN (Paper 46 descoped, §9.1). Strong-form Lorentzian propinquity — a Latrémolière-style metric on Krein spectral triples in their own right, without K^+ -restriction (Paper 45 §1.4 G1) — remains the principal open NCG-math problem of the Lorentzian arc. Paper 46 [41] proposed a closure via the operator-norm Lipschitz seminorm $L_{\text{op}}(a) = \|[D_L, a]\|_{\text{op}}$ on the chirality-doubled scalar-multiplier substrate of Paper 44, but that seminorm has kernel far exceeding the scalars (the temporal multipliers are L_{op} -invisible), so the construction is not a quantum-metric distance and is descoped pending repair; the “free-upgrade” bit-exact panel match is a re-evaluation of the spatial rate formula, not an independent Lorentzian convergence. The enlarged-substrate algebra of Paper 46 Appendix B (chirality-flipping generators with $\{J, M\} = 0$, where the commutators are non-zero) survives and is the named starting point for the repair. A genuine strong-form theorem requires an indefinite-inner-product Lipschitz seminorm compatible with Krein-self-adjoint D_L ; multi-month scale.

De-compactification $T \rightarrow \infty$ (G2) — CLOSED at norm-resolvent level by Paper 47 (§9.2); G2-metric open as multi-month frontier. The temporal factor $U(1)_T$ is finite; the standard physics applications (Hawking, Unruh) require non-compact $\mathbb{R}_t$. The rate $T/N_t \rightarrow 0$ in (22) is the joint mass-concentration on $U(1)_T$ as $N_t \rightarrow \infty$ at fixed T ; the $T \rightarrow \infty$ limit is closed at the *norm-resolvent* / *spectral* level by Paper 47 [42]’s two-rate hybrid composite (26). The remaining open question is the *metric* / *propinquity* level closure on a non-compact carrier (G2-metric or equivalently G1-genuine on the non-compact substrate), which requires a non-compact extension of Latrémolière’s propinquity not in the published literature as of May 2026. This is named multi-month frontier (Sprint L3e in our internal taxonomy), with the path-of-least-resistance being a pointed-propinquity bridge to the Mondino–Sämman synthetic Lorentzian GH program (§9.2).

Cross-manifold $\mathcal{T}_{S^3} \otimes \mathcal{T}_{\text{Hardy}(S^5)}$ (G3, Coulomb-side claim only). The natural tensor product of the Coulomb S^3 spectral triple with the Bargmann–Segal Hardy-sector spectral triple on S^5 [31] is blocked at the NCG-framework level by a four-layer asymmetry recorded in Paper 24 §V. The four layers are: (i) spectrum-computing role of L_0 (Coulomb-side only, by Paper 23 Fock rigidity); (ii) calibration π (Coulomb-side only); (iii) non-abelian Wilson gauge with natural matter coupling (both sides have Wilson, only Coulomb-side has natural matter coupling because the Bargmann tower $(N, 0)$ irreps have N -dependent dimensions and no uniform-rep matter sector); and (iv) modular-Hamiltonian structure of the wedge KMS state (Coulomb-side admits the Krein wedge HS-orthogonality of Corollary 6.4 with closed-form M_1 prefactor; the Hardy-sector side does not admit the construction at all). Closing G3 requires

new NCG-framework machinery to bridge Riemannian and Hardy-sector spectral triples in a single tensor product; we make no claim about whether this is achievable.

Higgs as inner fluctuation (G4). The GeoVac triple is naturally combined with a finite almost-commutative factor $\mathcal{A}_{\text{GV}} \otimes (\mathbb{C} \oplus \mathbb{H})$ to produce an electroweak-like sector. The Connes axiom audit at finite cutoff (Paper 32 §8.C) gives verdict POSITIVE-THIN: the construction *admits* a Higgs sector structurally (i.e. the inner fluctuation $D \mapsto D + \omega + \varepsilon' J\omega J^{-1}$ produces a non-trivial off-diagonal $\mathbb{C} \leftrightarrow \mathbb{H}$ block iff the finite Yukawa Y_F is non-zero), but *GeoVac does not autonomously select* Y_F from its internal data. In Marcolli-vS terms, GeoVac sits on the Yang–Mills-without-Higgs side of the Perez-Sanchez [23] clarification, with calibration data (the Yukawa) supplied externally. Paper 32 §8.C records this as the standing structural verdict.

State-side complement of the dictionary. Paper 44 highlights that the non-trivial K^+ -positivity content of the Lorentzian operator system sits at the *state* level, not the operator level. The natural object is the Wasserstein–Kantorovich metric on K^+ -states. Paper 45’s K^+ -weak-form analysis (now the annihilation theorem) is the first probe in this direction and shows the operator-level compression is degenerate; substantially more mathematics on Wasserstein–Kantorovich on Krein-positive *states* is required to complete the state-side dictionary.

Concluding remark. The structural identification of GeoVac as an almost-commutative spectral triple in the Marcolli–van Suijlekom lineage is proved at the state-space Gromov–Hausdorff level by Paper 38 (Riemannian, single factor), extended by Paper 39 (tensor product, state-space GH) and Paper 40 (all compact simple Lie groups). The Lorentzian K^+ -weak-form extension is *descoped*: Paper 45 proves a degeneracy theorem (the natural Lorentzian-proximity seminorm vanishes identically), and what survives on the temporal factor is Paper 45’s signature-agnostic product-carrier rate (not a Lorentzian convergence). The metric geometry of the truncated triples carries the same M_1 Hopf-base measure signature ($4/\pi$ in the Riemannian rate, $1/\pi^2$ in the Lorentzian Pythagorean closed form) that the abstract π -source case-exhaustion theorem (Theorem 8.1) identifies at the spectral-triple level. The cross-arc identification of these signatures is the deepest structural observation of the corpus.

References

- [1] J. J. Bisognano and E. H. Wichmann, *On the duality condition for quantum fields*, J. Math. Phys. **17** (1976), 303–321.
- [2] N. Bizi, C. Brouder, and F. Besnard, *Space and time dimensions of algebras with application to Lorentzian noncommutative geometry and quantum electrodynamics*, J. Math. Phys. **59** (2018), 062303; arXiv:1611.07062.
- [3] M. Bożejko and G. Fendler, *Herz–Schur multipliers and uniformly bounded representations of discrete groups*, Arch. Math. **57** (1991), 290–298.
- [4] R. Camporesi and A. Higuchi, *On the eigenfunctions of the Dirac operator on spheres and real hyperbolic spaces*, J. Geom. Phys. **20** (1996), 1–18; arXiv:gr-qc/9505009.
- [5] A. H. Chamseddine and A. Connes, *Noncommutative geometry as a framework for unification of all fundamental interactions including gravity. I*, Fortsch. Phys. **58** (2010), 553–600.
- [6] A. Connes, *Noncommutative Geometry*, Academic Press, 1995.
- [7] A. Connes and W. D. van Suijlekom, *Spectral truncations in noncommutative geometry and operator systems*, Comm. Math. Phys. **383** (2021), 2021–2067; arXiv:2004.14115.

- [8] C. Farsi and F. Latrémolière, *Continuity of spectral propinquity for spectrally truncated spheres*, preprint, 2024.
- [9] C. Farsi and F. Latrémolière, *Continuity for the spectral propinquity of the Dirac operators associated with an analytic path of Riemannian metrics*, arXiv:2504.11715, 2025.
- [10] J. B. Hartle and S. W. Hawking, *Path-integral derivation of black-hole radiance*, Phys. Rev. D **13** (1976), 2188–2203.
- [11] E. Hawkins, *Quantization of equivariant vector bundles, and the correspondence with deformation quantization*, Comm. Math. Phys. **215** (2000), 409–432.
- [12] E. Hekkelman, *Truncations of the circle and Connes’ geometric formula*, M.Sc. thesis, Radboud University, 2022; arXiv:2206.13744.
- [13] E. Hekkelman and E. McDonald, *Spectral truncations of $\mathbb{T}^d$ and quantum metric geometry*, J. Noncommut. Geom., to appear (2024); arXiv:2403.18619.
- [14] E. Hekkelman and E. McDonald, *A noncommutative integral on spectrally truncated spectral triples, and a link with quantum ergodicity*, J. Funct. Anal., to appear (2025); arXiv:2412.00628.
- [15] F. Latrémolière, *The Gromov–Hausdorff propinquity for metric spectral triples*, Adv. Math. **415** (2023), Paper No. 108876, 88pp; preprint arXiv:1811.10843, November 2018.
- [16] F. Latrémolière, *The Gromov–Hausdorff propinquity*, Trans. Amer. Math. Soc. **370** (2018), 365–411.
- [17] M. Leimbach and W. D. van Suijlekom, *Gromov–Hausdorff convergence of spectral truncations of the torus*, Adv. Math. **439** (2024), Paper No. 109496.
- [18] M. Marcolli and W. D. van Suijlekom, *Gauge networks in noncommutative geometry*, J. Geom. Phys. **75** (2014), 71–91; arXiv:1301.3480.
- [19] E. Minguzzi and S. Suhr, *Bounded Lorentzian metric spaces*, Lett. Math. Phys. **114** (2024), Paper No. 73; arXiv:2209.14384.
- [20] A. Mondino and C. Sämann, *Synthetic Lorentzian Gromov–Hausdorff convergence and pre-compactness*, arXiv:2504.10380, 2025.
- [21] V. I. Paulsen, *Completely Bounded Maps and Operator Algebras*, Cambridge Studies in Advanced Mathematics 78, Cambridge University Press, 2002.
- [22] C. I. Perez-Sanchez, *Bratteli networks and the Spectral Action on quivers*, arXiv:2401.03705, 2024.
- [23] C. I. Perez-Sanchez, *Comment on ‘Gauge networks in noncommutative geometry’*, arXiv:2508.17338, 2025.
- [24] J.-P. Pier, *Amenable Locally Compact Groups*, Wiley, 1984.
- [25] G. Pisier, *Similarity Problems and Completely Bounded Maps*, Lecture Notes in Math. 1618, Springer, 2nd ed., 2001.
- [26] G. L. Sewell, *Quantum fields on manifolds: PCT and gravitationally induced thermal states*, Ann. Phys. (NY) **141** (1982), 201–224.
- [27] E. M. Stein and G. Weiss, *Introduction to Fourier Analysis on Euclidean Spaces*, Princeton Math. Ser. 32, Princeton Univ. Press, 1971.

- [28] M. Toyota, *Quantum Gromov–Hausdorff convergence of spectral truncations for groups with polynomial growth*, arXiv:2309.13469, 2023.
- [29] W. G. Unruh, *Notes on black-hole evaporation*, Phys. Rev. D **14** (1976), 870–892.
- [30] K. van den Dungen, *Krein spectral triples and the fermionic action*, Math. Phys. Anal. Geom. **19** (2016), Art. 4, 22pp; arXiv:1505.01939.
- [31] J. Loutey, *Bargmann–Segal lattice for the 3D harmonic oscillator and the Coulomb/HO structural asymmetry*, GeoVac internal preprint (2026), DOI via Zenodo.
- [32] J. Loutey, *The GeoVac Hopf graphs are Ramanujan: Ihara zeta, integer-algebraic zeros, and the scope boundary on Selberg-on- hydrogen*, GeoVac internal preprint (2026), DOI via Zenodo.
- [33] J. Loutey, *The GeoVac spectral triple, the master Mellin engine case-exhaustion theorem, and the GH-convergence theorem on S^3* , GeoVac internal preprint (2026), DOI via Zenodo.
- [34] J. Loutey, *State-space Gromov–Hausdorff convergence of truncated Camporesi–Higuchi spectral triples on S^3* , GeoVac internal preprint (2026), DOI via Zenodo.
- [35] J. Loutey, *Tensor-product propinquity convergence for distinct-focal-length truncated Camporesi–Higuchi spectral triples on S^3* , GeoVac internal preprint (2026), DOI via Zenodo.
- [36] J. Loutey, *State-space Gromov–Hausdorff convergence of spectral truncations of compact Lie groups with bi-invariant metric*, GeoVac internal preprint (2026), DOI via Zenodo.
- [37] J. Loutey, *Tomita–Takesaki modular structure on truncated $SU(2)$ spectral triples: four-witness Wick-rotation literal identification at finite cutoff*, GeoVac internal preprint (2026), DOI via Zenodo.
- [38] J. Loutey, *Lorentzian extension of the four-witness Wick-rotation theorem at finite cutoff*, GeoVac internal preprint (2026), DOI via Zenodo.
- [39] J. Loutey, *Operator-system extension of the BBB Krein spectral triple at finite cutoff: propagation number and envelope dependence on the Camporesi–Higuchi spinor bundle*, GeoVac internal preprint (2026), DOI via Zenodo.
- [40] J. Loutey, *Lorentzian propinquity on truncated $SU(2) \times U(1)_T$ Krein spectral triples: a degeneracy theorem*, GeoVac internal preprint (2026), DOI via Zenodo.
- [41] J. Loutey, *Strong-form Lorentzian propinquity on truncated $SU(2) \times U(1)_T$ Krein spectral triples*, GeoVac internal preprint (2026), DOI via Zenodo.
- [42] J. Loutey, *Norm-resolvent convergence to the non-compact Lorentzian Krein spectral triple on $S^3 \times \mathbb{R}_t$ and the two-rate hybrid limit*, GeoVac internal preprint (2026), DOI via Zenodo.
- [43] M. Reed and B. Simon, *Methods of Modern Mathematical Physics, Vol. IV: Analysis of Operators*, Academic Press, 1978.
- [44] J. Loutey, *Krein-pointed proper quantum metric spaces and the bridge to Mondino–Sämman Lorentzian pre-length spaces, with application to the GeoVac hemispheric wedge*, GeoVac internal preprint (2026), DOI via Zenodo.
- [45] J. Loutey, *OSLPLS and the strong-form Krein–Mondino–Sämman bridge: twin-paradox-as-quantum-information via the Connes–Rovelli thermal-time stack across distinct KMS states*, GeoVac internal preprint (2026), DOI via Zenodo.

- [46] J. Loutey, *Bit-exact CFT-on-sphere partition functions on S^3 and S^5 from the master Mellin engine M_2/M_3 orthogonal decomposition*, GeoVac internal preprint (2026), DOI via Zenodo.
- [47] J. Loutey, *The foundations arc of the GeoVac framework: a reading guide for spectral graph theory, the dimensionless vacuum, and the natural geometry hierarchy*, GeoVac internal synthesis (2026).
- [48] I. R. Klebanov, S. S. Pufu, and B. R. Safdi, *F-theorem without supersymmetry*, J. High Energy Phys. **10** (2011), 038; arXiv:1105.4598.
- [49] M. Henningson and K. Skenderis, *The holographic Weyl anomaly*, J. High Energy Phys. **07** (1998), 023; arXiv:hep-th/9806087.
- [50] T. Hartman, J. Kruthoff, E. Shaghoulian, and A. Tajdini, *Holography at finite cutoff with a T^2 deformation*, J. High Energy Phys. **03** (2019), 004; arXiv:1807.11401.
- [51] R. Bousso, V. Chandrasekaran, P. Rath, and A. Shahbazi-Moghaddam, *Gravity dual of Connes cocycle flow*, Phys. Rev. D **102** (2020) 066008; arXiv:2007.00230.
- [52] A. Connes and C. Rovelli, *Von Neumann algebra automorphisms and time- thermodynamics relation in generally covariant quantum theories*, Class. Quantum Grav. **11** (1994), 2899–2917.
- [53] F. Latrémolière, *The hypertopology of the Gromov–Hausdorff propinquity for pointed proper quantum metric spaces*, arXiv:2512.03573, December 2025.
- [54] N. Datta, *Min- and max-relative entropies and a new entanglement monotone*, IEEE Trans. Inf. Theory **55** (2009), 2816–2826; arXiv:0803.2770.