

**Problem 1.** Verify that the functional

$$J[u] = \int_R \left[ \left( \frac{\partial u}{\partial x} \right)^2 + \left( \frac{\partial u}{\partial y} \right)^2 \right] dx dy \quad (1)$$

is invariant under a rotation of the coordinate axis:

$$\tilde{x} = x \cos \epsilon + y \sin \epsilon, \quad \tilde{y} = -x \sin \epsilon + y \cos \epsilon. \quad (2)$$

**Solution.** The functional is invariant if  $J[u(x, y)] = J[u(\tilde{x}, \tilde{y})]$ . By the chain rule,

$$\frac{\partial}{\partial x} = \frac{\partial}{\partial \tilde{x}} \frac{\partial \tilde{x}}{\partial x} + \frac{\partial}{\partial \tilde{y}} \frac{\partial \tilde{y}}{\partial x} = \cos \epsilon \frac{\partial}{\partial \tilde{x}} - \sin \epsilon \frac{\partial}{\partial \tilde{y}}, \quad \frac{\partial}{\partial y} = \frac{\partial}{\partial \tilde{x}} \frac{\partial \tilde{x}}{\partial y} + \frac{\partial}{\partial \tilde{y}} \frac{\partial \tilde{y}}{\partial y} = \sin \epsilon \frac{\partial}{\partial \tilde{x}} + \cos \epsilon \frac{\partial}{\partial \tilde{y}}.$$

The Jacobian for (2) is

$$A = \begin{bmatrix} \partial \tilde{x} / \partial x & \partial \tilde{x} / \partial y \\ \partial \tilde{y} / \partial x & \partial \tilde{y} / \partial y \end{bmatrix} = \begin{bmatrix} \cos \epsilon & \sin \epsilon \\ -\sin \epsilon & \cos \epsilon \end{bmatrix} \implies \det(A) = \cos^2 \epsilon + \sin^2 \epsilon = 1,$$

and therefore

$$\int_R dx dy \mapsto \int_{\tilde{R}} d\tilde{x} d\tilde{y}.$$

Making these substitutions into (1), we have

$$\begin{aligned} J[u(x, y)] &= \int_R \left[ \left( \cos \epsilon \frac{\partial u}{\partial \tilde{x}} - \sin \epsilon \frac{\partial u}{\partial \tilde{y}} \right)^2 + \left( \sin \epsilon \frac{\partial u}{\partial \tilde{x}} + \cos \epsilon \frac{\partial u}{\partial \tilde{y}} \right)^2 \right] dx dy \\ &= \int_R \left( \cos^2 \epsilon \frac{\partial^2 u}{\partial \tilde{x}^2} - 2 \cos \epsilon \sin \epsilon \frac{\partial^2 u}{\partial \tilde{x} \partial \tilde{y}} + \sin^2 \epsilon \frac{\partial^2 u}{\partial \tilde{y}^2} + \sin^2 \epsilon \frac{\partial^2 u}{\partial \tilde{x}^2} + 2 \cos \epsilon \sin \epsilon \frac{\partial^2 u}{\partial \tilde{x} \partial \tilde{y}} + \cos^2 \epsilon \frac{\partial^2 u}{\partial \tilde{y}^2} \right) dx dy \\ &= \int_R \left[ \left( \frac{\partial u}{\partial \tilde{x}} \right)^2 + \left( \frac{\partial u}{\partial \tilde{y}} \right)^2 \right] dx dy = \int_{\tilde{R}} \left[ \left( \frac{\partial u}{\partial \tilde{x}} \right)^2 + \left( \frac{\partial u}{\partial \tilde{y}} \right)^2 \right] d\tilde{x} d\tilde{y} \\ &= J[u(\tilde{x}, \tilde{y})] \end{aligned}$$

as desired.  $\square$

**Problem 2.** Consider the real-valued Lagrangian density  $\mathcal{L}$  depending on a complex-valued function  $\phi(t, x, y)$ :

$$\mathcal{L} = \frac{i}{2} \left( \phi^* \frac{d\phi}{dt} - \frac{d\phi^*}{dt} \phi \right) - \nabla \phi^* \cdot \nabla \phi - m^2 \phi^* \phi, \quad (3)$$

where  $*$  is complex conjugation, and  $\nabla \phi = (\partial \phi / \partial x, \partial \phi / \partial y)$ . Treating  $\phi$  and  $\phi^*$  as independent objects, derive the Euler-Lagrange equations.

**Solution.** We will have two Euler-Lagrange equations; one for  $\phi$  and one for  $\phi^*$ . In general, they are given by

$$0 = \frac{\partial \mathcal{L}}{\partial \phi} - \frac{d}{dt} \frac{\partial \mathcal{L}}{\partial \phi_t} - \frac{d}{dx} \frac{\partial \mathcal{L}}{\partial \phi_x} - \frac{d}{dy} \frac{\partial \mathcal{L}}{\partial \phi_y}, \quad 0 = \frac{\partial \mathcal{L}}{\partial \phi^*} - \frac{d}{dt} \frac{\partial \mathcal{L}}{\partial \phi_t^*} - \frac{d}{dx} \frac{\partial \mathcal{L}}{\partial \phi_x^*} - \frac{d}{dy} \frac{\partial \mathcal{L}}{\partial \phi_y^*}.$$

Expanding out  $\nabla \phi^* \cdot \nabla \phi$ , (3) becomes

$$\mathcal{L} = \frac{i}{2} \left( \phi^* \frac{d\phi}{dt} - \frac{d\phi^*}{dt} \phi \right) - \frac{\partial \phi^*}{\partial x} \frac{\partial \phi}{\partial x} - \frac{\partial \phi^*}{\partial y} \frac{\partial \phi}{\partial y} - m^2 \phi^* \phi.$$

Then

$$\begin{aligned}\frac{\partial \mathcal{L}}{\partial \phi} &= -\frac{i}{2} \frac{d\phi^*}{dt} - m^2 \phi^*, & \frac{\partial \mathcal{L}}{\partial \phi_t} &= \frac{i}{2} \phi^*, & \frac{\partial \mathcal{L}}{\partial \phi_x} &= -\frac{\partial \phi^*}{\partial x}, & \frac{\partial \mathcal{L}}{\partial \phi_y} &= -\frac{\partial \phi^*}{\partial y}, \\ \frac{\partial \mathcal{L}}{\partial \phi^*} &= \frac{i}{2} \frac{d\phi}{dt} - m^2 \phi, & \frac{\partial \mathcal{L}}{\partial \phi_t^*} &= -\frac{i}{2} \phi, & \frac{\partial \mathcal{L}}{\partial \phi_x^*} &= -\frac{\partial \phi}{\partial x}, & \frac{\partial \mathcal{L}}{\partial \phi_y^*} &= -\frac{\partial \phi}{\partial y},\end{aligned}$$

and

$$\begin{aligned}\frac{d}{dt} \frac{\partial \mathcal{L}}{\partial \phi_t} &= \frac{i}{2} \frac{d\phi^*}{dt}, & \frac{d}{dx} \frac{\partial \mathcal{L}}{\partial \phi_x} &= -\frac{\partial^2 \phi^*}{\partial x^2}, & \frac{d}{dy} \frac{\partial \mathcal{L}}{\partial \phi_y} &= -\frac{\partial^2 \phi^*}{\partial y^2}, \\ \frac{d}{dt} \frac{\partial \mathcal{L}}{\partial \phi_t^*} &= -\frac{i}{2} \frac{d\phi}{dt}, & \frac{d}{dx} \frac{\partial \mathcal{L}}{\partial \phi_x^*} &= -\frac{\partial^2 \phi}{\partial x^2}, & \frac{d}{dy} \frac{\partial \mathcal{L}}{\partial \phi_y^*} &= -\frac{\partial^2 \phi}{\partial y^2}.\end{aligned}$$

Then the Euler-Lagrange equations are

$$0 = -\frac{i}{2} \frac{d\phi^*}{dt} - m^2 \phi^* - \frac{i}{2} \frac{d\phi^*}{dt} + \frac{\partial^2 \phi^*}{\partial x^2} + \frac{\partial^2 \phi^*}{\partial y^2}, \quad 0 = \frac{i}{2} \frac{d\phi}{dt} - m^2 \phi + \frac{i}{2} \frac{d\phi}{dt} + \frac{\partial^2 \phi}{\partial x^2} + \frac{\partial^2 \phi}{\partial y^2},$$

which simplify to

$$0 = i \frac{d\phi^*}{dt} - \nabla^2 \phi^* + m^2 \phi^*, \quad 0 = i \frac{d\phi}{dt} + \nabla^2 \phi - m^2 \phi.$$

**Problem 3.** The nondimensionalized, multidimensional Sine-Gordon equation,

$$\theta_{xx} + \theta_{yy} - \theta_{tt} = \sin \theta$$

for  $\theta(x, y, t)$ , is the Euler-Lagrange equation for the action integral

$$S[\theta] = \int_R \left\{ \frac{1}{2} [\theta_t^2 - (\nabla \theta)^2] - \sin \theta \right\} dx dy dt$$

with  $\nabla \theta = (\partial \theta / \partial x, \partial \theta / \partial y)$ . The functional  $S[\theta]$  is invariant under translation of  $x$ ,  $y$ , and  $t$ . Find the associated energy-momentum tensor and energy-momentum vector.

**Solution.** Expanding out  $(\nabla \theta)^2$ , the Lagrangian density is

$$\mathcal{L} = \frac{1}{2} (\theta_t^2 - \theta_x^2 - \theta_y^2) - \sin \theta. \quad (4)$$

The energy-momentum tensor is defined by

$$T_{ij} = \frac{\partial \mathcal{L}}{\partial \theta_{x_i}} \frac{\partial \theta}{\partial x_j} - \mathcal{L} \delta_{ij},$$

where  $x_i \in \{x_0, x_1, x_2\} = \{t, x, y\}$ . The diagonal elements of  $T$  are then

$$\begin{aligned}T_{00} &= \frac{\partial \mathcal{L}}{\partial \theta_t} \frac{\partial \theta}{\partial t} - \mathcal{L} = \theta_t^2 - \frac{1}{2} (\theta_t^2 - \theta_x^2 - \theta_y^2) + \sin \theta = \frac{1}{2} (\theta_t^2 + \theta_x^2 + \theta_y^2) + \sin \theta, \\ T_{11} &= \frac{\partial \mathcal{L}}{\partial \theta_x} \frac{\partial \theta}{\partial x} - \mathcal{L} = -\theta_x^2 - \frac{1}{2} (\theta_t^2 - \theta_x^2 - \theta_y^2) + \sin \theta = -\frac{1}{2} (\theta_t^2 + \theta_x^2 - \theta_y^2) + \sin \theta, \\ T_{22} &= \frac{\partial \mathcal{L}}{\partial \theta_y} \frac{\partial \theta}{\partial y} - \mathcal{L} = -\theta_y^2 - \frac{1}{2} (\theta_t^2 - \theta_x^2 - \theta_y^2) + \sin \theta = -\frac{1}{2} (\theta_t^2 - \theta_x^2 + \theta_y^2) + \sin \theta,\end{aligned}$$

and the nondiagonal elements are

$$\begin{aligned} T_{01} &= \frac{\partial \mathcal{L}}{\partial \theta_t} \frac{\partial \theta}{\partial x} = \theta_t \theta_x, & T_{02} &= \frac{\partial \mathcal{L}}{\partial \theta_t} \frac{\partial \theta}{\partial y} = \theta_t \theta_y, & T_{12} &= \frac{\partial \mathcal{L}}{\partial \theta_x} \frac{\partial \theta}{\partial y} = -\theta_x \theta_y, \\ T_{10} &= \frac{\partial \mathcal{L}}{\partial \theta_x} \frac{\partial \theta}{\partial t} = -\theta_t \theta_x, & T_{20} &= \frac{\partial \mathcal{L}}{\partial \theta_y} \frac{\partial \theta}{\partial t} = -\theta_t \theta_y, & T_{21} &= \frac{\partial \mathcal{L}}{\partial \theta_y} \frac{\partial \theta}{\partial x} = -\theta_x \theta_y. \end{aligned}$$

In matrix form, we have

$$T = \begin{bmatrix} (\theta_t^2 + \theta_x^2 + \theta_y^2)/2 + \sin \theta & \theta_t \theta_x & \theta_t \theta_y \\ -\theta_t \theta_x & -(\theta_t^2 + \theta_x^2 - \theta_y^2)/2 + \sin \theta & -\theta_x \theta_y \\ -\theta_t \theta_y & -\theta_x \theta_y & -(\theta_t^2 - \theta_x^2 + \theta_y^2)/2 + \sin \theta \end{bmatrix}.$$

The energy-momentum vector is defined by

$$P_j = \int T_{0j} dx_1 dx_2.$$

Its components are then

$$P_0 = \int \left[ \frac{1}{2}(\theta_t^2 + \theta_x^2 + \theta_y^2) + \sin \theta \right] dx dy, \quad P_1 = \int \theta_t \theta_x dx dy, \quad P_2 = \int \theta_t \theta_y dx dy.$$

#### Problem 4. Extra credit

**4.a** Verify that the nondimensionalized, one-dimensional Sine-Gordon equation,

$$\theta_{xx} - \theta_{tt} = \sin \theta, \tag{5}$$

is also invariant under a Lorentz transformation on  $(x_0 = t, x_1 = x)$ . The transformation is given by

$$\begin{bmatrix} \gamma & -\gamma\nu \\ -\gamma\nu & \gamma \end{bmatrix},$$

where  $\gamma = 1/\sqrt{1 - \nu^2}$ .

**Solution.** Define  $(\tilde{t}, \tilde{x})$  as the transformed coordinates. (5) is invariant if it has the same form under the substitution  $\theta(t, x) \mapsto \theta(\tilde{t}, \tilde{x})$ . The new coordinates are given by

$$\begin{bmatrix} \tilde{t} \\ \tilde{x} \end{bmatrix} = \begin{bmatrix} \gamma & -\gamma\nu \\ -\gamma\nu & \gamma \end{bmatrix} \begin{bmatrix} t \\ x \end{bmatrix} = \begin{bmatrix} \gamma(t - \nu x) \\ \gamma(x - \nu t) \end{bmatrix},$$

or

$$\tilde{t} = \gamma(t - \nu x), \quad \tilde{x} = \gamma(x - \nu t).$$

Proceeding similarly to problem 1, the chain rule gives us

$$\frac{\partial}{\partial t} = \frac{\partial}{\partial \tilde{t}} \frac{\partial \tilde{t}}{\partial t} + \frac{\partial}{\partial \tilde{x}} \frac{\partial \tilde{x}}{\partial t} = \gamma \left( \frac{\partial}{\partial \tilde{t}} - \nu \frac{\partial}{\partial \tilde{x}} \right), \quad \frac{\partial}{\partial x} = \frac{\partial}{\partial \tilde{t}} \frac{\partial \tilde{t}}{\partial x} + \frac{\partial}{\partial \tilde{x}} \frac{\partial \tilde{x}}{\partial x} = \gamma \left( \frac{\partial}{\partial \tilde{x}} - \nu \frac{\partial}{\partial \tilde{t}} \right).$$

For the second derivatives,

$$\begin{aligned}\frac{\partial^2}{\partial t^2} &= \gamma^2 \left( \frac{\partial}{\partial \tilde{t}} - \nu \frac{\partial}{\partial \tilde{x}} \right)^2 = \gamma^2 \left( \frac{\partial^2}{\partial \tilde{t}^2} - 2\nu \frac{\partial^2}{\partial \tilde{t} \partial \tilde{x}} + \nu^2 \frac{\partial^2}{\partial \tilde{x}^2} \right), \\ \frac{\partial^2}{\partial x^2} &= \gamma^2 \left( \frac{\partial}{\partial \tilde{x}} - \nu \frac{\partial}{\partial \tilde{t}} \right)^2 = \gamma^2 \left( \frac{\partial^2}{\partial \tilde{x}^2} - 2\nu \frac{\partial^2}{\partial \tilde{t} \partial \tilde{x}} + \nu^2 \frac{\partial^2}{\partial \tilde{t}^2} \right).\end{aligned}$$

Making these substitutions, (5) becomes

$$\begin{aligned}\sin \theta &= \frac{\partial^2 \theta}{\partial x^2} - \frac{\partial^2 \theta}{\partial t^2} \\ &= \gamma^2 \left( \frac{\partial^2 \theta}{\partial \tilde{x}^2} - 2\nu \frac{\partial^2 \theta}{\partial \tilde{t} \partial \tilde{x}} + \nu^2 \frac{\partial^2 \theta}{\partial \tilde{t}^2} \right) - \gamma^2 \left( \frac{\partial^2 \theta}{\partial \tilde{t}^2} - 2\nu \frac{\partial^2 \theta}{\partial \tilde{t} \partial \tilde{x}} + \nu^2 \frac{\partial^2 \theta}{\partial \tilde{x}^2} \right) \\ &= \gamma^2 \left[ (1 - \nu^2) \frac{\partial^2 \theta}{\partial \tilde{x}^2} - (1 - \nu^2) \frac{\partial^2 \theta}{\partial \tilde{t}^2} \right] \\ &= \frac{\partial^2 \theta}{\partial \tilde{x}^2} - \frac{\partial^2 \theta}{\partial \tilde{t}^2},\end{aligned}$$

because  $\gamma^2 = 1/(1 - \nu^2)$ . Thus, we have demonstrated the invariance of (5).  $\square$

**4.b** Find the associated conserved quantity. Is it analogous to a common conserved quantity in classical mechanics?

**Solution.** By analogy to problem 3, the Lagrangian for this system is given by

$$\mathcal{L} = \frac{1}{2}(\theta_t^2 - \theta_x^2) - \sin \theta$$

which is like (4), but with only one spatial dimension. Continuing the analogy, the components of the energy-momentum vector are

$$P_0 = \int \left[ \frac{1}{2}(\theta_t^2 + \theta_x^2) + \sin \theta \right] dx, \quad P_1 = \int \theta_t \theta_x dx.$$

These are the conserved quantities, or “currents.” The component  $P_0$  is analogous to the classical Hamiltonian, or the total energy of the system. This corresponds to  $\mathcal{L}$ ’s having no explicit  $t$  dependence. The component  $P_1$  is like the momentum conjugate to  $x$ , since it corresponds to  $\mathcal{L}$ ’s having no explicit  $x$  dependence. Since we are concerned with only one spatial dimension,  $P_1$  is analogous to the classical total (linear) momentum of the system.

## Problem 5. Interacting line vortices

A system of  $n$  vortices moving on a two-dimensional plane has the Hamiltonian

$$H = \sum_{j=1}^n \sum_{i=1}^{j-1} -\gamma^{(i)} \gamma^{(j)} \ln |\mathbf{r}_i - \mathbf{r}_j|,$$

where  $\gamma^{(i)}$  is the strength of the  $i$ th line vortex, and  $\mathbf{r}_i = (x_i, y_i)$  its position in the plane. Using the Poisson bracket structure

$$[f, g] = \sum_{i=1}^n \frac{1}{\gamma^{(i)}} \left( \frac{\partial f}{\partial x_i} \frac{\partial g}{\partial y_i} - \frac{\partial f}{\partial y_i} \frac{\partial g}{\partial x_i} \right),$$

Hamilton's equations for the vortex system simplify to

$$\dot{x}_i = [x_i, H], \quad \dot{y}_i = [y_i, H].$$

Consider two vortices. Show that the equations of motion can be solved explicitly. Most importantly, show that the solution tells us the two vortices orbit each other with a frequency that is inversely proportional to the square of their separation.

**Solution.** For two vortices, the Hamiltonian reduces to

$$H = -\gamma^{(1)}\gamma^{(2)} \ln |\mathbf{r}_1 - \mathbf{r}_2|.$$

For  $i, j \in \{1, 2\}$ , note that

$$\frac{\partial x_i}{\partial x_j} = \frac{\partial y_i}{\partial y_j} = \delta_{ij}, \quad \frac{\partial x_i}{\partial y_j} = \frac{\partial y_i}{\partial x_j} = 0.$$

Note also that

$$|\mathbf{r}_1 - \mathbf{r}_2| = \sqrt{(x_1 - x_2)^2 + (y_1 - y_2)^2} = \sqrt{x_1^2 - 2x_1x_2 + x_2^2 + y_1^2 - 2y_1y_2 + y_2^2},$$

and define  $R \equiv |\mathbf{r}_1 - \mathbf{r}_2|$  as the separation of the vortices. Define also

$$u \equiv \ln R, \quad v \equiv (x_1 - x_2)^2 + (y_1 - y_2)^2 = R^2.$$

Then

$$\begin{aligned} \frac{\partial H}{\partial x_i} &= \frac{\partial H}{\partial u} \frac{\partial u}{\partial R} \frac{\partial R}{\partial v} \frac{\partial v}{\partial x_i} = -\gamma^{(i)}\gamma^{(j)} \frac{1}{R} \frac{1}{2R} (2x_i - 2x_j) = -\gamma^{(i)}\gamma^{(j)} \frac{x_i - x_j}{(x_1 - x_2)^2 + (y_1 - y_2)^2} = -\gamma^{(i)}\gamma^{(j)} \frac{x_i - x_j}{R^2}, \\ \frac{\partial H}{\partial y_i} &= \frac{\partial H}{\partial u} \frac{\partial u}{\partial R} \frac{\partial R}{\partial v} \frac{\partial v}{\partial y_i} = -\gamma^{(i)}\gamma^{(j)} \frac{y_i - y_j}{R^2}, \end{aligned}$$

where  $i \neq j$ .

Combining the above, and again fixing  $i \neq j$ ,

$$\begin{aligned} \dot{x}_i &= \frac{1}{\gamma^{(i)}} \left( \frac{\partial x_i}{\partial x_i} \frac{\partial H}{\partial y_i} - \frac{\partial x_i}{\partial y_i} \frac{\partial H}{\partial x_i} \right) + \frac{1}{\gamma^{(j)}} \left( \frac{\partial x_i}{\partial x_j} \frac{\partial H}{\partial y_j} - \frac{\partial x_i}{\partial y_j} \frac{\partial H}{\partial x_j} \right) = \frac{1}{\gamma^{(i)}} \frac{\partial H}{\partial y_i} = -\gamma^{(j)} \frac{y_i - y_j}{R^2}, \\ \dot{y}_i &= \frac{1}{\gamma^{(i)}} \left( \frac{\partial y_i}{\partial x_i} \frac{\partial H}{\partial y_i} - \frac{\partial y_i}{\partial y_i} \frac{\partial H}{\partial x_i} \right) + \frac{1}{\gamma^{(j)}} \left( \frac{\partial y_i}{\partial x_j} \frac{\partial H}{\partial y_j} - \frac{\partial y_i}{\partial y_j} \frac{\partial H}{\partial x_j} \right) = -\frac{1}{\gamma^{(i)}} \frac{\partial H}{\partial x_i} = \gamma^{(j)} \frac{x_i - x_j}{R^2}. \end{aligned}$$

Explicitly, the four equations of motion are

$$\dot{x}_1 = -\gamma^{(2)} \frac{y_1 - y_2}{R^2}, \quad \dot{x}_2 = \gamma^{(1)} \frac{y_1 - y_2}{R^2}, \quad \dot{y}_1 = \gamma^{(2)} \frac{x_1 - x_2}{R^2}, \quad \dot{y}_2 = -\gamma^{(1)} \frac{x_1 - x_2}{R^2}. \quad (6)$$

Note that

$$\frac{\partial R}{\partial x_i} = \frac{\partial R}{\partial v} \frac{\partial v}{\partial x_i} = \frac{x_i - x_j}{R^2}, \quad \frac{\partial R}{\partial y_i} = \frac{y_i - y_j}{R^2}$$

so

$$[H, R] = \sum_{i=1}^n \frac{1}{\gamma^{(i)}} \left( \frac{\partial H}{\partial x_i} \frac{\partial v}{\partial y_i} - \frac{\partial H}{\partial y_i} \frac{\partial v}{\partial x_i} \right) = \sum_{i=1}^n \frac{1}{\gamma^{(i)}} \left( -\gamma^{(i)}\gamma^{(j)} \frac{x_i - x_j}{R^2} \frac{y_i - y_j}{R^2} + \gamma^{(i)}\gamma^{(j)} \frac{y_i - y_j}{R^2} \frac{x_i - x_j}{R^2} \right) = 0.$$

This means  $R$  is a constant.

Define  $\mathbf{R} \equiv \mathbf{r}_1 - \mathbf{r}_2 = (X, Y)$ , where  $|\mathbf{R}| = R$ . Now we have two generalized coordinates  $X$  and  $Y$ , where  $X = x_1 - x_2$  and  $Y = y_1 - y_2$ . This gives us the two equations of motion

$$\dot{X} = \dot{x}_1 - \dot{x}_2 = -(\gamma^{(1)} + \gamma^{(2)}) \frac{y_1 - y_2}{R^2} = -\frac{\gamma^{(1)} + \gamma^{(2)}}{R^2} Y, \quad (7)$$

$$\dot{Y} = \dot{y}_1 - \dot{y}_2 = (\gamma^{(1)} + \gamma^{(2)}) \frac{x_1 - x_2}{R^2} = \frac{\gamma^{(1)} + \gamma^{(2)}}{R^2} X. \quad (8)$$

We can differentiate these to obtain two uncoupled second-order equations:

$$\ddot{X} = -\frac{\gamma^{(1)} + \gamma^{(2)}}{R^2} \dot{Y} = -\frac{(\gamma^{(1)} + \gamma^{(2)})^2}{R^4} X, \quad \ddot{Y} = \frac{\gamma^{(1)} + \gamma^{(2)}}{R^2} \dot{X} = -\frac{(\gamma^{(1)} + \gamma^{(2)})^2}{R^4} Y.$$

These equations have the solutions

$$X(t) = C_1 \cos\left(\frac{\gamma^{(1)} + \gamma^{(2)}}{R^2} t\right) + C_2 \sin\left(\frac{\gamma^{(1)} + \gamma^{(2)}}{R^2} t\right), \quad Y(t) = D_1 \cos\left(\frac{\gamma^{(1)} + \gamma^{(2)}}{R^2} t\right) + D_2 \sin\left(\frac{\gamma^{(1)} + \gamma^{(2)}}{R^2} t\right),$$

where  $C_1, C_2, D_1, D_2$  are constants. Define

$$\omega = \frac{\gamma^{(1)} + \gamma^{(2)}}{R^2}$$

as the angular frequency. Then

$$\dot{X}(t) = -C_1 \omega \sin \omega t + C_2 \omega \cos \omega t = -\omega Y, \quad \dot{Y}(t) = -D_1 \omega \sin \omega t + D_2 \omega \cos \omega t = \omega X,$$

where we have used (7) and (8). This fixes

$$C_1 = D_2, \quad C_2 = -D_1, \quad -D_1 = C_2, \quad D_2 = C_1$$

which implies  $C_1 = D_2 = 0$  and  $C_2 = D_1 \equiv K$ , where we have defined some new constant  $K$ . Finally, we have

$$X(t) = K \sin \omega t = x_1(t) - x_2(t), \quad Y(t) = K \cos \omega t = y_1(t) - y_2(t).$$

Thus, we have shown that the equations of motion (6) can be solved explicitly. Since  $\mathbf{R} = (X, Y)$  is the vector separating the vortices, we have also shown that they orbit each other. The orbital frequency

$$\omega = \frac{\gamma^{(1)} + \gamma^{(2)}}{R^2}$$

is clearly inversely proportional to  $R^2$ , where  $R = |\mathbf{R}|$  is the magnitude of the vortices' separation.

## Problem 6. Conserved quantities for a system of line vortices

Now consider the general case of  $n$  vortices.

**6.a** Verify that the “total linear momentum along  $x$ ” and the “total linear momentum along  $y$ ,”

$$P_x = \sum_{i=1}^n \gamma^{(i)} y_i, \quad P_y = \sum_{i=1}^n -\gamma^{(i)} x_i,$$

are conserved.

**6.b** Verify that  $[P_x, P_y]$  gives a conserved quantity.

### Problem 7. Charged particle in a magnetic field

Suppose a charged particle moves in a two-dimensional plane while experiencing a magnetic field  $\mathbf{B} = (0, 0, B)$ . Use the vector potential  $\mathbf{A} = (-By, 0, 0)$ . The Hamiltonian for the particle is

$$H = \frac{1}{2m} \left( p_x + \frac{eB}{c} y \right)^2 + \frac{p_y^2}{2m}.$$

**7.a** Write down Hamilton's equations. Verify that by appropriate manipulation we have

$$p_y + \frac{eB}{c} x = a, \quad p_x = m\dot{x} - \frac{eB}{c} y = b,$$

where  $a$  and  $b$  are constants.

**Solution.** Note that

$$H = \frac{1}{2m} \left( p_x^2 + 2\frac{eB}{c} p_x y + \frac{e^2 B^2}{c^2} y^2 \right) + \frac{p_y^2}{2m} = \frac{p_x^2}{2m} + \frac{eB}{c} \frac{p_x y}{m} + \frac{e^2 B^2}{c^2} \frac{y^2}{2m} + \frac{p_y^2}{2m}$$

Hamilton's equations are

$$\dot{x} = \frac{\partial H}{\partial p_x} = \frac{1}{m} \left( p_x + \frac{eB}{c} y \right), \quad (9)$$

$$\dot{p}_x = -\frac{\partial H}{\partial x} = 0, \quad (10)$$

$$\dot{y} = \frac{\partial H}{\partial p_y} = \frac{p_y}{m}, \quad (11)$$

$$\dot{p}_y = -\frac{\partial H}{\partial y} = -\frac{eB}{c} \frac{1}{m} \left( p_x + \frac{eB}{c} y \right). \quad (12)$$

Substituting (9) into (12),

$$\dot{p}_y = -\frac{eB}{c} \dot{x}.$$

By integrating with respect to  $t$ , we obtain

$$p_y = -\frac{eB}{c} \int \dot{x} dt = -\frac{eB}{c} x + a,$$

where  $a$  is some constant. Therefore, we have

$$p_y + \frac{eB}{c} x = a, \quad (13)$$

as desired.

From (9),

$$m\dot{x} = p_x + \frac{eB}{c} y \iff p_x = m\dot{x} - \frac{eB}{c} y,$$

and from (10),

$$p_x = \int 0 dt = b,$$

where  $b$  is some constant. Combining these, we have

$$p_x = m\dot{x} - \frac{eB}{c}y = b \quad (14)$$

as desired.  $\square$

**7.b** Using the relations above and the equations of motion, verify that the charged particle moves in a circle in the  $(x, y)$  plane and that the circling frequency  $\omega$  is given by

$$\omega = \frac{eB}{mc}.$$

This is called the *Larmor frequency*.

**Solution.** Substituting (14) into (9) yields

$$\dot{x} = \frac{1}{m} \left( b + \frac{eB}{c}y \right) = \frac{eB}{mc} \left( \frac{c}{eB}b + y \right). \quad (15)$$

Similarly, solving (13) for  $p_y$  and substituting into (11) gives us

$$\dot{y} = \frac{1}{m} \left( a - \frac{eB}{c}x \right) = \frac{eB}{mc} \left( \frac{c}{eB}a - x \right). \quad (16)$$

(15) and (16) are a system of two coupled first-order equations. Differentiating each by  $t$ , we obtain two uncoupled second-order equations:

$$\ddot{x} = \frac{eB}{mc}\dot{y} = -\frac{e^2B^2}{m^2c^2} \left( x - \frac{c}{eB}a \right), \quad \ddot{y} = -\frac{eB}{mc}\dot{x} = -\frac{e^2B^2}{m^2c^2} \left( y + \frac{c}{eB}b \right). \quad (17)$$

Define

$$\tilde{x} \equiv x - \frac{c}{eB}a, \quad \tilde{y} \equiv y + \frac{c}{eB}b.$$

Then

$$\frac{d\tilde{x}}{dt} = \dot{x}, \quad \frac{d^2\tilde{x}}{dt^2} = \ddot{x}, \quad \frac{d\tilde{y}}{dt} = \dot{y}, \quad \frac{d^2\tilde{y}}{dt^2} = \ddot{y},$$

and the equations (17) can be rewritten in terms of  $\tilde{x}$  and  $\tilde{y}$ :

$$\frac{d^2\tilde{x}}{dt^2} = -\frac{e^2B^2}{m^2c^2}\tilde{x}, \quad \frac{d^2\tilde{y}}{dt^2} = -\frac{e^2B^2}{m^2c^2}\tilde{y}.$$

These equations have the solutions

$$\begin{aligned} \tilde{x}(t) &= C_1 \cos\left(\frac{eB}{mc}t\right) + C_2 \sin\left(\frac{eB}{mc}t\right) \equiv C_1 \cos(\omega t) + C_2 \sin(\omega t), \\ \tilde{y}(t) &= D_1 \cos\left(\frac{eB}{mc}t\right) + D_2 \sin\left(\frac{eB}{mc}t\right) \equiv D_1 \cos(\omega t) + D_2 \sin(\omega t), \end{aligned}$$

where  $C_1, C_2, D_1, D_2$  are constants, and we have defined

$$\omega \equiv \frac{eB}{mc}.$$



Note that

$$\frac{d\tilde{x}}{dt} = -C_1\omega \sin(\omega t) + C_2\omega \cos(\omega t), \quad \frac{d\tilde{y}}{dt} = -D_1\omega \sin(\omega t) + D_2\omega \cos(\omega t).$$

Applying (15) and (16),

$$\begin{aligned} -C_1\omega \sin(\omega t) + C_2\omega \cos(\omega t) &= \omega [D_1 \cos(\omega t) + D_2 \sin(\omega t)], \\ -D_1\omega \sin(\omega t) + D_2\omega \cos(\omega t) &= \omega [C_1 \cos(\omega t) + C_2 \sin(\omega t)] \end{aligned}$$

which imply

$$-C_1 = D_2, \quad C_2 = D_1, \quad -D_1 = C_2, \quad D_2 = C_1$$

**7.c** Now consider the limit where the  $B$  field can be made arbitrarily strong. Compare the Poisson bracket  $[x, p_x]$  for the charged particle with the Poisson bracket relation

$$[x_i, y_i] = \frac{\delta_{ij}}{\gamma^{(i)}}$$

for the system of line vortices described in problems 5 and 6.

**Solution.** The Poisson bracket for the charged particle is given by

$$[x, p_x] = \frac{\partial x}{\partial x} \frac{\partial p_x}{\partial p_x} - \frac{\partial x}{\partial p_x} \frac{\partial p_x}{\partial x} + \frac{\partial x}{\partial y} \frac{\partial p_x}{\partial p_y} - \frac{\partial x}{\partial p_y} \frac{\partial p_x}{\partial y}.$$

Note that

$$\frac{\partial x}{\partial x} = \frac{\partial p_x}{\partial p_x} = 1, \quad \frac{\partial x}{\partial p_x} = \frac{\partial p_x}{\partial x} = \frac{\partial x}{\partial y} = \frac{\partial p_x}{\partial p_y} = 0, \quad \frac{\partial x}{\partial p_y} = -\frac{c}{eB}, \quad \frac{\partial p_x}{\partial y} = -\frac{eB}{c}.$$

Then

$$[x, p_x] = 1 - 1 = 0.$$

It's different?

While writing up these solutions, I consulted Gelfand and Fomin's *Calculus of Variations*, Goldstein's *Classical Mechanics*, Tong's *Classical Dynamics*, and Riley, Hobson, and Bence's *Mathematical Methods for Physics and Engineering*.