

Problem 1. Consider the following probabilistic game: There are four doors (Q, R, S, T). Behind each door is a device which displays ± 1 randomly according to the probability $P(Q = \pm 1, R = \pm 1, S = \pm 1, T = \pm 1)$. Alice and Bob are on the same team. Alice has to choose either Q and R , and then Bob has to choose either S and T . When the numbers match, they get $+1$ point; when the numbers do not match, they get -1 point. However, when they open Q and T , it's an exception. When the numbers (do not) match, they get -1 ($+1$).

1.1 Let's assume Alice and Bob open the doors completely randomly. When all numbers are $+1$ with probability 1, what is the expectation value of the point they get?

Solution. Let \mathbf{E} be the expectation value of the number of points. In this case, the numbers behind the two doors will always match. So

$$\mathbf{E} = \frac{QS + RS + RT - QT}{4} = \frac{1 + 1 + 1 - 1}{4} = \frac{1}{2}.$$

1.2 As it turns out, irrespective of how hard you fine tune the probability $P(Q = \pm 1, R = \pm 1, S = \pm 1, T = \pm 1)$, the expectation value of the point Alice and Bob get cannot exceed a certain value Max:

$$\frac{\mathbf{E}(QS) + \mathbf{E}(RS) + \mathbf{E}(RT) - \mathbf{E}(QT)}{4} \leq \text{Max}.$$

Here, $\mathbf{E}(QS)$, etc. is the expectation value of the point when Alice opens Q and Bob opens S . This is a Bell inequality. Determine Max.

Hint: For a given realization of the numbers $Q = \pm 1, R = \pm 1, S = \pm 1, T = \pm 1$, which occurs with probability $P(Q, R, S, T)$, note that $QS + RS + RT - QT = (Q + R)S + (R - Q)T$, where one of $\{(R + Q), (R - Q)\}$ is 2 and the other 0.

Solution. In addition to the information provided in the hint, both S and T must be ± 1 . This means the only possibilities for the number of points earned are

$$\frac{(Q + R)S + (R - Q)T}{4} = \begin{cases} \frac{(0)(-1) + (2)(1)}{4} = \frac{1}{2}, \\ \frac{(0)(1) + (2)(-1)}{4} = -\frac{1}{2}. \end{cases}$$

Thus,

$$\text{Max} = \frac{1}{2}.$$

1.3 Frustrated by the upper bound set by the Bell inequality, Bob decides to cheat. He now changes the value of T after Alice chooses Q or R . Assume Q, R, S are set to be $+1$ with probability 1. To make the expectation value of the point they get equal to $+1$, what values should Bob set after Alice chooses Q or R ?

Solution. If Alice chooses R , Bob should set $T = 1$. If Alice chooses Q , Bob should set $T = -1$. This way,

$$\frac{\mathbf{E}(QS) + \mathbf{E}(RS) + \mathbf{E}(RT) - \mathbf{E}(QT)}{4} = \frac{1 + 1 + 1 + 1}{4} = 1.$$

1.4 Now consider a quantum mechanical version of the game. There are quantum states of two spin-1/2 degrees of freedom shared by Alice and Bob. Alice can measure the z component or x components of the first spin \mathbf{S}^A . (This corresponds to $Q = \pm 1$ or $R = \pm 1$.) Bob can measure the $-(z + x)$ component or the $(z - x)$ component of the first spin \mathbf{S}^B . (This corresponds to $S = \pm 1$ or $T = \pm 1$.)

More specifically, Alice and Bob share the quantum state

$$|\psi\rangle = \frac{|\uparrow_z\rangle \otimes |\downarrow_z\rangle - |\downarrow_z\rangle \otimes |\uparrow_z\rangle}{\sqrt{2}}.$$

The operators to be measured are

$$Q = S_z^A, \quad R = S_x^A, \quad S = -\frac{S_z^B + S_x^B}{\sqrt{2}}, \quad T = \frac{S_x^B - S_z^B}{\sqrt{2}}.$$

Let us consider the case when Alice measures Q and Bob measures T . Calculate the probability $P(Q, T)$ for Alice and Bob getting the measurement outcomes $(Q, T) = (\pm 1, \pm 1)$.

Solution. All of the operators have eigenvalues $\pm\hbar/2$. In the S_z^A basis, Q and its eigenvectors are

$$Q = \frac{\hbar}{2} \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}, \quad |Q_+\rangle = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \quad |Q_-\rangle = \begin{bmatrix} 0 \\ 1 \end{bmatrix}.$$

In the S_z^B basis, T can be written

$$T = \frac{\hbar}{2\sqrt{2}} \left(\begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix} - \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \right) = \frac{\hbar}{2\sqrt{2}} \begin{bmatrix} 1 & -1 \\ -1 & -1 \end{bmatrix}.$$

The eigenvectors of T corresponding to eigenvalues $\pm\hbar/2$ can be found by

$$\begin{bmatrix} 1 & -1 \\ -1 & -1 \end{bmatrix} \begin{bmatrix} u \\ v \end{bmatrix} = \pm\sqrt{2} \begin{bmatrix} u \\ v \end{bmatrix},$$

which is satisfied when

$$v = (1 \mp \sqrt{2})u, \quad u = -(1 \pm \sqrt{2})v.$$

We will fix $u = 1$. Then the normalization constants A_{\pm} are found by

$$1 = |\langle T_{\pm} | T_{\pm} \rangle|^2 = A_{\pm}^2 \begin{bmatrix} 1 & 1 \mp \sqrt{2} \end{bmatrix} \begin{bmatrix} 1 \\ 1 \mp \sqrt{2} \end{bmatrix} = A_{\pm}^2 (4 \mp 2\sqrt{2}) \implies A_{\pm} = \frac{1}{\sqrt{4 \mp 2\sqrt{2}}}. \quad (1)$$

The normalized eigenvectors are

$$|T_+\rangle = \frac{1}{\sqrt{4 - 2\sqrt{2}}} \begin{bmatrix} 1 \\ 1 - \sqrt{2} \end{bmatrix}, \quad |T_-\rangle = \frac{1}{\sqrt{4 + 2\sqrt{2}}} \begin{bmatrix} 1 \\ 1 + \sqrt{2} \end{bmatrix}. \quad (2)$$

The probability that Alice and Bob obtain $(Q, T) = (+1, +1)$ is

$$\begin{aligned} P(+1, +1) &= |\langle Q_+, T_+ | \psi \rangle|^2 = \left| \begin{bmatrix} 1 & 0 \end{bmatrix} \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ -1 \end{bmatrix} \otimes \frac{1}{\sqrt{4 - 2\sqrt{2}}} \begin{bmatrix} 1 & 1 - \sqrt{2} \end{bmatrix} \frac{1}{\sqrt{2}} \begin{bmatrix} -1 \\ 1 \end{bmatrix} \right|^2 = \left| \frac{1}{\sqrt{2}} \otimes -\frac{1}{\sqrt{4 - 2\sqrt{2}}} \right|^2 \\ &= \frac{1}{8 - 4\sqrt{2}}, \end{aligned}$$

and the probability that they obtain $(Q, T) = (-1, -1)$ is

$$\begin{aligned} P(-1, -1) &= |\langle Q_-, T_- | \psi \rangle|^2 = \left| \begin{bmatrix} 0 & 1 \end{bmatrix} \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ -1 \end{bmatrix} \otimes \frac{1}{\sqrt{4+2\sqrt{2}}} \begin{bmatrix} 1 & 1+\sqrt{2} \end{bmatrix} \frac{1}{\sqrt{2}} \begin{bmatrix} -1 \\ 1 \end{bmatrix} \right|^2 = \left| -\frac{1}{\sqrt{2}} \otimes \frac{1}{\sqrt{4+2\sqrt{2}}} \right|^2 \\ &= \frac{1}{8+4\sqrt{2}}. \end{aligned}$$

1.5 Similarly, consider the case when Alice measures R and Bob measures T . Calculate the probability $P(R, T)$ for Alice and Bob getting the measurement outcomes $(R, T) = (\pm 1, \pm 1)$.

Solution. In the S_z^A basis, R and its eigenvectors are

$$R = \frac{\hbar}{2} \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \quad |R_+\rangle = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ 1 \end{bmatrix} \quad |R_-\rangle = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ -1 \end{bmatrix}.$$

Using (2), the probability for Alice and Bob obtain $(R, T) = (+1, +1)$ is

$$P(+1, +1) = |\langle R_+, T_+ | \psi \rangle|^2 = \left| \frac{1}{\sqrt{2}} \begin{bmatrix} 1 & 1 \end{bmatrix} \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ -1 \end{bmatrix} \otimes \frac{1}{\sqrt{4-2\sqrt{2}}} \begin{bmatrix} 1 & 1-\sqrt{2} \end{bmatrix} \frac{1}{\sqrt{2}} \begin{bmatrix} -1 \\ 1 \end{bmatrix} \right|^2 = 0,$$

and the probability that they obtain $(R, T) = (-1, -1)$ is

$$\begin{aligned} P(-1, -1) &= |\langle R_-, T_- | \psi \rangle|^2 = \left| \frac{1}{\sqrt{2}} \begin{bmatrix} 1 & -1 \end{bmatrix} \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ -1 \end{bmatrix} \otimes \frac{1}{\sqrt{4+2\sqrt{2}}} \begin{bmatrix} 1 & 1+\sqrt{2} \end{bmatrix} \frac{1}{\sqrt{2}} \begin{bmatrix} -1 \\ 1 \end{bmatrix} \right|^2 = \left| 1 \otimes \frac{1}{\sqrt{4+2\sqrt{2}}} \right|^2 \\ &= \frac{1}{4+2\sqrt{2}}. \end{aligned}$$

1.6 Compute the expectation values $\mathbf{E}(QS)$, $\mathbf{E}(RS)$, $\mathbf{E}(QT)$, and $\mathbf{E}(RT)$. Compute

$$\frac{\mathbf{E}(QS) + \mathbf{E}(RS) + \mathbf{E}(RT) - \mathbf{E}(QT)}{4}.$$

Solution. Firstly, in the S_z^B basis, S can be written

$$S = -\frac{\hbar}{2\sqrt{2}} \left(\begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} + \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix} \right) = \frac{\hbar}{2\sqrt{2}} \begin{bmatrix} -1 & -1 \\ -1 & 1 \end{bmatrix}.$$

Then the expectation values are

$$\begin{aligned} \mathbf{E}(QS) &= \langle \psi | QS | \psi \rangle = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 & -1 \end{bmatrix} \frac{\hbar}{2} \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix} \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ -1 \end{bmatrix} \otimes \frac{1}{\sqrt{2}} \begin{bmatrix} -1 & 1 \end{bmatrix} \frac{\hbar}{2\sqrt{2}} \begin{bmatrix} -1 & -1 \\ -1 & 1 \end{bmatrix} \frac{1}{\sqrt{2}} \begin{bmatrix} -1 \\ 1 \end{bmatrix} \\ &= \frac{\hbar^2}{4} \begin{bmatrix} 1 & -1 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \end{bmatrix} \otimes \frac{\hbar^2}{4\sqrt{2}} \begin{bmatrix} -1 & 1 \end{bmatrix} \begin{bmatrix} 0 \\ 2 \end{bmatrix} = 0, \end{aligned}$$

$$\begin{aligned} \mathbf{E}(RS) &= \langle \psi | RS | \psi \rangle = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 & -1 \end{bmatrix} \frac{\hbar}{2} \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ -1 \end{bmatrix} \otimes \frac{1}{\sqrt{2}} \begin{bmatrix} -1 & 1 \end{bmatrix} \frac{\hbar}{2\sqrt{2}} \begin{bmatrix} -1 & -1 \\ -1 & 1 \end{bmatrix} \frac{1}{\sqrt{2}} \begin{bmatrix} -1 \\ 1 \end{bmatrix} \\ &= \frac{\hbar}{4} \begin{bmatrix} 1 & -1 \end{bmatrix} \begin{bmatrix} -1 \\ 1 \end{bmatrix} \otimes \frac{\hbar}{4\sqrt{2}} \begin{bmatrix} -1 & 1 \end{bmatrix} \begin{bmatrix} 0 \\ 2 \end{bmatrix} = -\frac{\hbar}{2} \otimes \frac{\hbar}{2\sqrt{2}} = -\frac{\hbar^2}{4\sqrt{2}}, \end{aligned}$$

$$\begin{aligned}\mathbf{E}(QT) &= \langle \psi | QT | \psi \rangle = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 & -1 \end{bmatrix} \frac{\hbar}{2} \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix} \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ -1 \end{bmatrix} \otimes \frac{1}{\sqrt{2}} \begin{bmatrix} -1 & 1 \end{bmatrix} \frac{\hbar}{2\sqrt{2}} \begin{bmatrix} 1 & -1 \\ -1 & -1 \end{bmatrix} \frac{1}{\sqrt{2}} \begin{bmatrix} -1 \\ 1 \end{bmatrix} \\ &= \frac{\hbar}{4} \begin{bmatrix} 1 & -1 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \end{bmatrix} \otimes \frac{\hbar}{4\sqrt{2}} \begin{bmatrix} -1 & 1 \end{bmatrix} \begin{bmatrix} -2 \\ 0 \end{bmatrix} = 0,\end{aligned}$$

$$\begin{aligned}\mathbf{E}(RT) &= \langle \psi | RT | \psi \rangle = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 & -1 \end{bmatrix} \frac{\hbar}{2} \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ -1 \end{bmatrix} \otimes \frac{1}{\sqrt{2}} \begin{bmatrix} -1 & 1 \end{bmatrix} \frac{\hbar}{2\sqrt{2}} \begin{bmatrix} 1 & -1 \\ -1 & -1 \end{bmatrix} \frac{1}{\sqrt{2}} \begin{bmatrix} -1 \\ 1 \end{bmatrix} \\ &= \frac{\hbar}{4} \begin{bmatrix} 1 & -1 \end{bmatrix} \begin{bmatrix} -1 \\ 1 \end{bmatrix} \otimes \frac{\hbar}{4\sqrt{2}} \begin{bmatrix} -1 & 1 \end{bmatrix} \begin{bmatrix} -2 \\ 0 \end{bmatrix} = -\frac{\hbar}{2} \otimes \frac{\hbar}{2\sqrt{2}} = -\frac{\hbar^2}{4\sqrt{2}}.\end{aligned}$$

Then

$$\frac{\mathbf{E}(QS) + \mathbf{E}(RS) + \mathbf{E}(RT) - \mathbf{E}(QT)}{4} = \frac{1}{4} \left(-\frac{\hbar^2}{4\sqrt{2}} - \frac{\hbar^2}{4\sqrt{2}} \right) = -\frac{\hbar^2}{8\sqrt{2}},$$

which doesn't make sense. Any of it.

I consulted Sakurai's *Modern Quantum Mechanics*, Shankar's *Principles of Quantum Mechanics*, and Wolfram MathWorld while writing up these solutions.