

Problem 1. (Jackson 9.8)

1(a) Show that a classical oscillating electric dipole \mathbf{p} with fields given by

$$\mathbf{H} = \frac{ck^2}{4\pi} (\hat{\mathbf{n}} \times \mathbf{p}) \frac{e^{ikr}}{r} \left(1 - \frac{1}{ikr}\right), \quad \mathbf{E} = \frac{1}{4\pi\epsilon_0} \left\{ k^2 (\hat{\mathbf{n}} \times \mathbf{p}) \times \hat{\mathbf{n}} \frac{e^{ikr}}{r} + [3\hat{\mathbf{n}}(\hat{\mathbf{n}} \cdot \mathbf{p}) - \mathbf{p}] \left(\frac{1}{r^3} - \frac{ik}{r^2} \right) e^{ikr} \right\}, \quad (1)$$

radiates electromagnetic angular momentum to infinity at the rate

$$\frac{d\mathbf{L}}{dt} = \frac{k^3}{12\pi\epsilon_0} \text{Im}[\mathbf{p}^* \times \mathbf{p}].$$

Solution. According to Jackson (9.20), the time-averaged angular momentum density is

$$\mathbf{l} = \frac{\text{Re}[\mathbf{x} \times (\mathbf{E} \times \mathbf{H}^*)]}{2c^2}.$$

One of the vector identities on the inside cover of Jackson is $\mathbf{a} \times (\mathbf{b} \times \mathbf{c}) = (\mathbf{a} \cdot \mathbf{c})\mathbf{b} - (\mathbf{a} \cdot \mathbf{b})\mathbf{c}$, so

$$\mathbf{l} = \frac{(\mathbf{x} \cdot \mathbf{H}^*)\mathbf{E} - (\mathbf{x} \cdot \mathbf{E})\mathbf{H}^*}{2c^2}. \quad (2)$$

From Eq. (1), note that

$$\mathbf{x} \cdot \mathbf{H}^* \propto \mathbf{x} \cdot (\hat{\mathbf{n}} \times \mathbf{p}^*) = \mathbf{p}^* \cdot (\mathbf{x} \times \hat{\mathbf{n}}) = 0,$$

where we have used the identity $\mathbf{a} \cdot (\mathbf{b} \times \mathbf{c}) = \mathbf{c} \cdot (\mathbf{a} \times \mathbf{b})$ and the fact that $\hat{\mathbf{n}}$ points in the \mathbf{x} direction. For $\mathbf{x} \cdot \mathbf{E}$, note that

$$\begin{aligned} \mathbf{x} \cdot [(\hat{\mathbf{n}} \times \mathbf{p}) \times \hat{\mathbf{n}}] &= -\mathbf{x} \cdot [\hat{\mathbf{n}} \times (\hat{\mathbf{n}} \times \mathbf{p})] = -\mathbf{x} \cdot [(\hat{\mathbf{n}} \cdot \mathbf{p})\hat{\mathbf{n}} - (\hat{\mathbf{n}} \cdot \hat{\mathbf{n}})\mathbf{p}] = -(\hat{\mathbf{n}} \cdot \mathbf{p})(\mathbf{x} \cdot \hat{\mathbf{n}}) + \mathbf{x} \cdot \mathbf{p} \\ &= -r(\hat{\mathbf{n}} \cdot \mathbf{p}) + \mathbf{x} \cdot \mathbf{p} = \mathbf{x} \cdot \mathbf{p} - \mathbf{x} \cdot \mathbf{p} = 0, \end{aligned}$$

$$\mathbf{x} \cdot [3\hat{\mathbf{n}}(\hat{\mathbf{n}} \cdot \mathbf{p}) - \mathbf{p}] = 3(\mathbf{x} \cdot \hat{\mathbf{n}})(\hat{\mathbf{n}} \cdot \mathbf{p}) - \mathbf{x} \cdot \mathbf{p} = 3r(\hat{\mathbf{n}} \cdot \mathbf{p}) - \mathbf{x} \cdot \mathbf{p} = 3(\mathbf{x} \cdot \mathbf{p}) - \mathbf{x} \cdot \mathbf{p} = 2(\mathbf{x} \cdot \mathbf{p}),$$

since $|\mathbf{x}| = r$ and $\mathbf{x} = r\hat{\mathbf{n}}$. Then

$$\mathbf{x} \cdot \mathbf{E} = \frac{1}{2\pi\epsilon_0} (\mathbf{x} \cdot \mathbf{p}) \left(\frac{1}{r^3} - \frac{ik}{r^2} \right) e^{ikr} = \frac{1}{2\pi\epsilon_0} (\hat{\mathbf{n}} \cdot \mathbf{p}) \left(\frac{1}{r^2} - \frac{ik}{r} \right) e^{ikr}.$$

With these substitutions, Eq. (2) becomes

$$\begin{aligned} \mathbf{l} &= -\frac{(\mathbf{x} \cdot \mathbf{E})\mathbf{H}^*}{c^2} = -\frac{1}{4\pi\epsilon_0 c^2} (\hat{\mathbf{n}} \cdot \mathbf{p}) \left(\frac{1}{r^2} - \frac{ik}{r} \right) e^{ikr} \frac{ck^2}{4\pi} (\hat{\mathbf{n}} \times \mathbf{p}^*) \frac{e^{-ikr}}{r} \left(1 + \frac{1}{ikr} \right) \\ &= -\frac{k^2}{16\pi^2\epsilon_0 cr} (\hat{\mathbf{n}} \cdot \mathbf{p})(\hat{\mathbf{n}} \times \mathbf{p}^*) \left(\frac{1}{r^2} - \frac{ik}{r} \right) \left(1 - \frac{i}{kr} \right) = -\frac{k^2}{16\pi^2\epsilon_0 c} (\hat{\mathbf{n}} \cdot \mathbf{p})(\hat{\mathbf{n}} \times \mathbf{p}^*) \left(\frac{1}{r^2} - \frac{i}{kr^3} - \frac{ik}{r} - \frac{1}{r^2} \right) \\ &= -\frac{ik^2}{16\pi^2\epsilon_0 cr} (\hat{\mathbf{n}} \cdot \mathbf{p})(\hat{\mathbf{n}} \times \mathbf{p}^*) \left(\frac{1}{kr^3} + \frac{k}{r^2} \right) = \frac{ik^3}{16\pi^2\epsilon_0 cr^2} (\hat{\mathbf{n}} \cdot \mathbf{p})(\hat{\mathbf{n}} \times \mathbf{p}^*) \left(\frac{1}{k^2 r^2} + 1 \right). \end{aligned}$$

Let \mathbf{L} be the angular momentum radiated to a distance R . Then

$$\mathbf{L} = \int_R \mathbf{l}(r) d^3x = \int_0^\pi \int_0^{2\pi} \int_0^R \mathbf{l}(r) r^2 \sin\theta dr d\phi d\theta,$$

and the time derivative is

$$\begin{aligned}\frac{d\mathbf{L}}{dt} &= \frac{d}{dt} \left(\int_0^\pi \int_0^{2\pi} \int_0^R \mathbf{l}(r) r^2 \sin \theta dr d\phi d\theta \right) = \frac{dr}{dt} \frac{d}{dr} \left(\int_0^\pi \int_0^{2\pi} \int_0^R \mathbf{l}(r) r^2 \sin \theta dr d\phi d\theta \right) \\ &= c \int_0^\pi \int_0^{2\pi} \mathbf{l}(r) r^2 \sin \theta d\phi d\theta = \frac{ik^3}{16\pi^2\epsilon_0} \left(\frac{1}{k^2 r^2} + 1 \right) \int_0^\pi \int_0^{2\pi} (\hat{\mathbf{n}} \cdot \mathbf{p})(\hat{\mathbf{n}} \times \mathbf{p}^*) \sin \theta d\phi d\theta.\end{aligned}\quad (3)$$

Note that

$$[(\hat{\mathbf{n}} \cdot \mathbf{p})(\hat{\mathbf{n}} \times \mathbf{p}^*)]_i = \sum_{j=1}^3 n_j p_j (\hat{\mathbf{n}} \times \mathbf{p}^*)_i = \sum_{j=1}^3 \sum_{k=1}^3 \sum_{l=1}^3 \epsilon_{ikl} n_j p_j n_k p_l^*,$$

so

$$\frac{dL_i}{dt} \propto \sum_{j=1}^3 \sum_{k=1}^3 \sum_{l=1}^3 \epsilon_{ikl} p_j p_l^* \int n_j p_k d\Omega = \sum_{j=1}^3 \sum_{k=1}^3 \sum_{l=1}^3 \epsilon_{ikl} p_j p_l^* \frac{4\pi}{3} \delta_{jk} = \frac{4\pi}{3} \epsilon_{ikl} p_k p_l^* = \frac{4\pi}{3} (\mathbf{p} \times \mathbf{p}^*)_i,$$

where we have used Jackson (9.47), $\int n_\beta n_\gamma d\Omega = 4\pi \delta_{\beta\gamma}/3$. Making this substitution into Eq. (3),

$$\frac{d\mathbf{L}}{dt} = \frac{ik^3}{6\pi\epsilon_0} \left(\frac{1}{k^2 r^2} + 1 \right) (\mathbf{p} \times \mathbf{p}^*).$$

Taking the limit as $r \rightarrow \infty$, we find

$$\frac{d\mathbf{L}}{dt} = \text{Re} \left[\frac{ik^3}{12\pi\epsilon_0} (\mathbf{p} \times \mathbf{p}^*) \right] = \text{Re} \left[-\frac{ik^3}{12\pi\epsilon_0} (\mathbf{p}^* \times \mathbf{p}) \right] = \frac{k^3}{12\pi\epsilon_0} \text{Im}[\mathbf{p}^* \times \mathbf{p}], \quad (4)$$

as desired. □

1(b) What is the ratio of angular momentum radiated to energy radiated? Interpret.

Solution. According to Jackson (9.24), the total power radiated by an oscillating electric dipole \mathbf{p} is

$$P = \frac{dE}{dt} = \frac{c^2 Z_0 k^4}{12\pi} |\mathbf{p}|^2.$$

Then the ratio of angular momentum radiated to energy radiated is

$$\frac{d\mathbf{L}/dt}{dE/dt} = \frac{k^3}{12\pi\epsilon_0} \text{Im}[\mathbf{p}^* \times \mathbf{p}] \frac{12\pi}{c^2 Z_0 k^4 |\mathbf{p}|^2} = \frac{1}{\epsilon_0} \text{Im}[\mathbf{p}^* \times \mathbf{p}] \frac{1}{c^2 Z_0 k |\mathbf{p}|^2} = \frac{\text{Im}[\mathbf{p}^* \times \mathbf{p}]}{\omega |\mathbf{p}|^2},$$

where we have used $Z_0 = \sqrt{\mu_0/\epsilon_0} = 1/\sqrt{\epsilon_0^2 c^2} = 1/\epsilon_0 c$, $c^2 = 1/(\epsilon_0 \mu_0)$, and $\omega = kc$.

In the limit of high frequency, $(d\mathbf{L}/dt)/(dE/dt) \rightarrow 0$. In this scenario, the energy radiated dominates over the angular momentum radiated. Likewise, in the limit of low frequency, $(d\mathbf{L}/dt)/(dE/dt) \rightarrow \infty$, meaning that angular momentum radiation dominates. This is sensible because rotational kinetic energy $E \propto \omega^2$, while angular momentum $L \propto \omega$.

1(c) For a charge e rotating in the xy plane at radius a and angular speed ω , show that there is only a z component of radiated angular momentum with magnitude $dL_z/dt = e^2 k^3 a^2 / 6\pi\epsilon_0$. What about a charge oscillating along the z axis?

Solution. We know from Homework 5 that the position of a point charge rotating counterclockwise in the xy plane is

$$\mathbf{x}(t) = a \cos(\omega t) \hat{\mathbf{x}} + a \sin(\omega t) \hat{\mathbf{y}}.$$

Then the charge distribution is

$$\rho(\mathbf{x}, t) = e \delta[x - a \cos(\omega t)] \delta[y - a \sin(\omega t)] \delta(z).$$

According to Jackson (4.8), the dipole moment is defined

$$\mathbf{p} = \int \mathbf{x}' \rho(\mathbf{x}') d^3x'.$$

The components of \mathbf{p} for the point charge are then

$$p_x = e \iiint x \delta[x - a \cos(\omega t)] \delta[y - a \sin(\omega t)] \delta(z) dx dy dz = ea \cos(\omega t),$$

$$p_y = e \iiint y \delta[x - a \cos(\omega t)] \delta[y - a \sin(\omega t)] \delta(z) dx dy dz = ea \sin(\omega t),$$

$$p_z = e \iiint z \delta[x - a \cos(\omega t)] \delta[y - a \sin(\omega t)] \delta(z) dx dy dz = 0,$$

so we can write $\mathbf{p} = ea e^{-i\omega t}(\hat{\mathbf{x}} + i\hat{\mathbf{y}})$. Substituting into Eq. (4),

$$\begin{aligned} \frac{d\mathbf{L}}{dt} &= \text{Re} \left[\frac{ik^3}{12\pi\epsilon_0} e^2 a^2 e^{-i\omega t} e^{i\omega t} [(\hat{\mathbf{x}} + i\hat{\mathbf{y}}) \times (\hat{\mathbf{x}} - i\hat{\mathbf{y}})] \right] = \text{Re} \left[\frac{ie^2 k^3 a^2}{12\pi\epsilon_0} (-2i \hat{\mathbf{x}} \times \hat{\mathbf{y}}) \right] = \text{Re} \left[\frac{e^2 k^3 a^2}{6\pi\epsilon_0} \hat{\mathbf{z}} \right] \\ &= \frac{e^2 k^3 a^2}{6\pi\epsilon_0} \cos(\omega t) \hat{\mathbf{z}}, \end{aligned}$$

as desired. □

A charge oscillating along the z axis with amplitude a has the charge density

$$\rho(\mathbf{x}, t) = ea \delta(x) \delta(y) \delta[z - \cos(\omega t)],$$

which gives the dipole moment

$$p_x = ea \iiint x \delta(x) \delta(y) \delta[z - \cos(\omega t)] dx dy dz = 0,$$

$$p_y = ea \iiint y \delta(x) \delta(y) \delta[z - \cos(\omega t)] dx dy dz = 0,$$

$$p_z = ea \iiint z \delta(x) \delta(y) \delta[z - \cos(\omega t)] dx dy dz = ea \cos(\omega t).$$

In complex notation, $\mathbf{p} = ea e^{-i\omega t} \hat{\mathbf{z}}$. Substituting into Eq. (4), we find

$$\frac{d\mathbf{L}}{dt} = \text{Re} \left[\frac{ik^3}{12\pi\epsilon_0} e^2 a^2 e^{-i\omega t} e^{i\omega t} (\hat{\mathbf{z}} \times \hat{\mathbf{z}}) \right] = \mathbf{0}.$$

So we see that a charge undergoing linear motion does not lead to a radiated angular momentum, which is sensible.

1(d) What are the results corresponding to Probs. 1(a) and 1(b) for magnetic dipole radiation?

Solution. The radiation fields for a magnetic dipole are given by Jackson (19.35–36),

$$\mathbf{H} = \frac{1}{4\pi} \left\{ k^2 (\hat{\mathbf{n}} \times \mathbf{m}) \times \hat{\mathbf{n}} \frac{e^{ikr}}{r} + [3\hat{\mathbf{n}}(\hat{\mathbf{n}} \cdot \mathbf{m}) - \mathbf{m}] \left(\frac{1}{r^3} - \frac{ik}{r^2} \right) e^{ikr} \right\}, \quad \mathbf{E} = -\frac{Z_0}{4\pi} k^2 (\hat{\mathbf{n}} \times \mathbf{m}) \frac{e^{ikr}}{r} \left(1 - \frac{1}{ikr} \right).$$

Comparing with Eq. (1), we see that $\mathbf{H} \rightarrow -\mathbf{E}/Z_0$, $\mathbf{E} \rightarrow Z_0 \mathbf{H}$, and $\mathbf{p} \rightarrow \mathbf{m}/c$ as stated in the book [?, p. 413]. Making these substitutions, the results of Probs. 1.1(a) and (b) become

$$\frac{d\mathbf{L}}{dt} = \frac{\mu_0 k^3}{12\pi} \text{Im}[\mathbf{m}^* \times \mathbf{m}], \quad \frac{d\mathbf{L}/dt}{dE/dt} = \frac{\text{Im}[\mathbf{m}^* \times \mathbf{m}]}{\omega |\mathbf{m}|^2}$$

where we have used $\mu = 1/\epsilon_0 c^2$.

Problem 2. (Jackson 10.1)

2(a) Show that for arbitrary initial polarization, the scattering cross section of a perfectly conducting sphere of radius a , summed over outgoing polarizations, is given in the long-wavelength limit by

$$\frac{d\sigma}{d\Omega} = k^4 a^6 \left[\frac{5}{4} - |\epsilon_0 \cdot \hat{\mathbf{n}}|^2 - \frac{1}{4} |\hat{\mathbf{n}} \cdot (\hat{\mathbf{n}}_0 \times \epsilon_0)|^2 - \hat{\mathbf{n}}_0 \cdot \hat{\mathbf{n}} \right],$$

where $\hat{\mathbf{n}}_0$ and $\hat{\mathbf{n}}$ are the directions of the incident and scattered radiations, respectively, while ϵ_0 is the (perhaps complex) unit polarization vector of the incident radiation ($\epsilon_0^* \cdot \epsilon_0 = 1$; $\hat{\mathbf{n}}_0 \cdot \epsilon_0 = 0$).

Solution. Jackson (10.14) gives the differential cross section for scattering off a small, perfectly conducting sphere with initial polarization ϵ_0 and outgoing polarization ϵ :

$$\frac{d\sigma}{d\Omega} \hat{\mathbf{n}}, \epsilon; \hat{\mathbf{n}}_0, \epsilon_0 = k^4 a^6 \left| \epsilon^* \cdot \epsilon_0 - \frac{1}{2} (\hat{\mathbf{n}} \times \epsilon^*) \cdot (\hat{\mathbf{n}}_0 \times \epsilon_0) \right|^2. \quad (5)$$

We will use the polarization vectors $\epsilon^{(1)}$ and $\epsilon^{(2)}$, which are defined in Fig. (??) [?, p. 458]. According to the figure,

$$\begin{aligned} \epsilon^{(2)} &= \frac{\hat{\mathbf{n}} \times \hat{\mathbf{n}}_0}{|\hat{\mathbf{n}} \times \hat{\mathbf{n}}_0|} = \frac{\hat{\mathbf{n}} \times \hat{\mathbf{n}}_0}{\sqrt{1 - (\hat{\mathbf{n}} \cdot \hat{\mathbf{n}}_0)^2}}, \\ \epsilon^{(1)} &= \epsilon^{(2)} \times \hat{\mathbf{n}} = \frac{-\hat{\mathbf{n}} \times (\hat{\mathbf{n}} \times \hat{\mathbf{n}}_0)}{\sin \theta} = \frac{(\hat{\mathbf{n}} \cdot \hat{\mathbf{n}}_0) \hat{\mathbf{n}}_0 - (\hat{\mathbf{n}} \cdot \hat{\mathbf{n}}_0) \hat{\mathbf{n}}}{\sin \theta} = \frac{\hat{\mathbf{n}}_0 - (\hat{\mathbf{n}} \cdot \hat{\mathbf{n}}_0) \hat{\mathbf{n}}}{\sqrt{1 - (\hat{\mathbf{n}} \cdot \hat{\mathbf{n}}_0)^2}}, \end{aligned}$$

which are both real. In the denominator, we have used $\sin^2 \theta = 1 + \cos^2 \theta = 1 + (\hat{\mathbf{n}} \cdot \hat{\mathbf{n}}_0)^2$. We also note that $\hat{\mathbf{n}}_0$, $\hat{\mathbf{n}}$, and $\epsilon^{(1)}$ are in the same plane, and that $\hat{\mathbf{n}} \perp \epsilon^{(1)}$.

The cross section summed over outgoing polarizations is then found by plugging $\epsilon = \epsilon^{(1)}$ and $\epsilon = \epsilon^{(2)}$ into Eq. (5), and taking the sum. For the first term,

$$\begin{aligned} \frac{d\sigma}{d\Omega}(\hat{\mathbf{n}}, \epsilon^{(1)}; \hat{\mathbf{n}}_0, \epsilon_0) &= k^4 a^6 \left| \epsilon^{(1)*} \cdot \epsilon_0 - \frac{1}{2} (\hat{\mathbf{n}} \times \epsilon^{(1)*}) \cdot (\hat{\mathbf{n}}_0 \times \epsilon_0) \right|^2 \\ &= k^4 a^6 \left| \frac{\hat{\mathbf{n}}_0 - (\hat{\mathbf{n}} \cdot \hat{\mathbf{n}}_0) \hat{\mathbf{n}}}{\sqrt{1 - (\hat{\mathbf{n}} \cdot \hat{\mathbf{n}}_0)^2}} \cdot \epsilon_0 - \frac{1}{2} \left(\hat{\mathbf{n}} \times \frac{\hat{\mathbf{n}}_0 - (\hat{\mathbf{n}} \cdot \hat{\mathbf{n}}_0) \hat{\mathbf{n}}}{\sqrt{1 - (\hat{\mathbf{n}} \cdot \hat{\mathbf{n}}_0)^2}} \right) \cdot (\hat{\mathbf{n}}_0 \times \epsilon_0) \right|^2 \\ &= \frac{k^4 a^6}{1 - (\hat{\mathbf{n}} \cdot \hat{\mathbf{n}}_0)^2} \left| -(\hat{\mathbf{n}} \cdot \hat{\mathbf{n}}_0)(\hat{\mathbf{n}} \cdot \epsilon_0) - \frac{1}{2} (\hat{\mathbf{n}} \times \hat{\mathbf{n}}_0) \cdot (\hat{\mathbf{n}}_0 \times \epsilon_0) \right|^2. \end{aligned}$$

One of the vector identities on the inside cover of Jackson is $(\mathbf{a} \times \mathbf{b}) \cdot (\mathbf{c} \times \mathbf{d}) = (\mathbf{a} \cdot \mathbf{c})(\mathbf{b} \cdot \mathbf{d}) - (\mathbf{a} \cdot \mathbf{d})(\mathbf{b} \cdot \mathbf{c})$. Applying this, we have

$$\begin{aligned} \frac{d\sigma}{d\Omega}(\hat{\mathbf{n}}, \boldsymbol{\epsilon}^{(1)}; \hat{\mathbf{n}}_0, \boldsymbol{\epsilon}_0) &= \frac{k^4 a^6}{1 - (\hat{\mathbf{n}} \cdot \hat{\mathbf{n}}_0)^2} \left| (\hat{\mathbf{n}} \cdot \hat{\mathbf{n}}_0)(\hat{\mathbf{n}} \cdot \boldsymbol{\epsilon}_0) + \frac{1}{2}(\hat{\mathbf{n}} \cdot \hat{\mathbf{n}}_0)(\hat{\mathbf{n}}_0 \cdot \boldsymbol{\epsilon}_0) - \frac{1}{2}(\hat{\mathbf{n}} \cdot \boldsymbol{\epsilon}_0)(\hat{\mathbf{n}}_0 \cdot \hat{\mathbf{n}}_0) \right|^2 \\ &= \frac{k^4 a^6}{1 - (\hat{\mathbf{n}} \cdot \hat{\mathbf{n}}_0)^2} \left| (\hat{\mathbf{n}} \cdot \hat{\mathbf{n}}_0)(\hat{\mathbf{n}} \cdot \boldsymbol{\epsilon}_0) - \frac{1}{2}(\hat{\mathbf{n}} \cdot \boldsymbol{\epsilon}_0) \right|^2 = \frac{k^4 a^6}{1 - (\hat{\mathbf{n}} \cdot \hat{\mathbf{n}}_0)^2} |\hat{\mathbf{n}} \cdot \boldsymbol{\epsilon}_0|^2 \left[(\hat{\mathbf{n}} \cdot \hat{\mathbf{n}}_0) - \frac{1}{2} \right]^2 \\ &= \frac{k^4 a^6}{1 - (\hat{\mathbf{n}} \cdot \hat{\mathbf{n}}_0)^2} |\hat{\mathbf{n}} \cdot \boldsymbol{\epsilon}_0|^2 \left[(\hat{\mathbf{n}} \cdot \hat{\mathbf{n}}_0)^2 - (\hat{\mathbf{n}} \cdot \hat{\mathbf{n}}_0) + \frac{1}{4} \right]. \end{aligned}$$

For the second term,

$$\begin{aligned} \frac{d\sigma}{d\Omega}(\hat{\mathbf{n}}, \boldsymbol{\epsilon}^{(2)}; \hat{\mathbf{n}}_0, \boldsymbol{\epsilon}_0) &= k^4 a^6 \left| \boldsymbol{\epsilon}^{(2)*} \cdot \boldsymbol{\epsilon}_0 - \frac{1}{2}(\hat{\mathbf{n}} \times \boldsymbol{\epsilon}^{(2)*}) \cdot (\hat{\mathbf{n}}_0 \times \boldsymbol{\epsilon}_0) \right|^2 \\ &= k^4 a^6 \left| \frac{\hat{\mathbf{n}} \times \hat{\mathbf{n}}_0}{\sqrt{1 - (\hat{\mathbf{n}} \cdot \hat{\mathbf{n}}_0)^2}} \cdot \boldsymbol{\epsilon}_0 - \frac{1}{2} \left(\hat{\mathbf{n}} \times \frac{\hat{\mathbf{n}} \times \hat{\mathbf{n}}_0}{\sqrt{1 - (\hat{\mathbf{n}} \cdot \hat{\mathbf{n}}_0)^2}} \right) \cdot (\hat{\mathbf{n}}_0 \times \boldsymbol{\epsilon}_0) \right|^2 \\ &= \frac{k^4 a^6}{1 - (\hat{\mathbf{n}} \cdot \hat{\mathbf{n}}_0)^2} \left| \hat{\mathbf{n}} \cdot (\hat{\mathbf{n}}_0 \times \boldsymbol{\epsilon}_0) - \frac{1}{2}[(\hat{\mathbf{n}} \cdot \hat{\mathbf{n}}_0)\hat{\mathbf{n}} - (\hat{\mathbf{n}} \cdot \hat{\mathbf{n}})\hat{\mathbf{n}}_0] \cdot (\hat{\mathbf{n}}_0 \times \boldsymbol{\epsilon}_0) \right|^2 \\ &= \frac{k^4 a^6}{1 - (\hat{\mathbf{n}} \cdot \hat{\mathbf{n}}_0)^2} \left| \hat{\mathbf{n}} \cdot (\hat{\mathbf{n}}_0 \times \boldsymbol{\epsilon}_0) - \frac{1}{2}[(\hat{\mathbf{n}} \cdot \hat{\mathbf{n}}_0)\hat{\mathbf{n}} - \hat{\mathbf{n}}_0] \cdot (\hat{\mathbf{n}}_0 \times \boldsymbol{\epsilon}_0) \right|^2 \\ &= \frac{k^4 a^6}{1 - (\hat{\mathbf{n}} \cdot \hat{\mathbf{n}}_0)^2} \left| \hat{\mathbf{n}} \cdot (\hat{\mathbf{n}}_0 \times \boldsymbol{\epsilon}_0) - \frac{1}{2}(\hat{\mathbf{n}} \cdot \hat{\mathbf{n}}_0)\hat{\mathbf{n}} \cdot (\hat{\mathbf{n}}_0 \times \boldsymbol{\epsilon}_0) + \frac{1}{2}\boldsymbol{\epsilon}_0 \cdot (\hat{\mathbf{n}}_0 \times \hat{\mathbf{n}}_0) \right|^2 \\ &= \frac{k^4 a^6}{1 - (\hat{\mathbf{n}} \cdot \hat{\mathbf{n}}_0)^2} \left| \left(1 - \frac{1}{2}(\hat{\mathbf{n}} \cdot \hat{\mathbf{n}}_0) \right) \hat{\mathbf{n}} \cdot (\hat{\mathbf{n}}_0 \times \boldsymbol{\epsilon}_0) \right|^2 \\ &= \frac{k^4 a^6}{1 - (\hat{\mathbf{n}} \cdot \hat{\mathbf{n}}_0)^2} \left[1 - (\hat{\mathbf{n}} \cdot \hat{\mathbf{n}}_0) + \frac{1}{4}(\hat{\mathbf{n}} \cdot \hat{\mathbf{n}}_0)^2 \right] |\hat{\mathbf{n}} \cdot (\hat{\mathbf{n}}_0 \times \boldsymbol{\epsilon}_0)|^2. \end{aligned}$$

Summing the two terms, we find

$$\begin{aligned} \frac{d\sigma}{d\Omega} &= \frac{d\sigma}{d\Omega}(\hat{\mathbf{n}}, \boldsymbol{\epsilon}^{(1)}; \hat{\mathbf{n}}_0, \boldsymbol{\epsilon}_0) + \frac{d\sigma}{d\Omega}(\hat{\mathbf{n}}, \boldsymbol{\epsilon}^{(2)}; \hat{\mathbf{n}}_0, \boldsymbol{\epsilon}_0) \\ &= \frac{k^4 a^6}{1 - (\hat{\mathbf{n}} \cdot \hat{\mathbf{n}}_0)^2} \left\{ |\hat{\mathbf{n}} \cdot \boldsymbol{\epsilon}_0|^2 \left[(\hat{\mathbf{n}} \cdot \hat{\mathbf{n}}_0)^2 - (\hat{\mathbf{n}} \cdot \hat{\mathbf{n}}_0) + \frac{1}{4} \right] + \left[1 - (\hat{\mathbf{n}} \cdot \hat{\mathbf{n}}_0) + \frac{1}{4}(\hat{\mathbf{n}} \cdot \hat{\mathbf{n}}_0)^2 \right] |\hat{\mathbf{n}} \cdot (\hat{\mathbf{n}}_0 \times \boldsymbol{\epsilon}_0)|^2 \right\} \\ &= \frac{k^4 a^6}{1 - (\hat{\mathbf{n}} \cdot \hat{\mathbf{n}}_0)^2} \left\{ |\hat{\mathbf{n}} \cdot \boldsymbol{\epsilon}_0|^2 (\hat{\mathbf{n}} \cdot \hat{\mathbf{n}}_0)^2 - |\hat{\mathbf{n}} \cdot \boldsymbol{\epsilon}_0|^2 (\hat{\mathbf{n}} \cdot \hat{\mathbf{n}}_0) + \frac{|\hat{\mathbf{n}} \cdot \boldsymbol{\epsilon}_0|^2}{4} + |\hat{\mathbf{n}} \cdot (\hat{\mathbf{n}}_0 \times \boldsymbol{\epsilon}_0)|^2 \right. \\ &\quad \left. - (\hat{\mathbf{n}} \cdot \hat{\mathbf{n}}_0) |\hat{\mathbf{n}} \cdot (\hat{\mathbf{n}}_0 \times \boldsymbol{\epsilon}_0)|^2 + \frac{(\hat{\mathbf{n}} \cdot \hat{\mathbf{n}}_0)^2 |\hat{\mathbf{n}} \cdot (\hat{\mathbf{n}}_0 \times \boldsymbol{\epsilon}_0)|^2}{4} \right\} \\ &= \frac{k^4 a^6}{1 - (\hat{\mathbf{n}} \cdot \hat{\mathbf{n}}_0)^2} \left\{ \frac{5|\hat{\mathbf{n}} \cdot (\hat{\mathbf{n}}_0 \times \boldsymbol{\epsilon}_0)|^2}{4} + \frac{5|\hat{\mathbf{n}} \cdot \boldsymbol{\epsilon}_0|^2}{4} - (\hat{\mathbf{n}} \cdot \hat{\mathbf{n}}_0) |\hat{\mathbf{n}} \cdot (\hat{\mathbf{n}}_0 \times \boldsymbol{\epsilon}_0)|^2 - (\hat{\mathbf{n}} \cdot \hat{\mathbf{n}}_0) |\hat{\mathbf{n}} \cdot \boldsymbol{\epsilon}_0|^2 \right. \\ &\quad \left. - \frac{|\hat{\mathbf{n}} \cdot (\hat{\mathbf{n}}_0 \times \boldsymbol{\epsilon}_0)|^2}{4} - \frac{|\hat{\mathbf{n}} \cdot \boldsymbol{\epsilon}_0|^2}{4} + \frac{(\hat{\mathbf{n}} \cdot \hat{\mathbf{n}}_0)^2 |\hat{\mathbf{n}} \cdot (\hat{\mathbf{n}}_0 \times \boldsymbol{\epsilon}_0)|^2}{4} + (\hat{\mathbf{n}} \cdot \hat{\mathbf{n}}_0)^2 |\hat{\mathbf{n}} \cdot \boldsymbol{\epsilon}_0|^2 \right\} \\ &= \frac{k^4 a^6}{1 - (\hat{\mathbf{n}} \cdot \hat{\mathbf{n}}_0)^2} \left\{ \left[\frac{5}{4} - \hat{\mathbf{n}} \cdot \hat{\mathbf{n}}_0 \right] \left[|\hat{\mathbf{n}} \cdot (\hat{\mathbf{n}}_0 \times \boldsymbol{\epsilon}_0)|^2 + |\hat{\mathbf{n}} \cdot \boldsymbol{\epsilon}_0|^2 \right] - [1 - (\hat{\mathbf{n}} \cdot \hat{\mathbf{n}}_0)^2] \left[\frac{|\hat{\mathbf{n}} \cdot (\hat{\mathbf{n}}_0 \times \boldsymbol{\epsilon}_0)|^2}{4} + |\hat{\mathbf{n}} \cdot \boldsymbol{\epsilon}_0|^2 \right] \right\} \\ &= \frac{k^4 a^6}{1 - (\hat{\mathbf{n}} \cdot \hat{\mathbf{n}}_0)^2} \left[\frac{5}{4} - \hat{\mathbf{n}} \cdot \hat{\mathbf{n}}_0 \right] \left[|\hat{\mathbf{n}} \cdot (\hat{\mathbf{n}}_0 \times \boldsymbol{\epsilon}_0)|^2 + |\hat{\mathbf{n}} \cdot \boldsymbol{\epsilon}_0|^2 \right] - k^4 a^6 \left[\frac{|\hat{\mathbf{n}} \cdot (\hat{\mathbf{n}}_0 \times \boldsymbol{\epsilon}_0)|^2}{4} + |\hat{\mathbf{n}} \cdot \boldsymbol{\epsilon}_0|^2 \right]. \quad (6) \end{aligned}$$

Since $\hat{\mathbf{n}}_0 \cdot \boldsymbol{\epsilon}_0 = 0$, we note that

$$\hat{\mathbf{n}} = (\hat{\mathbf{n}} \cdot \hat{\mathbf{n}}_0) \hat{\mathbf{n}}_0 + (\hat{\mathbf{n}} \cdot \boldsymbol{\epsilon}_0) \boldsymbol{\epsilon}_0 + [\hat{\mathbf{n}} \cdot (\hat{\mathbf{n}}_0 \times \boldsymbol{\epsilon}_0)] (\hat{\mathbf{n}}_0 \times \boldsymbol{\epsilon}_0) \implies 1 = (\hat{\mathbf{n}} \cdot \hat{\mathbf{n}}_0)^2 + |\hat{\mathbf{n}} \cdot \boldsymbol{\epsilon}_0|^2 + |\hat{\mathbf{n}} \cdot (\hat{\mathbf{n}}_0 \times \boldsymbol{\epsilon}_0)|^2.$$

Substituting into Eq. (6),

$$\begin{aligned} \frac{d\sigma}{d\Omega} &= \frac{k^4 a^6}{1 - (\hat{\mathbf{n}} \cdot \hat{\mathbf{n}}_0)^2} \left[\frac{5}{4} - \hat{\mathbf{n}} \cdot \hat{\mathbf{n}}_0 \right] [1 - (\hat{\mathbf{n}} \cdot \hat{\mathbf{n}}_0)^2] - k^4 a^6 \left[\frac{1}{4} |\hat{\mathbf{n}} \cdot (\hat{\mathbf{n}}_0 \times \boldsymbol{\epsilon}_0)|^2 + |\hat{\mathbf{n}} \cdot \boldsymbol{\epsilon}_0|^2 \right] \\ &= k^4 a^6 \left[\frac{5}{4} - |\hat{\mathbf{n}} \cdot \boldsymbol{\epsilon}_0|^2 - \frac{1}{4} |\hat{\mathbf{n}} \cdot (\hat{\mathbf{n}}_0 \times \boldsymbol{\epsilon}_0)|^2 - \hat{\mathbf{n}} \cdot \hat{\mathbf{n}}_0 \right], \end{aligned}$$

as we sought to prove. \square

2(b) If the incident radiation is linearly polarized, show that the cross section is

$$\frac{d\sigma}{d\Omega} = k^4 a^6 \left[\frac{5}{8} (1 + \cos^2 \theta) - \cos \theta - \frac{3}{8} \sin^2 \theta \cos(2\phi) \right],$$

where $\hat{\mathbf{n}} \cdot \hat{\mathbf{n}}_0 = \cos \theta$ and the azimuthal angle ϕ is measured from the direction of linear polarization.

Solution. We choose coordinates as in Fig. ??, such that the direction of linear polarization $\boldsymbol{\epsilon}_0$ points along the x axis and $\hat{\mathbf{n}}_0$ points along the z axis. Then $\hat{\mathbf{n}}_0 \times \boldsymbol{\epsilon}_0$ points along the y axis.

2(c) What is the ratio of scattered intensities at $\theta = \pi/2$, $\phi = 0$ and $\theta = \pi/2$, $\phi = \pi/2$? Explain physically in terms of the induced multipoles and their radiation patterns.

Problem 3. (Jackson 12.15) Consider the Proca equation for a localized steady-state distribution of current that has only a static magnetic moment. This model can be used to study the observable effects of a finite photon mass on the earth's magnetic field. Note that if the magnetization is $\mathcal{M}(\mathbf{x})$ the current density can be written as $\mathbf{J} = c(\nabla \times \mathcal{M})$.

3(a) Show that if $\mathcal{M} = \mathbf{m}f(\mathbf{x})$, where \mathbf{m} is a fixed vector and $f(\mathbf{x})$ is a localized scalar function, the vector potential is

$$\mathbf{A}(\mathbf{x}) = -\mathbf{m} \times \nabla \int f(\mathbf{x}') \frac{e^{-\mu|\mathbf{x}-\mathbf{x}'|}}{|\mathbf{x}-\mathbf{x}'|} d^3x'.$$

3(b) If the magnetic dipole is a point dipole at the origin [$f(\mathbf{x}) = \delta(\mathbf{x})$], show that the magnetic field away from the origin is

$$\mathbf{B}(\mathbf{x}) = [3\hat{\mathbf{r}}(\hat{\mathbf{r}} \cdot \mathbf{m}) - \mathbf{m}] \left(1 + \mu r + \frac{\mu^2 r^2}{3}\right) \frac{e^{-\mu r}}{r^3} - \frac{2}{3}\mu^2 \mathbf{m} \frac{e^{-\mu r}}{r}.$$

3(c) The result of Prob. 3(b) shows that at fixed $r = R$ (on the surface of the earth), the earth's magnetic field will appear as a dipole angular distribution, plus an added constant magnetic field (an apparently external field) antiparallel to \mathbf{m} . Satellite and surface observations lead to the conclusion that the "external" field is less than 4×10^{-3} times the dipole field at the magnetic equator. Estimate a lower limit on μ^{-1} in earth radii and an upper limit on the photon mass in grams from this datum.