

Problem 1. Show that for an arbitrary spatially bound charge-current source, the electric dipole moment \mathbf{p} satisfies

$$\frac{d\mathbf{p}}{dt} = \int \mathbf{J} d^3x.$$

Solution. The electric dipole moment \mathbf{p} is defined by Eq. (2.36),

$$\mathbf{p} = \int \mathbf{x} \rho(x) d^3x. \quad (1)$$

Differentiating both sides with respect to t , we find

$$\frac{d\mathbf{p}}{dt} = \frac{d}{dt} \int \mathbf{x} \rho d^3x = \int \frac{d}{dt} d^3x = \int \mathbf{x} \frac{\partial \rho}{\partial t} d^3x, \quad (2)$$

because \mathbf{x} is simply the point at which we are evaluating the potential, and is therefore independent of time.

The charge-current conservation law is given by Eq. (5.8),

$$\frac{\partial \rho}{\partial t} + \nabla \cdot \mathbf{J} = 0. \quad (3)$$

Multiplying by \mathbf{x} on both sides and integrating over all space, we obtain

$$\int \mathbf{x} \frac{\partial \rho}{\partial t} d^3x + \int \mathbf{x} (\nabla \cdot \mathbf{J}) d^3x = 0.$$

Applying (2), we have

$$\frac{d\mathbf{p}}{dt} = - \int \mathbf{x} (\nabla \cdot \mathbf{J}) d^3x. \quad (4)$$

It remains to be shown that the right side is equal to the integral of \mathbf{J} over all space.

Vector identity (5) in Griffiths is

$$\nabla \cdot (f\mathbf{a}) = f(\nabla \cdot \mathbf{a}) + \mathbf{a} \cdot (\nabla f).$$

Writing the right side of (4) in component notation and applying the identity gives us

$$- \int x_i (\nabla \cdot \mathbf{J}) d^3x = \int \mathbf{J} \cdot (\nabla x_i) d^3x - \int \nabla \cdot (x_i \mathbf{J}) d^3x. \quad (5)$$

Gauss's theorem is given by Eq. (2.6),

$$\int_{\mathcal{V}} \nabla \cdot \mathbf{v} d^3x = \int_S \mathbf{v} \cdot \hat{\mathbf{n}} dS,$$

Here, let \mathcal{V} be a ball of radius R , with R large enough that the entire charge-current source is enclosed. Then S is the surface of \mathcal{V} , and $\hat{\mathbf{n}} = \hat{\mathbf{r}}$. Applying Gauss's theorem to the second integral on the right side of (5), we have

$$\int \nabla \cdot (x_i \mathbf{J}) d^3x = \lim_{R \rightarrow \infty} \int_{\mathcal{V}} \nabla \cdot (x_i \mathbf{J}) d^3x = \lim_{R \rightarrow \infty} \int_S x_i \mathbf{J} \cdot \hat{\mathbf{r}} dS = 0,$$

since \mathbf{J} is bounded, and therefore \mathbf{J} evaluated on S reaches zero well before x_i becomes very large.

Returning to (5), we now have

$$- \int x_i (\nabla \cdot \mathbf{J}) d^3x = \int \mathbf{J} \cdot (\nabla x_i) d^3x = \sum_j \int J_j \partial_j x_i d^3x = \sum_j \int J_j \delta_{ij} d^3x = \int J_i d^3x,$$

where we have followed the proof in Eq. (4.24) of the course notes. Finally, (4) becomes

$$\frac{d\mathbf{p}}{dt} = \int \mathbf{J} d^3x \quad (6)$$

as desired. \square

Problem 2. A particle of charge q_1 moves with velocity v in a circular orbit of radius R about the origin in the xy plane, such that its φ coordinate varies as $\varphi = \omega t$, with $\omega = v/R$. Assume that $v \ll c$. Another particle of charge q_2 is at rest at point \mathbf{x} , where $|\mathbf{x}| \gg R$. To order $1/|\mathbf{x}|$, find the force \mathbf{F} on the particle of charge q_2 at time t .

Solution. The Lorentz force equation, Eq. (1.25), is written

$$\mathbf{F} = q \left(\mathbf{E} + \frac{\mathbf{v}}{c} \times \mathbf{B} \right), \quad (7)$$

where \mathbf{v} is the velocity of the charge q on which the force is exerted, and \mathbf{E} and \mathbf{B} are the total electric and magnetic fields. For this problem, we are interested in the force acting on a stationary point charge q_2 , so $\mathbf{v}_2 = 0$. Additionally, we do not have to consider the self-field contribution to \mathbf{E} , since static charge distributions do not experience any self force. Thus we need only find the electric field due to q_1 , \mathbf{E}_1 . The multipole expansion of the electric field in electrodynamics is given by Eq. (5.70),

$$\mathbf{E}(t, \mathbf{x}) = \frac{1}{c^2 |\mathbf{x}|} \left[\left(\hat{\mathbf{x}} \cdot \frac{d^2 \mathbf{p}}{dt^2} \right) \hat{\mathbf{x}} - \frac{d^2 \mathbf{p}}{dt^2} \right]_{\text{ret}} + \mathcal{O}\left(\frac{1}{|\mathbf{x}|^2}\right), \quad (8)$$

where $\hat{\mathbf{x}} = \mathbf{x}/|\mathbf{x}|$ is the unit vector in the direction of the point at which we are evaluating the field, and \mathbf{p} is the dipole moment defined by (1). In addition, (8) relies upon the assumption that the velocity of q_1 , v , satisfies $v \ll c$.

The position of q_1 at time t can be expressed as

$$\mathbf{x}_1(t) = R \cos(\omega t) \hat{\mathbf{x}} + R \sin(\omega t) \hat{\mathbf{y}},$$

so the charge density for q_1 everywhere is

$$\rho_1(t, \mathbf{x}) = q_1 \delta(\mathbf{x} - \mathbf{x}_1(t)).$$

Then the dipole moment $\mathbf{p}_1(t, \mathbf{x})$ is

$$\mathbf{p}_1(t, \mathbf{x}) = \int \mathbf{x} \rho_1(t, \mathbf{x}) d^3x = q_1 \int \mathbf{x} \delta(\mathbf{x} - \mathbf{x}_1(t)) d^3x = q_1 \mathbf{x}_1(t) = q_1 R \cos(\omega t) \hat{\mathbf{x}} + q_1 R \sin(\omega t) \hat{\mathbf{y}},$$

and so its second time derivative is

$$\frac{d^2 \mathbf{p}_1(t)}{dt^2} = \frac{d}{dt} \left(\frac{d\mathbf{p}_1}{dt} \right) = \frac{d}{dt} \left(-q_1 R \omega \sin(\omega t) \hat{\mathbf{x}} + q_1 R \omega \cos(\omega t) \hat{\mathbf{y}} \right) = -q_1 R \omega^2 \cos(\omega t) \hat{\mathbf{x}} - q_1 R \omega^2 \sin(\omega t) \hat{\mathbf{y}}.$$

To this order, the retarded time t' is defined

$$t' = t - \frac{|\mathbf{x}|}{c}. \quad (9)$$

In (8), let $\mathbf{x} \rightarrow \mathbf{r}$. Then $\hat{\mathbf{x}} \rightarrow \hat{\mathbf{r}}$, which is the radial unit vector, and $|\mathbf{x}| \rightarrow r$. To first order in $1/|\mathbf{x}|$, we get

$$\begin{aligned}\mathbf{E}(t, \mathbf{x}) &= \frac{1}{c^2 r} \left[-\frac{q_1 R \omega^2}{r} [\cos(\omega t') (\hat{\mathbf{r}} \cdot \hat{\mathbf{x}}) + \sin(\omega t') (\hat{\mathbf{r}} \cdot \hat{\mathbf{y}})] \mathbf{r} + q_1 R \omega^2 [\cos(\omega t') \hat{\mathbf{x}} + \sin(\omega t') \hat{\mathbf{y}}] \right]_{\text{ret}} \\ &= q_1 \frac{R \omega^2}{c^2 r} \left[\cos(\omega t') [\hat{\mathbf{x}} - (\hat{\mathbf{r}} \cdot \hat{\mathbf{x}}) \hat{\mathbf{r}}] + \sin(\omega t') [\hat{\mathbf{y}} - (\hat{\mathbf{r}} \cdot \hat{\mathbf{y}}) \hat{\mathbf{r}}] \right]_{\text{ret}} \\ &= q_1 \frac{R \omega^2}{c^2 r} \left\{ \cos\left(\omega t - \frac{\omega |\mathbf{x}|}{c}\right) [\hat{\mathbf{x}} - (\hat{\mathbf{r}} \cdot \hat{\mathbf{x}}) \hat{\mathbf{r}}] + \sin\left(\omega t - \frac{\omega |\mathbf{x}|}{c}\right) [\hat{\mathbf{y}} - (\hat{\mathbf{r}} \cdot \hat{\mathbf{y}}) \hat{\mathbf{r}}] \right\}.\end{aligned}$$

Note that

$$\hat{\mathbf{x}} = \sin \theta \cos \varphi \hat{\mathbf{r}} + \cos \theta \cos \varphi \hat{\boldsymbol{\theta}} - \sin \varphi \hat{\boldsymbol{\phi}}, \quad \hat{\mathbf{y}} = \sin \theta \sin \varphi \hat{\mathbf{r}} + \cos \theta \sin \varphi \hat{\boldsymbol{\theta}} + \cos \varphi \hat{\boldsymbol{\phi}},$$

so

$$\begin{aligned}\hat{\mathbf{x}} - (\hat{\mathbf{r}} \cdot \hat{\mathbf{x}}) \hat{\mathbf{r}} &= \hat{\mathbf{x}} - \sin \theta \cos \varphi \hat{\mathbf{r}} = \cos \theta \cos \varphi \hat{\boldsymbol{\theta}} - \sin \varphi \hat{\boldsymbol{\phi}}, \\ \hat{\mathbf{y}} - (\hat{\mathbf{r}} \cdot \hat{\mathbf{y}}) \hat{\mathbf{r}} &= \hat{\mathbf{y}} - \sin \theta \sin \varphi \hat{\mathbf{r}} = \cos \theta \sin \varphi \hat{\boldsymbol{\theta}} + \cos \varphi \hat{\boldsymbol{\phi}},\end{aligned}$$

and then

$$\mathbf{E}(t, \mathbf{x}) = q_1 \frac{R \omega^2}{c^2 r} \left[\cos\left(\omega t - \frac{\omega r}{c}\right) (\cos \theta \cos \varphi \hat{\boldsymbol{\theta}} - \sin \varphi \hat{\boldsymbol{\phi}}) + \sin\left(\omega t - \frac{\omega r}{c}\right) (\cos \theta \sin \varphi \hat{\boldsymbol{\theta}} + \cos \varphi \hat{\boldsymbol{\phi}}) \right].$$

Applying (7) with $\mathbf{v} = 0$, we have

$$\mathbf{F}(t, \mathbf{x}) = q_1 q_2 \frac{R \omega^2}{c^2 r} \left[\cos\left(\omega t - \frac{\omega r}{c}\right) (\cos \theta \cos \varphi \hat{\boldsymbol{\theta}} - \sin \varphi \hat{\boldsymbol{\phi}}) + \sin\left(\omega t - \frac{\omega r}{c}\right) (\cos \theta \sin \varphi \hat{\boldsymbol{\theta}} + \cos \varphi \hat{\boldsymbol{\phi}}) \right].$$

Problem 3. An “antenna” is a segment of conducting wire in which a current flows (driven by an external power supply). Suppose an antenna of length L is placed on the z axis between $z = 0$ and $z = L$, and suppose that the current in the antenna is

$$\mathbf{J}(t, z) = I_0 \sin\left(\frac{\pi z}{L}\right) \cos(\omega t) \delta(x) \delta(y) \hat{\mathbf{z}}. \quad (10)$$

3.a Find the charge density $\rho(t, z)$ in the antenna.

Solution. From the charge-current conservation law (3), we have

$$\rho(t, z) = - \int \nabla \cdot \mathbf{J} dt.$$

For \mathbf{J} given by (10),

$$\nabla \cdot \mathbf{J} = \frac{\partial J_z}{\partial z} = \frac{\pi}{L} I_0 \cos\left(\frac{\pi z}{L}\right) \cos(\omega t) \delta(x) \delta(y),$$

and so, discarding the constant of integration,

$$\rho(t, z) = -\frac{\pi}{L} I_0 \cos\left(\frac{\pi z}{L}\right) \delta(x) \delta(y) \int \cos(\omega t) dt = -\frac{\pi}{L} \frac{I_0}{\omega} \cos\left(\frac{\pi z}{L}\right) \sin(\omega t) \delta(x) \delta(y)$$

for $0 \leq z \leq L$.

3.b Assume that $\omega L \ll c$. Find the electric and magnetic fields, $\mathbf{E}(t, z)$ and $\mathbf{B}(t, z)$, at large distances from the antenna (valid to order $1/|\mathbf{x}|$).

Solution. We will use (8) to find $\mathbf{E}(t, z)$. From Eq. (5.68), we know

$$\int \mathbf{J}(\mathbf{x}) d^3x = \frac{d\mathbf{p}}{dt},$$

so from (10) we have

$$\begin{aligned} \frac{d\mathbf{p}}{dt} &= I_0 \cos(\omega t) \hat{\mathbf{z}} \int_0^L \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \sin\left(\frac{\pi z}{L}\right) \delta(x) \delta(y) dx dy dz = I_0 \cos(\omega t) \hat{\mathbf{z}} \int_0^L \sin\left(\frac{\pi z}{L}\right) dz \\ &= I_0 \cos(\omega t) \hat{\mathbf{z}} \left[-\frac{L}{\pi} \cos\left(\frac{\pi z}{L}\right) \right]_0^L = \frac{2L}{\pi} I_0 \cos(\omega t) \hat{\mathbf{z}}. \end{aligned}$$

Then

$$\frac{d^2\mathbf{p}}{dt^2} = -\frac{2L}{\pi} I_0 \omega \sin(\omega t) \hat{\mathbf{z}}.$$

Using the retarded time (9), to first order in $1/|\mathbf{x}|$ we obtain

$$\begin{aligned} \mathbf{E}(t, \mathbf{x}) &= \frac{1}{c^2 r} \left[\left(-\hat{\mathbf{r}} \cdot \frac{2L}{\pi} I_0 \omega \sin(\omega t) \hat{\mathbf{z}} \right) \hat{\mathbf{r}} + \frac{2L}{\pi} I_0 \omega \sin(\omega t) \hat{\mathbf{z}} \right]_{\text{ret}} = \frac{2L}{\pi c^3} \frac{I_0 \omega}{r} \left[\sin(\omega t) [\hat{\mathbf{z}} - (\hat{\mathbf{r}} \cdot \hat{\mathbf{z}}) \hat{\mathbf{r}}] \right]_{\text{ret}} \\ &= \frac{2L}{\pi c^2} \frac{I_0 \omega}{r} \sin\left(\omega t - \frac{\omega r}{c}\right) [\hat{\mathbf{z}} - (\hat{\mathbf{r}} \cdot \hat{\mathbf{z}}) \hat{\mathbf{r}}]. \end{aligned}$$

Note that $\hat{\mathbf{z}} = \cos \theta \hat{\mathbf{r}} - \sin \theta \hat{\boldsymbol{\theta}}$, so

$$\hat{\mathbf{z}} - (\hat{\mathbf{r}} \cdot \hat{\mathbf{z}}) \hat{\mathbf{r}} = \hat{\mathbf{z}} - \cos \theta \hat{\mathbf{r}} = -\sin \theta \hat{\boldsymbol{\theta}},$$

and then

$$\mathbf{E}(t, \mathbf{x}) = -\frac{2L}{\pi c^2} \frac{I_0 \omega}{r} \sin\left(\omega t - \frac{\omega r}{c}\right) \sin \theta \hat{\boldsymbol{\theta}}.$$

The multipole expansion of the magnetic field in electrodynamics is given by Eq. (5.73),

$$\mathbf{B}(t, \mathbf{x}) = -\frac{1}{c^2 |\mathbf{x}|} \hat{\mathbf{x}} \times \left[\frac{d^2 \mathbf{p}}{dt^2} \right]_{\text{ret}} + \mathcal{O}\left(\frac{1}{|\mathbf{x}|^2}\right). \quad (11)$$

To first order in $1/|\mathbf{x}|$, we obtain

$$\mathbf{B}(t, \mathbf{x}) = \frac{1}{c^2 r} \hat{\mathbf{r}} \times \left[\frac{2L}{\pi} I_0 \omega \sin(\omega t) \hat{\mathbf{z}} \right]_{\text{ret}} = \frac{2L}{\pi c^2} \frac{I_0 \omega}{r} \hat{\mathbf{r}} \times \left[\sin(\omega t) \hat{\mathbf{z}} \right]_{\text{ret}} = \frac{2L}{\pi c^2} \frac{I_0 \omega}{r} \sin\left(\omega t - \frac{\omega r}{c}\right) (\hat{\mathbf{r}} \times \hat{\mathbf{z}}).$$

Again using spherical coordinates, $r(\hat{\mathbf{r}} \times \hat{\mathbf{z}}) = -r \sin \theta \hat{\boldsymbol{\phi}}$, and so

$$\mathbf{B}(t, \mathbf{x}) = -\frac{2L}{\pi c^2} \frac{I_0 \omega}{r} \sin\left(\omega t - \frac{\omega r}{c}\right) \sin \theta \hat{\boldsymbol{\phi}}.$$

In addition to the course lecture notes, I consulted Griffiths's *Introduction to Electrodynamics* while writing up these solutions.