

**Problem 1. (Jackson 14.1)** Verify by explicit calculation that the Liénard-Wiechert expressions for *all* components of  $\mathbf{E}$  and  $\mathbf{B}$  for a particle moving with constant velocity agree with the ones obtained in the text by means of a Lorentz transformation. Follow the general method at the end of Section 14.1.

**Solution.** The Liénard-Wiechert expressions for the fields are given by Jackson (14.13–14):

$$\mathbf{B} = [\hat{\mathbf{n}} \times \mathbf{E}]_{\text{ret}}, \quad \mathbf{E}(\mathbf{x}, t) = e \left[ \frac{\hat{\mathbf{n}} - \boldsymbol{\beta}}{\gamma^2(1 - \boldsymbol{\beta} \cdot \hat{\mathbf{n}})^3 R^2} \right]_{\text{ret}} + \frac{e}{c} \left[ \frac{\hat{\mathbf{n}} \times \{(\hat{\mathbf{n}} - \boldsymbol{\beta}) \times \dot{\boldsymbol{\beta}}\}}{(1 - \boldsymbol{\beta} \cdot \hat{\mathbf{n}})^3 R} \right]_{\text{ret}}, \quad (1)$$

where  $\boldsymbol{\beta} = \mathbf{v}/c$  with  $\mathbf{v}$  being the particle's velocity,  $R$  is the distance from the observation point to the particle's position, and  $\hat{\mathbf{n}}$  is a unit vector defined by  $\mathbf{x} - \mathbf{r}(\tau) = R \hat{\mathbf{n}}$ . Here,  $\mathbf{r}(\tau)$  is the particle's present position and  $\tau$  the proper time.

The expressions for the components of  $\mathbf{E}$  and  $\mathbf{B}$  obtained by a Lorentz transformation are given by Jackson (11.152):

$$E_1 = -\frac{e\gamma vt}{(b^2 + \gamma^2 v^2 t^2)^{3/2}}, \quad E_2 = \frac{e\gamma b}{(b^2 + \gamma^2 v^2 t^2)^{3/2}}, \quad E_3 = B_1 = B_2 = 0, \quad B_3 = \beta E_2, \quad (2)$$

where the particle is moving in the  $x_1$  direction at impact parameter  $b$  on the  $x_2$  axis, as shown in Fig. (1).

For a particle moving with constant velocity in the  $x_1$  direction with velocity  $v$  as shown in Fig. (1),  $\boldsymbol{\beta} = \beta \hat{\mathbf{x}}_1$  and  $\dot{\boldsymbol{\beta}} = 0$ . From Jackson (14.16), note that

$$(1 - \boldsymbol{\beta} \cdot \hat{\mathbf{n}})^2 R^2 = b^2 + v^2 t^2 - \beta^2 b^2 = \frac{b^2 + \gamma^2 v^2 t^2}{\gamma^2} \implies (1 - \boldsymbol{\beta} \cdot \hat{\mathbf{n}})^3 R^2 = \frac{(b^2 + \gamma^2 v^2 t^2)^{3/2}}{R\gamma^3}.$$

This calculation comes from Fig. (2), where  $O$  is the observation point,  $P$  is the present position of the particle, and  $P'$  its retarded position. Also from Fig. 2,

$$\hat{\mathbf{n}} = \cos \theta \hat{\mathbf{x}}_1 + \sin \theta \hat{\mathbf{x}}_2 = \frac{\beta R - vt}{R} \hat{\mathbf{x}}_1 + \frac{b}{R} \hat{\mathbf{x}}_2.$$

Making these substitutions in the expression for  $\mathbf{E}(\mathbf{x}, t)$  in Eq. (1),

$$\mathbf{E}(\mathbf{x}, t) = e \left[ \frac{\hat{\mathbf{n}} - \boldsymbol{\beta}}{\gamma^2(1 - \boldsymbol{\beta} \cdot \hat{\mathbf{n}})^3 R^2} \right]_{\text{ret}} = e \left[ \frac{(\beta - vt/R - \beta) \hat{\mathbf{x}}_1 + (b/R) \hat{\mathbf{x}}_2}{\gamma^2(b^2 + \gamma^2 v^2 t^2)^{3/2}} R\gamma^3 \right]_{\text{ret}} = e\gamma \frac{-vt \hat{\mathbf{x}}_1 + b \hat{\mathbf{x}}_2}{(b^2 + \gamma^2 v^2 t^2)^{3/2}}. \quad (3)$$




Figure 1: (Jackson Fig. 11.8) Particle of charge  $q$  moving at constant velocity  $\mathbf{v}$  passes an observation point  $P$  at impact parameter  $b$ .




Figure 2: (Jackson Fig. 14.2) Present and retarded positions of a charge in uniform motion.

For  $\mathbf{B}(\mathbf{x}, t)$ , note that

$$\hat{\mathbf{n}} \times \mathbf{E} \propto \left( \frac{\beta R - vt}{R} \hat{\mathbf{x}}_1 + \frac{b}{R} \hat{\mathbf{x}}_2 \right) \times (-vt \hat{\mathbf{x}}_1 + b \hat{\mathbf{x}}_2) = \left( b \frac{\beta R - vt}{R} + \frac{bvt}{R} \right) \hat{\mathbf{x}}_3 = \beta b,$$

so

$$\mathbf{B}(\mathbf{x}, t) = e\gamma \frac{\beta b \hat{\mathbf{x}}_3}{(b^2 + \gamma^2 v^2 t^2)^{3/2}}. \quad (4)$$

Writing Eqs. (3) and (4) in component notation, we find

$$\begin{aligned} E_1 &= -\frac{e\gamma vt}{(b^2 + \gamma^2 v^2 t^2)^{3/2}}, & E_2 &= \frac{e\gamma b}{(b^2 + \gamma^2 v^2 t^2)^{3/2}}, & E_3 &= 0, \\ B_1 &= 0, & B_2 &= 0, & B_3 &= \frac{e\gamma \beta b}{(b^2 + \gamma^2 v^2 t^2)^{3/2}} = \beta E_2, \end{aligned}$$

which are identical to Eq. (2) as desired.  $\square$

**Problem 2. (Jackson 14.3)** The Heaviside-Feynman expression for the electric field of a particle of charge  $e$  in arbitrary motion, an alternative to the Liénard-Wiechert expression (14.14) is

$$\mathbf{E} = e \left[ \frac{\hat{\mathbf{n}}}{R^2} \right]_{\text{ret}} + e \left[ \frac{R}{c} \right]_{\text{ret}} \frac{d}{dt} \left[ \frac{\hat{\mathbf{n}}}{R^2} \right]_{\text{ret}} + \frac{e^2}{c^2} \frac{d^2}{dt^2} [\hat{\mathbf{n}}]_{\text{ret}},$$

where the time derivatives are with respect to the time at the observation point. The magnetic fields are given by (14.3).

Using the fact that the retarded time is  $t' = t - R(t')/c$  and that, as a result,

$$\frac{dt}{dt'} = 1 - \beta(t') \cdot \hat{\mathbf{n}}(t'),$$

show that the form above yields (14.14) when the time differentiations are performed.

**Problem 3. (Jackson 14.4)** Using the Liénard-Wiechart fields, discuss the time-averaged power radiated per unit solid angle in nonrelativistic motion of a particle with charge  $e$ . Sketch the angular distribution of the radiation and determine the total power radiated in each case.

**3(a)** The particle is moving along the  $z$  axis with instantaneous position  $z(t) = \alpha \cos \omega_0 t$ .

**3(b)** The particle is moving in a circle of radius  $R$  in the  $xy$  plane with constant angular frequency  $\omega_0$ .