

---

# Models

Justin Salmon<sup>1</sup> and George Lancaster<sup>2</sup>

<sup>1</sup>*wr18313*

<sup>2</sup>*qv18258*

---

October 15, 2018

**L**orem ipsum dolor sit amet, consectetur adipiscing elit. Fusce maximus nisi ligula. Morbi laoreet ex ligula, vitae lobortis purus mattis vel. Vestibulum ante ipsum primis in faucibus orci luctus et ultrices posuere cubilia Curae; Donec ac metus ut turpis mollis placerat et nec enim. Duis tristique nibh maximus faucibus facilisis. Praesent in consequat leo. Maecenas condimentum ex rhoncus, elementum diam vel, malesuada ante.

## 1 The Prior

Q1: Most probabilistic processes in nature tend to follow Gaussian distributions, hence this is generally a good place to start.

Noise in input data - assume noise is gaussian -> implies gaussian likelihood

Q2; Equally likely to deviate from the mean in all directions. Again a good place to start

Q3: The covariance matrix would not be in terms of the identity matrix (for some reason). We would have non-zero values in the offset diagonals which correspond to the correlations between different things.

Q5: Euclidean distance

## 2 Posterior

$$f \sim \mathcal{N}(\mathbf{0}, \beta^{-1} \mathbf{I}) \quad (1)$$

$$\beta \sim \Gamma(a) \quad (2)$$

## 3 Evidence