

Lecture 2: Asymptotic Analysis, Recurrences

Michael Dinitz

September 3, 2026

601.433/633 Introduction to Algorithms

Course changes and updates

Homeworks now released and due on Thursdays!

- ▶ HW1 released Sep 10, due Sep 17 at 11:59pm
- ▶ Makes more sense with office hours and lecture schedule.
- ▶ Note: probably HW due on fall break (Oct 22)!
- ▶ Oral presentations will be on Th/F

Office hours will be finalized soon (early next week)

- ▶ Prof. Dinitz office hours: Th 11am-12pm
- ▶ Prof. Sorrell office hours: sometime Wednesday (probably 11am-12pm)
- ▶ CA office hours: likely M-Th, 6pm-9pm
- ▶ TA Shruthi Prusty office hours: TBD

Today

Should be review, some might be new.

See math background in CLRS.

Asymptotics: $O(\cdot)$, $\Omega(\cdot)$, and $\Theta(\cdot)$ notation.

- ▶ Should know from Data Structures / MFCS. We'll be a bit more formal.
- ▶ Intuitively: hide constants and lower-order terms, since we only care what happens “at scale” (asymptotically).

Recurrences: How to solve recurrence relations.

- ▶ Should know from MFCS / Discrete Math.

Asymptotic Notation

$O(\cdot)$

Definition

$g(n) \in O(f(n))$ if there exist constants $c, n_0 > 0$ such that $g(n) \leq c \cdot f(n)$ for all $n > n_0$.

$O(\cdot)$

Definition

$g(n) \in O(f(n))$ if there exist constants $c, n_0 > 0$ such that $g(n) \leq c \cdot f(n)$ for all $n > n_0$.

Technically $O(f(n))$ is a set.

Abuse notation: “ $g(n)$ is $O(f(n))$ ” or $g(n) = O(f(n))$.

$O(\cdot)$

Definition

$g(n) \in O(f(n))$ if there exist constants $c, n_0 > 0$ such that $g(n) \leq c \cdot f(n)$ for all $n > n_0$.

Technically $O(f(n))$ is a set.

Abuse notation: “ $g(n)$ is $O(f(n))$ ” or $g(n) = O(f(n))$.

Examples:

- ▶ $2n^2 + 27 = O(n^2)$: set $n_0 = 6$ and $c = 3$
- ▶ $2n^2 + 27 = O(n^3)$: same values, or $n_0 = 4$ and $c = 1$
- ▶ $n^3 + 2000n^2 + 2000n = O(n^3)$: set $n_0 = 10000$ and $c = 2$

$O(\cdot)$

Definition

$g(n) \in O(f(n))$ if there exist constants $c, n_0 > 0$ such that $g(n) \leq c \cdot f(n)$ for all $n > n_0$.

Technically $O(f(n))$ is a set.

Abuse notation: “ $g(n)$ is $O(f(n))$ ” or $g(n) = O(f(n))$.

Examples:

- ▶ $2n^2 + 27 = O(n^2)$: set $n_0 = 6$ and $c = 3$
- ▶ $2n^2 + 27 = O(n^3)$: same values, or $n_0 = 4$ and $c = 1$
- ▶ $n^3 + 2000n^2 + 2000n = O(n^3)$: set $n_0 = 10000$ and $c = 2$

About *functions*, not algorithms!

Expresses an *upper* bound.

Example

Definition

$g(n) \in O(f(n))$ if there exist constants $c, n_0 > 0$ such that $g(n) \leq c \cdot f(n)$ for all $n > n_0$.

Theorem

$$2n^2 + 27 = O(n^2)$$

Example

Definition

$g(n) \in O(f(n))$ if there exist constants $c, n_0 > 0$ such that $g(n) \leq c \cdot f(n)$ for all $n > n_0$.

Theorem

$$2n^2 + 27 = O(n^2)$$

Proof.

Set $c = 3$. Suppose $2n^2 + 27 > cn^2 = 3n^2$

Example

Definition

$g(n) \in O(f(n))$ if there exist constants $c, n_0 > 0$ such that $g(n) \leq c \cdot f(n)$ for all $n > n_0$.

Theorem

$$2n^2 + 27 = O(n^2)$$

Proof.

Set $c = 3$. Suppose $2n^2 + 27 > cn^2 = 3n^2$
 $\implies n^2 < 27$

Example

Definition

$g(n) \in O(f(n))$ if there exist constants $c, n_0 > 0$ such that $g(n) \leq c \cdot f(n)$ for all $n > n_0$.

Theorem

$$2n^2 + 27 = O(n^2)$$

Proof.

Set $c = 3$. Suppose $2n^2 + 27 > cn^2 = 3n^2$
 $\implies n^2 < 27 \implies n < 6$

Example

Definition

$g(n) \in O(f(n))$ if there exist constants $c, n_0 > 0$ such that $g(n) \leq c \cdot f(n)$ for all $n > n_0$.

Theorem

$$2n^2 + 27 = O(n^2)$$

Proof.

Set $c = 3$. Suppose $2n^2 + 27 > cn^2 = 3n^2$

$$\implies n^2 < 27 \implies n < 6$$

$$\implies 2n^2 + 27 \leq 3n^2 \text{ for all } n \geq 6.$$

Example

Definition

$g(n) \in O(f(n))$ if there exist constants $c, n_0 > 0$ such that $g(n) \leq c \cdot f(n)$ for all $n > n_0$.

Theorem

$$2n^2 + 27 = O(n^2)$$

Proof.

Set $c = 3$. Suppose $2n^2 + 27 > cn^2 = 3n^2$

$$\implies n^2 < 27 \implies n < 6$$

$$\implies 2n^2 + 27 \leq 3n^2 \text{ for all } n \geq 6.$$

Set $n_0 = 6$. Then $2n^2 + 27 \leq cn^2$ for all $n > n_0$. □

Example

Definition

$g(n) \in O(f(n))$ if there exist constants $c, n_0 > 0$ such that $g(n) \leq c \cdot f(n)$ for all $n > n_0$.

Theorem

$$2n^2 + 27 = O(n^2)$$

Proof.

Set $c = 3$. Suppose $2n^2 + 27 > cn^2 = 3n^2$

$$\implies n^2 < 27 \implies n < 6$$

$$\implies 2n^2 + 27 \leq 3n^2 \text{ for all } n \geq 6.$$

Set $n_0 = 6$. Then $2n^2 + 27 \leq cn^2$ for all $n > n_0$. □

Many other ways to prove this!

$\Omega(\cdot)$

Counterpart to $O(\cdot)$: *lower* bound rather than upper bound.

Definition

$g(n) \in \Omega(f(n))$ if there exist constants $c, n_0 > 0$ such that $g(n) \geq c \cdot f(n)$ for all $n > n_0$.

$\Omega(\cdot)$

Counterpart to $O(\cdot)$: *lower* bound rather than upper bound.

Definition

$g(n) \in \Omega(f(n))$ if there exist constants $c, n_0 > 0$ such that $g(n) \geq c \cdot f(n)$ for all $n > n_0$.

Examples:

- ▶ $2n^2 + 27 = \Omega(n^2)$: set $n_0 = 1$ and $c = 1$
- ▶ $2n^2 + 27 = \Omega(n)$: set $n_0 = 1$ and $c = 1$
- ▶ $\frac{1}{100}n^3 - 1000n^2 = \Omega(n^3)$: set $n_0 = 1000000$ and $c = 1/1000$

$\Theta(\cdot)$

Combination of $O(\cdot)$ and $\Omega(\cdot)$.

Definition

$g(n) \in \Theta(f(n))$ if $g(n) \in O(f(n))$ and $g(n) \in \Omega(f(n))$.

Note: constants n_0, c can be different in the proofs for $O(f(n))$ and $\Omega(f(n))$

$\Theta(\cdot)$

Combination of $O(\cdot)$ and $\Omega(\cdot)$.

Definition

$g(n) \in \Theta(f(n))$ if $g(n) \in O(f(n))$ and $g(n) \in \Omega(f(n))$.

Note: constants n_0, c can be different in the proofs for $O(f(n))$ and $\Omega(f(n))$

Equivalently:

Definition

$g(n) \in \Theta(f(n))$ if there are constants $c_1, c_2, n_0 > 0$ such that $c_1 f(n) \leq g(n) \leq c_2 f(n)$ for all $n > n_0$.

Both lower bound and upper bound, so asymptotic equality.

Little notation

Strict versions of $\mathcal{O}$ and Ω :

Definition

$g(n) \in \mathcal{o}(f(n))$ if for every constant $c > 0$ there exists a constant $n_0 > 0$ such that $g(n) < c \cdot f(n)$ for all $n > n_0$.

Definition

$g(n) \in \omega(f(n))$ if for every constant $c > 0$ there exists a constant $n_0 > 0$ such that $g(n) > c \cdot f(n)$ for all $n > n_0$.

Examples:

- ▶ $2n^2 + 27 = \mathcal{o}(n^2 \log n)$
- ▶ $2n^2 + 27 = \omega(n)$

Recurrence Relations

Sorting

Many algorithms are recursive, so their running time is naturally a recurrence relation (Karatsuba, Strassen).

Sorting

Many algorithms are recursive, so their running time is naturally a recurrence relation (Karatsuba, Strassen).

Sorting: Let $T(n)$ denote the (worst-case) running time on an array of size n .

Sorting

Many algorithms are recursive, so their running time is naturally a recurrence relation (Karatsuba, Strassen).

Sorting: Let $T(n)$ denote the (worst-case) running time on an array of size n .

- ▶ Selection Sort

Sorting

Many algorithms are recursive, so their running time is naturally a recurrence relation (Karatsuba, Strassen).

Sorting: Let $T(n)$ denote the (worst-case) running time on an array of size n .

- ▶ Selection Sort
 - ▶ Find smallest unsorted element, put it just after sorted elements. Repeat.

Sorting

Many algorithms are recursive, so their running time is naturally a recurrence relation (Karatsuba, Strassen).

Sorting: Let $T(n)$ denote the (worst-case) running time on an array of size n .

- ▶ Selection Sort

- ▶ Find smallest unsorted element, put it just after sorted elements. Repeat.
- ▶ Running time: Takes $O(n)$ time to find smallest unsorted element, decreases remaining unsorted by 1.

$$\implies T(n) = T(n-1) + cn$$

Sorting

Many algorithms are recursive, so their running time is naturally a recurrence relation (Karatsuba, Strassen).

Sorting: Let $T(n)$ denote the (worst-case) running time on an array of size n .

- ▶ Selection Sort
 - ▶ Find smallest unsorted element, put it just after sorted elements. Repeat.
 - ▶ Running time: Takes $O(n)$ time to find smallest unsorted element, decreases remaining unsorted by 1.
 $\implies T(n) = T(n-1) + cn$
- ▶ Mergesort

Sorting

Many algorithms are recursive, so their running time is naturally a recurrence relation (Karatsuba, Strassen).

Sorting: Let $T(n)$ denote the (worst-case) running time on an array of size n .

- ▶ Selection Sort

- ▶ Find smallest unsorted element, put it just after sorted elements. Repeat.
- ▶ Running time: Takes $O(n)$ time to find smallest unsorted element, decreases remaining unsorted by 1.
 $\implies T(n) = T(n-1) + cn$

- ▶ Mergesort

- ▶ Split array into left and right halves. Recursively sort each half, then merge.

Sorting

Many algorithms are recursive, so their running time is naturally a recurrence relation (Karatsuba, Strassen).

Sorting: Let $T(n)$ denote the (worst-case) running time on an array of size n .

- ▶ Selection Sort

- ▶ Find smallest unsorted element, put it just after sorted elements. Repeat.
- ▶ Running time: Takes $O(n)$ time to find smallest unsorted element, decreases remaining unsorted by 1.
 $\implies T(n) = T(n-1) + cn$

- ▶ Mergesort

- ▶ Split array into left and right halves. Recursively sort each half, then merge.
- ▶ Running time: Merging takes $O(n)$ time. Two recursive calls on half the size.
 $\implies T(n) = T(n/2) + T(n/2) + cn = 2T(n/2) + cn$

Sorting

Many algorithms are recursive, so their running time is naturally a recurrence relation (Karatsuba, Strassen).

Sorting: Let $T(n)$ denote the (worst-case) running time on an array of size n .

- ▶ Selection Sort

- ▶ Find smallest unsorted element, put it just after sorted elements. Repeat.
- ▶ Running time: Takes $O(n)$ time to find smallest unsorted element, decreases remaining unsorted by 1.
 $\implies T(n) = T(n-1) + cn$

- ▶ Mergesort

- ▶ Split array into left and right halves. Recursively sort each half, then merge.
- ▶ Running time: Merging takes $O(n)$ time. Two recursive calls on half the size.
 $\implies T(n) = T(n/2) + T(n/2) + cn = 2T(n/2) + cn$

Also need base case. For algorithms, constant-size input takes constant time.

$\implies T(n) \leq c$ for all $n \leq n_0$, for some constants $n_0, c > 0$.

Guess and Check

$$T(n) = 3T(n/3) + n$$

$$T(1) = 1$$

Guess and Check

$$T(n) = 3T(n/3) + n$$

$$T(1) = 1$$

Guess: $T(n) \leq cn$.

Guess and Check

$$T(n) = 3T(n/3) + n$$

$$T(1) = 1$$

Guess: $T(n) \leq cn$.

Check: assume true for $n' < n$, prove true for n (induction).

Guess and Check

$$T(n) = 3T(n/3) + n$$

$$T(1) = 1$$

Guess: $T(n) \leq cn$.

Check: assume true for $n' < n$, prove true for n (induction).

$$T(n) = 3T(n/3) + n \leq 3cn/3 + n = (c + 1)n$$

Guess and Check

$$T(n) = 3T(n/3) + n$$

$$T(1) = 1$$

Guess: $T(n) \leq cn$.

Check: assume true for $n' < n$, prove true for n (induction).

$$T(n) = 3T(n/3) + n \leq 3cn/3 + n = (c + 1)n$$

Failure! Wanted $T(n) \leq cn$, got $T(n) \leq (c + 1)n$. Guess was wrong.

Guess and Check

$$T(n) = 3T(n/3) + n$$

$$T(1) = 1$$

Guess: $T(n) \leq cn$.

Check: assume true for $n' < n$, prove true for n (induction).

$$T(n) = 3T(n/3) + n \leq 3cn/3 + n = (c + 1)n$$

Failure! Wanted $T(n) \leq cn$, got $T(n) \leq (c + 1)n$. Guess was wrong.

Better guess? What goes up by 1 when n goes up by a factor of 3?

Guess and Check

$$T(n) = 3T(n/3) + n$$

$$T(1) = 1$$

Guess: $T(n) \leq cn$.

Check: assume true for $n' < n$, prove true for n (induction).

$$T(n) = 3T(n/3) + n \leq 3cn/3 + n = (c + 1)n$$

Failure! Wanted $T(n) \leq cn$, got $T(n) \leq (c + 1)n$. Guess was wrong.

Better guess? What goes up by 1 when n goes up by a factor of 3? $\log_3 n$

Guess and Check

$$T(n) = 3T(n/3) + n$$

$$T(1) = 1$$

Guess: $T(n) \leq cn$.

Check: assume true for $n' < n$, prove true for n (induction).

$$T(n) = 3T(n/3) + n \leq 3cn/3 + n = (c + 1)n$$

Failure! Wanted $T(n) \leq cn$, got $T(n) \leq (c + 1)n$. Guess was wrong.

Better guess? What goes up by 1 when n goes up by a factor of 3? $\log_3 n$

Guess: $T(n) \leq n \log_3(3n)$

Guess and Check

$$T(n) = 3T(n/3) + n$$

$$T(1) = 1$$

Guess: $T(n) \leq cn$.

Check: assume true for $n' < n$, prove true for n (induction).

$$T(n) = 3T(n/3) + n \leq 3cn/3 + n = (c + 1)n$$

Failure! Wanted $T(n) \leq cn$, got $T(n) \leq (c + 1)n$. Guess was wrong.

Better guess? What goes up by 1 when n goes up by a factor of 3? $\log_3 n$

Guess: $T(n) \leq n \log_3(3n)$

Check: assume true for $n' < n$, prove true for n (induction).

$$\begin{aligned} T(n) &= 3T(n/3) + n \leq 3(n/3) \log_3(3(n/3)) + n = n \log_3(n) + n \\ &= n(\log_3(n) + \log_3 3) = n \log_3(3n). \end{aligned}$$

Unrolling

Example: selection sort. $T(n) = T(n-1) + cn$

Idea: “unroll” the recurrence.

Unrolling

Example: selection sort. $T(n) = T(n-1) + cn$

Idea: “unroll” the recurrence.

$$T(n) = cn + T(n-1)$$

Unrolling

Example: selection sort. $T(n) = T(n-1) + cn$

Idea: “unroll” the recurrence.

$$\begin{aligned} T(n) &= cn + T(n-1) \\ &= cn + c(n-1) + T(n-2) \end{aligned}$$

Unrolling

Example: selection sort. $T(n) = T(n-1) + cn$

Idea: “unroll” the recurrence.

$$\begin{aligned}T(n) &= cn + T(n-1) \\&= cn + c(n-1) + T(n-2) \\&= cn + c(n-1) + c(n-2) + T(n-3)\end{aligned}$$

Unrolling

Example: selection sort. $T(n) = T(n-1) + cn$

Idea: “unroll” the recurrence.

$$\begin{aligned}T(n) &= cn + T(n-1) \\&= cn + c(n-1) + T(n-2) \\&= cn + c(n-1) + c(n-2) + T(n-3) \\&\quad \vdots\end{aligned}$$

Unrolling

Example: selection sort. $T(n) = T(n-1) + cn$

Idea: “unroll” the recurrence.

$$\begin{aligned}T(n) &= cn + T(n-1) \\&= cn + c(n-1) + T(n-2) \\&= cn + c(n-1) + c(n-2) + T(n-3) \\&\quad \vdots \\&= cn + c(n-1) + c(n-2) + \cdots + c\end{aligned}$$

Unrolling

Example: selection sort. $T(n) = T(n-1) + cn$

Idea: “unroll” the recurrence.

$$\begin{aligned}T(n) &= cn + T(n-1) \\&= cn + c(n-1) + T(n-2) \\&= cn + c(n-1) + c(n-2) + T(n-3) \\&\quad \vdots \\&= cn + c(n-1) + c(n-2) + \cdots + c\end{aligned}$$

n terms, each of which is at most $cn \implies T(n) \leq cn^2 = O(n^2)$

Unrolling

Example: selection sort. $T(n) = T(n-1) + cn$

Idea: “unroll” the recurrence.

$$\begin{aligned}T(n) &= cn + T(n-1) \\&= cn + c(n-1) + T(n-2) \\&= cn + c(n-1) + c(n-2) + T(n-3) \\&\quad \vdots \\&= cn + c(n-1) + c(n-2) + \cdots + c\end{aligned}$$

n terms, each of which is at most $cn \implies T(n) \leq cn^2 = O(n^2)$

At least $n/2$ terms are each at least $cn/2 \implies T(n) \geq c\frac{n^2}{4} = \Omega(n^2)$

Unrolling

Example: selection sort. $T(n) = T(n-1) + cn$

Idea: “unroll” the recurrence.

$$\begin{aligned}T(n) &= cn + T(n-1) \\&= cn + c(n-1) + T(n-2) \\&= cn + c(n-1) + c(n-2) + T(n-3) \\&\quad \vdots \\&= cn + c(n-1) + c(n-2) + \cdots + c\end{aligned}$$

n terms, each of which is at most $cn \implies T(n) \leq cn^2 = O(n^2)$

At least $n/2$ terms are each at least $cn/2 \implies T(n) \geq c\frac{n^2}{4} = \Omega(n^2)$
 $\implies T(n) = \Theta(n^2)$.

Recursion Tree: Mergesort


Generalizes unrolling: draw out full tree of “recursive calls”.

Mergesort: $T(n) = 2T(n/2) + cn$.

Recursion Tree: Mergesort

Generalizes unrolling: draw out full tree of “recursive calls”.


Mergesort: $T(n) = 2T(n/2) + cn$.



Recursion Tree: Mergesort

Generalizes unrolling: draw out full tree of “recursive calls”.

Mergesort: $T(n) = 2T(n/2) + cn$.




levels:

Recursion Tree: Mergesort

Generalizes unrolling: draw out full tree of “recursive calls”.

Mergesort: $T(n) = 2T(n/2) + cn$.




levels: $\log_2 n$

Recursion Tree: Mergesort

Generalizes unrolling: draw out full tree of “recursive calls”.

Mergesort: $T(n) = 2T(n/2) + cn$.




levels: $\log_2 n$

Contribution of level i :

Recursion Tree: Mergesort

Generalizes unrolling: draw out full tree of “recursive calls”.

Mergesort: $T(n) = 2T(n/2) + cn$.




levels: $\log_2 n$

Contribution of level i : $2^{i-1} cn / 2^{i-1} = cn$

Recursion Tree: Mergesort

Generalizes unrolling: draw out full tree of “recursive calls”.

Mergesort: $T(n) = 2T(n/2) + cn$.



levels: $\log_2 n$

Contribution of level i : $2^{i-1} cn/2^{i-1} = cn$


$\implies T(n) = \Theta(n \log n)$

Recursion Tree: Strassen

$$T(n) = 7T(n/2) + cn^2$$


Recursion Tree: Strassen

$$T(n) = 7T(n/2) + cn^2$$



Recursion Tree: Strassen


$$T(n) = 7T(n/2) + cn^2$$



Level i : $7^{i-1}c(n/2^{i-1})^2 = (7/4)^{i-1}cn^2$

Recursion Tree: Strassen

$$T(n) = 7T(n/2) + cn^2$$




Level i : $7^{i-1}c(n/2^{i-1})^2 = (7/4)^{i-1}cn^2$

$$T(n) = \sum_{i=1}^{\log n+1} \left(\frac{7}{4}\right)^{i-1} cn^2 = cn^2 \sum_{i=1}^{\log n+1} \left(\frac{7}{4}\right)^{i-1}$$

Total:

Recursion Tree: Strassen

$$T(n) = 7T(n/2) + cn^2$$



Level i : $7^{i-1}c(n/2^{i-1})^2 = (7/4)^{i-1}cn^2$

$$T(n) = \sum_{i=1}^{\log n+1} \left(\frac{7}{4}\right)^{i-1} cn^2 = cn^2 \sum_{i=1}^{\log n+1} \left(\frac{7}{4}\right)^{i-1}$$

Total:

$$\begin{aligned} \implies T(n) &= O(n^2(7/4)^{\log n}) = O(n^2 n^{\log(7/4)}) = O(n^2 n^{\log 7 - 2}) \\ &= O(n^{\log 7}) \end{aligned}$$

Master Theorem

$$T(n) = aT(n/b) + cn^k \qquad T(1) = c$$


a, b, c, k constants with **$a \geq 1$** , **$b > 1$** , **$c > 0$** , and **$k \geq 0$**

Master Theorem

$$T(n) = aT(n/b) + cn^k$$

$$T(1) = c$$

a, b, c, k constants with $a \geq 1$, $b > 1$, $c > 0$, and $k \geq 0$




Master Theorem

$$T(n) = aT(n/b) + cn^k$$

$$T(1) = c$$

a, b, c, k constants with $a \geq 1$, $b > 1$, $c > 0$, and $k \geq 0$




levels: $\log_b n + 1$

Master Theorem

$$T(n) = aT(n/b) + cn^k$$

$$T(1) = c$$

a, b, c, k constants with $a \geq 1$, $b > 1$, $c > 0$, and $k \geq 0$



levels: $\log_b n + 1$

Level i : $a^{i-1}c(n/b^{i-1})^k = cn^k(a/b^k)^{i-1}$

Master Theorem II

Let $\alpha = (a/b^k)$

$$\implies T(n) = cn^k \sum_{i=1}^{\log_b n+1} (a/b^k)^{i-1} = cn^k \sum_{i=1}^{\log_b n+1} \alpha^{i-1}$$

Master Theorem II

Let $\alpha = (a/b^k)$

$$\implies T(n) = cn^k \sum_{i=1}^{\log_b n+1} (a/b^k)^{i-1} = cn^k \sum_{i=1}^{\log_b n+1} \alpha^{i-1}$$

- ▶ Case 1: $\alpha = 1$. All levels the same. $T(n) = cn^k \sum_{i=1}^{\log_b n+1} 1 = \Theta(n^k \log n)$

Master Theorem II

Let $\alpha = (a/b^k)$

$$\implies T(n) = cn^k \sum_{i=1}^{\log_b n+1} (a/b^k)^{i-1} = cn^k \sum_{i=1}^{\log_b n+1} \alpha^{i-1}$$

- ▶ Case 1: $\alpha = 1$. All levels the same. $T(n) = cn^k \sum_{i=1}^{\log_b n+1} 1 = \Theta(n^k \log n)$
- ▶ Case 2: $\alpha < 1$. Dominated by top level.

Master Theorem II

Let $\alpha = (a/b^k)$

$$\implies T(n) = cn^k \sum_{i=1}^{\log_b n+1} (a/b^k)^{i-1} = cn^k \sum_{i=1}^{\log_b n+1} \alpha^{i-1}$$

► Case 1: $\alpha = 1$. All levels the same. $T(n) = cn^k \sum_{i=1}^{\log_b n+1} 1 = \Theta(n^k \log n)$

► Case 2: $\alpha < 1$. Dominated by top level.

$$\implies \sum_{i=1}^{\log_b n+1} \alpha^{i-1} \leq \sum_{i=1}^{\infty} \alpha^{i-1} = \frac{1}{1-\alpha}.$$

$$\implies T(n) = O(n^k)$$

Master Theorem II

Let $\alpha = (a/b^k)$

$$\implies T(n) = cn^k \sum_{i=1}^{\log_b n+1} (a/b^k)^{i-1} = cn^k \sum_{i=1}^{\log_b n+1} \alpha^{i-1}$$

► Case 1: $\alpha = 1$. All levels the same. $T(n) = cn^k \sum_{i=1}^{\log_b n+1} 1 = \Theta(n^k \log n)$

► Case 2: $\alpha < 1$. Dominated by top level.

$$\implies \sum_{i=1}^{\log_b n+1} \alpha^{i-1} \leq \sum_{i=1}^{\infty} \alpha^{i-1} = \frac{1}{1-\alpha}.$$

$$\implies T(n) = O(n^k)$$

$$T(n) \geq cn^k \implies T(n) = \Omega(n^k)$$

Master Theorem II

Let $\alpha = (a/b^k)$

$$\implies T(n) = cn^k \sum_{i=1}^{\log_b n+1} (a/b^k)^{i-1} = cn^k \sum_{i=1}^{\log_b n+1} \alpha^{i-1}$$

- ▶ Case 1: $\alpha = 1$. All levels the same. $T(n) = cn^k \sum_{i=1}^{\log_b n+1} 1 = \Theta(n^k \log n)$
- ▶ Case 2: $\alpha < 1$. Dominated by top level.

$$\implies \sum_{i=1}^{\log_b n+1} \alpha^{i-1} \leq \sum_{i=1}^{\infty} \alpha^{i-1} = \frac{1}{1-\alpha}.$$

$$\implies T(n) = O(n^k)$$

$$T(n) \geq cn^k \implies T(n) = \Omega(n^k) \implies T(n) = \Theta(n^k)$$

Master Theorem II

Let $\alpha = (a/b^k)$

$$\implies T(n) = cn^k \sum_{i=1}^{\log_b n+1} (a/b^k)^{i-1} = cn^k \sum_{i=1}^{\log_b n+1} \alpha^{i-1}$$

► Case 1: $\alpha = 1$. All levels the same. $T(n) = cn^k \sum_{i=1}^{\log_b n+1} 1 = \Theta(n^k \log n)$

► Case 2: $\alpha < 1$. Dominated by top level.

$$\implies \sum_{i=1}^{\log_b n+1} \alpha^{i-1} \leq \sum_{i=1}^{\infty} \alpha^{i-1} = \frac{1}{1-\alpha}.$$

$$\implies T(n) = O(n^k)$$

$$T(n) \geq cn^k \implies T(n) = \Omega(n^k) \implies T(n) = \Theta(n^k)$$

► Case 3: $\alpha > 1$. Dominated by bottom level.

Master Theorem II

Let $\alpha = (a/b^k)$

$$\implies T(n) = cn^k \sum_{i=1}^{\log_b n+1} (a/b^k)^{i-1} = cn^k \sum_{i=1}^{\log_b n+1} \alpha^{i-1}$$

► Case 1: $\alpha = 1$. All levels the same. $T(n) = cn^k \sum_{i=1}^{\log_b n+1} 1 = \Theta(n^k \log n)$

► Case 2: $\alpha < 1$. Dominated by top level.

$$\implies \sum_{i=1}^{\log_b n+1} \alpha^{i-1} \leq \sum_{i=1}^{\infty} \alpha^{i-1} = \frac{1}{1-\alpha}.$$

$$\implies T(n) = O(n^k)$$

$$T(n) \geq cn^k \implies T(n) = \Omega(n^k) \implies T(n) = \Theta(n^k)$$

► Case 3: $\alpha > 1$. Dominated by bottom level.

$$\begin{aligned} \implies \sum_{i=1}^{\log_b n+1} \alpha^{i-1} &= \alpha^{\log_b n} \sum_{i=1}^{\log_b n+1} \left(\frac{1}{\alpha}\right)^{i-1} \leq \alpha^{\log_b n} \frac{1}{1-(1/\alpha)} \\ &= O(\alpha^{\log_b n}) \end{aligned}$$

Master Theorem II

Let $\alpha = (a/b^k)$

$$\implies T(n) = cn^k \sum_{i=1}^{\log_b n+1} (a/b^k)^{i-1} = cn^k \sum_{i=1}^{\log_b n+1} \alpha^{i-1}$$

► Case 1: $\alpha = 1$. All levels the same. $T(n) = cn^k \sum_{i=1}^{\log_b n+1} 1 = \Theta(n^k \log n)$

► Case 2: $\alpha < 1$. Dominated by top level.

$$\implies \sum_{i=1}^{\log_b n+1} \alpha^{i-1} \leq \sum_{i=1}^{\infty} \alpha^{i-1} = \frac{1}{1-\alpha}.$$

$$\implies T(n) = O(n^k)$$

$$T(n) \geq cn^k \implies T(n) = \Omega(n^k) \implies T(n) = \Theta(n^k)$$

► Case 3: $\alpha > 1$. Dominated by bottom level.

$$\implies \sum_{i=1}^{\log_b n+1} \alpha^{i-1} = \alpha^{\log_b n} \sum_{i=1}^{\log_b n+1} \left(\frac{1}{\alpha}\right)^{i-1} \leq \alpha^{\log_b n} \frac{1}{1-(1/\alpha)}$$

$$= O(\alpha^{\log_b n})$$

$$\begin{aligned} \implies T(n) &= \Theta(n^k \alpha^{\log_b n}) = \Theta(n^k (a/b^k)^{\log_b n}) = \Theta(a^{\log_b n}) \\ &= \Theta(n^{\log_b a}) \end{aligned}$$

Master Theorem III

Theorem (“Master Theorem”)

The recurrence

$$T(n) = aT(n/b) + cn^k$$

$$T(1) = c$$

where a, b, c , and k are constants with $a \geq 1$, $b > 1$, $c > 0$, and $k \geq 0$, satisfies

$$T(n) = \begin{cases} \Theta(n^k) & \text{if } a < b^k, \\ \Theta(n^k \log n) & \text{if } a = b^k, \\ \Theta(n^{\log_b a}) & \text{if } a > b^k. \end{cases}$$