

Lecture 2: Asymptotic Analysis, Recurrences

Jessica Sorrell

Sep 3, 2026

601.433/633 Introduction to Algorithms

Slides from Mike Dinitz, modified for our section

Administrative

- ▶ Schedule change! See course website
- ▶ Oral Homeworks: The oral-presentation homeworks will be done in groups of three. Each of these assignments will consist of three problems. The members of your group will work together to solve the problems (so, unlike written homeworks, here collaboration is required). You will then present your solutions, as a group, to one of the instructors or TAs.
- ▶ Presentations will be given in 1-hour time slots (there will be an electronic sign-up sheet reachable from the course home page). At the presentation, each member of the group will spend 15 minutes presenting one of the problems. The instructor (or TA) will decide who presents which problem, but when one member is presenting, other members are allowed to chime in too. In the end, the three presentations together will determine the grade for the group.
- ▶ If you are nervous about your presentation, you may in addition hand in a written sketch of your solution as well. We will then take this writeup into consideration in determining your grade on the assignment.

Today

Should be review, some might be new.

See math background in CLRS

Asymptotics: $O(\cdot)$, $\Omega(\cdot)$, and $\Theta(\cdot)$ notation.

- ▶ Should know from Data Structures / MFCS. We'll be a bit more formal.
- ▶ Intuitively: hide constants and lower order terms, since we only care what happens “at scale” (asymptotically)

Recurrences: How to solve recurrence relations.

- ▶ Should know from MFCS / Discrete Math.

Asymptotic Notation

$O(\cdot)$

Definition

$g(n) \in O(f(n))$ if there exist constants $c, n_0 > 0$ such that $g(n) \leq c \cdot f(n)$ for all $n > n_0$.

$O(\cdot)$

Definition

$g(n) \in O(f(n))$ if there exist constants $c, n_0 > 0$ such that $g(n) \leq c \cdot f(n)$ for all $n > n_0$.

Technically $O(f(n))$ is a set.

Abuse notation: “ $g(n)$ is $O(f(n))$ ” or $g(n) = O(f(n))$.

$O(\cdot)$

Definition

$g(n) \in O(f(n))$ if there exist constants $c, n_0 > 0$ such that $g(n) \leq c \cdot f(n)$ for all $n > n_0$.

Technically $O(f(n))$ is a set.

Abuse notation: “ $g(n)$ is $O(f(n))$ ” or $g(n) = O(f(n))$.

Examples:

- ▶ $2n^2 + 27 = O(n^2)$: set $n_0 = 6$ and $c = 3$
- ▶ $2n^2 + 27 = O(n^3)$: same values, or $n_0 = 4$ and $c = 1$
- ▶ $n^3 + 2000n^2 + 2000n = O(n^3)$: set $n_0 = 10000$ and $c = 2$

$O(\cdot)$

Definition

$g(n) \in O(f(n))$ if there exist constants $c, n_0 > 0$ such that $g(n) \leq c \cdot f(n)$ for all $n > n_0$.

Technically $O(f(n))$ is a set.

Abuse notation: “ $g(n)$ is $O(f(n))$ ” or $g(n) = O(f(n))$.

Examples:

- ▶ $2n^2 + 27 = O(n^2)$: set $n_0 = 6$ and $c = 3$
- ▶ $2n^2 + 27 = O(n^3)$: same values, or $n_0 = 4$ and $c = 1$
- ▶ $n^3 + 2000n^2 + 2000n = O(n^3)$: set $n_0 = 10000$ and $c = 2$

About *functions* not algorithms!


Expresses an *upper* bound

Example

Definition

$g(n) \in O(f(n))$ if there exist constants $c, n_0 > 0$ such that $g(n) \leq c \cdot f(n)$ for all $n > n_0$.

Example: $2n^2 + 27 = O(n^2)$



Example

Definition

$g(n) \in O(f(n))$ if there exist constants $c, n_0 > 0$ such that $g(n) \leq c \cdot f(n)$ for all $n > n_0$.

Theorem

$$2n^2 + 27 = O(n^2)$$

Example

Definition

$g(n) \in O(f(n))$ if there exist constants $c, n_0 > 0$ such that $g(n) \leq c \cdot f(n)$ for all $n > n_0$.

Theorem

$$2n^2 + 27 = O(n^2)$$

Proof.

Set $c = 3$. Suppose $2n^2 + 27 > cn^2 = 3n^2$

Example

Definition

$g(n) \in O(f(n))$ if there exist constants $c, n_0 > 0$ such that $g(n) \leq c \cdot f(n)$ for all $n > n_0$.

Theorem

$$2n^2 + 27 = O(n^2)$$

Proof.

Set $c = 3$. Suppose $2n^2 + 27 > cn^2 = 3n^2$
 $\implies n^2 < 27$

Example

Definition

$g(n) \in O(f(n))$ if there exist constants $c, n_0 > 0$ such that $g(n) \leq c \cdot f(n)$ for all $n > n_0$.

Theorem

$$2n^2 + 27 = O(n^2)$$

Proof.

Set $c = 3$. Suppose $2n^2 + 27 > cn^2 = 3n^2$
 $\implies n^2 < 27 \implies n < 6$

Example

Definition

$g(n) \in O(f(n))$ if there exist constants $c, n_0 > 0$ such that $g(n) \leq c \cdot f(n)$ for all $n > n_0$.

Theorem

$$2n^2 + 27 = O(n^2)$$

Proof.

Set $c = 3$. Suppose $2n^2 + 27 > cn^2 = 3n^2$

$$\implies n^2 < 27 \implies n < 6$$

$$\implies 2n^2 + 27 \leq 3n^2 \text{ for all } n \geq 6.$$

Example

Definition

$g(n) \in O(f(n))$ if there exist constants $c, n_0 > 0$ such that $g(n) \leq c \cdot f(n)$ for all $n > n_0$.

Theorem

$$2n^2 + 27 = O(n^2)$$

Proof.

Set $c = 3$. Suppose $2n^2 + 27 > cn^2 = 3n^2$

$$\implies n^2 < 27 \implies n < 6$$

$$\implies 2n^2 + 27 \leq 3n^2 \text{ for all } n \geq 6.$$

Set $n_0 = 6$. Then $2n^2 + 27 \leq cn^2$ for all $n > n_0$. □

Example

Definition

$g(n) \in O(f(n))$ if there exist constants $c, n_0 > 0$ such that $g(n) \leq c \cdot f(n)$ for all $n > n_0$.

Theorem

$$2n^2 + 27 = O(n^2)$$

Proof.

Set $c = 3$. Suppose $2n^2 + 27 > cn^2 = 3n^2$

$$\implies n^2 < 27 \implies n < 6$$

$$\implies 2n^2 + 27 \leq 3n^2 \text{ for all } n \geq 6.$$

Set $n_0 = 6$. Then $2n^2 + 27 \leq cn^2$ for all $n > n_0$. □

Many other ways to prove this!

$\Omega(\cdot)$

Counterpart to $O(\cdot)$: *lower* bound rather than upper bound.

Definition

$g(n) \in \Omega(f(n))$ if there exist constants $c, n_0 > 0$ such that $g(n) \geq c \cdot f(n)$ for all $n > n_0$.

$\Omega(\cdot)$

Counterpart to $O(\cdot)$: *lower* bound rather than upper bound.

Definition

$g(n) \in \Omega(f(n))$ if there exist constants $c, n_0 > 0$ such that $g(n) \geq c \cdot f(n)$ for all $n > n_0$.

Examples:

- ▶ $2n^2 + 27 = \Omega(n^2)$: set $n_0 = 1$ and $c = 1$
- ▶ $2n^2 + 27 = \Omega(n)$: set $n_0 = 1$ and $c = 1$
- ▶ $\frac{1}{100}n^3 - 1000n^2 = \Omega(n^3)$: set $n_0 = 1000000$ and $c = 1/1000$

$\Theta(\cdot)$

Combination of $O(\cdot)$ and $\Omega(\cdot)$.

Definition

$g(n) \in \Theta(f(n))$ if $g(n) \in O(f(n))$ and $g(n) \in \Omega(f(n))$.

Note: constants n_0, c can be different in the proofs for $O(f(n))$ and $\Omega(f(n))$

$\Theta(\cdot)$

Combination of $O(\cdot)$ and $\Omega(\cdot)$.

Definition

$g(n) \in \Theta(f(n))$ if $g(n) \in O(f(n))$ and $g(n) \in \Omega(f(n))$.

Note: constants n_0, c can be different in the proofs for $O(f(n))$ and $\Omega(f(n))$

Equivalent:

Definition

$g(n) \in \Theta(f(n))$ if there are constants $c_1, c_2, n_0 > 0$ such that $c_1 f(n) \leq g(n) \leq c_2 f(n)$ for all $n > n_0$.

Both lower bound and upper bound, so asymptotic equality.


$\Theta(\cdot)$

Definition

$g(n) \in \Theta(f(n))$ if $g(n) \in O(f(n))$ and $g(n) \in \Omega(f(n))$.

Note: constants n_0, c can be different in the proofs for $O(f(n))$ and $\Omega(f(n))$

Example: $\frac{1}{100}n^3 - 1000n^2 = \Omega(n^3)$: set $n_0 = 1000000$ and $c = 1/1000$



Both lower bound and upper bound, so asymptotic equality.

Little notation

Strict versions of O and Ω :

Definition

$g(n) \in o(f(n))$ if for every constant $c > 0$ there exists a constant $n_0 > 0$ such that $g(n) < c \cdot f(n)$ for all $n > n_0$.

Definition

$g(n) \in \omega(f(n))$ if for every constant $c > 0$ there exists a constant $n_0 > 0$ such that $g(n) > c \cdot f(n)$ for all $n > n_0$.

Examples:

- ▶ $2n^2 + 27 = o(n^2 \log n)$
- ▶ $2n^2 + 27 = \omega(n)$

Recurrence Relations

Sorting

Many algorithms recursive so running time naturally a recurrence relation (Karatsuba, Strassen).

Sorting

Many algorithms recursive so running time naturally a recurrence relation (Karatsuba, Strassen).

Sorting: Let $T(n)$ denote (worst-case) running time on an array of size n .

Sorting

Many algorithms recursive so running time naturally a recurrence relation (Karatsuba, Strassen).

Sorting: Let $T(n)$ denote (worst-case) running time on an array of size n .

- ▶ Selection Sort

Sorting

Many algorithms recursive so running time naturally a recurrence relation (Karatsuba, Strassen).

Sorting: Let $T(n)$ denote (worst-case) running time on an array of size n .

- ▶ Selection Sort
 - ▶ Find smallest unsorted element, put it just after sorted elements. Repeat.

Sorting

Many algorithms recursive so running time naturally a recurrence relation (Karatsuba, Strassen).

Sorting: Let $T(n)$ denote (worst-case) running time on an array of size n .

- ▶ Selection Sort
 - ▶ Find smallest unsorted element, put it just after sorted elements. Repeat.
 - ▶ Running time: Takes $O(n)$ time to find smallest unsorted element, decreases remaining unsorted by 1.
 $\implies T(n) = T(n-1) + cn$

Sorting

Many algorithms recursive so running time naturally a recurrence relation (Karatsuba, Strassen).

Sorting: Let $T(n)$ denote (worst-case) running time on an array of size n .

- ▶ Selection Sort
 - ▶ Find smallest unsorted element, put it just after sorted elements. Repeat.
 - ▶ Running time: Takes $O(n)$ time to find smallest unsorted element, decreases remaining unsorted by 1.
 $\implies T(n) = T(n-1) + cn$
- ▶ Mergesort

Sorting

Many algorithms recursive so running time naturally a recurrence relation (Karatsuba, Strassen).

Sorting: Let $T(n)$ denote (worst-case) running time on an array of size n .

- ▶ Selection Sort

- ▶ Find smallest unsorted element, put it just after sorted elements. Repeat.
- ▶ Running time: Takes $O(n)$ time to find smallest unsorted element, decreases remaining unsorted by 1.
 $\implies T(n) = T(n-1) + cn$

- ▶ Mergesort

- ▶ Split array into left and right halves. Recursively sort each half, then merge.

Sorting

Many algorithms recursive so running time naturally a recurrence relation (Karatsuba, Strassen).

Sorting: Let $T(n)$ denote (worst-case) running time on an array of size n .

- ▶ Selection Sort

- ▶ Find smallest unsorted element, put it just after sorted elements. Repeat.
- ▶ Running time: Takes $O(n)$ time to find smallest unsorted element, decreases remaining unsorted by 1.

$$\implies T(n) = T(n-1) + cn$$

- ▶ Mergesort

- ▶ Split array into left and right halves. Recursively sort each half, then merge.
- ▶ Running time: Merging takes $O(n)$ time. Two recursive calls on half the size.

$$\implies T(n) = T(n/2) + T(n/2) + cn = 2T(n/2) + cn$$

Sorting

Many algorithms recursive so running time naturally a recurrence relation (Karatsuba, Strassen).

Sorting: Let $T(n)$ denote (worst-case) running time on an array of size n .

- ▶ Selection Sort

- ▶ Find smallest unsorted element, put it just after sorted elements. Repeat.
- ▶ Running time: Takes $O(n)$ time to find smallest unsorted element, decreases remaining unsorted by 1.

$$\implies T(n) = T(n-1) + cn$$

- ▶ Mergesort

- ▶ Split array into left and right halves. Recursively sort each half, then merge.
- ▶ Running time: Merging takes $O(n)$ time. Two recursive calls on half the size.

$$\implies T(n) = T(n/2) + T(n/2) + cn = 2T(n/2) + cn$$

Also need base case. For algorithms, constant size input takes constant time.

$$\implies T(n) \leq c \text{ for all } n \leq n_0, \text{ for some constants } n_0, c > 0.$$

Guess and Check

$$T(n) = 3T(n/3) + n$$

$$T(1) = 1$$

Guess and Check

$$T(n) = 3T(n/3) + n$$

$$T(1) = 1$$

Guess: $T(n) \leq cn$.

Guess and Check

$$T(n) = 3T(n/3) + n$$

$$T(1) = 1$$

Guess: $T(n) \leq cn$.

Check: assume true for $n' < n$, prove true for n (induction).

Guess and Check

$$T(n) = 3T(n/3) + n$$

$$T(1) = 1$$

Guess: $T(n) \leq cn$.

Check: assume true for $n' < n$, prove true for n (induction).

$$T(n) = 3T(n/3) + n \leq 3cn/3 + n = (c + 1)n$$

Guess and Check

$$T(n) = 3T(n/3) + n$$

$$T(1) = 1$$

Guess: $T(n) \leq cn$.

Check: assume true for $n' < n$, prove true for n (induction).

$$T(n) = 3T(n/3) + n \leq 3cn/3 + n = (c + 1)n$$

Failure! Wanted $T(n) \leq cn$, got $T(n) \leq (c + 1)n$. Guess was wrong.

Guess and Check

$$\begin{array}{r} 13 \\ \hline 3T(3) + 9 \end{array} \quad + 5 = 21$$

$$\underline{T(n) = 3T(n/3) + n}$$

$$\begin{aligned} T(1) &= 1 \\ T(3) &= 6 \\ T(9) &= 21 \end{aligned}$$

Guess: $T(n) \leq cn$.

Check: assume true for $n' < n$, prove true for n (induction).

$$T(n) = 3T(n/3) + n \leq 3cn/3 + n = (c+1)n$$

Failure! Wanted $T(n) \leq cn$, got $T(n) \leq (c+1)n$. Guess was wrong.

Better guess? What goes up by 1 when n goes up by a factor of 3?

Guess and Check

$$T(n) = 3T(n/3) + n$$

$$T(1) = 1$$

Guess: $T(n) \leq cn$.

Check: assume true for $n' < n$, prove true for n (induction).

$$T(n) = 3T(n/3) + n \leq 3cn/3 + n = (c + 1)n$$

Failure! Wanted $T(n) \leq cn$, got $T(n) \leq (c + 1)n$. Guess was wrong.

Better guess? What goes up by 1 when n goes up by a factor of 3? $\log_3 n$

Guess and Check

$$T(n) = 3T(n/3) + n$$

$$T(1) = 1$$

Guess: $T(n) \leq cn$.

Check: assume true for $n' < n$, prove true for n (induction).

$$T(n) = 3T(n/3) + n \leq 3cn/3 + n = (c + 1)n$$

Failure! Wanted $T(n) \leq cn$, got $T(n) \leq (c + 1)n$. Guess was wrong.

Better guess? What goes up by 1 when n goes up by a factor of 3? $\log_3 n$

Guess: $T(n) \leq n \log_3(3n)$

Guess and Check

$$T(n) = 3T(n/3) + n$$

$$T(1) = 1$$

Guess: $T(n) \leq cn$.

Check: assume true for $n' < n$, prove true for n (induction).

$$T(n) = 3T(n/3) + n \leq 3cn/3 + n = (c + 1)n$$

Failure! Wanted $T(n) \leq cn$, got $T(n) \leq (c + 1)n$. Guess was wrong.

Better guess? What goes up by 1 when n goes up by a factor of 3? $\log_3 n$

Guess: $T(n) \leq n \log_3(3n)$

Check: assume true for $n' < n$, prove true for n (induction).

$$\begin{aligned} T(n) &= 3T(n/3) + n \leq 3(n/3) \log_3(3n/3) + n = n \log_3(n) + n \\ &= n(\log_3(n) + \log_3 3) = n \log_3(3n). \end{aligned}$$

Unrolling

Example: selection sort. $T(n) = T(n - 1) + cn$

Idea: “unroll” the recurrence.

Unrolling

Example: selection sort. $T(n) = T(n - 1) + cn$

Idea: “unroll” the recurrence.

$$T(n) = cn + T(n - 1)$$

Unrolling

Example: selection sort. $T(n) = T(n - 1) + cn$

Idea: “unroll” the recurrence.

$$\begin{aligned} T(n) &= cn + T(n - 1) \\ &= cn + c(n - 1) + T(n - 2) \end{aligned}$$

Unrolling

Example: selection sort. $T(n) = T(n - 1) + cn$

Idea: “unroll” the recurrence.

$$\begin{aligned}T(n) &= cn + T(n - 1) \\&= cn + c(n - 1) + T(n - 2) \\&= cn + c(n - 1) + c(n - 2) + T(n - 3)\end{aligned}$$

Unrolling

Example: selection sort. $T(n) = T(n-1) + cn$

Idea: “unroll” the recurrence.

$$\begin{aligned}T(n) &= cn + T(n-1) \\&= cn + c(n-1) + T(n-2) \\&= cn + c(n-1) + c(n-2) + T(n-3) \\&\vdots\end{aligned}$$

Unrolling

Example: selection sort. $T(n) = T(n-1) + cn$

Idea: “unroll” the recurrence.

$$\begin{aligned}T(n) &= cn + T(n-1) \\&= cn + c(n-1) + T(n-2) \\&= cn + c(n-1) + c(n-2) + T(n-3) \\&\quad \vdots \\&= cn + c(n-1) + c(n-2) + \cdots + c\end{aligned}$$

Unrolling

Example: selection sort. $T(n) = T(n-1) + cn$

Idea: “unroll” the recurrence.

$$\begin{aligned}T(n) &= cn + T(n-1) \\&= cn + c(n-1) + T(n-2) \\&= cn + c(n-1) + c(n-2) + T(n-3) \\&\quad \vdots \\&= cn + c(n-1) + c(n-2) + \cdots + c\end{aligned}$$

n terms, each of which at most $cn \implies T(n) \leq cn^2 = O(n^2)$

Unrolling

Example: selection sort. $T(n) = T(n-1) + cn$

Idea: “unroll” the recurrence.

$$\begin{aligned}T(n) &= cn + T(n-1) \\&= cn + c(n-1) + T(n-2) \\&= cn + c(n-1) + c(n-2) + T(n-3) \\&\quad \vdots \\&= cn + c(n-1) + c(n-2) + \cdots + c\end{aligned}$$

n terms, each of which at most $cn \implies T(n) \leq cn^2 = O(n^2)$

At least $n/2$ terms which are at least $cn/2 \implies T(n) \geq c\frac{n^2}{4} = \Omega(n^2)$

Unrolling

Example: selection sort. $T(n) = T(n-1) + cn$

Idea: “unroll” the recurrence.

$$\begin{aligned}T(n) &= cn + T(n-1) \\&= cn + c(n-1) + T(n-2) \\&= cn + c(n-1) + c(n-2) + T(n-3) \\&\quad \vdots \\&= cn + c(n-1) + c(n-2) + \cdots + c\end{aligned}$$

n terms, each of which at most $cn \implies T(n) \leq cn^2 = O(n^2)$

At least $n/2$ terms which are at least $cn/2 \implies T(n) \geq c\frac{n^2}{4} = \Omega(n^2)$
 $\implies T(n) = \Theta(n^2)$.

Recursion Tree: Mergesort


Generalizes unrolling: draw out full tree of “recursive calls”.

Mergesort: $T(n) = 2T(n/2) + cn$.

Recursion Tree: Mergesort

Generalizes unrolling: draw out full tree of “recursive calls”.


Mergesort: $T(n) = 2T(n/2) + cn$.



Recursion Tree: Mergesort

Generalizes unrolling: draw out full tree of “recursive calls”.

Mergesort: $T(n) = 2T(n/2) + cn$.




levels:

Recursion Tree: Mergesort

Generalizes unrolling: draw out full tree of “recursive calls”.

Mergesort: $T(n) = 2T(n/2) + cn$.




levels: $\log_2 n$

Recursion Tree: Mergesort

Generalizes unrolling: draw out full tree of “recursive calls”.

Mergesort: $T(n) = 2T(n/2) + cn$.




levels: $\log_2 n$

Contribution of level i :

Recursion Tree: Mergesort

Generalizes unrolling: draw out full tree of “recursive calls”.

Mergesort: $T(n) = 2T(n/2) + cn$.




levels: $\log_2 n$

Contribution of level i : $2^{i-1} cn/2^{i-1} = cn$

Recursion Tree: Mergesort

Generalizes unrolling: draw out full tree of “recursive calls”.

Mergesort: $T(n) = 2T(n/2) + cn$.



levels: $\log_2 n$

Contribution of level i : $2^{i-1} cn/2^{i-1} = cn$


$\implies T(n) = \Theta(n \log n)$

Recursion Tree: Strassen

$$T(n) = 7T(n/2) + cn^2$$


Recursion Tree: Strassen

$$T(n) = 7T(n/2) + cn^2$$



Recursion Tree: Strassen


$$T(n) = 7T(n/2) + cn^2$$



$$\text{Level } i: 7^{i-1}c(n/2^{i-1})^2 = (7/4)^{i-1}cn^2$$

Recursion Tree: Strassen

$$T(n) = 7T(n/2) + cn^2$$




Level i : $7^{i-1}c(n/2^{i-1})^2 = (7/4)^{i-1}cn^2$

Total:

$$T(n) = \sum_{i=1}^{\log n+1} \left(\frac{7}{4}\right)^{i-1} cn^2 = cn^2 \sum_{i=1}^{\log n+1} \left(\frac{7}{4}\right)^{i-1}$$

Recursion Tree: Strassen

$$T(n) = 7T(n/2) + cn^2$$



$$\text{Level } i: 7^{i-1}c(n/2^{i-1})^2 = (7/4)^{i-1}cn^2$$

Total:

$$T(n) = \sum_{i=1}^{\log n+1} \left(\frac{7}{4}\right)^{i-1} cn^2 = cn^2 \sum_{i=1}^{\log n+1} \left(\frac{7}{4}\right)^{i-1}$$

$$\begin{aligned} \implies T(n) &= O(n^2(7/4)^{\log_2 n}) = O(n^2 2^{\log_2(7/4) \cdot \log_2 n}) = O(n^2 n^{\log_2(7/4)}) = O(n^2 n^{\log_2 7 - 2}) \\ &= O(n^{\log_2 7}) \end{aligned}$$

Master Theorem

$$T(n) = aT(n/b) + cn^k \qquad T(1) = c$$


a, b, c, k constants with **$a \geq 1$** , **$b > 1$** , **$c > 0$** , and **$k \geq 0$**

Master Theorem

$$T(n) = aT(n/b) + cn^k$$

$$T(1) = c$$

a, b, c, k constants with $a \geq 1$, $b > 1$, $c > 0$, and $k \geq 0$




Master Theorem

$$T(n) = aT(n/b) + cn^k$$

$$T(1) = c$$

a, b, c, k constants with $a \geq 1$, $b > 1$, $c > 0$, and $k \geq 0$




levels: $\log_b n + 1$

Master Theorem

$$T(n) = aT(n/b) + cn^k$$

$$T(1) = c$$

a, b, c, k constants with $a \geq 1$, $b > 1$, $c > 0$, and $k \geq 0$



levels: $\log_b n + 1$

Level i : $a^{i-1}c(n/b^{i-1})^k = cn^k(a/b^k)^{i-1}$

Master Theorem II

Let $\alpha = (a/b^k)$

$$\implies T(n) = cn^k \sum_{i=1}^{\log_b n+1} (a/b^k)^{i-1} = cn^k \sum_{i=1}^{\log_b n+1} \alpha^{i-1}$$

Master Theorem II

Let $\alpha = (a/b^k)$

$$\implies T(n) = cn^k \sum_{i=1}^{\log_b n+1} (a/b^k)^{i-1} = cn^k \sum_{i=1}^{\log_b n+1} \alpha^{i-1}$$

- ▶ Case 1: $\alpha = 1$. All levels the same. $T(n) = cn^k \sum_{i=1}^{\log_b n+1} 1 = \Theta(n^k \log n)$

Master Theorem II

Let $\alpha = (a/b^k)$

$$\implies T(n) = cn^k \sum_{i=1}^{\log_b n+1} (a/b^k)^{i-1} = cn^k \sum_{i=1}^{\log_b n+1} \alpha^{i-1}$$

- ▶ Case 1: $\alpha = 1$. All levels the same. $T(n) = cn^k \sum_{i=1}^{\log_b n+1} 1 = \Theta(n^k \log n)$
- ▶ Case 2: $\alpha < 1$. Dominated by top level.

Master Theorem II

Let $\alpha = (a/b^k)$

$$\implies T(n) = cn^k \sum_{i=1}^{\log_b n+1} (a/b^k)^{i-1} = cn^k \sum_{i=1}^{\log_b n+1} \alpha^{i-1}$$

▶ Case 1: $\alpha = 1$. All levels the same. $T(n) = cn^k \sum_{i=1}^{\log_b n+1} 1 = \Theta(n^k \log n)$

▶ Case 2: $\alpha < 1$. Dominated by top level.

$$\implies \sum_{i=1}^{\log_b n+1} \alpha^{i-1} \leq \sum_{i=1}^{\infty} \alpha^{i-1} = \frac{1}{1-\alpha}.$$

$$\implies T(n) = O(n^k)$$

Master Theorem II

Let $\alpha = (a/b^k)$

$$\implies T(n) = cn^k \sum_{i=1}^{\log_b n+1} (a/b^k)^{i-1} = cn^k \sum_{i=1}^{\log_b n+1} \alpha^{i-1}$$

- ▶ Case 1: $\alpha = 1$. All levels the same. $T(n) = cn^k \sum_{i=1}^{\log_b n+1} 1 = \Theta(n^k \log n)$
- ▶ Case 2: $\alpha < 1$. Dominated by top level.

$$\implies \sum_{i=1}^{\log_b n+1} \alpha^{i-1} \leq \sum_{i=1}^{\infty} \alpha^{i-1} = \frac{1}{1-\alpha}.$$

$$\implies T(n) = O(n^k)$$

$$T(n) \geq cn^k \implies T(n) = \Omega(n^k)$$

Master Theorem II

Let $\alpha = (a/b^k)$

$$\implies T(n) = cn^k \sum_{i=1}^{\log_b n+1} (a/b^k)^{i-1} = cn^k \sum_{i=1}^{\log_b n+1} \alpha^{i-1}$$

- ▶ Case 1: $\alpha = 1$. All levels the same. $T(n) = cn^k \sum_{i=1}^{\log_b n+1} 1 = \Theta(n^k \log n)$
- ▶ Case 2: $\alpha < 1$. Dominated by top level.

$$\implies \sum_{i=1}^{\log_b n+1} \alpha^{i-1} \leq \sum_{i=1}^{\infty} \alpha^{i-1} = \frac{1}{1-\alpha}.$$

$$\implies T(n) = O(n^k)$$

$$T(n) \geq cn^k \implies T(n) = \Omega(n^k) \implies T(n) = \Theta(n^k)$$

Master Theorem II

Let $\alpha = (a/b^k)$

$$\implies T(n) = cn^k \sum_{i=1}^{\log_b n+1} (a/b^k)^{i-1} = cn^k \sum_{i=1}^{\log_b n+1} \alpha^{i-1}$$

▶ Case 1: $\alpha = 1$. All levels the same. $T(n) = cn^k \sum_{i=1}^{\log_b n+1} 1 = \Theta(n^k \log n)$

▶ Case 2: $\alpha < 1$. Dominated by top level.

$$\implies \sum_{i=1}^{\log_b n+1} \alpha^{i-1} \leq \sum_{i=1}^{\infty} \alpha^{i-1} = \frac{1}{1-\alpha}.$$

$$\implies T(n) = O(n^k)$$

$$T(n) \geq cn^k \implies T(n) = \Omega(n^k) \implies T(n) = \Theta(n^k)$$

▶ Case 3: $\alpha > 1$. Dominated by bottom level

Master Theorem II

Let $\alpha = (a/b^k)$

$$\implies T(n) = cn^k \sum_{i=1}^{\log_b n+1} (a/b^k)^{i-1} = cn^k \sum_{i=1}^{\log_b n+1} \alpha^{i-1}$$

► Case 1: $\alpha = 1$. All levels the same. $T(n) = cn^k \sum_{i=1}^{\log_b n+1} 1 = \Theta(n^k \log n)$

► Case 2: $\alpha < 1$. Dominated by top level.

$$\implies \sum_{i=1}^{\log_b n+1} \alpha^{i-1} \leq \sum_{i=1}^{\infty} \alpha^{i-1} = \frac{1}{1-\alpha}.$$

$$\implies T(n) = O(n^k)$$

$$T(n) \geq cn^k \implies T(n) = \Omega(n^k) \implies T(n) = \Theta(n^k)$$

► Case 3: $\alpha > 1$. Dominated by bottom level

$$\begin{aligned} \implies \sum_{i=1}^{\log_b n+1} \alpha^{i-1} &= \alpha^{\log_b n} \sum_{i=1}^{\log_b n+1} \left(\frac{1}{\alpha}\right)^{i-1} \leq \alpha^{\log_b n} \frac{1}{1 - (1/\alpha)} \\ &= O(\alpha^{\log_b n}) \end{aligned}$$

Master Theorem II

Let $\alpha = (a/b^k)$

$$\implies T(n) = cn^k \sum_{i=1}^{\log_b n+1} (a/b^k)^{i-1} = cn^k \sum_{i=1}^{\log_b n+1} \alpha^{i-1}$$

► Case 1: $\alpha = 1$. All levels the same. $T(n) = cn^k \sum_{i=1}^{\log_b n+1} 1 = \Theta(n^k \log n)$

► Case 2: $\alpha < 1$. Dominated by top level.

$$\implies \sum_{i=1}^{\log_b n+1} \alpha^{i-1} \leq \sum_{i=1}^{\infty} \alpha^{i-1} = \frac{1}{1-\alpha}.$$

$$\implies T(n) = O(n^k)$$

$$T(n) \geq cn^k \implies T(n) = \Omega(n^k) \implies T(n) = \Theta(n^k)$$

► Case 3: $\alpha > 1$. Dominated by bottom level

$$\implies \sum_{i=1}^{\log_b n+1} \alpha^{i-1} = \alpha^{\log_b n} \sum_{i=1}^{\log_b n+1} \left(\frac{1}{\alpha}\right)^{i-1} \leq \alpha^{\log_b n} \frac{1}{1 - (1/\alpha)}$$

$$= O(\alpha^{\log_b n})$$

$$\begin{aligned} \implies T(n) &= \Theta(n^k \alpha^{\log_b n}) = \Theta(n^k (a/b^k)^{\log_b n}) = \Theta(a^{\log_b n}) \\ &= \Theta(n^{\log_b a}) \end{aligned}$$

Master Theorem III

Theorem (“Master Theorem”)

The recurrence

$$T(n) = aT(n/b) + cn^k$$

$$T(1) = c$$

where a, b, c , and k are constants with $a \geq 1$, $b > 1$, $c > 0$, and $k \geq 0$, is equal to

$$T(n) = \Theta(n^k) \text{ if } a < b^k,$$

$$T(n) = \Theta(n^k \log n) \text{ if } a = b^k,$$

$$T(n) = \Theta(n^{\log_b a}) \text{ if } a > b^k.$$