

High temperatures predict gut microbiome composition, metabolic capacity, and immune interactions in a wild primate species

Running title: Heat shapes primate gut microbiomes

Jordan M. Lucore^{1,2*}, Marcela E. Benítez^{2,3}, Amy F. White^{2,4}, Juan Carlos Ordoñez², Jacinta C. Beehner^{1,2,5}, Thore J. Bergman^{2,5,6}, Maria Luisa Savo Sardaro^{7,8}, Katherine R. Amato^{7†}, Andrew J. Marshall^{1,6,9, 10, 11†}

¹Department of Anthropology, University of Michigan, Ann Arbor, MI, United States,

²Capuchinos de Taboga Research Project, Taboga Forest Reserve, Guanacaste, Costa Rica,

³Department of Anthropology, Emory University, Atlanta, GA, United States, ⁴Department of

Anthropology, Durham University, Durham, UK, ⁵Department of Psychology, University of

Michigan, Ann Arbor, MI, United States, ⁶Department of Ecology and Evolutionary Biology,

University of Michigan, Ann Arbor, MI, United States, ⁷Department of Anthropology,

Northwestern University, Evanston, Illinois, ⁸Department for the Promotion of Human Sciences

and Quality of Life, San Raffaele University, 00166 Rome, Italy, ⁹Program in the Environment,

University of Michigan, Ann Arbor, MI, United States, ¹⁰School for Environment and

Sustainability, University of Michigan, Ann Arbor, MI, United States, ¹¹Program in Computing

for the Arts and Sciences, University of Michigan, Ann Arbor, MI, United States

* Corresponding author jlucore@umich.edu

† These authors contributed equally to this work.

Abstract: Rising temperatures due to climate change are impacting the function of critical physiological processes in wild animals. Heat stress may compromise the gut microbiome's ability to support essential host functions, highlighting an important pathway by which climate change could negatively affect numerous species. However, the relationship between the microbiome and heat stress has been studied primarily in experimental contexts and remains poorly understood in ecologically relevant environments. Here, we examine how natural thermal variation influences gut microbiome taxonomic composition and functional capacity in a wild population of white-face capuchin monkeys. We find that high temperatures are associated with changes in microbiome community-level diversity, taxonomic composition, and metabolic capacity consistent with physiological responses to heat stress. We also show that the microbiome's response to high temperatures is linked to host immunity and may affect microbiome-immune interactions. These results suggest that rising temperatures may compromise essential microbiome functions and highlight a potential mechanism by which climate change could negatively impact species through their microbiomes.

Keywords: gut microbiome, neopterin, heat stress, endotherm, climate change, wild primate

Introduction:

Rising temperatures due to climate change are predicted to push species beyond their evolved thermal tolerances and challenge them to adapt to hotter climates (1, 2). Efforts to evaluate the physiological consequences of heat stress in wild animals have shown that high temperatures can impair aerobic capacity, reducing oxygen availability to essential systems and compromising

function (3, 4). Much of this research has focused on ectotherms, whose lack of physiological thermoregulation makes them particularly vulnerable to high temperatures (4–6). However, mounting evidence suggests that endotherms are also physiologically susceptible to heat (7, 8), and evolved thermoregulatory mechanisms may be increasingly insufficient to cope with prolonged periods of extreme heat. Recently, research has turned to assessing the role of host-associated microbiomes in shaping how species respond to heat stress. Animals have evolved complex relationships with the microbial communities (i.e., microbiomes) that live on and inside them to the extent that hosts rely on their microbiomes to support diverse aspects of their physiology. Accordingly, microbiomes may play a key role in determining the thermal tolerances of their host species (9–12) because heat stress can deplete microbiome communities, and is associated with negative phenotypic, survivorship, and fitness outcomes (13, 14). While this research highlights that heat is an important determinant of host-microbiome relationships, it has largely been conducted in laboratory settings and offers limited insight into how these dynamics function in ecologically relevant contexts.

The gastrointestinal tract is home to a particularly dense microbial community, the gut microbiome, that plays a crucial role in regulating host responses to environmental change (15), making it an ideal system to explore how species may respond to rising temperatures. High temperatures have been shown to alter community composition (16) and are linked to changes in host digestion, metabolism, and immune function (17–20), but these relationships have yet to be demonstrated in wild populations. A common consequence of heat stress is redox imbalance—the accumulation of reactive oxygen species (ROS) that can damage cells (21, 22), compromise the epithelial barrier (23), and trigger immune activation (24). Gut microbes typically help maintain redox balance (25–27), but heat-induced changes in community composition have been linked to reduced antioxidant metabolism (28, 29) and increased epithelial damage (30). These redox shifts may disrupt oxygen gradients, favoring facultative anaerobes over obligate anaerobes. This is supported across a variety of vertebrate species where high temperatures are consistently associated with declines in Firmicutes and Bacteroidota (20, 31, 32), phyla composed primarily of obligate anaerobes lacking robust ROS defenses (33, 34), and increases in Proteobacteria (16), a phylum dominated by facultative anaerobes that thrive in higher-oxygen conditions (35, 36).

The loss of obligate anaerobes during heat stress could affect the relationship between the gut microbiome and the immune system. Many obligate anaerobes ferment dietary fiber and produce short-chain fatty acids (SCFAs) (37–39), which support anti-inflammatory mechanisms that promote immune tolerance of the microbiome (40–43). A loss of these obligate anaerobes could result in reduced SCFAs and impact the immune system. Experimental studies evaluating the effects of acute heat stress in livestock show that exposure to high temperatures is associated with reduced SCFA concentrations, (19) changes in the levels of circulating cytokines, and altered gut microbiome composition (17, 44), suggesting that heat-driven compositional shifts may disrupt microbiome-immune interactions and increase the risk of gut inflammation. These dynamics are likely to play an essential role in shaping animal responses to changing climates. However, interactions between heat stress and the microbiome have been studied almost exclusively in experimental contexts that examine the effect of acute periods of moderate heat stress [e.g., ~8–10 hours of exposure to temperatures ~32–34 °C, (45, 46)] and have focused either on ectotherms [e.g., amphibians (18, 47), fish (28), reptiles (32, 48)] or on domestic livestock [e.g., pigs (30), cows (17, 19, 44), poultry (29, 49)]. Therefore, we have little

understanding of how this system functions in wild populations in response to natural thermal variation.

To address this knowledge gap, here, we assess how temperature predicts gut microbiome composition and metabolic function in wild capuchin monkeys, *Cebus imitator*, in Costa Rica. We focus on evaluating the response to cumulative thermal load by averaging ambient temperature two weeks before sample collection, rather than attempting to measure short-term responses that are difficult to isolate in wild populations. Importantly, this predictor is also associated with thermal responses in immune performance in this population, supporting its relevance for studying the consequences of high temperatures on physiology (8). We also evaluate the interplay between temperature, the gut microbiome, and host immunity. Specifically, we measure host immunity using urinary neopterin, a byproduct of macrophage activity during the cellular immune response (50). Macrophages play a central role in host containment of the gut microbiome (51), and microbial metabolites, including SCFAs and saturated fatty acids, regulate macrophage activity to maintain host tolerance and prevent chronic inflammatory disorders (40, 52). Therefore, neopterin is a valuable tool for examining microbiome-immune homeostasis.

We analyzed 11 months of paired fecal and urine samples from 36 individually identified white-faced capuchin monkeys living in tropical dry forest fragments in northwestern Costa Rica at the Taboga Forest Reserve. Tropical dry forests are highly seasonal and defined by marked changes in ecology between the dry and wet seasons (53). During the dry season, lack of rainfall leads to food and water scarcity that has physiological consequences for capuchins (54) and can influence gut microbiome composition (55). Because the dry season is also characterized by high temperatures, this can make it difficult to disentangle the effects of heat from those of resource scarcity. However, our study population differs from most others in that we find no evidence of seasonal differences in physiological stress responses (56) or immunity (8), suggesting that these monkeys are not exposed to the same seasonal stressors. This is likely because our study site is highly disturbed and bordered by secondary forest and agriculture, where capuchins can exploit anthropogenic food and water sources even during the dry season (57). Consequently, studying this population provides a rare opportunity to isolate temperature-driven changes in the microbiome from those driven by broader changes in seasonal ecology. We characterized microbiome composition using 16S rRNA gene amplicon sequencing (n = 277), and characterized microbiome functional capacity using shotgun metagenomic sequencing (n = 71). Our results highlight the crucial role of temperature in shaping essential gut microbiome functions in a wild endotherm species and demonstrate how temperature is related to microbiome-immune interactions.

Methods:

Site information and data collection

We collected data from a wild population of *C. imitator* living in the Taboga Forest reserve in Guanacaste, Northwestern Costa Rica. The Taboga Forest Reserve is a 516-ha tropical dry forest fragment and forms an important piece of a fragmented biological corridor connecting the Tempisque River Basin to the Guanacaste Mountains (57). The Capuchinos de Taboga Research project has collected daily behavioral and endocrine data on the three groups of white-faced capuchins since 2018. All individuals are individually identified and habituated to human presence. Individual age is calculated by birth date for individuals younger than 7 years of age; for individuals older than 7 years, age was estimated based on body condition by a field team

member (J.C.O.) with more than two decades of experience observing wild white-faced capuchins. A team of two or more trained field assistants collected data between 5:00 AM and 6:30 PM, including detailed observations of feeding and foraging behavior using scan sampling. Scans occurred every 30 minutes for 10 minutes, where feeding and foraging observations of individually identified monkeys were recorded. Individuals were recorded multiple times in one scan if they were observed engaging with different food items. Plant food items were identified on the family or genus level.

Between September 2022 and July 2023, we opportunistically collected fecal and urine samples in conjunction with behavioral data. All samples were collected from identified individuals who appeared healthy upon visual inspection. We placed fecal samples into 4 mL of 98% ethanol immediately after defecation (58). To avoid contamination, we wore gloves and only collected feces that had not contacted the soil. We discarded feces contaminated with urine to prevent contamination from the urogenital microbiome (59). We collected urine samples using a clean catch method (54) to avoid rainfall and fecal contamination. We also collected urine from leaves or the ground if it appeared free of soil or fecal contamination. We immediately stored fecal and urine samples in a portable cooler and transferred them to a -20°F freezer at the end of each field day. We kept fecal samples under refrigeration until shipment to the Amato Microbiome Lab at Northwestern University in Evanston IL for DNA extraction and sequencing. Urine samples remained in the freezer until we conducted assay analysis onsite in the Capuchinos de Taboga field laboratory (56).

Temperature observations for this analysis came from the Palo Verde meteorological station, located at the Organization for Tropical Studies (OTS) Palo Verde research station (10.345°N, 85.339°E), approximately 20.81 km southwest of the *Taboga Forest Reserve*. Like Taboga, Palo Verde is one of Central America's last remaining tropical dry forest fragments (60). Palo Verde temperature data have been used in previous studies of the Taboga white-faced capuchin population (8), given the habitat similarity and proximity to Taboga. The station records maximum and minimum air temperature every 15 minutes.

Sample processing for 16S rRNA amplicon and shotgun metagenomic sequencing

DNA extraction

We conducted all lab work at the Amato Microbiome Lab at Northwestern University, Evanston IL. We extracted DNA from fecal samples using the DNeasy PowerLyzer PowerSoil Kit (Qiagen, Germantown, MD, USA), with minor modifications to manufacturer protocol. Specifically, we incubated samples at 65°C for 10 minutes after adding the C1 and PowerBead solutions, refrigerated the C2 and C3 incubation steps, and prewarmed the C6 solution before the final elution. These modifications align with other protocols designed for fecal samples containing high levels of plant fiber (61–63).

16s rRNA amplicon sequencing and data processing

We used 16s rRNA amplicon sequencing to describe the gut microbiome taxonomic composition. We amplified the hypervariable V4-V5 region of the 16s rRNA gene using the 515F/926R primers from the Earth Microbiome Project (www.earthmicrobiome.org) and a dual-indexing approach (64, 65). We used negative controls to monitor contamination during the DNA extraction and PCR amplification. We purified and normalized PCR products using SequalPrep Normalization and sequenced them on the Illumina MiSeq V4 platform at the Rush Genomics and Microbiome Core Facility.

We analyzed the resulting data using the Quantitative Insights into Microbial Ecology (QIIME2) platform (66), v2023.9. Pair-end sequences from 900 samples yielded 28,869,202 raw sequence reads (average 32,077 per sample, range of 7 to 64851 reads). We trimmed forward and reverse reads to 216 bases and 206 bases, respectively, to remove low-quality base pairs using a minimum Phred score of 33. We quality-filtered and denoised sequences using the Divisive Amplicon Denoising Algorithm 2 (DADA2)(67) via the q2-dada2 pipeline, which infers Amplicon Sequence Variants (ASVs) by identifying exact sequences. We assigned taxonomy to ASV's using the q2-feature classifier (68) against version 2022.10 of the Greengenes2 database (69, 70) by applying the classify-sklearn model. We removed sequences assigned to chloroplasts and mitochondria using the q2-taxa-filter-table to retain only microbial sequences for downstream analysis. After filtering, we retained 797 samples, which contained an average of 16,898 reads (range: 4,944–38,941). We imported the ASV count and taxonomy files generated by QIIME2 into R and removed ASV's found in less than 0.5% of total samples to avoid analyzing low-frequency taxa that are artifacts of the sequencing process (71). After filtering, we removed 102 additional ASV's that were not assigned at the phylum level, resulting in 1159 total remaining ASV's.

Shotgun metagenomic sequencing and data processing

We used metagenomic sequencing to characterize the functional capacity of the gut microbiome in a subset of 95 samples with sufficient DNA concentration for library preparation. We prepared sequencing libraries using the NEBNext Ultra™ II FS DNA PCR-free Library Prep Kit for Illumina (New England Biolabs, Ipswich, MA, USA). We followed the manufacturer protocol for low inputs (50ng to 99ng) and selected library inserts around 350bp. We pooled the libraries and sequenced them on the Illumina NovaSeq platform at the Rush Genomics and Microbiome Core Facility. Pair-end sequences from 95 samples yielded 2,902,714,548 raw sequence reads (average 15,277,445 per sample, range of 531,829 to 25,348,795 reads).

We quality-filtered the raw metagenomic sequences with KneadData (v0.12.0) using the standard parameters SLIDINGWINDOW:4:20 and MINLEN:90 (72). To remove host-associated reads we used the GRCh38.p13 human reference genome (RefSeq accession: GCF_009914755.1) which has been used to filter host genome data in previous capuchin metagenome studies (55). KneadData yielded 931,326,495 cleaned reads (average 98,034,376 per sample, range of 178,711 to 22,386,906 reads). We concatenated the forward and reverse reads and used HUMAnN3 (v3.9) for functional profiling. Gene families were assigned using the Uniref50 database and regrouped into MetaCyc reactions. The resulting table reported the abundance of each unique metabolic pathway normalized by the number of gene families contributing to that pathway. We imported the pathway abundance table into R and removed the unmapped and unintegrated values, leaving us with 548 unique metabolic pathways. We performed all subsequent analyses using R v4.3.1.

Metadata assembly

We paired 16s rRNA and shotgun metagenomic samples, the average neopterin value from a 7-day window, encompassing three days before, the day of, and three days after sample collection. Our time window provides a general measure of immune performance associated with the microbiome composition, enabling us to assess the relationship between the two. We averaged the maximum temperature (collected every 15 minutes) 15 days before sample collection, to assess response to cumulative thermal load (8). We also paired each sample with all diet

observations from the same individual within two weeks before sample collection, following time frames used in previous white-faced capuchin gut microbiome studies (55, 73). We quantified diet diversity by modeling diet richness (i.e., the total number of unique food items recorded) as a function of the total number of feeding observations and extracting the residuals (74). We then calculated the percent consumption of each food category (animal, flower, fruit, fungi, insect, leaves, nectar, nurse, pith, seeds) and plotted each against temperature to identify diet categories that vary with temperature. We found that percent fruit consumption has a strong, positive relationship with temperature (Fig 1S) and controlled for this in subsequent analyses. After matching each sample with the appropriate metadata, we were left with 277 16s rRNA samples and 71 shotgun metagenomic samples.

Community-level composition analysis

To evaluate how gut microbiome taxonomic composition and functional capacity are related to temperature, we generated metrics of community composition including alpha and beta diversity. To measure alpha diversity, we calculated two widely used indices: Shannon diversity index and observed richness. To measure beta diversity, we generated Aitchison distances (75) (76) by computing the Euclidean distance between pairwise samples after performing a centered log-ratio (clr) transformation on the raw count data (77). We added a pseudo count to all zeros using Bayesian-multiplicative replacement to preserve the structure of the data and minimize distortion (78). We then performed a Principal Components Analysis (PCA) on the Aitchison distance matrix and extracted the scores for the first three principal components (PC1–PC3) to capture major axes of variation in community composition. Components one through three were selected because they explained greater than 5% of the variation (Fig 2S).

To evaluate the relationship between temperature and community composition, we built mixed models using alpha diversity indices and the first three principal components as the outcome variables. We modeled each community metric twice, and compared model fit using Akaike information criterion adjusted for small sample sizes (AICc) (79, 80). The baseline model, *m_main*, included predictors for social group, age, sex, rainfall, diet diversity and fruit consumption. The temperature model, *m_full*, included all of the same predictors with an additional predictor for temperature (Fig 3S). Both models included a random intercept for individual to control for repeated sampling and an offset for the log-transformed number of reads to control for differences in sequencing depth between samples. We included fruit consumption as a predictor because it increased with rising temperatures (Fig 1S), and we wanted to control for this effect to better isolate the relationship between temperature and community features. We scaled all continuous predictors for ease of coefficient interpretation. We fit all models as linear mixed models (LMMs) with a Gaussian distribution except for observed taxonomic richness, which was fit using a Gamma distribution with a log link function because the variable was continuous and positive with a long right tail. We used the model weights from the AICc comparison to evaluate the importance of temperature as a predictor of community composition.

Temperature differential abundance analysis

We also used a model selection approach to identify taxa and metabolic pathways associated with temperature. For the taxonomic analysis, we aggregated the number of reads per taxa on the family and genus levels, calculated the relative abundance, and removed taxa with less than 0.2% average relative abundance across all samples to avoid analyzing extremely low-abundance taxa. This resulted in 134 genera and 79 families to be used for differential abundance testing. We

chose to use an Information-theoretic approach for our differential abundance analysis because, unlike null-hypothesis significance testing, it allows for the comparison of multiple alternative hypotheses to assess biological significance (81, 82). This allowed us to evaluate if adding temperature to our model improved the explanation of the abundance of specific taxa or metabolic pathways compared to other known predictors of microbiome composition.

We built two generalized linear mixed models (GLMMs) using a negative binomial distribution with a log link function (83) and used individual family and genera abundances as our outcome variables. We modeled each taxon twice and compared model fit using AICc. The baseline model, *m_main*, included predictors for social group, age, sex, rainfall, and fruit consumption. The temperature model, *m_full*, included all of the same predictors with an additional predictor for temperature (Fig 3S). Both models included a random intercept for individuals to control for repeated sampling and an offset for the log-transformed number of reads to control for differences in sequencing depth between samples. Similar to our community feature analysis, fruit consumption was included to better isolate the relationship between temperature and individual taxa while accounting for diet components also related to temperature. However, we did not include diet diversity as a predictor to reduce model complexity and facilitate model convergence. We scaled all continuous predictors. Model weights were used to evaluate the importance of temperature as a predictor of individual taxon abundance. Taxa where *m_full* held greater than 50% of the model weight were considered temperature sensitive. We then used the *m_full* model weights of all genera included in the analysis to perform ancestral state reconstruction. Internal node values represent the maximum likelihood estimates of the temperature model weight under a Brownian motion model of trait evolution.

The modeling approach for the metabolic pathway data mirrored that used for the taxonomic data, except we modeled clr-transformed abundances instead of raw counts. This allowed us to use a Gaussian distribution and omit the offset term for log-transformed read depth, as the clr transformation already accounts for differences in sequencing depth. We used clr-transformed abundances because pathway abundance values are normalized by the number of contributing gene families, meaning they are no longer raw counts. We modeled each of the 547 pathways twice, once with the baseline model, *m_main*, and once with the temperature model, *m_full* (Fig 3S) and compared the models using AICc. Pathways for which *m_full* accounted for more than 50% of the model weight were considered temperature sensitive.

Immune and temperature interaction analysis

To assess whether the relationship between temperature and community composition is associated with immune performance, we employed a second comparative modeling approach and used alpha diversity and the first three principal components as outcome variables. Similar to our previous approach, we modeled each community feature twice, once with each model, and compared model fit using AICc. The baseline model, *m_immune_main*, included the same predictors as *m_full* (i.e., social group, age, sex, rainfall, fruit consumption, temperature, random effect for ID, log total reads offset). The interaction model, *m_immune_full*, included all of the same predictors with an additional interaction effect between temperature and neopterin (Fig 3S). The interaction effect allowed us to test if the relationship between community composition and temperature changes depending on neopterin levels. We scaled all continuous predictors, including neopterin. All models were fit as linear mixed models (LMMs) with a Gaussian distribution except for observed taxonomic richness, which was fit using a Gamma distribution

with a log link function because the variable was continuous, positive, and left-skewed with a long right tail.

To assess whether the relationship between temperature and individual taxa or pathways is associated with immune performance, we employed the same model comparison approach but used taxon and pathway abundances as outcome variables. First, we filtered for taxa and pathways that were temperature-sensitive (i.e., m_full held greater than 50% of the model weight). Then, we modeled each feature twice, once with m_immune_main and once with m_immune_full , and compared model fit using AICc. Importantly, we fit the models with taxa as outcome variables as GLMMs with a negative binomial distribution, log link function, and included a log total reads offset. We fit the models with pathways as outcome variables as LMMs with a Gaussian distribution and did not include a log total reads offset. Features where m_immune_full held greater than 50% of the model weight were considered to have a relationship between temperature and abundance that changes depending on neopterin levels.

Results:

Temperature predicts gut microbiome taxonomic composition and functional capacity

Community-level results

We found a strong relationship between temperature and both taxonomic and functional community-level diversity in the gut microbiome. For taxonomic community composition, the model including temperature provided the best fit (controlling for model complexity) across all community-level indices, accounting for 100% of the model weight for every principal component and alpha diversity metric (Fig. 1B). The strong relationship between temperature and principal components 1–3 suggests that temperature drives shifts in taxonomic composition that are substantial enough to affect community-level beta diversity. Both alpha diversity indices showed a reliable negative relationship with temperature (Shannon $\beta_{temp} = -0.49$, Observed $\beta_{temp} = -0.49$)(Fig. 1A). Even though the effect sizes were small compared to the principal components, alpha diversity reliably declined with increasing temperatures (Fig. 4S). This suggests that higher temperatures are associated with reductions in both taxonomic richness and evenness.

For functional capacity, the temperature model provided the best fit for every principal component and the total number of observed metabolic pathways, accounting for at least 96% of the model weight. In contrast, the baseline model provided the best fit for Shannon diversity, accounting for 77% of the model weight (Fig. 1D). Therefore, temperature was associated with shifts in individual metabolic pathways substantial enough to affect community-level beta diversity, but the relationship between temperature and functional alpha diversity was less clear. Temperature was negatively associated with the total number of unique pathways (Observed: $\beta_{temp} = -35.05$), but the magnitude and direction of this effect were uncertain due to the wide confidence intervals that include zero. The effect size for Shannon diversity ($\beta_{temp} = -0.27$) was much smaller, and the confidence intervals also included zero (Fig. 1C). Together, these results suggest that higher temperatures may reduce the number of unique metabolic pathways but are not associated with changes in pathway evenness.

Individual taxon and metabolic pathway results

Temperature was also related to the abundance of individual taxa at both the family and genus levels. The temperature model provided the best fit (controlling for complexity) for 93 out of 163 taxa, accounting for at least 50% of the model weight. The majority of taxa, 82 out of 93,

declined in abundance with rising temperatures (Table 1S). Ancestral state reconstruction performed using the temperature model weights suggests that this is a conserved microbial trait for *C. imitator*, particularly in key phyla such as Actinobacteria, Bacteroidota, and Firmicutes. In contrast, phyla such as Campylobacterota and Proteobacteria appear less sensitive to temperature over evolutionary time (Fig 2).

Temperature was also related to the abundance of microbial metabolic pathways. The temperature model provided the best fit for 169 out of 383 total pathways, accounting for at least 50% of the model weight (Table 2S). To facilitate interpretation, we grouped 103 of these pathways into five functional categories based on shared biological roles and their apparent relationship with temperature: lipid metabolism, oxygen-driven metabolism and fermentation, redox maintenance, nucleotide mobilization, and immune regulation (Fig 3). Patterns in these categories suggest a metabolic restructuring that may help microbes respond to rising temperatures. For example, we found an increase in fatty acid synthesis pathways like oleate, cis-vaccenate, palmitoleate, stearate, petroselinic acid, and gondoate, and lipid remodeling pathways such as the mevalonate pathway and downstream isoprenoid lipid synthesis (Fig 3A), that may help maintain cytoplasmic membrane integrity at high temperatures (84, 85). However, we also found a decrease in pathways involved in lipid catabolism and recycling, including fatty acid β -oxidation, fatty acid salvage, and phospholipid turnover, which likely impairs membrane repair mechanisms (85, 86) and may compromise integrity during extended periods of high temperatures. Further, we observed an increase in anaerobic pathways, including the reductive TCA cycle and several involved in glycolysis, alongside a decrease in pathways that support aerobic metabolism, including the pyruvate dehydrogenase complex, the TCA cycle, and glyoxylate cycle (Fig 3B). This may reflect a response to changing oxygen gradients with rising temperatures to maintain energy production (87). The anaerobic pathways primarily metabolized simple carbohydrates and coincided with a decline in complex polysaccharide fermentation, likely reflecting taxonomic shifts with rising temperatures (37).

We also observed declines in pathways that support redox maintenance and cycling (Fig 3C), including the synthesis of cofactors such as B vitamins (88, 89), coenzymes (90), antioxidant precursors to glutathione (91), NADPH production, quinone synthesis, siderophore-mediated iron transport (92, 93), and detoxification pathways for aromatic compounds and aldehydes (94). In conjunction, we observed an increase in nucleotide and translation-associated metabolism pathways (Fig 3D), such as purine and pyrimidine de novo synthesis, nucleotide salvage and recycling (95, 96), queuosine synthesis, and S-adenosylmethionine salvage (97). This coordinated increase in nucleotide and translation-associated metabolism may reflect the consequences of redox imbalance at high temperatures because the accumulation of ROS can lead to DNA damage (98). Some pathways that support redox maintenance increased with rising temperatures (Fig 3D), including the synthesis of antioxidant precursors and vitamin B12 (99), coenzyme metabolism (100), and degradation of vitamin-like compounds (101), which may reflect a compensatory effort to restore redox balance.

Rising temperatures were also associated with a decline in pathways that produce metabolites known to support anti-inflammatory signaling (Fig 3E), including synthesis of the SCFAs butyrate and propanoate (42, 102), catabolism of monounsaturated fatty acids like oleate (103), and polyamine production from arginine degradation and ornithine synthesis (104). However, these shifts were accompanied by an increase in select pathways that may help compensate for the loss of microbial immune regulation, including the synthesis of the saturated

fatty acids (5Z)-dodecenoate and L-tryptophan biosynthesis, which leads to the production of indole derivatives that suppress pro-inflammatory cytokine signaling (105).

Neopterin is related to the gut microbiome's response to temperature

Community-level results

For taxonomic composition, the model including the interaction effect between temperature and neopterin concentration provided the best fit (controlling for complexity) for every principal component, accounting for at least 60% of the model weight (Fig 5S). Principal component 1 was the only component with confidence intervals that did not include zero. This result, along with the high model weight for every principal component, suggests that the relationship between temperature and beta diversity is associated with neopterin. In contrast, the temperature model provided the best fit for both alpha diversity metrics, accounting for at least 86% of the model weight. These results, combined with the small effect sizes (Shannon $\beta_{\text{temp}*\text{neopterin}} = -0.10$, Observed $\beta_{\text{temp}*\text{neopterin}} = -0.13$) and confidence intervals that include zero (Fig 5S), suggest that the relationship between taxonomic alpha diversity and temperature is independent of neopterin concentration.

For metabolic functional composition, the interaction model provided the best fit for every principal component and observed pathways, accounting for at least 98% of the model weight, while the temperature model provided the best fit for Shannon diversity, accounting for 63% of the model weight (Fig 5S). Even though principal component 3 was the only component where the confidence intervals did not include zero, the large effect sizes for all three components (PC1 $\beta_{\text{temp}*\text{neopterin}} = 17.60$, PC2 $\beta_{\text{temp}*\text{neopterin}} = -7.66$, PC3 $\beta_{\text{temp}*\text{neopterin}} = -11.44$) plus the model weights suggest that the interaction between neopterin and temperature has a meaningful influence on metabolic functional composition. The effect size and model weight for observed pathways (Observed $\beta_{\text{temp}*\text{neopterin}} = 72.78$) suggest that neopterin may be associated with the relationship between the total number of distinct metabolic pathways and temperature. However, it is unlikely that neopterin is associated with the relationship between temperature and pathway evenness due to the small effect size (Shannon $\beta_{\text{temp}*\text{neopterin}} = 0.46$), confidence intervals that include zero, and model weight for Shannon diversity.

Individual taxon and metabolic pathway results

Neopterin concentration was also associated with the relationship between temperature and the abundance of individual taxa. The interaction model provided the best fit for 20 of the 93 temperature-associated taxa, accounting for at least 50% of the model weight in each case (Table 3S). To interpret the interaction effect, we plotted the relationship between temperature and abundance across three discrete ranges of neopterin values (Fig 4). These results indicated two contrasting effects: (1) the relationship between temperature and abundance became more negative (i.e., the slopes decrease) at higher neopterin concentrations (Fig 4A), and (2) the relationship between temperature and abundance became more positive (i.e., the slopes increase) at higher neopterin concentrations (Fig 4B & C). The majority of taxa (12 out of 20) exhibited decreasing slopes at higher neopterin concentrations. These taxa were almost exclusively from the phyla Bacteroidota (5 taxa) and Firmicutes (6 taxa), and included two known SCFA producers, *Parabacteroides* and *Alistipes* (106, 107). The taxa that exhibited increasing slopes (8 out of 20) showed two distinct patterns: one where a positive slope became steeper when neopterin was high, and another where a negative slope changed to positive when neopterin was

high. These taxa included two SCFA producers, *Phascolarctobacterium* and *Eubacterium* (108, 109), and the family *Helicobacteraceae*, which commonly includes facultative anaerobes (110).

Neopterin concentration was also associated with the relationship between temperature and the abundance of individual metabolic pathways. The interaction model provided a better fit for 65 out of 116 pathways tested, accounting for at least 50% of the model weight in each case (Table 4S). To interpret the interaction effect, we plotted the relationship between abundance and temperature across two discrete ranges of neopterin values. These results indicated the same two effects as the taxonomic analysis (1) where slopes decreased at higher neopterin (Fig 5A-C) and (2) where slopes increased at higher neopterin (Fig 5D & E). The pathways that exhibited decreasing slopes showed three distinct patterns: one where a positive slope changed to negative, another where a positive slope became less positive, and one where a negative slope became steeper. This group included pathways involved in anti-inflammatory signaling, including synthesis of monounsaturated fatty acids (103) and tryptophan, as well as those typically associated with obligate anaerobes, including complex carb fermentation (Flint et al., 2012) and Gram-positive cell wall biosynthesis characteristic of obligate anaerobes from the Firmicutes phylum (111). The majority of pathways (37 out of 65) exhibited increasing slopes at higher neopterin concentrations. There were two patterns of increasing slopes: one where a negative slope changed to positive when neopterin was high, and another where a positive slope became steeper when neopterin was high. These groups included pathways commonly associated with facultative anaerobes, including the glyoxylate cycle (112), Rubisco shunt (113), ubiquinone biosynthesis (114), enterobacterial common antigen biosynthesis (115), and Lipid A biosynthesis (116). They also included pathways involved in redox maintenance, including the synthesis of B vitamins (88, 89), menaquinol (90), and NADPH (93).

Discussion:

Our study had two main goals: (1) to assess how temperature influences gut microbiome composition and function in wild, white-faced capuchins, and (2) to evaluate whether temperature-driven changes to the gut microbiome are related to host immunity. We found strong evidence that temperature is associated with community-level diversity, taxonomic composition, and metabolic capacity that may indicate the capuchins are undergoing periods of heat stress. Moreover, the microbiome's taxonomic and functional responses to temperature are linked to differences in urinary neopterin concentrations, suggesting that immunity is related to the gut microbiome's response to heat.

Gut microbiome response to temperature

Our data suggest that the gut microbiome's response to high temperatures may affect its role in conferring colonization resistance—the ability to protect the host from exogenous pathogens. Colonization resistance arises from direct competition of commensals with pathogens for resources and microbial support for host immunity (117). Community-level analysis reveals that taxonomic alpha diversity declines with increasing temperatures (Fig 2S), suggesting that heat may impair the gut microbiome's ability to resist colonization by reducing the diversity of taxa that are available to compete with pathogens (118). High temperatures are also associated with the decline in the Bacteroidota and Firmicutes phyla, which contain taxa known to support host immunity such as *Clostridiaceae*, *Lactococcus*, and *Barnesiella* (Fig 2)(119), which may leave hosts more vulnerable to infection. However, functional alpha diversity appears more resilient to increasing temperatures (Fig 1), likely due to functional redundancy—when

taxonomically distinct microbes perform similar functional roles (120). This redundancy may buffer the loss of taxonomic diversity and help preserve colonization resistance at high temperatures because remaining taxa fill enough functional niches to outcompete pathogens. Although we did not observe an increase in potentially pathogenic taxa, the phyla Proteobacteria and Campylobacteria, which are known to contain gut pathogens (36, 121), are less sensitive to heat compared to other phyla. Therefore, sustained high temperatures may continue to erode both taxonomic and functional diversity, eventually increasing the host's vulnerability to infection

Metabolic restructuring in response to rising temperatures suggests that the gut microbiome undergoes functional shifts that may support community stability. For example, the increase in lipid synthesis and remodeling pathways (Fig 3A) could reflect cytoplasmic membrane restructuring, a process bacteria use to preserve membrane stability at high temperatures (84, 122). However, these changes may be insufficient to maintain membrane integrity during extended periods of high temperatures because the observed decline in lipid catabolism and recycling pathways likely compromises membrane repair mechanisms (85). Limited repair capacity could leave gut microbes more vulnerable to peroxidative damage, a common consequence of heat stress driven by an accumulation of ROS that damages the lipid components of cell membranes (21, 123). Similarly, the observed increase in anaerobic metabolic pathways and the decline in aerobic pathways (Fig 3B) likely reflect a response to maintain energy production under changing oxygen gradients due to redirected blood flow during heat exposure (87). However, these anaerobic pathways are primarily involved in simple carbohydrate metabolism typical of facultative taxa (124, 125), and coincide with a decline in complex polysaccharide fermentation typical of obligate taxa (37–39). Therefore, this shift is likely driven by the relative persistence of facultative versus obligate anaerobes at high temperatures. Since fiber fermentation creates SCFAs, which are crucial energy sources (126) and immunomodulators (41, 42), extended periods of high temperatures may eventually cause a decline in overall energy availability and affect immune function. While these metabolic shifts suggest that capuchin gut microbes actively respond to heat in ways that may support community stability, the decline in membrane repair and inferred loss of obligate fermenters indicate that this response is unlikely to be sustainable over longer periods of high temperatures.

The taxonomic response to temperature suggests that changes in gut physiology at high temperatures affect the persistence of obligate anaerobes compared to facultative anaerobes. There are broad declines in two phyla known to include obligate anaerobes, Bacteroidota and Firmicutes, including essential obligate taxa such as *Bacteroides*, *Parabacteroides*, *Clostridium*, *Ruminococcus*, *Eubacterium*, *Blautia*, *Anaerostipes*, and *Alistipes* (Fig 2). In contrast, Proteobacteria, comprised of primarily facultative taxa (35, 36), are relatively stable in response to rising temperatures. These patterns mirror compositional changes in response to heat stress in a wide range of vertebrates [e.g., fish (127, 128), reptiles (32, 48), amphibians, birds (20), and mammals (129)]. The decrease in obligate anaerobes is likely due to redox imbalance induced by high temperatures, where the production of reactive oxygen species (ROS) exceeds the antioxidant capacity of the gut environment. Obligate anaerobes possess limited enzymatic defenses against ROS, and are therefore less able to tolerate changes in the oxygen environment compared to facultative taxa (33, 34). Although direct markers of redox balance are needed to assess the gut's redox environment, the observed decline in redox-maintenance pathways (Fig 3C) suggests that rising temperatures impair microbial antioxidant capacity, which can disrupt the redox environment (25–27). We also observe an increase in DNA repair pathways (Fig 3D) that suggests cellular damage from the accumulation of ROS (98). Together, these results

provide compelling evidence that capuchins experience periods of redox imbalance at high temperatures that may affect the relative abundance of facultative versus obligate anaerobes.

Further, a decline in obligate anaerobes during heat stress may disrupt the gut microbiome's ability to regulate host immunity by impairing key immunomodulatory functions. Obligate anaerobes are essential SCFA producers that help regulate intestinal immunity and maintain host tolerance towards gut microbiomes, including the promotion of anti-inflammatory regulatory T-cell activity (41, 102) and the suppression of proinflammatory cytokine signaling (42, 103). The decrease in SCFA-producing pathways, particularly butyrate and pyruvate (Fig 3E), likely reflects the loss of obligate anaerobes at high temperatures due to redox imbalance and may decrease host tolerance towards gut microbiomes. The resulting increase in immune activity may exacerbate redox imbalance because activated immune cells, particularly phagocytes, release ROS as part of the inflammatory response (130).

These results may be driven by confounding variables related to seasonal ecology, such as changes in diet, ranging patterns, social structure, activity budgets, or water sources that can influence microbiome composition and function(55, 73, 131). However, many of our results mirror those from experimental studies evaluating the effects of high temperature on the microbiome, indicating that we have likely captured a true temperature effect rather than an indirect signature of seasonal ecology. This is likely because we assessed cumulative exposure to high temperatures rather than short-term fluctuations that are more likely to covary with other ecological variables. We also focused on interpreting broad patterns across taxonomic groups and metabolic pathways (Figs. 2 & 3) rather than interpreting individual features that may be more sensitive to ecological confounds.

Importantly, our results resemble those associated with heat stress in experimental work, including a decrease in obligate anaerobes relative to facultative anaerobes (16), evidence of redox imbalance (28, 29), and changes in immunomodulatory function (17, 19, 44). Together, these findings suggest that the capuchins may be experiencing heat stress at the upper limit of their thermal range that could compromise essential microbiome functions. These effects also likely reinforce one another during extended periods of high temperatures via reciprocal interactions linking redox balance, microbial composition, and immune function (132, 133)(Fig 3F).

Gut microbiome response to temperature is related to microbiome-immune interactions

Declines in the abundance of obligate anaerobes with temperature may impact host immunity. For example, stronger declines in obligate anaerobes at high temperatures, such as *Parabacteroides* and *Alistipes*, are associated with higher neopterin levels (Fig 4A). These taxa contribute to anti-inflammatory signaling through the production of SCFAs (106, 107), suggesting that their loss may affect immune tolerance towards gut microbes. Similarly, stronger declines in metabolic pathways that produce anti-inflammatory immunomodulators are also associated with higher neopterin. This includes the synthesis of monounsaturated fatty acids (103) and tryptophan, a compound primarily synthesized by obligate anaerobes (134, 135) (Fig 5A & B). Conversely, high neopterin could also contribute to a stronger decline in obligate anaerobes at high temperatures, likely because increased macrophage activity exacerbates heat-driven redox imbalance. Taxa that exhibit stronger declines when neopterin is high are primarily from the Bacteroidota and Firmicutes phyla, which include many obligate anaerobes (Fig 4A). We also observed a stronger decline in pathways commonly associated with obligate anaerobes

including complex carb fermentation (37) and Gram-positive cell wall biosynthesis characteristic of obligates from the Firmicutes phylum (111)(Fig 5A).

In contrast, facultative anaerobes may be more resilient to these conditions. For example, *Helicobacteraceae* exhibits a stronger increase in abundance when neopterin is high (Fig 4B), suggesting they are better able to tolerate redox imbalance, and may exploit niches vacated by the loss of obligates. Similarly, several pathways commonly associated with facultative anaerobes exhibit a stronger increase at high neopterin levels and may help these taxa actively respond to changes in the gut environment (Fig 5D). For example, the glyoxylate cycle and Rubisco shunt both increase under high neopterin, which likely supports energy conservation under oxygen and energy-limited conditions (112, 113). In addition, redox maintenance pathways such as ubiquinone biosynthesis also increase which helps facultative anaerobes neutralize ROS (114). Synthesis of enterobacterial common antigen (ECA) also increases, which is characteristic of *Enterobacteriaceae* (115). Conversely, an increase in facultative anaerobes at high temperatures could also affect host immunity. For example, lipid A synthesis, which is characteristic of many facultative taxa, increases at higher neopterin and forms pathogen-associated molecular patterns (PAMPs) that are recognized by Toll-like receptors (TLRs) and can initiate pro-inflammatory signaling (116)(Fig 5D). *Helicobacteraceae* is also associated with inflammatory diseases and may promote macrophage activation by damaging the gut epithelium and disrupting the mucosal barrier (136).

Therefore, the decline of obligate anaerobes relative to facultative taxa at high temperatures may drive, and be driven by, increased immune activity in the gut. Although we cannot conclude directionality, these findings suggest that gut-immune interactions are an integral component of the gut microbiome's response to heat and may increase susceptibility to gut inflammation during prolonged high temperatures. These results may also reflect physiological heat stress in the capuchins, given the observed relationship between taxonomic shifts and immune activity that mirrors experimental findings, and the potential inflammatory consequences of long-term heat exposure (16, 17, 19, 44). Future work should incorporate additional immune biomarkers that directly assess inflammation, as neopterin primarily reflects macrophage activation and represents only one of the many mechanisms linking immunity and the gut microbiome. Continued research should also assess whether these effects have measurable impacts on host health and fitness, providing critical insight into how climate warming may influence wildlife through gut-immune pathways.

Conclusion:

Our results highlight the pivotal role of temperature in shaping the gut microbiome and its relationship with host immunity in a wild primate species. Many of the observed taxonomic and metabolic shifts resemble those characteristic of experimentally induced heat stress, including declines in obligate anaerobes relative to facultative anaerobes, evidence of redox imbalance, and altered immunomodulatory functions, suggesting that the capuchins may be experiencing heat stress that could compromise essential microbiome functions. We also found that gut-immune interactions are linked to the gut microbiome's response to heat and may increase susceptibility to gut inflammation during prolonged high temperatures. These results highlight the crucial role of temperature in shaping essential gut microbiome functions in a wild endotherm species and offer insight into how rising temperatures due to climate change may threaten species resilience through their microbiomes.

Data availability statement: All the data and computer code used for this manuscript have been deposited in a GitHub repository that will be made public upon acceptance. Raw sequence data generated by 16s rRNA amplicon sequencing and shotgun metagenomics will be deposited on NCBI upon publication.

Author contributions statement:

Conception and design of the manuscript: J.M.L. and A.J.M.

Drafting of the manuscript: J.M.L., A.J.M., and K.R.A.

Statistical analysis and figures: J.M.L. and A.J.M.

R code: J.M.L.

Sample acquisition: J.M.L., A.F.W., and J.C.O.

Sample processing: J.M.L., M.L.S.S., M.E.B., and A.F.W.

Study funding: J.M.L., M.E.B., and A.J.M.


Field site funding, logistics, and management: M.E.B., J.C.B., T.J.B., and J.C.O.

Competing interests: The authors declare that they have no competing interests.

Funding information: This research was supported by National Science Foundation grant BA-DDRIG 2236061 awarded to J.M.L. and A.J.M.; National Science Foundation grant BCS 1945121 awarded to T.J.B. (PI), M.E.B (Co-PI), and J.C.B. (Co-PI); National Science Foundation grant BCS 2127373 awarded to M.E.B.; a Leakey Foundation Research grant awarded to J.M.L. and A.J.M.; an International Primatological Society Research Award awarded to J.M.L., a University of Michigan International Institute grant awarded to J.M.L.; and a University of Michigan Rackham Graduate School grant awarded to J.M.L.

Acknowledgements: We would like to thank the Costa Rican institutions el Sistema Nacional de Áreas de Conservación (SINAC), the Comisión Nacional para la Gestión de la Biodiversidad (CONAGEBIO), the Ministerio de Ambiente y Energía (MINAE), and the Universidad Técnica Nacional (UTN) for supporting our research. We acknowledge the Capuchinos de Taboga field team for contributing to site logistics and data collection. We specifically want to thank L. F. Sinclair, V. A. Martins, S. A. Kovalaskas, W. E. Acevedo, E. M. Saborío, M. Chaves, C. I. Wright, C. Berry, and N. Guisneuf for help with sample collection and V. Viard for assistance in the laboratory. We would also like to thank Amato lab member A. Pulvino, University of Michigan Bioinformatics Core director C. Gates, and University of Michigan High Performance Computing Consultant C. Antonelli for their assistance with bioinformatics. All protocols involved in this study were previously approved by the University of Michigan IACUC (no. PRO00011898).

685 Figures:



686
687 **Figure 1:** Model comparison results assessing whether adding temperature to a base model of
688 known microbiome predictors improves the explanation of community-level taxonomic and
689 metabolic diversity. Panels **A** and **C** show the coefficients of the temperature predictor for
690 community diversity metrics, including beta diversity (measured by principal components 1-3)
691 and alpha diversity (measured by the Shannon Index and the observed number of ASVs or
692 metabolic pathways). The thick bars represent 50% confidence intervals (CI), while the thin bars
693 represent 95% CI. Panels **B** and **D** compare the model weights of the base model (in green) to the
694 temperature model (in yellow). The weights of the base model plus the temperature model equal
695 100%. The top panels **A** and **B** show the results for community-level taxonomic diversity, while
696 the bottom panels **C** and **D** show the results for metabolic pathway diversity. Results show a
697 strong association between temperature and beta diversity for both taxonomic composition and
698 metabolic capacity. In contrast, the association between temperature and alpha diversity differs
699 by data type: taxonomic alpha diversity declines with rising temperature, whereas metabolic
700 alpha diversity indicates a possible reduction in the number of unique pathways at higher
701 temperatures but no relationship with pathway evenness.
702





Figure 2: Cladogram showing ancestral state reconstruction of temperature sensitivity for 97 genera with $\geq 0.2\%$ average relative abundance. Branch color indicates model weight for the temperature model (includes base predictors plus temperature), with yellow representing high model weight (greater temperature sensitivity) and green representing low model weight. Genera where the temperature model accounts for $\geq 50\%$ of the model weight are classified as temperature sensitive and are depicted with red (negative association) or blue (positive association) circles at the branch tips. Genera with $< 50\%$ model weight are considered not temperature sensitive and are shown with grey circles. Most temperature-sensitive taxa decline in abundance with rising temperatures, especially within the Firmicutes and Bacteroidota phyla. In contrast, Proteobacteria and Campylobacterota appear less temperature sensitive compared to the other phyla.



Figure 3: Model results for the microbial metabolic pathways associated with temperature. Pathways are grouped into five functional categories (Panels A-E) based on shared biological roles and their associations with temperature. The thick bars represent 50% confidence intervals (CI), while the thin bars represent 95% CI. Summary annotations to the right of each panel describe the broad functional roles of the pathways within each category. Panel F visualizes the effects consistent with experimentally induced heat stress and how they may reinforce one another during extended periods of high temperatures.

Decreasing slopes



Increasing slopes





Figure 4: The relationship between taxa abundance and temperature plotted across three distinct ranges of neopterin concentration. There are two main effects. Row **A** includes taxa for which the relationship between temperature and abundance becomes increasingly negative at higher neopterin concentrations. Rows **B** and **C** include taxa for which the relationship becomes increasingly positive, either through an increasingly positive slope (Row **B**) or a slope that switches from negative to positive (Row **C**). Taxa highlighted in the results and discussion are labeled on the right side of each row. A complete list of all taxa included in the plots is available in Supplementary Table 3.

Decreasing slopes



Increasing slopes






Figure 5: The relationship between metabolic pathway abundance and temperature plotted across two distinct ranges of neopterin concentration. Pathway abundances are shown as centered log-ratio (clr)–transformed values. There are two main effects. Rows A–C include

pathways for which the relationship becomes increasingly negative at higher neopterin, either through a slope that switches from positive to negative (Row **A**), a slope that becomes less positive (Row **B**), or an increasingly negative slope (Row **C**). Rows **D** and **E** include pathways for which the relationship between temperature and abundance becomes increasingly positive at higher neopterin, either through a slope that switches from negative to positive (Row **D**) or an increasingly positive slope (Row **E**). Pathways highlighted in the results and discussion are labeled on the right side of each row. A complete list of all pathways included in the plots is available in Supplementary Table 4.

826
827 Supplemental information:



828 **Figure 1S:** Scatter plot of the count of fruit feeding observations two weeks before sample
829 collection and its relationship with temperature. The plot demonstrates a clear positive
830 relationship where fruit consumption increases with rising temperatures.
831



832 **Figure 2S:** Visualization of the eigenvalues of the taxonomic data for each principal component
833 in descending order. Principal components 1 through 3 were used for analysis because they
834 explain >5% of the variation.




Figure 3S: Flowchart outlining the model selection approach for each component of the analysis. Dark grey boxes represent the two data types: 16S rRNA amplicon sequences and shotgun metagenomic sequences. Light grey boxes specify the model formulas applied to each analysis. While the community-level modeling approach used the same model structure for both data types, the models for the temperature and interaction differential abundance analyses differed. Only features for which *m_full* accounted for more than 50% of the model weight were included in the neopterin–temperature interaction analysis.




Figure 4S: Plotted model results showing the relationship between taxonomic alpha diversity and temperature. Panel (A) measures alpha diversity using Shannon Index, while panel (B) uses the total number of ASVs. Both metrics demonstrate a reliable decrease in taxonomic alpha diversity as temperature increases.




Figure 5S: Model comparison results assessing whether adding an interaction between temperature and neopterin to the temperature model (model with base predictors plus temperature) improves the explanation of community-level taxonomic and metabolic diversity. Panels **A** and **C** show the coefficients of the interaction effect for community diversity metrics including beta diversity (measured by principal components 1-3) and alpha diversity (measured by the Shannon Index and the observed number of ASVs or metabolic pathways). The thick bars represent 50% confidence intervals (CI), while the thin bars represent 95% CI. Panels **B** and **D** compare the model weights of the temperature model (in green) to the interaction model (in yellow). The weights of the temperature model plus the interaction model equal 100%. The top panels **A** and **B** show the results for community-level taxonomic diversity while the bottom panels **C** and **D** show the results for metabolic pathway diversity. Results show that the relationship between temperature and beta diversity is associated with neopterin concentration for both taxonomic and metabolic community composition, but the relationship between alpha diversity and temperature is independent of neopterin concentration.

Table 1S: Csv file containing results from the differential abundance analysis of 163 microbial taxa to evaluate their relationship with temperature. The analysis compares a baseline model, *m_main*, which includes predictors for social group, age, sex, rainfall, and fruit consumption, to a full model, *m_full*, that includes all the same predictors plus temperature. The column contents are as follows:

- Column 1: Taxon name
- Column 2: Temperature coefficient from the *m_full* model
- Columns 3-7: Standard error and 50% and 95% confidence intervals for the temperature coefficient
- Columns 8-9: Taxonomic classification
- Column 10: Model weight for the *m_full* model
- Column 11: Direction of the temperature coefficient (positive or negative)
- Column 12: Classification of the taxon as temperature-sensitive (model weight > 0.5) or neutral (model weight ≤ 0.5)

Table 2S: Csv file containing results from the differential abundance analysis of 383 metabolic pathways to evaluate their relationship with temperature. The analysis compares a baseline model, *m_main*, which includes predictors for social group, age, sex, rainfall, and fruit consumption, to a full model, *m_full*, that includes all the same predictors plus temperature. The column contents are as follows:

- Column 1: pathway Metacyc ID
- Column 2: Temperature coefficient from the *m_full* model
- Columns 3–7: Standard error and 50% and 95% confidence intervals for the temperature coefficient
- Column 8: pathway name
- Column 9: Model weight for the *m_full* model
- Column 10: Direction of the temperature coefficient (positive or negative)
- Column 12: Classification of the pathway as temperature-sensitive (model weight > 0.5) or neutral (model weight ≤ 0.5)
- Columns 13-14: Functional category assignments for each pathway, or an indication if the pathway was not assigned to any functional category.

Table 3S: Csv file containing the results of the second comparative modeling approach containing 20 taxa whose relationship with temperature is associated with neopterin. The analysis compares a baseline model, *m_immune_main*, which includes predictors for social group, age, sex, rainfall, and fruit consumption, and temperature to a full model, *m_immune_full*, that includes all the same predictors plus an interaction effect between neopterin and temperature. The column contents are as follows:

- Column 1: Taxon name
- Column 2: Temperature*neopterin coefficient from the *m_immune_full* model
- Columns 3-7: Standard error and 50% and 95% confidence intervals for the interaction coefficient
- Columns 8-9: Taxonomic classification

- Column 10: Model weight for the m_immune_full model
- Column 11: Category of the interaction effect, whether the relationship between temperature and abundance becomes more negative (i.e. i.e., the slopes decrease) at higher neopterin concentrations or the relationship between temperature and abundance becomes more positive (i.e. i.e., the slopes increase) at higher neopterin concentrations

Table 4S: Csv file containing the results of the second comparative modeling approach containing 65 metabolic pathways whose relationship with temperature is associated with neopterin. The analysis compares a baseline model, m_immune_main, which includes predictors for social group, age, sex, rainfall, and fruit consumption, and temperature to a full model, m_immune_full, that includes all the same predictors plus an interaction effect between neopterin and temperature. The column contents are as follows:

- Column 1: pathway Metacyc ID
- Column 2: Temperature*neopterin coefficient from the m_immune_full model
- Columns 3-7: Standard error and 50% and 95% confidence intervals for the interaction coefficient
- Columns 8: Pathway name
- Column 9: Model weight for the m_immune_full model

Column 10: Category of the interaction effect, whether the relationship between temperature and abundance becomes more negative (i.e. i.e., the slopes decrease) at higher neopterin concentrations or the relationship between temperature and abundance becomes more positive (i.e. i.e., the slopes increase) at higher neopterin concentrations.

References:

1. C. Bellard, C. Bertelsmeier, P. Leadley, W. Thuiller, F. Courchamp, Impacts of climate change on the future of biodiversity. *Ecology Letters* **15**, 365–377 (2012).
2. J. M. Bennett, J. Sunday, P. Calosi, F. Villalobos, B. Martínez, R. Molina-Venegas, M. B. Araújo, A. C. Algar, S. Clusella-Trullas, B. A. Hawkins, S. A. Keith, I. Kühn, C. Rahbek, L. Rodríguez, A. Singer, I. Morales-Castilla, M. Á. Olalla-Tárraga, The evolution of critical thermal limits of life on Earth. *Nat Commun* **12**, 1198 (2021).
3. W. W. Dowd, F. A. King, M. W. Denny, Thermal variation, thermal extremes and the physiological performance of individuals. *Journal of Experimental Biology* **218**, 1956–1967 (2015).
4. H. O. Pörtner, R. Knust, Climate Change Affects Marine Fishes Through the Oxygen Limitation of Thermal Tolerance. *Science* **315**, 95–97 (2007).
5. B. J. Sinclair, K. E. Marshall, M. A. Sewell, D. L. Levesque, C. S. Willett, S. Slotsbo, Y. Dong, C. D. G. Harley, D. J. Marshall, B. S. Helmuth, R. B. Huey, Can we predict ectotherm responses to climate change using thermal performance curves and body temperatures? *Ecology Letters* **19**, 1372–1385 (2016).
6. R. S. Telemeco, M. J. Elphick, R. Shine, Nesting lizards (*Bassiana duperreyi*) compensate partly, but not completely, for climate change. *Ecology* **90**, 17–22 (2009).
7. B. G. Lovegrove, C. Canale, D. Levesque, G. Fluch, M. Řeháková-Petrů, T. Ruf, Are Tropical Small Mammals Physiologically Vulnerable to Arrhenius Effects and Climate Change? *Physiological and Biochemical Zoology* **87**, 30–45 (2014).
8. J. M. Lucore, J. C. Beehner, A. F. White, L. F. Sinclair, V. A. Martins, S. A. Kovalaskas, J. C. Ordoñez, T. J. Bergman, M. E. Benítez, A. J. Marshall, High temperatures are associated with decreased immune system performance in a wild primate. *Science Advances* **10**, eadq6629 (2024).
9. N. N. Moghadam, Thorshauge ,Pia Mai, Kristensen ,Torsten N., de Jonge ,Nadieh, Bahrndorff ,Simon, Kjeldal ,Henrik, J. L. and Nielsen, Strong responses of *Drosophila melanogaster* microbiota to developmental temperature. *Fly* **12**, 1–12 (2018).
10. M. Ziegler, F. O. Seneca, L. K. Yum, S. R. Palumbi, C. R. Voolstra, Bacterial community dynamics are linked to patterns of coral heat tolerance. *Nat Commun* **8**, 14213 (2017).
11. L. Baldassarre, H. Ying, A. M. Reitzel, S. Franzenburg, S. Fraune, Microbiota mediated plasticity promotes thermal adaptation in the sea anemone *Nematostella vectensis*. *Nat Commun* **13**, 3804 (2022).
12. S. S. Fontaine, K. D. Kohl, Ectotherm heat tolerance and the microbiome: current understanding, future directions and potential applications. *Journal of Experimental Biology* **226**, jeb245761 (2023).

- 995 13. A. Jaramillo, L. E. Castañeda, Gut Microbiota of *Drosophila subobscura* Contributes to Its
996 Heat Tolerance and Is Sensitive to Transient Thermal Stress. *Front. Microbiol.* **12** (2021).
- 997 14. B. Zhang, S. P. Leonard, Y. Li, N. A. Moran, Obligate bacterial endosymbionts limit
998 thermal tolerance of insect host species. *Proceedings of the National Academy of Sciences*
999 **116**, 24712–24718 (2019).
- 1000 15. K. R. Amato, Incorporating the gut microbiota into models of human and non-human
1001 primate ecology and evolution. *American Journal of Physical Anthropology* **159**, 196–215
1002 (2016).
- 1003 16. J. Sepulveda, A. H. Moeller, The Effects of Temperature on Animal Gut Microbiomes.
1004 *Front. Microbiol.* **11**, 384 (2020).
- 1005 17. S. Chen, J. Wang, D. Peng, G. Li, J. Chen, X. Gu, Exposure to heat-stress environment
1006 affects the physiology, circulation levels of cytokines, and microbiome in dairy cows. *Sci*
1007 *Rep* **8**, 14606 (2018).
- 1008 18. S. S. Fontaine, A. J. Novarro, K. D. Kohl, Environmental temperature alters the digestive
1009 performance and gut microbiota of a terrestrial amphibian. *Journal of Experimental*
1010 *Biology* **221**, jeb187559 (2018).
- 1011 19. K. Tajima, I. Nonaka, K. Higuchi, N. Takusari, M. Kurihara, A. Takenaka, M. Mitsumori,
1012 H. Kajikawa, R. I. Aminov, Influence of high temperature and humidity on rumen
1013 bacterial diversity in Holstein heifers. *Anaerobe* **13**, 57–64 (2007).
- 1014 20. L. Zhu, R. Liao, N. Wu, G. Zhu, C. Yang, Heat stress mediates changes in fecal
1015 microbiome and functional pathways of laying hens. *Appl Microbiol Biotechnol* **103**, 461–
1016 472 (2019).
- 1017 21. I. Belhadj Slimen, N. Taha, G. Abdeljelil, D. Hajer, B. M. Moncef, A. Manef, Reactive
1018 oxygen species, heat stress and oxidative-induced mitochondrial damage. A review.
1019 *International Journal of Hyperthermia* **30**, 513–523 (2014).
- 1020 22. M. Dodson, R. Castro-Portuguez, D. D. Zhang, NRF2 plays a critical role in mitigating
1021 lipid peroxidation and ferroptosis. *Redox Biology* **23**, 101107 (2019).
- 1022 23. G. P. Lambert, C. V. Gisolfi, D. J. Berg, P. L. Moseley, L. W. Oberley, K. C. Kregel,
1023 Selected Contribution: Hyperthermia-induced intestinal permeability and the role of
1024 oxidative and nitrosative stress. *Journal of Applied Physiology* **92**, 1750–1761 (2002).
- 1025 24. F. Koch, U. Thom, E. Albrecht, R. Weikard, W. Nolte, B. Kuhla, C. Kuehn, Heat stress
1026 directly impairs gut integrity and recruits distinct immune cell populations into the bovine
1027 intestine. *Proceedings of the National Academy of Sciences* **116**, 10333–10338 (2019).
- 1028 25. C. Kunst, S. Schmid, M. Michalski, D. Tümen, J. Buttenschön, M. Müller, K. Gülow, The
1029 Influence of Gut Microbiota on Oxidative Stress and the Immune System. *Biomedicines*
1030 **11**, 1388 (2023).

- 1031 26. Y. Sun, X. Wang, L. Li, C. Zhong, Y. Zhang, X. Yang, M. Li, C. Yang, The role of gut
1032 microbiota in intestinal disease: from an oxidative stress perspective. *Front. Microbiol.* **15**
1033 (2024).
- 1034 27. T. Yardeni, C. E. Tanes, K. Bittinger, L. M. Mattei, P. M. Schaefer, L. N. Singh, G. D.
1035 Wu, D. G. Murdock, D. C. Wallace, Host mitochondria influence gut microbiome
1036 diversity: A role for ROS. *Science Signaling* **12**, eaaw3159 (2019).
- 1037 28. M. Fang, Z. Lei, M. Ruilin, W. Jing, D. Leqiang, High temperature stress induced
1038 oxidative stress, gut inflammation and disordered metabolome and microbiome in tsinling
1039 lenok trout. *Ecotoxicology and Environmental Safety* **266**, 115607 (2023).
- 1040 29. X. Liu, Z. Ma, Y. Wang, H. Jia, Z. Wang, L. Zhang, Heat stress exposure cause alterations
1041 in intestinal microbiota, transcriptome, and metabolome of broilers. *Front. Microbiol.* **14**
1042 (2023).
- 1043 30. B. Xia, W. Wu, W. Fang, X. Wen, J. Xie, H. Zhang, Heat stress-induced mucosal barrier
1044 dysfunction is potentially associated with gut microbiota dysbiosis in pigs. *Animal*
1045 *Nutrition* **8**, 289–299 (2022).
- 1046 31. X. Bai, Y. Shi, L. Tang, L. Chen, H. Fan, H. Wang, J. Wang, X. Jia, S. Chen, S. Lai, Heat
1047 Stress Affects Faecal Microbial and Metabolic Alterations of Rabbits. *Front. Microbiol.*
1048 **12** (2022).
- 1049 32. E. Bestion, S. Jacob, L. Zinger, L. Di Gesu, M. Richard, J. White, J. Cote, Climate
1050 warming reduces gut microbiota diversity in a vertebrate ectotherm. *Nat Ecol Evol* **1**, 1–3
1051 (2017).
- 1052 33. Z. Lu, R. Sethu, J. A. Imlay, Endogenous superoxide is a key effector of the oxygen
1053 sensitivity of a model obligate anaerobe. *Proceedings of the National Academy of Sciences*
1054 **115**, E3266–E3275 (2018).
- 1055 34. J. M. McCord, B. B. Keele, I. Fridovich, An Enzyme-Based Theory of Obligate
1056 Anaerobiosis: The Physiological Function of Superoxide Dismutase. *Proceedings of the*
1057 *National Academy of Sciences* **68**, 1024–1027 (1971).
- 1058 35. L. Albenberg, T. V. Esipova, C. P. Judge, K. Bittinger, J. Chen, A. Laughlin, S. Grunberg,
1059 R. N. Baldassano, J. D. Lewis, H. Li, S. R. Thom, F. D. Bushman, S. A. Vinogradov, G.
1060 D. Wu, Correlation Between Intraluminal Oxygen Gradient and Radial Partitioning of
1061 Intestinal Microbiota. *Gastroenterology* **147**, 1055–1063.e8 (2014).
- 1062 36. N.-R. Shin, T. W. Whon, J.-W. Bae, *Proteobacteria*: microbial signature of dysbiosis in
1063 gut microbiota. *Trends in Biotechnology* **33**, 496–503 (2015).
- 1064 37. H. J. Flint, Scott ,Karen P., Duncan ,Sylvia H., Louis ,Petra, E. and Forano, Microbial
1065 degradation of complex carbohydrates in the gut. *Gut Microbes* **3**, 289–306 (2012).

- 1066 38. L. V. Hooper, T. Midtvedt, J. I. Gordon, How host-microbial interactions shape the
1067 nutrient environment of the mammalian intestine. *Annual Review of Nutrition* **22**, 283–307
1068 (2002).
- 1069 39. E. Maier, R. C. Anderson, N. C. Roy, Understanding How Commensal Obligate
1070 Anaerobic Bacteria Regulate Immune Functions in the Large Intestine. *Nutrients* **7**, 45–73
1071 (2014).
- 1072 40. R. Corrêa-Oliveira, J. L. Fachi, A. Vieira, F. T. Sato, M. A. R. Vinolo, Regulation of
1073 immune cell function by short-chain fatty acids. *Clinical & Translational Immunology* **5**,
1074 e73 (2016).
- 1075 41. Y. Furusawa, Y. Obata, S. Fukuda, T. A. Endo, G. Nakato, D. Takahashi, Y. Nakanishi, C.
1076 Uetake, K. Kato, T. Kato, M. Takahashi, N. N. Fukuda, S. Murakami, E. Miyauchi, S.
1077 Hino, K. Atarashi, S. Onawa, Y. Fujimura, T. Lockett, J. M. Clarke, D. L. Topping, M.
1078 Tomita, S. Hori, O. Ohara, T. Morita, H. Koseki, J. Kikuchi, K. Honda, K. Hase, H. Ohno,
1079 Commensal microbe-derived butyrate induces the differentiation of colonic regulatory T
1080 cells. *Nature* **504**, 446–450 (2013).
- 1081 42. L. Klampfer, J. Huang, T. Sasazuki, S. Shirasawa, L. Augenlicht, Inhibition of Interferon γ
1082 Signaling by the Short Chain Fatty Acid Butyrate¹. *Molecular Cancer Research* **1**, 855–
1083 862 (2003).
- 1084 43. X. Liu, J. Shao, Y.-T. Liao, L.-N. Wang, Y. Jia, P. Dong, Z. Liu, D. He, C. Li, X. Zhang,
1085 Regulation of short-chain fatty acids in the immune system. *Front Immunol* **14**, 1186892
1086 (2023).
- 1087 44. Z. Yu, J. M. Cantet, H. A. Paz, J. D. Kaufman, M. S. Orellano, I. R. Ipharraguerre, A. G.
1088 Rius, Heat stress–associated changes in the intestinal barrier, inflammatory signals, and
1089 microbiome communities in dairy calves. *Journal of Dairy Science* **107**, 1175–1196
1090 (2024).
- 1091 45. J. He, Y. He, D. Pan, J. Cao, Y. Sun, X. Zeng, Associations of Gut Microbiota With Heat
1092 Stress-Induced Changes of Growth, Fat Deposition, Intestinal Morphology, and
1093 Antioxidant Capacity in Ducks. *Front. Microbiol.* **10** (2019).
- 1094 46. Y.-J. Seo, L. Chiwoong, L. Byeonghwi, K. Jun-Mo, Microbial-transcriptome integrative
1095 analysis of heat stress effects on amino acid metabolism and lipid peroxidation in poultry
1096 jejunum. *Animal Biotechnology* **35**, 2331179 (2024).
- 1097 47. K. J. Emerson, S. S. Fontaine, K. D. Kohl, S. K. Woodley, Temperature and the microbial
1098 environment alter brain morphology in a larval amphibian. *Journal of Experimental*
1099 *Biology* **226**, jeb245333 (2023).
- 1100 48. A. H. Moeller, K. Ivey, M. B. Cornwall, K. Herr, J. Rede, E. N. Taylor, A. R. Gunderson,
1101 The Lizard Gut Microbiome Changes with Temperature and Is Associated with Heat
1102 Tolerance. *Applied and Environmental Microbiology* **86**, e01181-20 (2020).

- 1103 49. G. Wang, X. Li, Y. Zhou, J. Feng, M. Zhang, Effects of Heat Stress on Gut-Microbial
1104 Metabolites, Gastrointestinal Peptides, Glycolipid Metabolism, and Performance of
1105 Broilers. *Animals* **11**, 1286 (2021).
- 1106 50. C. Murr, B. Widner, B. Wirleitner, D. Fuchs, Neopterin as a marker for immune system
1107 activation. *Curr Drug Metab* **3**, 175–187 (2002).
- 1108 51. B. Kelsall, Recent progress in understanding the phenotype and function of intestinal
1109 dendritic cells and macrophages. *Mucosal Immunology* **1**, 460–469 (2008).
- 1110 52. J. Wang, W.-D. Chen, Y.-D. Wang, The Relationship Between Gut Microbiota and
1111 Inflammatory Diseases: The Role of Macrophages. *Front. Microbiol.* **11** (2020).
- 1112 53. C. A. Portillo-Quintero, G. A. Sánchez-Azofeifa, Extent and conservation of tropical dry
1113 forests in the Americas. *Biological Conservation* **143**, 144–155 (2010).
- 1114 54. M. L. Bergstrom, M. Emery Thompson, A. D. Melin, L. M. Fedigan, Using urinary
1115 parameters to estimate seasonal variation in the physical condition of female white-faced
1116 capuchin monkeys (*Cebus capucinus imitator*). *American Journal of Physical*
1117 *Anthropology* **163**, 707–715 (2017).
- 1118 55. J. D. Orkin, F. A. Campos, M. S. Myers, S. E. Cheves Hernandez, A. Guadamuz, A. D.
1119 Melin, Seasonality of the gut microbiota of free-ranging white-faced capuchins in a
1120 tropical dry forest. *The ISME Journal* **13**, 183–196 (2019).
- 1121 56. J. C. Beehner, J. Alfaro, C. Allen, M. E. Benítez, T. J. Bergman, M. S. Buehler, S. C.
1122 Carrera, E. M. Chester, T. Deschner, A. Fuentes, C. M. Gault, I. Godoy, K. M. Jack, J. D.
1123 Kim, L. Kolinski, N. K. Kulick, T. Losch, J. C. Ordoñez, S. E. Perry, F. Pinto, O. T.
1124 Reilly, E. T. Johnson, M. D. Wasserman, Using an on-site laboratory for fecal steroid
1125 analysis in wild white-faced capuchins. *General and Comparative Endocrinology* **329**,
1126 114109 (2022).
- 1127 57. E. Tinsley Johnson, M. E. Benítez, A. Fuentes, C. R. McLean, A. B. Norford, J. C.
1128 Ordoñez, J. C. Beehner, T. J. Bergman, High density of white-faced capuchins (*Cebus*
1129 *capucinus*) and habitat quality in the Taboga Forest of Costa Rica. *American Journal of*
1130 *Primatology* **82**, e23096 (2020).
- 1131 58. K. R. Amato, C. J. Yeoman, A. Kent, N. Righini, F. Carbonero, A. Estrada, H. Rex
1132 Gaskins, R. M. Stumpf, S. Yildirim, M. Torralba, M. Gillis, B. A. Wilson, K. E. Nelson,
1133 B. A. White, S. R. Leigh, Habitat degradation impacts black howler monkey (*Alouatta*
1134 *pigra*) gastrointestinal microbiomes. *ISME J* **7**, 1344–1353 (2013).
- 1135 59. K. Thomas-White, S. C. Forster, N. Kumar, M. Van Kuiken, C. Putonti, M. D. Stares, E.
1136 E. Hilt, T. K. Price, A. J. Wolfe, T. D. Lawley, Culturing of female bladder bacteria
1137 reveals an interconnected urogenital microbiota. *Nat Commun* **9**, 1557 (2018).

- 1138 60. M. Stern, M. Quesada, K. E. Stoner, Changes in composition and structure of a tropical
1139 dry forest following intermittent Cattle grazing. *Revista de Biología Tropical* **50**, 1021–
1140 1034 (2002).
- 1141 61. K. R. Amato, S. Kuthyar, M. Ekanayake-Weber, R. Salmi, N. Snyder-Mackler, L.
1142 Wijayathunga, R. Vandercone, A. Lu, Gut microbiome, diet, and conservation of
1143 endangered langurs in Sri Lanka. *Biotropica* **52**, 981–990 (2020).
- 1144 62. E. k. Mallott, R. s. Malhi, P. a. Garber, High-throughput sequencing of fecal DNA to
1145 identify insects consumed by wild Weddell’s saddleback tamarins (*aguinus weddelli*,
1146 Cebidae, Primates) in Bolivia. *American Journal of Physical Anthropology* **156**, 474–481
1147 (2015).
- 1148 63. S. B. Pickett, C. M. Bergey, A. Di Fiore, A Metagenomic Study of Primate Insect Diet
1149 Diversity. *American Journal of Primatology* **74**, 622–631 (2012).
- 1150 64. D. M. Gohl, P. Vangay, J. Garbe, A. MacLean, A. Hauge, A. Becker, T. J. Gould, J. B.
1151 Clayton, T. J. Johnson, R. Hunter, D. Knights, K. B. Beckman, Systematic improvement
1152 of amplicon marker gene methods for increased accuracy in microbiome studies. *Nat*
1153 *Biotechnol* **34**, 942–949 (2016).
- 1154 65. E. K. Mallott, K. R. Amato, The microbial reproductive ecology of white-faced capuchins
1155 (*Cebus capucinus*). *American Journal of Primatology* **80**, e22896 (2018).
- 1156 66. J. G. Caporaso, J. Kuczynski, J. Stombaugh, K. Bittinger, F. D. Bushman, E. K. Costello,
1157 N. Fierer, A. G. Peña, J. K. Goodrich, J. I. Gordon, G. A. Huttley, S. T. Kelley, D.
1158 Knights, J. E. Koenig, R. E. Ley, C. A. Lozupone, D. McDonald, B. D. Muegge, M.
1159 Pirrung, J. Reeder, J. R. Sevinsky, P. J. Turnbaugh, W. A. Walters, J. Widmann, T.
1160 Yatsunenko, J. Zaneveld, R. Knight, QIIME allows analysis of high-throughput
1161 community sequencing data. *Nat Methods* **7**, 335–336 (2010).
- 1162 67. B. J. Callahan, P. J. McMurdie, M. J. Rosen, A. W. Han, A. J. A. Johnson, S. P. Holmes,
1163 DADA2: High-resolution sample inference from Illumina amplicon data. *Nat Methods* **13**,
1164 581–583 (2016).
- 1165 68. N. A. Bokulich, B. D. Kaehler, J. R. Rideout, M. Dillon, E. Bolyen, R. Knight, G. A.
1166 Huttley, J. Gregory Caporaso, Optimizing taxonomic classification of marker-gene
1167 amplicon sequences with QIIME 2’s q2-feature-classifier plugin. *Microbiome* **6**, 90
1168 (2018).
- 1169 69. D. McDonald, M. N. Price, J. Goodrich, E. P. Nawrocki, T. Z. DeSantis, A. Probst, G. L.
1170 Andersen, R. Knight, P. Hugenholtz, An improved Greengenes taxonomy with explicit
1171 ranks for ecological and evolutionary analyses of bacteria and archaea. *ISME J* **6**, 610–618
1172 (2012).
- 1173 70. D. McDonald, Y. Jiang, M. Balaban, K. Cantrell, Q. Zhu, A. Gonzalez, J. T. Morton, G.
1174 Nicolaou, D. H. Parks, S. M. Karst, M. Albertsen, P. Hugenholtz, T. DeSantis, S. J. Song,
1175 A. Bartko, A. S. Havulinna, P. Jousilahti, S. Cheng, M. Inouye, T. Niiranen, M. Jain, V.

- 1176 Salomaa, L. Lahti, S. Mirarab, R. Knight, Greengenes2 unifies microbial data in a single
1177 reference tree. *Nat Biotechnol* **42**, 715–718 (2024).
- 1178 71. Q. Cao, X. Sun, K. Rajesh, N. Chalasani, K. Gelow, B. Katz, V. H. Shah, A. J. Sanyal, E.
1179 Smirnova, Effects of Rare Microbiome Taxa Filtering on Statistical Analysis. *Front.*
1180 *Microbiol.* **11** (2021).
- 1181 72. F. Beghini, L. J. McIver, A. Blanco-Míguez, L. Dubois, F. Asnicar, S. Maharjan, A.
1182 Mailyan, P. Manghi, M. Scholz, A. M. Thomas, M. Valles-Colomer, G. Weingart, Y.
1183 Zhang, M. Zolfo, C. Huttenhower, E. A. Franzosa, N. Segata, Integrating taxonomic,
1184 functional, and strain-level profiling of diverse microbial communities with bioBakery 3.
1185 *eLife* **10**, e65088 (2021).
- 1186 73. E. K. Mallott, K. R. Amato, P. A. Garber, R. S. Malhi, Influence of fruit and invertebrate
1187 consumption on the gut microbiota of wild white-faced capuchins (*Cebus capucinus*).
1188 *American Journal of Physical Anthropology* **165**, 576–588 (2018).
- 1189 74. A. J. Marshall, E. Meijaard, M. Leighton, Extreme ecological specialisation in a rainforest
1190 mammal, the Bornean tufted ground squirrel, *Rheithrosciurus macrotis* Gray, 1857. *Raffles*
1191 *Bulletin of Zoology* **69**, 497506 (2021).
- 1192 75. G. B. Gloor, G. Reid, Compositional analysis: a valid approach to analyze microbiome
1193 high-throughput sequencing data. *Can. J. Microbiol.* **62**, 692–703 (2016).
- 1194 76. G. B. Gloor, J. M. Macklaim, V. Pawlowsky-Glahn, J. J. Egozcue, Microbiome Datasets
1195 Are Compositional: And This Is Not Optional. *Front. Microbiol.* **8** (2017).
- 1196 77. J. Aitchison, M. Greenacre, Biplots of Compositional Data. *Journal of the Royal*
1197 *Statistical Society Series C: Applied Statistics* **51**, 375–392 (2002).
- 1198 78. J. Palarea-Albaladejo, J. A. Martín-Fernández, zCompositions — R package for
1199 multivariate imputation of left-censored data under a compositional approach.
1200 *Chemometrics and Intelligent Laboratory Systems* **143**, 85–96 (2015).
- 1201 79. H. Akaike, “Information Theory and an Extension of the Maximum Likelihood Principle”
1202 in *Selected Papers of Hirotugu Akaike*, E. Parzen, K. Tanabe, G. Kitagawa, Eds. (Springer
1203 New York, New York, NY, 1998; [http://link.springer.com/10.1007/978-1-4612-1694-](http://link.springer.com/10.1007/978-1-4612-1694-0_15)
1204 [0_15](http://link.springer.com/10.1007/978-1-4612-1694-0_15)) *Springer Series in Statistics*, pp. 199–213.
- 1205 80. C. M. Hurvich, C.-L. Tsai, Regression and time series model selection in small samples.
- 1206 81. D. R. Anderson, K. P. Burnham, W. L. Thompson, Null Hypothesis Testing: Problems,
1207 Prevalence, and an Alternative. *The Journal of Wildlife Management* **64**, 912–923 (2000).
- 1208 82. K. P. Burnham, D. R. Anderson, K. P. Huyvaert, AIC model selection and multimodel
1209 inference in behavioral ecology: some background, observations, and comparisons. *Behav*
1210 *Ecol Sociobiol* **65**, 23–35 (2011).

- 1211 83. X. Zhang, Y.-F. Pei, L. Zhang, B. Guo, A. H. Pendegraft, W. Zhuang, N. Yi, Negative
1212 Binomial Mixed Models for Analyzing Longitudinal Microbiome Data. *Front. Microbiol.*
1213 **9** (2018).
- 1214 84. M. C. Mansilla, L. E. Cybulski, D. Albanesi, D. de Mendoza, Control of Membrane Lipid
1215 Fluidity by Molecular Thermosensors. *Journal of Bacteriology* **186**, 6681–6688 (2004).
- 1216 85. Y.-M. Zhang, C. O. Rock, Membrane lipid homeostasis in bacteria. *Nat Rev Microbiol* **6**,
1217 222–233 (2008).
- 1218 86. E. A. Dennis, Diversity of group types, regulation, and function of phospholipase A2.
1219 *Journal of Biological Chemistry* **269**, 13057–13060 (1994).
- 1220 87. D. M. Hall, K. R. Baumgardner, T. D. Oberley, C. V. Gisolfi, Splanchnic tissues undergo
1221 hypoxic stress during whole body hyperthermia. *American Journal of Physiology-*
1222 *Gastrointestinal and Liver Physiology* **276**, G1195–G1203 (1999).
- 1223 88. H. Hellmann, S. Mooney, Vitamin B6: A Molecule for Human Health? *Molecules* **15**,
1224 442–459 (2010).
- 1225 89. H. J. Powers, Riboflavin (vitamin B-2) and health¹². *The American Journal of Clinical*
1226 *Nutrition* **77**, 1352–1360 (2003).
- 1227 90. I. Gout, Coenzyme A, protein CoAlation and redox regulation in mammalian cells.
1228 *Biochemical Society Transactions* **46**, 721–728 (2018).
- 1229 91. S. C. Lu, Glutathione synthesis. *Biochimica et Biophysica Acta (BBA) - General Subjects*
1230 **1830**, 3143–3153 (2013).
- 1231 92. S. Dhungana, A. L. and Crumbliss, Coordination Chemistry and Redox Processes in
1232 Siderophore-Mediated Iron Transport. *Geomicrobiology Journal* **22**, 87–98 (2005).
- 1233 93. W. Ying, NAD⁺ /NADH and NADP⁺ /NADPH in Cellular Functions and Cell Death:
1234 Regulation and Biological Consequences. *Antioxidants & Redox Signaling* **10**, 179–206
1235 (2008).
- 1236 94. M. P. Kalapos, Methylglyoxal in living organisms: Chemistry, biochemistry, toxicology
1237 and biological implications. *Toxicology Letters* **110**, 145–175 (1999).
- 1238 95. M. Kilstrup, K. Hammer, P. Ruhdal Jensen, J. Martinussen, Nucleotide metabolism and its
1239 control in lactic acid bacteria. *FEMS Microbiol Rev* **29**, 555–590 (2005).
- 1240 96. A. N. Lane, T. W.-M. Fan, Regulation of mammalian nucleotide metabolism and
1241 biosynthesis. *Nucleic Acids Research* **43**, 2466–2485 (2015).
- 1242 97. M. Vinayak, C. Pathak, Queuosine modification of tRNA: its divergent role in cellular
1243 machinery. *Bioscience Reports* **30**, 135–148 (2009).

- 1244 98. J. A. Imlay, The molecular mechanisms and physiological consequences of oxidative
1245 stress: lessons from a model bacterium. *Nat Rev Microbiol* **11**, 443–454 (2013).
- 1246 99. J.-H. Martens, H. Barg, M. Warren, D. Jahn, Microbial production of vitamin B12. *Appl*
1247 *Microbiol Biotechnol* **58**, 275–285 (2002).
- 1248 100. P. Belenky, K. L. Bogan, C. Brenner, NAD⁺ metabolism in health and disease. *Trends in*
1249 *Biochemical Sciences* **32**, 12–19 (2007).
- 1250 101. C. L. Linster, E. Van Schaftingen, Vitamin C. *The FEBS Journal* **274**, 1–22 (2007).
- 1251 102. N. Arpaia, C. Campbell, X. Fan, S. Dikiy, J. van der Veeken, P. deRoos, H. Liu, J. R.
1252 Cross, K. Pfeffer, P. J. Coffey, A. Y. Rudensky, Metabolites produced by commensal
1253 bacteria promote peripheral regulatory T-cell generation. *Nature* **504**, 451–455 (2013).
- 1254 103. H.-C. Hung, S.-F. Tsai, H.-W. Chou, M.-J. Tsai, P.-L. Hsu, Y.-M. Kuo, Dietary fatty acids
1255 differentially affect secretion of pro-inflammatory cytokines in human THP-1 monocytes.
1256 *Sci Rep* **13**, 5511 (2023).
- 1257 104. A. Nakamura, M. Matsumoto, Role of polyamines in intestinal mucosal barrier function.
1258 *Semin Immunopathol* **47**, 9 (2025).
- 1259 105. P. Konopelski, I. Mogilnicka, Biological Effects of Indole-3-Propionic Acid, a Gut
1260 Microbiota-Derived Metabolite, and Its Precursor Tryptophan in Mammals' Health and
1261 Disease. *International Journal of Molecular Sciences* **23**, 1222 (2022).
- 1262 106. K. Hiippala, V. Kainulainen, M. Suutarinen, T. Heini, J. R. Bowers, D. Jasso-Selles, D.
1263 Lemmer, M. Valentine, R. Barnes, D. M. Engelthaler, R. Satokari, Isolation of Anti-
1264 Inflammatory and Epithelium Reinforcing Bacteroides and Parabacteroides Spp. from A
1265 Healthy Fecal Donor. *Nutrients* **12**, 935 (2020).
- 1266 107. B. J. Parker, P. A. Wearsch, A. C. M. Veloo, A. Rodriguez-Palacios, The Genus Alistipes:
1267 Gut Bacteria With Emerging Implications to Inflammation, Cancer, and Mental Health.
1268 *Front. Immunol.* **11** (2020).
- 1269 108. P. Louis, H. J. Flint, Formation of propionate and butyrate by the human colonic
1270 microbiota. *Environmental Microbiology* **19**, 29–41 (2017).
- 1271 109. F. Wu, X. Guo, J. Zhang, M. Zhang, Z. Ou, Y. Peng, Phascolarctobacterium faecium
1272 abundant colonization in human gastrointestinal tract. *Experimental and Therapeutic*
1273 *Medicine* **14**, 3122–3126 (2017).
- 1274 110. K. O. Alfarouk, A. H. H. Bashir, A. N. Aljarbou, A. M. Ramadan, A. K. Muddathir, S. T.
1275 S. AlHoufie, A. Hifny, G. O. Elhassan, M. E. Ibrahim, S. S. Alqahtani, S. D. AlSharari, C.
1276 T. Supuran, C. Rauch, R. A. Cardone, S. J. Reshkin, S. Fais, S. Harguindey, The Possible
1277 Role of Helicobacter pylori in Gastric Cancer and Its Management. *Front. Oncol.* **9**
1278 (2019).

- 1279 111. M. Rajilić-Stojanović, W. M. de Vos, The first 1000 cultured species of the human
1280 gastrointestinal microbiota. *FEMS Microbiology Reviews* **38**, 996–1047 (2014).
- 1281 112. S. Ahn, J. Jung, I.-A. Jang, E. L. Madsen, W. Park, Role of Glyoxylate Shunt in Oxidative
1282 Stress Response. *Journal of Biological Chemistry* **291**, 11928–11938 (2016).
- 1283 113. M. R. Badger, E. J. Bek, Multiple Rubisco forms in proteobacteria: their functional
1284 significance in relation to CO₂ acquisition by the CBB cycle. *Journal of Experimental*
1285 *Botany* **59**, 1525–1541 (2008).
- 1286 114. M. Bekker, G. Kramer, A. F. Hartog, M. J. Wagner, C. G. De Koster, K. J. Hellingwerf,
1287 M. J. Teixeira De Mattos, Changes in the redox state and composition of the quinone pool
1288 of *Escherichia coli* during aerobic batch-culture growth. *Microbiology* **153**, 1974–1980
1289 (2007).
- 1290 115. H.-M. Kuhn, U. Meier-Dieter, H. Mayer, ECA, the enterobacterial common antigen*.
1291 *FEMS Microbiology Reviews* **4**, 195–222 (1988).
- 1292 116. C. R. H. Raetz, C. Whitfield, Lipopolysaccharide endotoxins. *Annual Review of*
1293 *Biochemistry* **71**, 635 (2002).
- 1294 117. G. Caballero-Flores, J. M. Pickard, G. Núñez, Microbiota-mediated colonization
1295 resistance: mechanisms and regulation. *Nat Rev Microbiol* **21**, 347–360 (2023).
- 1296 118. F. Spragge, E. Bakkeren, M. T. Jahn, E. B. N. Araujo, C. F. Pearson, X. Wang, L.
1297 Pankhurst, O. Cunrath, K. R. Foster, Microbiome diversity protects against pathogens by
1298 nutrient blocking. *Science* **382**, eadj3502 (2023).
- 1299 119. C. G. Buffie, E. G. Pamer, Microbiota-mediated colonization resistance against intestinal
1300 pathogens. *Nat Rev Immunol* **13**, 790–801 (2013).
- 1301 120. S. Louca, M. F. Polz, F. Mazel, M. B. N. Albright, J. A. Huber, M. I. O’Connor, M.
1302 Ackermann, A. S. Hahn, D. S. Srivastava, S. A. Crowe, M. Doebeli, L. W. Parfrey,
1303 Function and functional redundancy in microbial systems. *Nat Ecol Evol* **2**, 936–943
1304 (2018).
- 1305 121. N. O. Kaakoush, N. Castaño-Rodríguez, H. M. Mitchell, S. M. Man, Global Epidemiology
1306 of *Campylobacter* Infection. *Clinical Microbiology Reviews* **28**, 687–720 (2015).
- 1307 122. J. Sáenz, Hopanoids as functional analogues of cholesterol in bacterial membranes. *PNAS*
1308 **112**, 11971–11976 (2015).
- 1309 123. H. Lin, E. Decuypere, J. Buyse, Acute heat stress induces oxidative stress in broiler
1310 chickens. *Comparative Biochemistry and Physiology Part A: Molecular & Integrative*
1311 *Physiology* **144**, 11–17 (2006).
- 1312 124. K. M. Ng, J. A. Ferreyra, S. K. Higginbottom, J. B. Lynch, P. C. Kashyap, S. Gopinath, N.
1313 Naidu, B. Choudhury, B. C. Weimer, D. M. Monack, J. L. Sonnenburg, Microbiota-

1314 liberated host sugars facilitate post-antibiotic expansion of enteric pathogens. *Nature* **502**,
1315 96–99 (2013).

1316 125. M. Y. Zeng, N. Inohara, G. Nuñez, Mechanisms of inflammation-driven bacterial
1317 dysbiosis in the gut. *Mucosal Immunology* **10**, 18–26 (2017).

1318 126. D. R. Donohoe, N. Garge, X. Zhang, W. Sun, T. M. O’Connell, M. K. Bunger, S. J.
1319 Bultman, The Microbiome and Butyrate Regulate Energy Metabolism and Autophagy in
1320 the Mammalian Colon. *Cell Metabolism* **13**, 517–526 (2011).

1321 127. C. Neuman, E. Hatje, K. Z. Zarkasi, R. Smullen, J. P. Bowman, M. Katouli, The effect of
1322 diet and environmental temperature on the faecal microbiota of farmed Tasmanian
1323 Atlantic Salmon (almo salar L.). *Aquaculture Research* **47**, 660–672 (2016).

1324 128. E. L. Soriano, D. T. Ramírez, D. R. Araujo, B. Gómez-Gil, L. I. Castro, C. G. Sánchez,
1325 Effect of temperature and dietary lipid proportion on gut microbiota in yellowtail kingfish
1326 *Seriola lalandi* juveniles. *Aquaculture* **497**, 269–277 (2018).

1327 129. K. D. Kohl, J. Yahn, Effects of environmental temperature on the gut microbial
1328 communities of tadpoles. *Environmental Microbiology* **18**, 1561–1565 (2016).

1329 130. C. Nathan, A. Cunningham-Bussel, Beyond oxidative stress: an immunologist’s guide to
1330 reactive oxygen species. *Nat Rev Immunol* **13**, 349–361 (2013).

1331 131. J. D. Orkin, S. E. Webb, A. D. Melin, Small to modest impact of social group on the gut
1332 microbiome of wild Costa Rican capuchins in a seasonal forest. *American Journal of*
1333 *Primatology* **81**, e22985 (2019).

1334 132. L. Li, P. Peng, N. Ding, W. Jia, C. Huang, Y. Tang, Oxidative Stress, Inflammation, Gut
1335 Dysbiosis: What Can Polyphenols Do in Inflammatory Bowel Disease? *Antioxidants* **12**,
1336 967 (2023).

1337 133. H. Mostafavi Abdolmaleky, J.-R. Zhou, Gut Microbiota Dysbiosis, Oxidative Stress,
1338 Inflammation, and Epigenetic Alterations in Metabolic Diseases. *Antioxidants* **13**, 985
1339 (2024).

1340 134. D. Dodd, M. H. Spitzer, W. Van Treuren, B. D. Merrill, A. J. Hryckowian, S. K.
1341 Higginbottom, A. Le, T. M. Cowan, G. P. Nolan, M. A. Fischbach, J. L. Sonnenburg, A
1342 gut bacterial pathway metabolizes aromatic amino acids into nine circulating metabolites.
1343 *Nature* **551**, 648–652 (2017).

1344 135. T. Zelante, R. G. Iannitti, C. Cunha, A. De Luca, G. Giovannini, G. Pieraccini, R. Zecchi,
1345 C. D’Angelo, C. Massi-Benedetti, F. Fallarino, A. Carvalho, P. Puccetti, L. Romani,
1346 Tryptophan Catabolites from Microbiota Engage Aryl Hydrocarbon Receptor and Balance
1347 Mucosal Reactivity via Interleukin-22. *Immunity* **39**, 372–385 (2013).

1348 136. S. H. Chu, J. W. Lim, K. H. Kim, H. Kim, NF- κ B and Bcl-2 in Helicobacter pylori-
1349 Induced Apoptosis in Gastric Epithelial Cells. *Annals of the New York Academy of*
1350 *Sciences* **1010**, 568–572 (2003).

1351

1352