




predicted Abundance for chorismate biosynthesis I




predicted Abundance for L-arginine degradation II (AST pathway)




predicted Abundance for enterobacterial common antigen biosynthesis




predicted Abundance for glucose and glucose-1-phosphate degradation




predicted Abundance for superpathway of glycolysis, pyruvate dehydrogenase, TCA, and glyo




predicted Abundance for glyoxylate cycle




predicted Abundance for superpathway of (Kdo)2-lipid A biosynthesis




predicted Abundance for methylerythritol phosphate pathway I




predicted Abundance for superpathway of ornithine degradation




predicted Abundance for L-glutamate degradation V (via hydroxyglutarate)




predicted Abundance for pyruvate fermentation to acetate and (S)-lactate I




predicted Abundance for incomplete reductive TCA cycle




predicted Abundance for peptidoglycan biosynthesis I (meso-diaminopimelate containing)




predicted Abundance for folate transformations I




predicted Abundance for L-lysine biosynthesis II




predicted Abundance for 4-aminobutanoate degradation V




predicted Abundance for pyruvate fermentation to acetate and lactate II




predicted Abundance for 2-oxobutanoate degradation I




predicted Abundance for fatty acid β-oxidation IV (unsaturated, even number)




predicted Abundance for L-arginine biosynthesis III (via N-acetyl-L-citrulline)




predicted Abundance for tetrapyrrole biosynthesis I (from glutamate)




predicted Abundance for mannosylglycerate biosynthesis I




predicted Abundance for Rubisco shunt




predicted Abundance for ubiquinol-7 biosynthesis (early decarboxylation)




predicted Abundance for stearate biosynthesis II (bacteria and plants)




predicted Abundance for S-adenosyl-L-methionine salvage I




predicted Abundance for superpathway of adenosylcobalamin salvage from cobinamide II




predicted Abundance for superpathway of fatty acids biosynthesis (E. coli)




predicted Abundance for L-lysine degradation X




predicted Abundance for peptidoglycan biosynthesis III (mycobacteria)




predicted Abundance for UDP-N-acetylmuramoyl-pentapeptide biosynthesis II (lysine-containing)




predicted Abundance for UDP-N-acetylmuramoyl-pentapeptide biosynthesis I (meso-diaminopimelate)




predicted Abundance for stachyose degradation




predicted Abundance for chondroitin sulfate degradation I (bacterial)




predicted Abundance for preQ0 biosynthesis




predicted Abundance for ubiquinol-8 biosynthesis (early decarboxylation)




predicted Abundance for L-N Δ -acetylornithine biosynthesis




predicted Abundance for dTDP-3-acetamido- α -D-fucose biosynthesis




predicted Abundance for fatty acid salvage




predicted Abundance for pyridoxal 5'-phosphate salvage II (plants)




predicted Abundance for guanosine ribonucleotides de novo biosynthesis




predicted Abundance for mitochondrial NADPH production (yeast)




predicted Abundance for 1,4-dihydroxy-6-naphthoate biosynthesis II




predicted Abundance for 1,3-propanediol biosynthesis (engineered)




predicted Abundance for L-arginine biosynthesis IV (archaeobacteria)




predicted Abundance for sulfoquinovose degradation I




predicted Abundance for methylerythritol phosphate pathway II




predicted Abundance for oleate biosynthesis IV (anaerobic)




predicted Abundance for (aminomethyl)phosphonate degradation




predicted Abundance for glyphosate degradation III




predicted Abundance for L-threonate degradation




predicted Abundance for superpathway of menaquinol-8 biosynthesis III




predicted Abundance for 5'-deoxyadenosine degradation II




predicted Abundance for L-glutamate degradation XI (reductive Stickland reaction)




predicted Abundance for oleate β-oxidation




predicted Abundance for polymyxin resistance




predicted Abundance for methylphosphonate degradation I




predicted Abundance for L-ascorbate degradation I (bacterial, anaerobic)




predicted Abundance for L-lysine degradation I




predicted Abundance for (5Z)-dodecenoate biosynthesis I




predicted Abundance for fatty acid β-oxidation VI (mammalian peroxisome)




predicted Abundance for NAD de novo biosynthesis I (from aspartate)



predicted Abundance for superpathway of L-threonine metabolism



predicted Abundance for L-tryptophan biosynthesis



predicted Abundance for superpathway of ubiquinol-8 biosynthesis (early decarboxylation)

