


eFigure 2: **<Subgroup> estimate beside the whole sample.** Each point is the same model's estimate with its 95% interval, in everyone and in <the subgroup>; the dashed line marks zero. Placeholder figure of Tools' Paper-Example; no real data.

What this display says

In <the subgroup> the association with <the policy> is larger than in the whole sample, and its interval is wider, so the paper's answer to RQ2, for whom it changes, rests on fewer units.

How this display was made

Data

<The data source>, years <YYYY to YYYY>: the work Runs b90.j02.t01.r01 and b90.j02.t02.r01, each its result/values.csv (placeholder).

Sample

- the whole sample, 12480 units, as in Display A
- <the subgroup>, 3120 units

Method

The main model, fitted once in everyone and once in <the subgroup>; each estimate with its 95% interval.

Made by

- `r01_example_main/result/values.csv` (main.b, main.ci_lo, main.ci_hi, sample.n_units), the Run of `work/b90_example_paper_results/j02_example_estimates/t01_example_main/`
- `r01_example_subgroup/result/values.csv` (sub.b, sub.ci_lo, sub.ci_hi, sub.n_units), the Run of `work/b90_example_paper_results/j02_example_estimates/t02_example_subgroup/`
- `scripts/draw_subgroup_estimate.py`, the worker of `t51_display-d-subgroup-estimate/`

Numbers

Every point and interval is read from the two values.csv files above; nothing is typed in.