lxNavbar
lx-
Header
lx-
Footer

Hyperconvexity and domains of discontinuity for surface groups in $SL_3\mathbb{C}$

BY

Jennifer Vaccaro
BS, Olin College of Engineering, 2017
MS, University of Illinois at Chicago, 2022

THESIS [PhD candidates may use DISSERTATION here instead]

Submitted as partial fulfillment of the requirements
for the degree of Doctorate of Philosophy in Mathematics
in the Graduate College of the
University of Illinois Chicago, 2026

Chicago, Illinois

Defense Committee:
Emily Dumas
Name
Name
Name, Outside Members Need Affiliation Here

Accessibility Statement

An accessible EPUB version of this document is available at [URL] or can be obtained by contacting [contact details].

Dedication

In memory of my grandmother Martha, who shared her love of mathematics with me and many others.

Acknowledgements

Emily (integrity, knowledge, high standards)
Sam (love, support, meals, no pressure)
Family (you don't know why I wanted to do this, but that's ok)
Friends (kept me company, kept me happy)
Other math teachers and mentors
Gertrude (tiniest companion)

Preface

Duis aute irure dolor in reprehenderit in voluptate velit esse cillum dolore eu fugiat nulla pariatur. Excepteur sint occaecat cupidatat non proident, sunt in culpa qui officia deserunt mollit anim id est laborum.

Curabitur pretium tincidunt lacus. Nulla gravida orci a odio. Nullam varius, turpis et commodo pharetra, est eros bibendum elit, nec luctus magna felis sollicitudin mauris.

Integer in mauris eu nibh euismod gravida. Duis ac tellus et risus vulputate vehicula. Donec lobortis risus a elit. Etiam tempor. Ut ullamcorper, ligula eu tempor congue, eros est euismod turpis, id tincidunt sapien risus a quam.

Contents

Accessibility Statement ii

Dedication iii

Acknowledgements iv

Preface vi

List of Figures x

List of Tables xi

List of Abbreviations xii

Summary xiv

Chapter 1. Introduction 1
1. Motivation 2

Chapter 2. Background 4
1. Anosov representations in $SL_3\mathbb{C}$ 5
2. The subvarieties of $\mathcal{F}^3$ 5
3. Richardson varieties in $\mathcal{F}^n$ for $n \geq 4$ 6

Chapter 3. Hyperconvexity is closed under topological conjugacy 8
1. Assumptions and notation 9
2. Hyperconvexity is closed under topological conjugacy 9
3. Proof of Theorem 3.1. 10
4. Action of $\rho(g)$ on $\mathcal{F}^3$ 10
5. F preserves the Richardson varieties of pairs $\xi(g^+), \xi(g^-)$ 11
6. F preserves Richardson varieties of Λ 12
7. F matches or switches Richardson varieties 13
8. Random remarks 14

Chapter 4. Anosov components have one topological conjugacy class 15
1. Assumptions and notation 16
2. Anosov components have one topological conjugacy class 16
3. Key ingredients for proof 16
4. Proof of Theorem 4.1 17
5. Topological conjugacies of reducible representations 18

Chapter 5. Vitae 20

Appendix A. Technical Lemmas 22

Bibliography 24

List of Figures

1 This is a torus 2

2 The Snake Lemma 2

1 Graph representing the cyclic connectivity of Richardson 0- and 1-varieties for x and y transverse. 7

List of Tables

1 Example Table 3

List of Abbreviations

- $\mathbb{R}$ The real numbers
- $\mathbb{C}$ The complex numbers

Summary

Cras mollis scelerisque nunc. Nullam arcu. Aliquam consequat. Curabitur augue lorem, dapibus quis, laoreet et, pretium ac, nisi. Aenean magna nisl, mollis quis, molestie eu, feugiat in, orci. In hac habitasse platea dictumst.

Maecenas fermentum consequat mi. Donec fermentum. Pellentesque malesuada nulla a mi. Duis sapien sem, aliquet nec, commodo eget, consequat quis, neque. Aliquam faucibus, elit ut dictum aliquet, felis nisl adipiscing sapien, sed malesuada diam lacus eget erat.

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Vivamus luctus urna sed urna ultricies ac tempor dui sagittis. In condimentum facilisis porta.

Sed nec diam eu diam mattis viverra. Nulla fringilla, orci ac euismod semper, magna diam porttitor mauris, quis sollicitudin sapien justo in libero.

Fusce lacinia quam nisi, euismod ultrices nulla malesuada ut. Curabitur sit amet magna quam. Praesent in libero vel turpis pellentesque egestas sit amet vel nunc.

Morbi lectus risus, iaculis vel, suscipit quis, luctus non, massa. Fusce ac turpis quis ligula lacinia aliquet. Mauris ipsum. Nulla metus metus, ullamcorper vel, tincidunt sed, euismod in, nibh.

Quisque volutpat condimentum velit. Class aptent taciti sociosqu ad litora torquent per conubia nostra, per inceptos himenaeos. Nam nec ante. Sed lacinia, urna non tincidunt mattis, tortor neque adipiscing diam, a cursus ipsum ante quis turpis.

Cras elementum ultrices diam. Maecenas ligula massa, varius a, semper congue, euismod non, mi. Proin porttitor, orci nec nonummy molestie, enim est eleifend mi, non fermentum diam nisl sit amet erat.

Duis semper. Duis arcu massa, scelerisque vitae, consequat in, pretium a, enim. Pellentesque congue. Ut in risus volutpat libero pharetra tempor. Cras vestibulum bibendum augue.

Praesent egestas leo in pede. Praesent blandit odio eu enim. Pellentesque sed dui ut augue blandit sodales. Vestibulum ante ipsum primis in faucibus orci luctus et ultrices posuere cubilia Curae.

Aliquam nibh. Mauris ac mauris sed pede pellentesque fermentum. Maecenas adipiscing ante non diam sodales hendrerit.

Ut velit mauris, egestas sed, gravida nec, ornare ut, mi. Aenean ut orci vel massa suscipit pulvinar. Nulla sollicitudin. Fusce varius, ligula non tempus aliquam, nunc turpis ullamcorper nibh, in tempus sapien eros vitae ligula.

Pellentesque rhoncus nunc et augue. Integer id felis. Curabitur aliquet pellentesque diam. Integer quis metus vitae elit lobortis egestas.

Lorem ipsum dolor sit amet, consectetuer adipiscing elit. Morbi vel erat non mauris convallis vehicula. Nulla et sapien. Integer tortor tellus, aliquam faucibus, convallis id, congue eu, quam.

Mauris ullamcorper felis vitae erat. Proin feugiat, augue non elementum posuere, metus purus iaculis lectus, et tristique ligula justo vitae magna.

Aliquam convallis sollicitudin purus. Praesent aliquam, enim at fermentum mollis, ligula massa adipiscing nisl, ac euismod nibh nisl eu lectus.

Fusce vulputate sem at sapien. Vivamus leo. Aliquam euismod libero eu enim. Nulla nec felis sed leo placerat imperdiet. Aenean suscipit nulla in justo.

Suspendisse cursus rutrum augue. Nulla tincidunt tincidunt mi. Curabitur iaculis, lorem vel rhoncus faucibus, felis magna fermentum augue, et ultricies lacus lorem varius purus.

Sed adipiscing ornare risus. Morbi est est, blandit sit amet, sagittis vel, euismod vel, velit. Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas.

Nullam nonummy metus. Vestibulum volutpat pretium libero. Cras id dui. Aenean ut eros et nisl sagittis vestibulum.

Nulla facilisi. Integer lacinia sollicitudin massa. Cras metus. Sed aliquet risus a tortor. Integer id quam.

CHAPTER 1

Introduction



FIGURE 1. This is a torus

$$\begin{tikzcd}
 & 0 \arrow[d] & 0 \arrow[d] & 0 \arrow[d] & \\
0 \arrow[r] & A' \arrow[r, "f'"] \arrow[d, "a"'] & B' \arrow[r, "g'"] \arrow[d, "b"'] & C' \arrow[r] \arrow[d, "c"'] & 0 \\
0 \arrow[r] & A \arrow[r, "f"] \arrow[d, "a''"'] & B \arrow[r, "g"] \arrow[d, "b''"'] & C \arrow[r] \arrow[d, "c''"'] & 0 \\
0 \arrow[r] & A'' \arrow[r, "f''"] \arrow[d] & B'' \arrow[r, "g''"] \arrow[d] & C'' \arrow[r] \arrow[d] & 0 \\
 & 0 & 0 & 0 &
\end{tikzcd}$$

FIGURE 2. The Snake Lemma

This is the introduction chapter. We cite some classic works [**3, 4**].

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Morbi tincidunt nisi sed neque volutpat, et imperdiet metus congue. Donec interdum, neque at varius vehicula, elit odio tempus sapien, sed finibus lorem felis nec turpis. Aliquam erat volutpat. Mauris vehicula imperdiet erat, non aliquet magna commodo id.

Sed porttitor urna ac ipsum rhoncus, vel dictum risus commodo. Integer ultrices dui sit amet pulvinar tempor. Phasellus nec tellus eu turpis sodales accumsan. Proin ultr

THEOREM 1.1. *This is a theorem*

We reference Theorem 1.1.

University of Illinois Chicago.

1. Motivation

1.1. Historical context. A brief overview of how the problem developed.

$$\int_0^1 f(x)dx = 2 \tag{1}$$

How to solve (1)

1.2. Open questions. Some questions remain open for future work.
Note: I have not tested the accessibility of this table.

Monkeys	Lions
100	200

Table 1. Example Table

CHAPTER 2

Background

1. Anosov representations in $SL_3\mathbb{C}$

Consider the collection of Anosov representations

$$\mathcal{A}_{\mathbb{R}}(\Gamma) := \{\rho : \Gamma \to SL_3\mathbb{R} \text{ Anosov}\}, \text{ and } \mathcal{A}(\Gamma) := \{\rho : \Gamma \to SL_3\mathbb{C} \text{ Anosov}\}. \tag{2}$$

Let $\rho_0 : \Gamma \to PSL_2\mathbb{R}$ Fuchsian and $\rho_i : \Gamma \to PSL_2\mathbb{C}$ quasi-Fuchsian. To create Anosov representations in $SL_3\mathbb{C}$, we can compose with an irreducible representation $\iota : PSL_2\mathbb{R} \to SL_3\mathbb{R}$:

$$\rho_F = \iota \circ \rho_0 : \Gamma \to SL_3\mathbb{R}, \text{ and } \rho_{QF} = \iota \circ \rho_i : \Gamma \to SL_3\mathbb{C}. \tag{3}$$

Similarly, (assuming that ρ_0, ρ_i lift to $SL_2\mathbb{R}$) we can compose with a reducible representation $\Delta : SL_2\mathbb{C} \to SL_3\mathbb{C}$:

$$\rho_R = \Delta \circ \rho_0 : \Gamma \to SL_3\mathbb{R}, \text{ and } \rho_{QR} = \Delta \circ \rho_i : \Gamma \to SL_3\mathbb{C}. \tag{4}$$

The representations ρ_{Fuchs} and ρ_{Red} are Anosov. We can then define the following subsets of $\mathcal{A}$:

- $SL_3\mathbb{C}$ quasi-Fuchsian representations $\mathcal{H}(\Gamma) := \{\rho \in \mathcal{A}(\Gamma) : \rho = \iota\circ$in the same Anosov component a
- Hitchin representations $\mathcal{H}(\Gamma) := \{\rho \in \mathcal{A}_{\mathbb{R}}(\Gamma) : \rho$in the same Anosov component as $\rho_0\}$
- Quasi-Hitchin representations
- $SL_3\mathbb{C}$-reducible representations
- Barbot representations
- Quasi-Barbot representations

2. The subvarieties of $\mathcal{F}^3$

2.1. Richardson varieties in $\mathcal{F}^3$. For any flag $x \in \mathcal{F}^3$, we define $Sch(x)$ to be the *principle Schubert varieties* centered at x. These are the closure of the principle Schubert cells, each identified by an element of the Weyl group $W(G) = A_2 = Sym(3)$. Let $x = (l, H) \in \mathcal{F}^3$ where $l \in \mathbb{CP}^2$ and $H \in (\mathbb{CP}^2)^*$ such that $l \subset H$. Then $Sch(x)$ consists of:

- $\alpha_{Id}(x) = \{(l, H)\} = \{x\}$,
- $\alpha_1(x) = \{(l, H_l) : l \subset H_l\}$,
- $\alpha_2(x) = \{(l_H, H) : l_H \subset H\}$,
- $\alpha_{12}(x) = \{(k, H_l) : l \subset H_l\}$,
- $\alpha_{21}(x) = \{(l_H, G) : l_H \subset H\}$, and
- $\alpha_{121}(x) = \alpha_{212}(x) = \mathcal{F}^3$.

We also define Sch as the *types* of principle Schubert varieties, e.g. α_1 or α_2, without specifying a basepoint x. For $g \in W(G)$, the complex dimension of $\alpha_g \in Sch$ is length of g in the Bruhat order. More explicitly, $dim(\alpha_{Id}) = 0$, $dim(\alpha_1) = dim(\alpha_2) = 1$, $dim(\alpha_{12}) = dim(\alpha_{21}) = 2$, and $dim(\alpha_{121}) = 3$.

We call a pair of flags $x, y \in \mathcal{F}^3$ *transverse* if they are in general position, meaning that for any proper, principle Schubert variety α, then $x \notin \alpha(y)$ and $y \notin \alpha(x)$.

For distinct flags $x, y \in \mathcal{F}^3$ such that either $l_x = l_y$ or $H_x = H_y$, then we define the *span* of x and y as follows:

$$span(x, y) := \begin{cases} \alpha_1(x) = \alpha_1(y) : l_x = l_y \\ \alpha_2(x) = \alpha_2(y) : H_x = H_y. \end{cases} \tag{5}$$

2.2. Richardson 0-varieties. *TODO: define Richardson variety more generally, then specialize to irreducible 0-varieties and irreducible 1-varieties.*

Let $x, y \in \mathcal{F}^3$ transverse. Define $R^0_{x,y}$ to be their irreducible *0-dimensional Richardson varieties*, which are the nonempty intersections between principle Schubert varieties $\alpha(x) \cap \beta(y)$ for $\alpha, \beta \in Sch$ such that $dim(\alpha) + dim(\beta) = dim(\mathcal{F}^3)$. There are exactly six such varieties $\{x, x_y, x^y, y_x, y^x, y\}$, where:

- $x := \alpha_{Id}(x) \cap \alpha_{121}(y)$,
- $x_y := \alpha_1(x) \cap \alpha_{12}(y)$,
- $x^y := \alpha_2(x) \cap \alpha_{21}(y)$,
- $y_x := \alpha_{12}(x) \cap \alpha_1(y)$,
- $y^x := \alpha_{21}(x) \cap \alpha_2(y)$, and
- $y := \alpha_{121}(x) \cap \alpha_{Id}(y)$.

Given two tranverse flags, the Richardson 0-varieties may be computed explicitly.

Example 2.1. *Choose x and y are the attracting and repelling flags of $\rho(g) = diag(\lambda, \mu, \nu)$ where $|\lambda| > |\mu| > |\nu|$. Then the six Richardson 0-varieties of x and y are $x = (e_1, e_1 + e_2)$, $x_y = (e_1, e_1 + e_3)$, $x^y = (e_2, e_1 + e_2)$, $y_x = (e_3, e_1 + e_3)$, $y^x = (e_2, e_2 + e_3)$, and $y = (e_3, e_2 + e_3)$.*

2.3. Richardson 1-varieties. Let $x, y \in \mathcal{F}^3$ transverse, and define $R^1_{x,y}$ to be their irreducible *1-dimensional Richardson varieties*. These are nonempty intersections between principle Schubert varieties $\alpha(x) \cap \beta(y)$ for $\alpha, \beta \in Sch$ such that $dim(\alpha) + dim(\beta) = dim(\mathcal{F}^3) + 1$. There are six such 1-varieties, each homeomorphic to $\mathbb{CP}^1$, which we distinguish using pairs of elements of $R^0_{x,y}$.

- $span(x, x_y) := \alpha_1(x) \cap \alpha_{121}(y)$,
- $span(x, x^y) := \alpha_2(x) \cap \alpha_{121}(y)$,
- $span(x_y, y_x) := \alpha_{12}(x) \cap \alpha_{12}(y)$,
- $span(x^y, y^x) := \alpha_{21}(x) \cap \alpha_{21}(y)$,
- $span(y_x, y) := \alpha_{121}(x) \cap \alpha_1(y)$, and
- $span(y^x, y) := \alpha_{121}(x) \cap \alpha_2(y)$.

The 1-varieties are not pairwise disjoint, e.g. $\{x\} = span(x, x_y) \cap span(x, x^y)$. The intersections of adjacent Richardson 1-varieties are exactly the Richardson 0-varieties. The connectivity of the 0- and 1-varieties is cyclic in the following way:

$$x \in span(x, x_y) \ni x_y \in span(x_y, y_x) \ni y_x \in span(y, y_x) \ni y \tag{6}$$

$$x \in span(x, x^y) \ni x^y \in span(x^y, y^x) \ni y^x \in span(y, y^x) \ni y. \tag{7}$$

Equivalently, in Figure 2.3 we describe the connectivity using a graph with six vertices representing 0-varieties and six edges representing 1-varieties. Two vertices are connected if the span of their 0-varieties is an irreducible 1-variety.

Remark 2.1. *If $x, y \in \mathcal{F}^3$ transverse and $\alpha, \beta \in Sch$ such that*

$$dim(\alpha) + dim(\beta) < dim(\mathcal{F}^3), \tag{8}$$

then $\alpha(x) \cap \beta(y) = \emptyset$.

3. Richardson varieties in $\mathcal{F}^n$ for $n \geq 4$

Conjectural picture for Richardson varieties in higher dimensions:



FIGURE 1. Graph representing the cyclic connectivity of Richardson 0- and 1-varieties for x and y transverse.

I suspect that topological conjugacies of Borel Anosov representations will preserve Richardson varieties up to $R^{n-2}_{x,y}$ organized into a cell decomposition with data an $(n-1)$-valent graph where vertices correspond to 0-varieties, edges correspond to 1-varieties, fully connected triples correspond to 2-varieties, etc. Matching/switching Richardson varieties will mean sending Richardson varieties to their dual.

I think balanced ideals I correspond to subgraphs of the Richardson variety graph containing exactly half of the vertices. In other words, $Th^I(x) \cap Th^I(y) = \emptyset$, and

$$(Th^I(x) \cap R^0_{x,y}) \cup (Th^I(y) \cap R^0_{x,y}) = R^0_{x,y}. \tag{9}$$

However, topological conjugacies should not depend on the choice of balanced ideal, so then they determine diffeomorphisms between all types of quotient manifolds. In other words, if $F : \mathcal{F}^n \to \mathcal{F}^n$ a topological conjugacy between ρ, ρ', then:

$$F(M^I_\rho) = M^I_{\rho'} \text{ for all balanced ideals } I \subset W(SL_n\mathbb{C}). \tag{10}$$

Most importantly, I believe the conclusions of Theorem 3.1 will apply in higher rank:

CONJECTURE 2.1. *For $\rho, \rho' : \Gamma \to SL_n\mathbb{C}$ Borel Anosov and topologically conjugate. Then ρ hyperconvex if and only if ρ' hyperconvex.*

CHAPTER 3

Hyperconvexity is closed under topological conjugacy

1. Assumptions and notation

Let Γ be the fundamental group of a closed hyperbolic surface, and let

$$\rho, \rho' : \Gamma \to SL_3\mathbb{C} \tag{11}$$

Anosov representations. We use the following notation related to ρ and ρ'.

- $\xi, \xi' : \partial\Gamma \to \mathcal{F}^3$ are the continuous, transverse, ρ- or ρ'- equivariant, dynamics-preserving boundary maps.
- $\Lambda, \Lambda' \subset \mathcal{F}^3$ are the limit curves, which are the images of ξ, ξ'.
- $Th(x) = \alpha_1(x) \cup \alpha_2(x)$ is the balanced ideal thickening of some $x \in \mathcal{F}^3$.
- $Th(\Lambda) = \cup_{x\in\Lambda} Th(x)$ is the thickening of the limit curve.

We call ρ, ρ' *topologically conjugate* if there exists a homeomorphism $F : \mathcal{F}^3 \to \mathcal{F}^3$ such that for all $g \in \Gamma$ and all $x \in \mathcal{F}^3$, then $F(\rho(g) \cdot x) = \rho'(g) \cdot F(x)$. We call F the *topological conjugacy* between ρ and ρ'.

Recall that ρ is *hyperconvex* if for all $x, y, z \in \Lambda^{(3)}$ then $l_z \not\subset l_x + l_y$ and $(H_x \cap H_y) \not\subset H_z$.

For $x, y \in \mathcal{F}^3$ transverse, recall the Richardson varieties of x and y, denoted by $R^0_{x,y}$ and $R^1_{x,y}$.

2. Hyperconvexity is closed under topological conjugacy

Theorem 3.1. *If ρ and ρ' are topologically conjugate, then ρ is hyperconvex if and only if ρ' is hyperconvex.*

We use two supporting lemmas to prove this theorem. First, we show that the relative position of 0-dimensional Richardson varieties of Λ determines whether ρ is hyperconvex.

Lemma 3.1.1. *An Anosov representation ρ is hyperconvex* if and only if *for all $x, y, z \in \Lambda^{(3)}$ then $R^0_{x,y} \cap R^0_{x,z} = \{x\}$.*

For the second lemma, we assume that $F : \mathcal{F}^3 \to \mathcal{F}^3$ is a topological conjugacy between ρ and ρ'. Clearly $F(\Lambda) = \Lambda'$. This mapping is pointwise, i.e. for all $\alpha \in \partial\Gamma$, then $F(\xi(\alpha)) = \xi'(\alpha)$. So for $x \in \Lambda$, we can use the notation $x' := F(x) \in \Lambda'$.

We say F *preserves* the Richardson varieties of Λ if for all $x, y \in \Lambda^{(2)}$, then $F(R^0_{x,y}) = R^0_{x',y'}$ and $F(R^1_{x,y}) = R^1_{x',y'}$.

We will show that F preserves the Richardson varieties of Λ in one of two ways:

- We say F *matches* Richardson varieties if for all $x, y \in \Lambda^{(2)}$, then $F(x_y) = x'_{y'}$ and $F(span(x, x_y)) = span(x, x'_{y'})$.
- We say F *switches* Richardson varieties if for all $x, y \in \Lambda^{(2)}$, then $F(x_y) = x'^{y'}$ and $F(span(x, x_y)) = span(x, x'^{y'})$.

Lemma 3.1.2. *If F is a topological conjugacy between ρ and ρ', then F preserves the Richardson varieties of Λ. Furthermore, F either matches or switches Richardson varieties.*

We also highlight the following consequence of the argument in Lemma 3.1.2:

Corollary 3.1.1. *For all $x \in \Lambda$, then $F(Th(x)) = Th(x')$.*

3. Proof of Theorem 3.1.

We first prove our main theorem using our supporting lemmas.

Proof. (Theorem 3.1) Suppose $F : \mathcal{F}^3 \to \mathcal{F}^3$ is a topological conjugacy between Anosov representations $\rho, \rho' :\to SL_3\mathbb{C}$.

We begin with the forward direction. Assume that ρ is hyperconvex. Let $x, y, z \in \Lambda^{(3)}$. By Lemma 3.1.1, then $R^0_{x,y} \cap R^0_{x,z} = \{x\}$. By Lemma 3.1.2, then

$$F(R^0_{x,y}) = R^0_{x',y'}, \tag{12}$$

$$F(R^0_{x,z}) = R^0_{x',z'}. \tag{13}$$

Since F is a homeomorphism, then

$$R^0_{x',y'} \cap R^0_{x',z'} = F(R^0_{x,y} \cap R^0_{x,z}) \tag{14}$$

$$= F(x) \tag{15}$$

$$= \{x\}. \tag{16}$$

Since x, y, z were general, then this holds for all triples in $x', y', z' \in \Lambda'$, and ρ' is hyperconvex.

The other direction proceeds similarly. Assume that ρ is *not* hyperconvex. By Lemma 3.1.1, there exist $x, y, z \in \Lambda^{(3)}$ such that $R^0_{x,y} \cap R^0_{x,z} \neq \{x\}$. By Lemma 3.1.2, then

$$R^0_{x',y'} \cap R^0_{x',z'} = F(R^0_{x,y}) \cap F(R^0_{x,z}) \tag{17}$$

$$= F(R^0_{x,y} \cap F^(R^0_{x,z})) \qquad \neq F(x). \tag{18}$$

So in particular $R^0_{x',y'} \cap R^0_{x',z'} \neq \{x'\}$, so ρ' is *not* hyperconvex. □

We also prove our lemma that hyperconvexity is equivalent to the relative position of the Richardson 0-varieties of Λ.

Proof. (Lemma 3.1.1) Suppose ρ is Anosov. Let $x, y, z \in \Lambda^{(3)}$. Since x, y, z are pairwise transverse, then clearly

$$y_x, y^x, z_x, z^x \notin R^0_{x,y} \cap R^0_{x,z}. \tag{19}$$

To show that $R^0_{x,y} \cap R^0_{x,z} = \{x\}$, it suffices to check the pairs $x_y \neq x_z \in \alpha_1(x)$ and $x^y \neq x^z \in \alpha_2(x)$.

By definition, $x_y = (l_x, l_x + l_y)$ and $x_z = (l_x, l_x + l_z)$, which are distinct if and only if $l_x + l_y \neq l_x + l_z$, or equivalently if and only if $l_z \not\subset l_x + l_y$.

Similarly, $x^y = (H_x \cap H_y, H_x)$ and $x^z = (H_x \cap H_z, H_x)$, which are distinct if and only if $H_x \cap H_y \neq H_x \cap H_z$, or equivalently if and only if $H_x \cap H_y \not\subset H_z$.

So $R^0_{x,y} \cap R^0_{x,z} = \{x\}$ for all $x, y, z \in \Lambda^{(3)}$ is equivalent to ρ hyperconvex. □

The remainder of the chapter is devoted to the proof of Lemma 3.1.2, which says that a topological conjugacy F preserves irreducible Richardson varities of Λ.

4. Action of $\rho(g)$ on $\mathcal{F}^3$

In this section we describe the action of $\rho(g)$ on $\mathcal{F}^3$ for $g \in \Gamma$ infinite order. Then g has attracting and repelling group boundary points which we call $g^+, g^- \in \partial\Gamma$, respectively. Since ρ is Anosov, then $\rho(g)$ is strongly loxodromic, i.e. its eigenvalues have distinct magnitude. That means $\rho(g)$ has attracting and repelling limit curve points which we call $x := \xi(g^+)$ and $y := \xi(g^-)$, respectively.

Under the action $\rho(g)$, most of $\mathcal{F}^3$ is attracted to x and repelled by y. We define the *uphill* of x as the subset attracted to x under $\rho(g)$:

$$U_{\rho(g)}(x) := \{f \in F^3 : \rho(g^n) \cdot f \to x\}. \tag{20}$$

Similarly, we define the *downhill* of y as the subset repelled by y under $\rho(g)$, or equivalently the subset attracted by y under $\rho(g^{-1})$:

$$D_{\rho(g)}(y) = U_{rho(g^{-1})}(y) := \{f \in \mathcal{F}^3 : \rho(g^{-n}) \cdot f \to y\}. \tag{21}$$

We then define the *exceptional set* of $\rho(g)$ as the subset *not* attracted to x or *not* repelled by y under $\rho(g)$:

$$E_{\rho(g)} = \mathcal{F}^3 \setminus (U_{\rho(g)}(x) \cap D_{\rho(g)}(y)). \tag{22}$$

The set $E_{\rho(g)}$ is nonempty and closed, since it is complement of an open set. It contains the fixed points of $\rho(g) \cdot \mathcal{F}^3$, which are precisely $R^0_{x,y}$. It suffices to check this for $x, y, \rho(g)$ as in Example 2.1, since $\rho(g)$ is conjugate to $diag(\lambda, \mu, \nu)$.

We can partition $E_{\rho(g)} \setminus R^0_{x,y}$ into connected subsets using the attracting and repelling fixed points. By taking the closure of these subsets, we recover the 1-dimensional Richardson varieties $R^1_{x,y}$.

- $span(x, x_y) = \alpha_1(x) = \overline{U(x) \cap D(x_y)}$
- $span(x, x^y) = \alpha_2(x) = \overline{U(x) \cap D(x^y)}$
- $span(x_y, y_x) = \overline{U(x_y) \cap D(y_x)}$
- $span(x^y, y^x) = \overline{U(x^y) \cap D(y^x)}$
- $span(y_x, y) = \alpha_1(y) = \overline{U(y_x) \cap D(y)}$
- $span(y^x, y) = \alpha_2(y) = \overline{U(y^x) \cap D(y)}$

So $E_{\rho(g)}$ is exactly the union of the Richardson 1-varieties. The cyclic connectivity of the Richardson 0- and 1-varieties, and therefore that of the exceptional set, is described explicitly Equation 6 and Figure 2.3.

5. F preserves the Richardson varieties of pairs $\xi(g^+), \xi(g^-)$

Assume $F : \mathcal{F}^3 \to \mathcal{F}^3$ is a topological conjugacy between $\rho, \rho' : \Gamma \to SL_3\mathbb{C}$. We use the notation $x' := F(x), y' := F(y)$.

First, we simplify the conclusions that we need to check.

Proposition 3.1. *Suppose that F a topological conjugacy such that F preserves the Richardson 1-varieties of a pair x and y in Λ. In other words, for some pair $x, y \in \Lambda^{(2)}$, we assume $F(R^1_{x,y}) = R^1_{x',y'}$. Then*

1. *F preserves the Richardson 0-varieties of x and y, i.e. $F(R^0_{x,y}) = R^0_{x',y'}$.*
2. *F either matches or switches the Richardson varieties of x and y.*

Proof. Let F be a homeomorphism that preserves the Richardson 1-varieties of x and y. First, let $z \in R^0_{x,y}$. Then $z = U \cap V$ for distinct $U, V \in R^1_{x,y}$. Since F is a homeomorphism, then

$$F(z) = F(U \cap V) \tag{23}$$

$$= F(U) \cap F(V). \tag{24}$$

By our assumption, then there exist $U', V' \in R^1_{x',y'}$ such that $U' = F(U)$ and $V' = F(V)$. Observe that $z' = U' \cap V'$ is therefore a Richardson 0-variety, so then $F(R^0_{x,y}) \subset F(R^0_{x',y'})$. Since F is a homeomorphism, then it restricts to a bijection on finite sets and $F(R^0_{x,y}) = R^0_{x',y'}$.

Recall that $x' = F(x), y' = F(y) \in \Lambda'$. Since $x \in span(x, x_y) \cap span(x, x^y)$, then $x' \in F(span(x, x_y)) \cap F(span(x, x^y))$, so either $F(span(x, x_y)) = span(x', x'_{y'})$ or $F(span(x, x_y)) = span(x, x^y)$. In the first case, F matches the Richardson varieties of x, y and $F(x_y) = x'_{y'}$, and in the second case F switches the Richardson varieties of x, y and $F(x_y) = x'^{y'}$. □

Next, we show that F preserves Richardson 1-varieties. We consider the case where x, y are the attracting and repelling flags of an infinite order element $g \in \Gamma$.

Proposition 3.2. *Let $x, y \in \Lambda$ such that $x = \xi(g^+)$ and $y = \xi(g^-)$. Then F preserves the Richardson 1-varieties of x and y.*

Proof. Since F is a topological conjugacy, it must preserve the exceptional set of $\rho(g)$, so $F(E_{\rho(g)}) = E_{\rho'(g)}$. Recall that the exceptional set of $\rho(g)$ is the union of the Richardson 1-varieties of x and y, that is:

$$E_{\rho(g)} = \cup_{V \in R^1_{x,y}} V. \tag{25}$$

Since $F|_{E_{\rho(g)}}$ is continuous and injective, then it must send 1-varieties to 1-varieties, so $F(R^1_{x,y}) = R^1_{x',y'}$. □

6. F preserves Richardson varieties of Λ

In this section, we extend our argument in Prop 3.2 from pairs $\xi(g^+), \xi(g^-)$ to all distinct pairs $\Lambda^{(2)}$. The main idea is that pairs of the form $\xi(g^+), \xi(g^-)$ are dense in Λ^2.

We use the following result from group theory:

Fact 3.1 (TODO REFERENCE). *Let Γ hyperbolic with boundary $\partial\Gamma \equiv S^1$, and let $x, z \in \partial\Gamma$. Then there exists a quasigeodesic sequence $\{g_i\}_{i=1} \in \Gamma^{\mathbb{N}}$ such that for all $y \in \partial\Gamma$:*

$$lim_{i\to\infty} g_i \cdot y = \begin{cases} x : y = x \\ z : y \neq x \end{cases} . \tag{26, 27}$$

We also observe that Richardson 1-varieties vary continuously on transverse pairs of flags:

Remark 3.1. *Let $\{x_i\}_{i=1}, \{y_i\}_{i=1} \in (\mathcal{F}^3)^{\mathbb{N}}$ such that for all $i \in \mathbb{N}$, the pair (x_i, y_i) are transverse. Furthermore, assume $x_i \to x$ and $y_i \to y$, and that the pair (x, y) is transverse. Then $lim_{i\to\infty} R^1_{x_i,y_i} = R^1_{x,y}$.*

The continuinity of Richardson 1-varieties also respects the type, e.g. $lim_{i\to\infty} span(x_i, x_i^{y_i}) = span(x, x^y)$.

So then we consider the Richardson varieties of pairs x, z where $x = \xi(g^+)$ and z is a (distinct) generic point on Λ.

Proposition 3.3. *Let $g \in \Gamma$ infinite order, and define $x := \xi(g^+)$. Let $z \in \Lambda \setminus \{x\}$. Then F preserves the Richardson 1-varieties of x and z.*

PROOF. (Proposition 3.3) If $z = \xi(g^-)$ then we are done by Prop 3.2, so assume not. Set $y := \xi(g^-)$.

Let $\{g_i\}_{i=1} \in \Gamma^{\mathbb{N}}$ as in Fact 3.1, such that $\lim_{i=1} \rho(g_i) \cdot y = z$ and $lim_{i=1}\rho(g_i) \cdot x = x$. For all $i \in \mathbb{N}$, define the following:

$$x_i := \rho(g_i) \cdot x, \tag{28}$$

$$y_i := \rho(g_i) \cdot y, \tag{29}$$

$$x_i' := F(x_i), \text{ and} \tag{30}$$

$$y_i' := F(y_i). \tag{31}$$

By Prop. 3.2, then for all $i \in \mathbb{N}$, F preserves the Richardson 1-varieties of x_i and y_i. By Rem. 3.1, then $\lim_{i\to\infty} R^1_{x_i,y_i} = R^1_{x,z}$. By continuity of F, then $\lim_{i\to\infty} x_i' = x'$ and $\lim_{i\to\infty} y_i' = z'$. We apply Rem. 3.1 again to the transverse sequence (x_i', y_i'), so then

$$F(R^1_{x,z}) = \lim_{i\to\infty} F(R^1_{x_i,y_i}) \tag{32}$$

$$= \lim_{i\to\infty} R^1_{x_i',y_i'} \tag{33}$$

$$= R^1_{x',z'}. \tag{34}$$

□

Next, we extend Proposition 3.3 to Richardson varieties of pairs w, z where both w and z are distinct generic points of Λ.

PROPOSITION 3.4. *Let $w, z \in \Lambda^{(2)}$. Then F preserves the Richardson 1-varieties of w and z.*

PROOF. If w or z is the attracting fixed point of an infinite order element $g \in \Gamma$, then we are done by Prop 3.3. Assume not.

Let $\{x_i\}_{i=1} \in \Lambda^{\mathbb{N}}$ such that each $x_i = \xi(g_i)$ is the attracting fixed point of an infinite order word, and $\lim_{i\to\infty} x_i = w$. This sequence exists because attracting fixed points are dense in Λ. Define $x_i' = F(x_i)$ for all $i \in \mathbb{N}$. By continuity of F, then $\lim_{i\to\infty} x_i' = w'$.

The argument proceeds similarly to that of Prop. 3.3. By Prop. 3.3, then for all $i \in \mathbb{N}$, F preserves the Richardson 1-varieties of x_i and z. By Rem. 3.1, then $\lim_{i\to\infty} R^1_{x_i,z} = R^1_{w,z}$. We apply Prop. 3.1 again to the transverse sequence $(x_i', z') \to (w', z')$, so then:

$$F(R^1_{w,z}) = \lim_{i\to\infty} F(R^1_{x_i,z}) \tag{35}$$

$$= \lim_{i\to\infty} R^1_{x_i',z'} \tag{36}$$

$$= R^1_{w',z'}. \tag{37}$$

□

7. F matches or switches Richardson varieties

In this section, we show that F either matches all Richardson varieties of Λ or switches all Richardson varieties of Λ, which concludes the proof of Lemma 3.1.2. We also highlight that F preserves the balanced ideal thickening of each $x \in \Lambda$.

Recall that $Th(x) = \alpha_1(x) \cup \alpha_2(x)$, so $Th(x) = span(x, x_y) \cup span(x, x^y)$ for any $y \in \Lambda$ distinct from x.

Proof. (Corollary 3.1.1) Let $x \in \Lambda$, and let $y \in \Lambda \setminus \{x\}$. By Prop. 3.4, F preserves the Richardson 1-varieties of x and y. By Prop. 3.1, F either matches or switches the Richardson varieties of each pair x and y. So then

$$F(Th(x)) = F(span(x, x_y) \cup span(x, x^y)) \tag{38}$$
$$= F(span(x, x_y)) \cup F(span(x, x^y)) \tag{39}$$
$$= span(x', x'_{y'}) \cup span(x', x'^{y'}) \tag{40}$$
$$= Th(x'). \tag{41}$$

□

If F matches (resp. switches) the Richardson varieties of some pair of $x, y \in \Lambda^{(2)}$, then F matches (resp. switches) the Richardson varieties of all such pairs.

Proof. Observe that

$$\alpha_1(\Lambda) = \cup_{x \in \Lambda} \alpha_1(x) \tag{42}$$
$$= \cup_{x,y \in \Lambda^{(2)}} span(x, x_y) \tag{43}$$
$$\cong S^1 \times \mathbb{CP}^1 \subset \mathcal{F}^3. \tag{44}$$

Similarly, $\alpha_2(\Lambda) \cong S^1 \times \mathbb{CP}^1$. In particular, $\alpha_1(\Lambda)$ and $\alpha_2(\Lambda)$ are connected subsets of $\mathcal{F}^3$ whose intersection is Λ.

By Cor. 3.1.1, clearly

$$F(\alpha_1(\Lambda)) \cup F(\alpha_2(\Lambda)) = F(\alpha_1(\Lambda) \cup \alpha_2(\Lambda)) \tag{45}$$
$$= F(Th(\Lambda)) \tag{46}$$
$$= Th(\Lambda') \tag{47}$$
$$= \alpha_1(\Lambda') \cup \alpha_2(\Lambda'). \tag{48}$$

Since F is a homeomorphism which sends Λ to Λ', then either $F(\alpha_1(\Lambda)) = \alpha_1(\Lambda')$, or $F(\alpha_1(\Lambda)) = \alpha_2(\Lambda)$.

In other words, if F matches the Richardson varieties of one pair in $\Lambda^{(2)}$, then it matches the Richardson varieties of all such pairs. Otherwise, F switches the Richardson varieties of all such pairs. □

This completes the proof of Lemma 3.1.2.

8. Random remarks

This might belong better in a different chapter.

Remark 3.2. *For distinct $x, y \in \mathcal{F}^3 \setminus \Lambda$, F will not necessarily preserve the 0-dimensional Richardson varieties.*

Remark 3.3. *Theorem 3.1 applies to virtual hyperbolic surface groups as well, e.g. $\Gamma = \pi_1(\mathcal{O}^2)$ where $\mathcal{O}^2$ is a hyperbolic 2-orbifold.*

CHAPTER 4

Anosov components have one topological conjugacy class

1. Assumptions and notation

TODO assumptions on ρ, ρ'.

Recall the set of Anosov representations $\mathcal{A}(\Gamma)$, the quasi-Hitchin representations $\mathcal{QH}(\Gamma)$ and the quasi-Barbot representations $\mathcal{QB}(\Gamma)$.

Recall irreducible representations.

2. Anosov components have one topological conjugacy class

THEOREM 4.1. *Suppose $\rho, \rho' : \Gamma \to SL_3\mathbb{C}$ are irreducible representations in the same Anosov component. Then ρ, ρ' are topologically conjugate.*

Combined with Theorem 3.1, we get the following corollaries:

COROLLARY 4.1.1. *All quasi-Hitchin representations are hyperconvex. All quasi-Barbot representations are* not *hyperconvex.*

Furthermore:

COROLLARY 4.1.2. *The quasi-Hitchin and quasi-Barbot Anosov components are disjoint.*

COROLLARY 4.1.3. *The collection of hyperconvex representations is open and closed in $\mathcal{A}(\Gamma)$.*

We observe that distinct Anosov components may share a single topological conjugacy class. For example, the quasi-Barbot components in $Hom(T(p,q,r), SL_3\mathbb{C})$ are disjoint. However, they are dual, so have the same conjugacy class:

PROPOSITION 4.1. *Let $\rho : \Gamma \to SL_3\mathbb{C}$ Anosov and let ρ^* its dual. Then ρ, ρ^* are topologically conjugate.*

We also describe the topological conjugacy classes of the connected components of reducible Anosov representations $\mathcal{A}_{red}(\Gamma) \subsetneq QB(\Gamma)$.

THEOREM 4.2. *There are exactly three topological conjugacy classes of $\mathcal{A}_{red}(\Gamma)$:*

- *The class of semisimple representations,*
- *The class of line- or plane-reducible representations, and*
- *The class of reducible, but neither line- nor plane-reducible representations.*

Representations in the last class are topologically conjugate to the irreducible representations in the same component of $QB(\Gamma)$.

3. Key ingredients for proof

In this section, we highlight prior results that are used in the proof of Theorem 4.1. First, we use the following theorem of Stecker:

THEOREM 4.3 (**??**, Thm 3.24). *Let $\rho : \Gamma \to G$ Borel Anosov. If $dim(\Lambda) \leq mbic(G) - 2$, then every non-empty cocompact domain of discontinuity in $\mathcal{F}^3$ is dense and connected, and there is a bijection:*

$$\{\textit{balanced ideals in } W(G)\} \leftrightarrow \{\textit{nonempty cocompact domains of discontinuity in } \mathcal{F}(G)\} \tag{49}$$

We tailor Stecker's theorem to our assumptions. Recall that $G = SL_3\mathbb{C}$, so $mbic(G) = 4$ and $\mathcal{F}(G) = \mathcal{F}^3$. Then $W(G) = A_2$ which has exactly one balanced ideal. In $SL_3\mathbb{C}$, all Anosov representations are Borel Anosov. Furthermore, we assume that Γ is a hyperbolic surface group, so $\partial\Gamma \cong S^1$. Then $1 \leq dim(\Lambda_\rho) \leq 2$, because ξ_ρ is an injective and continuous function of S^1. So the condition in Theorem 4.3 is met, and:

Remark 4.1. *For Γ a hyperbolic surface group and $\rho : \Gamma \to SL_3\mathbb{C}$ Anosov, then ρ has a* unique *nonempty cocompact domain of discontinuity $\Omega_\rho \subset \mathcal{F}^3$. Furthermore, $\Omega_r ho$ is connected and dense in $\mathcal{F}^3$.*

We also use the following lemma, proved in the following section:

Lemma 4.3.1. *For Γ a hyperbolic surface group and $\rho : \Gamma \to SL_3\mathbb{C}$ Anosov, then Ω_ρ is simply connected.*

Finally, we use the following theorem of Dumas and Sanders:

Theorem 4.4 (**??**, Thm 5.1). *Let (P, Q) be parabolic subgroups of G, with P symmetric, and let $I \subset W(G)$ be a balanced ideal of type (P, Q). Let $\rho : \Gamma \to G$ be a P-Anosov representation. Then for any $\rho' \in \mathcal{A}(\rho, P)$, the quotient manifolds M^I_ρ and $M^I_{\rho'}$ are diffeomorphic.*

We again tailor this theorem to our assumptions that $G = SL_3\mathbb{C}$, where there is only symmetric parabolic subgroup and one balanced ideal in $W(G)$. Then:

Remark 4.2. *For Γ a hyperbolic surface group and $\rho, \rho' : \Gamma \to SL_3\mathbb{C}$ in the same Anosov component, then*

$$M_\rho \equiv_{diffeo} M_{\rho'}. \tag{50}$$

4. Proof of Theorem 4.1

Proof. (Theorem 4.1) Let $\rho, \rho' : \Gamma \to SL_3\mathbb{C}$ irreducible and in the same Anosov component. We want to show the existence of $F : \mathcal{F}^3 \to \mathcal{F}^3$ a topological connjugacy of ρ, ρ'.

Then by Thm. 4.4, there exists a diffeomorphism $D : M_\rho \to M_{\rho'}$. This lifts to a diffeomorphism of the universal cover $\tilde{D} : \tilde{M}_\rho \to \tilde{M}_{\rho'}$. Then by Lemma 4.3.1, the domains of discontinuity Ω_ρ, $\Omega_{\rho'}$ are simply connected, so $\tilde{M}_\rho \equiv \Omega_\rho$ and $\tilde{M}_{\rho'} \equiv \Omega_{\rho'}$. So we can consider $\tilde{D} : \Omega_\rho \to \Omega_{\rho'}$.

Clearly $\tilde{D}$ is a topological conjugacy of Ω_ρ and $\Omega_{\rho'}$, since it is a diffeomorphism and for all $f \in \Omega_\rho$ and $g \in \Gamma$ we have $\tilde{D}(\rho(g) \cdot f) = \rho'(g) \cdot \tilde{D}(f)$.

Next, we want to extend $\tilde{D}$ to a homeomorphism $F : \mathcal{F}^3 \to \mathcal{F}^3$ such that $F|_{\Omega_\rho} \equiv \tilde{D}$. By Theorem 4.3, Ω_ρ is an open, connected dense subset of $\mathcal{F}^3$, so $\tilde{D}$ has a unique continuous extension to $\mathcal{F}^3$ using sequences in Ω_ρ which limit to points in $Th(\Gamma)$.

TODO check thet F is a topological conjugacy. TODO where do we use irreducible? Probably in the checking F top conj □

Remark 4.3. *The topological conjugacy F may not be unique, however it is uniquely determined by the choice of the diffeomorphism D and its lift $\tilde{D}$.*

Proof. (Lemma 4.3.1) Fact: $\pi_1\mathcal{F}^3$ is trivial. Fact: The odd homology groups $H^{2n+1}(\Omega_\rho)$ vanish. Fact: Then $\pi_1(\Omega_\rho)$ is perfect, since $\pi_1(\Omega_\rho)^{ab} = H^1(\Omega_\rho) = \{Id\}$.

Claim: First consider $\Omega_{\rho_{Fuchs}}$ where ρ_F is $SL_3\mathbb{C}$-Fuchsian. Then $\Omega_{\rho_{Fuchs}}$ is simply connnected.

Claim: If there is a transverse homotopy between $\Lambda_{\rho_{Fuchs}}$ and Λ_ρ, then $\pi_1(\Omega_{\rho_{Fuchs}}) = \pi_1(\Omega_\rho)$.

Claim: There is a transverse homotopy between $\Lambda_{\rho_{Fuchs}}$ and Λ_ρ. □

5. Topological conjugacies of reducible representations

This argument for Theorem 4.2 hinges on whether ρ fixes a line and/or a plane in $\mathbb{C}^3$. Let $\rho : \Gamma \to SL_3\mathbb{C}$ reducible. We say that ρ is

(1) *semisimple* if ρ fixes a line $l \in CP^2$ and a plane $H \in (CP^2)^*$,
(2) *line-reducible* if ρ fixes a line $l \in CP^2$, but no plane,
(3) *plane-reducible* if ρ fixes a plane $H \in (CP^2)^*$, but no line, and
(4) *neither line- nor plane-reducible* otherwise.

Given $l \in \mathbb{CP}^2$, we use the abuse of notation

$$\alpha_1(l) := \alpha_1(l, H) \tag{51}$$

for any $H \supset l$ because $\alpha_1(l, H)$ does not depend on choice of H. Similarly, for $H \in (\mathbb{CP}^2)^*$, we use the analogous abuse of notation

$$\alpha_2(H) := \alpha_2(l, H) \text{ for some } l \subset H. \tag{52}$$

If ρ is semisimple, then for all $x, y, z \in \Lambda_\rho^{(3)}$ we have $x_y = x_z$, $y_x = y_z$, $z_x = z_y$, and $x^y = x^z$, $y^x = y^z$, $z^x = z^y$. So ρ fixes the line $l = l_{x^y}$ and the plane $H = H_{x_y}$. This means that the following sets are invariant under ρ:

$$\alpha_1(l) = span(x_y, y_x) = span(x_z, z_x) = span(y_z, z_y), \text{ and} \tag{53}$$

$$\alpha_2(H) = span(x^y, y^x) = span(x^z, z^x) = span(y^z, z^y). \tag{54}$$

Similarly, if ρ is line-reducible (but not semisimmple), then all $x^y = x^z$, $y^x = y^z$, $z^x = z^y$ but not all $x_y = x_z$. So ρ also fixes the line $l = l_{x^y}$, so the following set is invariant under ρ:

$$\alpha_1(l) = span(x^y, y^x) = span(x^z, z^x) = span(y_z, z_y). \tag{55}$$

If ρ is plane-reducible (but not semisimple), then for all $x, y, z \in \Lambda_\rho^{(3)}$ we have $x_y = x_z$, $y_x = y_z$, $z_y = z_x$, but not all $x^y = x^z$. So ρ fixes the plane $H = H_{x_y}$, so the following set is invariant under ρ:

$$\alpha_2(H) = span(x_y, y_x) = span(x_z, z_x) = span(y_z, z_y). \tag{56}$$

Finally, if ρ is neither line- nor plane-reducible, then ρ fixes no complex line or plane in $\mathbb{C}^3$.

Proof. (Theorem 4.2) If ρ, ρ' are in the same representation conjugacy class and have the same reducible type, then they are topologically conjugate.

Now we consider pairs $\rho, \rho' \in \overline{[\rho]}$ with different reducible types.

Semisimple representations are only topologically conjugate to other semisimple representations.

Since line-reducible and plane-reducible representations but fix a complex line in Ω_ρ, then they can be topologically conjugate.

Finally, if ρ is neither line- nor plane-reducible, then it does not fix any complex line in Ω_ρ, and is therefore not topologically conjugate to other reducible types. However, it is topologically conjugate to other irreducible representations in the same Anosov component. □

We also gain the following corollary, which is immediate from the proof of the Theorem and Lemma 3.1.2:

COROLLARY 4.4.1. *Suppose F a topological conjugacy between ρ line-reducible and some reducible ρ'. If ρ' line-irreducible, then F matches Richardson varieties. Otherwise ρ' is plane-irreducible and F switches Richardson varieties.*

CHAPTER 5

Vitae

For UIC this can be a full CV or a short version (e.g. half a page) with previoius degrees, work experience and publications.

APPENDIX A

Technical Lemmas

Here we collect some supporting lemmas.

Bibliography

[1] E. Dumas, and A. Sanders "Geometry of compact complex manifolds associated to generalized quasi-Fuchsian varieties" (2017)

[2] F. Stecker, "Balanced ideals and domains of discontinuity of Anosov representations" (2022)

[3] R. Hartshorne, *Algebraic Geometry*, Springer-Verlag, New York, 1977.

[4] D. Mumford, *Abelian Varieties*, Oxford University Press, 1970.

[5] J. Draisma, E. Horobet, G. Ottaviani, B. Sturmfels, and R. R. Thomas, "The Euclidean distance degree of an algebraic variety," *arXiv:1309.0049* (2013). Available at: https://arxiv.org/abs/1309.0049