tellmeGen™

Name: Walid Soaida
Date: 10-05-2026

Name	Pharmacological action	Results
Amitriptyline (Dosage)	Antidepressants	Use an alternative drug or reduce the dose by 50%, depending on the case.
Brivaracetam (Dosage)	Antiepileptics	Poor CYP2C19 metabolizer. Consider dose reduction
Carbamazepine (Dosage)	Antiepileptics	A higher dose of carbamazepine would be necessary.
Citalopram (Dosage)	Antidepressants	Poor CYP2C19 metabolizer. Evaluate dose or consider alternative drug
Clobazam (Adverse Reactions)	Muscle relaxants	Poor CYP2C19 metabolizer. Increased likelihood of adverse effects
Clomipramine (Dosage)	Antidepressants	Use an alternative drug or reduce the dose by 50%, depending on the case.
Clopidogrel (Dosage)	Antithrombotics	Poor CYP2C19 metabolizer. Avoid clopidogrel if possible.
Diazepam (Adverse Reactions)	Anxiolytics	Poor CYP2C19 metabolizer. Increased likelihood of adverse effects
Docetaxel (Adverse effects)	Antineoplastics	Decreased risk of adverse effects, leukopenia and neutropenia (Asians)
Doxepin (Dosage)	Antidepressants	Use an alternative drug or reduce the dose by 50%, depending on the case.
Escitalopram (Dosage)	Antidepressants	Poor CYP2C19 metabolizer. Evaluate dose or consider alternative drug
Imipramine (Dosage)	Antidepressants	Use an alternative drug or reduce the dose by 50%, depending on the case.
Ivacaftor (Efficacy)	Other drugs for the respiratory system	In case of cystic fibrosis, the use of ivacaftor is not recommended.
Lansoprazole, Dexlansoprazole (Dosage)	Anti-acid treatment	Poor CYP2C19 metabolizer. Consider dose reduction in chronic therapy
Methotrexate (Adverse Reactions)	Immunomodulators	Slightly increased toxicity risk
Methotrexate in rheumatoid arthritis (Efficacy)	Immunomodulators	Decreased response to treatment
Olanzapine (Adverse effects)	Antipsychotics	Increased likelihood of certain adverse effects
Olanzapine (Efficacy)	Antipsychotics	Probable worse response to treatment

Omeprazole (Dosage)	**Anti-acid treatment**	**Poor CYP2C19 metabolizer. Consider dose reduction in chronic therapy**
Pantoprazole (Dosage)	**Anti-acid treatment**	**Poor CYP2C19 metabolizer. Consider dose reduction in chronic therapy**
Peginterferons alfa-2a -2b and ribavirin (Efficacy)	**Antivirals**	**Lower probability of response**
Sertraline (Dosage)	**Antidepressants**	**Poor CYP2C19 metabolizer. Decrease dose or evaluate alternative drug.**
Tacrolimus (Dosage)	**Immunosuppressants**	**Poor CYP3A5 metabolizer. Initiate treatment with standard dose and monitor**
Trimipramine (Dosage)	**Antidepressants**	**Use an alternative drug or reduce the dose by 50%, depending on the case.**
Voriconazole (Dosage)	**Antifungals**	**Poor CYP2C19 metabolizer. Evaluate dose or consider alternative drug**
Abacavir (Adverse effects)	Antivirals	Reduced risk of hypersensitivity
Acenocoumarol, Fenprocoumon (Adverse effects)	Antithrombotics	Does not present c.-1639G>A in VKORC1. Risk of adverse effects unchanged
Allopurinol (Dosage)	Antigout	Standard dose treatment recommended.
Amifampridine (Dosage)	Cholinergics	Probably NAT2 intermediate acetylator. Usual dose
Aminoglycoside antibiotics (Adverse effects)	Antibiotics	No variation in the risk of developing hearing loss
Aripiprazole (Dosage)	Antipsychotics	Normal CYP2D6 metabolizer. Use the recommended dose
Atazanavir (Adverse reactions)	Antivirals	Intermediate UGT1A1 metabolizer. Low probability of hyperbilirubinemia.
Atomoxetine (Dosage)	Centrally acting psychostimulants	Normal CYP2D6 metabolizer. Consider increasing initial dose if there is no response or adverse effects.
Atorvastatin (Dosage)	Hypolipidemic agents	Normal SLCO1B1 activity. Treatment with standard doses
Brexpiprazole (Dosage)	Antipsychotics	Normal CYP2D6 metabolizer. Use the recommended dose
Celecoxib (Dosage)	Non-steroidal anti-inflammatory drugs	Normal CYP2C9 metabolizer. Start treatment with the recommended dose
Cisplatin (Adverse Reactions)	Antineoplastics	Typical risk of ototoxicity
Codeine (Dosage)	Opioid analgesics	Normal CYP2D6 metabolizer. Use the recommended dose
Daunorubicin, Doxorubicin (Adverse Reactions)	Antineoplastics	Moderate risk of cardiotoxicity
Desipramine (Dosage)	Antidepressants	Normal CYP2D6 metabolizer. Use the recommended dose

Efavirenz (Adverse effects)	Antivirals	Low risk of adverse reactions
Eliglustat (Adverse Reactions)	Treatment of metabolic diseases	CYP2D6 metabolizer normal. Typical risk of adverse effects
Fentanyl (Efficacy)	Opioid analgesics	No variation in response
Flecainide (Dosage)	Antiarrhythmics	Normal CYP2D6 metabolizer. Use the recommended dose
Floxacillin (Adverse effects)	Antibiotics	No variations in the risk of developing severe liver injury
Flucytosine (Adverse Reactions)	Antifungals	Normal metabolizer DPYD. Normal toxicity risk
Fluindione (Adverse Reactions)	Antithrombotics	Lower probability of bleeding at standard doses
Fluorouracil, Capecitabine (Dosage)	Antineoplastics	Normal DPYD metabolizer. Normal risk of fluoropyrimidine toxicity.
Flurbiprofen (Dosage)	Anti-inflammatory	Normal CYP2C9 metabolizer. Start treatment with the recommended dose
Fluvastatin (Dosage)	Hypolipidemic agents	Treatment with standard doses is recommended
Fluvoxamine (Dosage)	Antidepressants	Normal CYP2D6 metabolizer. Initiate therapy with standard doses
G6PD Deficiency and Adverse Drug Reactions	Multiple pharmacological actions	Low risk of hemolytic anemia for certain drugs
Haloperidol (Dosage)	Antipsychotics	Normal CYP2D6 metabolizer. Use the recommended dose
Hydrochlorothiazide (Efficacy)	Antihypertensives	No variation in response
Hydrocodone (Dosage)	Opioid analgesics	Normal CYP2D6 metabolizer. Standard dose recommended
Ibuprofen (Dosage)	Anti-inflammatory	Normal CYP2C9 metabolizer. Start treatment with the recommended dose
Iloperidone (Dosage)	Antipsychotics	Normal CYP2D6 metabolizer. Use the recommended dose
Inhalation anesthetics and succinylcholine (Adverse Reactions)	General anesthetics	Low risk of developing malignant hyperthermia (see report)
Irinotecan (Adverse effects)	Antineoplastics	Intermediate UGT1A1 metabolizer. Lower probability of adverse effects
Isoniazid (Adverse effects)	Antibiotics	Probably NAT2 intermediate acetylator. Lower risk of liver toxicity
Lornoxicam (Dosage)	Anti-inflammatory	Normal CYP2C9 metabolizer. Start treatment with the recommended dose
Lovastatin (Dosage)	Hypolipidemic agents	Normal SLCO1B1 activity. Treatment with standard doses
Lumacaftor + Ivacaftor (Efficacy)	Other drugs for the respiratory system	Cystic Fibrosis treatment with lumacaftor/ivacaftor not recommended

Meloxicam (Dosage)	Anti-inflammatory	Normal CYP2C9 metabolizer. Start treatment with the recommended dose
Metoprolol (Dosage)	Beta-blockers cardiovascular system	Normal CYP2D6 metabolizer. Use the recommended dose
Mirtazapine (Adverse Reactions)	Antidepressants	Normal CYP2D6 metabolizer. Standard risk of adverse effects
Mivacurium and succinylcholine (Adverse Reactions)	Muscle relaxants	Low risk of prolonged neuromuscular blockade
Nilotinib (Adverse Reactions)	Antineoplastics	Lower risk of adverse reactions
Nortriptyline (Dosage)	Antidepressants	Normal CYP2D6 metabolizer. Use the recommended dose
Paroxetin (Dosage)	Antidepressants	Normal CYP2D6 metabolizer. Use the recommended dose
Phenytoin (Dosage)	Antiepileptics	Normal CYP2C9 metabolizer. Initiate treatment at the recommended dose. Subsequent doses should be adjusted according to response and side effects.
Pimozide (Dosage)	Antipsychotics	Normal CYP2D6 metabolizer. Standard dose is recommended
Piroxicam (Dosage)	Anti-inflammatory	Normal CYP2C9 metabolizer. Start treatment with the recommended dose
Pitavastatin (Dosage)	Hypolipidemic agents	Normal SLCO1B1 activity. Treatment with standard doses
Pitolisant (Dosage)	Other nervous system drugs	Normal CYP2D6 metabolizer. Initiate therapy with standard dose
Pravastatin (Dosage)	Hypolipidemic agents	Normal SLCO1B1 activity. Treatment with standard doses
Propafenone (Adverse Reactions)	Antiarrhythmics	Normal CYP2D6 metabolizer. Normal risk of adverse effects
Quetiapine (Dosage)	Antipsychotics	Probably normal CYP3A4 metabolizer. It is recommended to use standard doses
Risperidone (Dosage)	Antipsychotics	Normal CYP2D6 metabolizer. Use the recommended dose
Rosuvastatin (Dosage)	Hypolipidemic agents	Treatment with standard doses is recommended
Selective Serotonin Reuptake Inhibitors (SSRIs) (Adverse effects)	Antidepressants	No variations in the likelihood of sexual dysfunction associated with SSRI treatment
Simvastatin (Dosage)	Hypolipidemic agents	Normal SLCO1B1 activity. Treatment with standard doses
Siponimod (Dosage)	Immunosuppressants	Normal CYP2C9 metabolizer. Initiate treatment at the recommended dose. Subsequent doses should be adjusted according to response and side effects.
Statins (Dosage and risk of myopathy)	Hypolipidemic agents	Normal risk of myopathy. Treatment with standard doses
Tamoxifen (Efficacy)	Antineoplastics	Normal CYP2D6 metabolizer. Standard dosage is recommended. Avoid CYP2D6 inhibitors.

Tenoxicam (Dosage)	Anti-inflammatory	Normal CYP2C9 metabolizer. Start treatment with the recommended dose
Thioguanine, Azathioprine, Mercaptopurine (Dosage)	Immunosuppressants	Normal TPMT metabolizer and normal NUDT15. Use standard doses
Tramadol (Dosage)	Opioid analgesics	Normal CYP2D6 metabolizer. Use the recommended dose
Valproic Acid (Adverse effects)	Antiepileptics	No variations in liver toxicity risk
Venlafaxine (Dosage)	Antidepressants	Normal CYP2D6 metabolizer. Use the recommended dose
Vortioxetine (Dosage)	Antidepressants	Normal CYP2D6 metabolizer. Initiate therapy with standard doses
Warfarin (Adverse Reactions)	Antithrombotics	Lower probability of bleeding at standard doses
Zuclopenthixol (Dosage)	Antipsychotics	Normal CYP2D6 metabolizer. Use the recommended dose