

**Title: Potential gains in life expectancy by reducing inequality of lifespans in Denmark: An international comparison and cause of death analysis**

**Authors:** José Manuel Aburto <sup>a\*</sup>, Maarten Wensink <sup>a\*</sup>, Alyson van Raalte <sup>b</sup> & Rune Lindahl-Jacobsen <sup>a</sup>

**Author affiliations:**

<sup>a</sup> Institute of Public Health- Unit of Biodemography, University of Southern Denmark, Odense 5000, Denmark.

<sup>b</sup> Max Planck Institute for Demographic Research, Rostock 18057, Germany

\* These authors contributed equally to the paper

**Corresponding author:**

José Manuel Aburto

Institute of Public Health- Unit of Biodemography, University of Southern Denmark

J.B. Winsløws Vej 9, DK-5000 Odense C

T: +45 65509416

[jmaburto@health.sdu.dk](mailto:jmaburto@health.sdu.dk)

**Email addresses:**

JMA: [jmaburto@health.sdu.dk](mailto:jmaburto@health.sdu.dk)

MW: [mwensink@health.sdu.dk](mailto:mwensink@health.sdu.dk)

AvR: [vanraalte@demogr.mpg.de](mailto:vanraalte@demogr.mpg.de)

RLJ: [rjacobsen@health.sdu.dk](mailto:rjacobsen@health.sdu.dk)

**Keywords:** Demography, lifespan variability, aging, mortality, public health

**Word count:** 2949

## Abstract

**Background** Reducing lifespan inequality is increasingly recognized as a health policy objective. Whereas lifespan inequality declined with rising longevity in most developed countries, Danish life expectancy stagnated between 1975 and 1995 for females and progressed slowly for males. It is unknown how Danish lifespan inequality changed, which causes of death drove these developments, and where the opportunities for further improvements lie now.

**Methods** We present an analytical strategy based on cause-by-age decompositions to simultaneously analyze changes in Danish life expectancy and lifespan inequality from 1960 to 2014, as well as current Swedish-Danish differences.

**Results** Stagnation in Danish life expectancy coincided with a shorter period of stagnation in lifespan inequality (1975-1990). The stagnation in life expectancy was mainly driven by increases in cancer and non-infectious respiratory mortality at higher ages (-.63 years), offsetting a reduction in cardiovascular and infant mortality (+1.52 years). Lifespan inequality stagnated because most causes of death did not show compression over the time period. Both these observations were consistent with higher smoking-related mortality in Danes born in 1919-1939. After 1995, life expectancy and lifespan equality increased in lockstep, but still lag behind Sweden, mainly due to infant mortality and cancer.

**Conclusions** Since 1960, Danish improvements in life expectancy and lifespan equality were halted by smoking-related mortality in those born 1919-1939, while also reductions in old-age cardiovascular mortality held back lifespan equality. The comparison with Sweden suggests that Denmark can reduce inequality in lifespans and increase life expectancy through a consistent policy target: reducing cancer and infant mortality.

## Background

Life expectancy is the most commonly used measure of the health status of a population and the performance of the healthcare system.(1) Another important dimension is the uncertainty around that expectation: lifespan inequality.(2) Lifespan inequality has become relevant for policy makers with the growing interest in economic and health inequalities, (3, 4) in particular because: (1) it is a marker of heterogeneity in age at death at the macro level, and (2) it is a marker of uncertainty in the timing of death at the micro level.(5-7) Typically, early deaths are more common in underprivileged groups, simultaneously reducing life expectancy and increasing lifespan inequality.(8-11) Both indicators may have implications for individuals' decisions over their life course. For instance, when to invest in education or when to retire are decisions based on life expectancy but also on the uncertainty surrounding the eventual time of death.(10)

Life expectancy is lower in Denmark than in Norway and Sweden for females and males. From 1975 to 1995, while their Scandinavian counterparts showed continuous improvement, life expectancy stagnated among Danish women, while Danish men experienced only slow progress. For both sexes, life expectancy improved after 1995, but remains lower than in Sweden and Norway.(12) Differences between Denmark and Sweden in life expectancy have been thoroughly documented.(13, 14) Among females, the stagnation in life expectancy resulted mainly from the increased mortality of those born from 1919 to 1939, cohorts with high levels of smoking and alcohol consumption compared to their Swedish contemporaries.(13, 14) Similarly, smoking-related mortality was considerably higher in Danish compared to Swedish males because of the widespread use of snus instead of cigarettes in

Sweden.(15) While these factors are a known contributor to life expectancy differences,(16) their effect on lifespan inequality differences is unknown. Previous evidence has shown mixed results for the effects of smoking on lifespan inequality: little to no effect on the Finnish population,(17) while it increased lifespan inequality in some European countries.(18)

The Danish case, juxtaposed with its Scandinavian counterparts, is interesting given the shared history, culture and similarities in their healthcare systems.(19) It is unknown how the different age and cause-of-death mortality trends in the three countries would extend to lifespan inequality patterns.

Because life expectancy and lifespan inequality tend to be positively correlated (5, 7) we hypothesize that 1) during the last decades, Denmark experienced higher lifespan inequality relative to Sweden and Norway in females and males; 2) the 1975-1995 stagnation in life expectancy of Danish women was accompanied by stagnation in lifespan inequality; 3) the slow increase in life expectancy for males in the same period was accompanied by slow reduction of lifespan inequality. Because it is well-documented that smoking in the interwar Danish female cohorts was a major cause of the 1975-1995 stagnation in Danish female life expectancy,(14) we hypothesize that 4) any 1975-1995 stagnation in lifespan inequality can also be attributed to smoking-related deaths in these cohorts.

Hence, we analyze data since 1960 for Denmark, Sweden and Norway to make a cause-by-age analysis of changes in life expectancy and lifespan inequality for both sexes.

## **Methods**

### ***Mortality and cause of death data***

Period lifetables by sex and single year of age (0-110+) were retrieved from the Human Mortality Database (12) for Denmark, Sweden and Norway for the period 1960 to 2014. Cause-of-death data were taken from the WHO Mortality Database to compute the proportion of deaths by cause, age, and sex in a given year.(20) Cause-of-death data are available in 5-year age and single year categories. To increase the accuracy of the resulting estimates,(21) causes of death were ungrouped into single years of age using efficient estimation of smooth distributions.(22)

### ***Cause-of-death classification***

Data on causes of death were classified using the seventh, eighth, ninth and tenth revisions of the International Classification of Diseases (ICD) for the period studied.(23) Deaths were grouped in seven major cause-of-death categories aimed at capturing conditions that might have affected mortality in these countries. We considered that smoking prevalence was comparatively high among women (and still remains higher) in Denmark;(14, 24) that the decrease in mortality from heart conditions (cardiovascular revolution) took place during the studied period;(25) and that the management of infectious diseases has improved greatly over the past half century.(26) Hence, we grouped causes of death up to age 84 as follows: 1) Cancers sensitive to smoking,(27) 2) Cancers not sensitive to smoking, 3) Cardiovascular diseases, 4) Non-infectious respiratory diseases, 5) Infectious respiratory diseases, 6) External causes and 7) Rest of causes. For ICD codes and details on the classification see Supplementary Table 1. Causes of death above age 85 were not decomposed, because of lower reliability in the presence of multi-morbidities.(28) Our groupings over the various ICD revisions were cross-checked with other coding practices across ICD versions in the literature.(29) We also

checked for discontinuities in death counts for each of the seven causes of death over ICD transition years (Supplementary Figures 2-4).

### ***Lifespan inequality measure***

Several dispersion measures have been proposed to analyze lifespan inequality.<sup>(30)</sup> Here, we use the coefficient of variation (CoV), which is the standard deviation divided by the mean of the lifetable age-at-death distribution, i.e. life expectancy (Supplementary Section 1 for a brief description). CoV has been found to be a good indicator to measure lifespan inequality.<sup>(31)</sup> The strong correlation between dispersion indicators suggests that main conclusions and results would not differ much between measures used.<sup>(30, 32, 33)</sup> Life expectancy and lifespan inequality (CoV) were calculated for Denmark, Norway and Sweden throughout 1960-2014.

### ***Decomposition techniques***

Cause-by-age decompositions of the changes in lifespan and lifespan inequality in Denmark, Norway and Sweden were made from 1960 to 2014 using standard decomposition techniques.<sup>(34)</sup> These decompositions allow singling out any period during those years, for instance 1975-1995<sup>1</sup>. We quantified the cause-by-age contributions to the current differences in life expectancy and lifespan inequality between Denmark and Sweden for females and males.

## **Results**

---

<sup>1</sup> We have created an interactive app where the reader can analyze any period he/she might be interested in for any sex. Available [online app](#).

### ***Trends in lifespan inequality and life expectancy 1960-2014***

The 1975-1995 stagnation in life expectancy for Danish females was accompanied by a shorter period of stagnation in lifespan inequality (Figure 1A). Swedish and Norwegian females experienced a decrease in inequality and increase in life expectancy throughout the period (Figure 1A). For males in all three countries, life expectancy increase was slow in 1960-1980, but accelerated thereafter, while the decrease in lifespan inequality was more monotonic (Figure 1B).

[Figure 1 about here]

### ***Decomposition of changes in life expectancy and lifespan inequality for Denmark***

Between 1960 and 1975, Danish female life expectancy increased from 74 to 77 years mainly due to reductions in infant mortality and mid- and old-age cardiovascular mortality (Figure 2). For males<sup>1</sup>, infant mortality was also reduced, but the contribution from cardiovascular diseases was absent (see Supplementary Figure 1), resulting in a small increase in life expectancy from 70.4 to 71.3 years. For both sexes, lifespan inequality was reduced mainly because of the reduction in infant mortality.

Between 1975 and 1995, Danish female life expectancy stagnated at about 77 years because a continued reduction in infant mortality and old-age cardiovascular mortality was offset by an increase in (mainly smoking-related) cancer and non-infectious respiratory mortality between ages 55 and 85 (Figure 2). Also, reduction in cardiovascular mortality was lower in Danish females relative to Danish males and Swedish females (Supplementary Figure 1).

The impact of mortality change on lifespan inequality is more complicated: at younger ages mortality reduction results in deaths being compressed into a narrower age range, reducing lifespan inequalities. At older ages mortality reduction stretches out the right tail of the age-at-death distribution, increasing lifespan inequality. Overall, lifespan inequality was mostly unchanged among Danish females because there was little compression of mortality for most causes. Increases in smoking-related cancer and non-infectious respiratory diseases were apparent over both these ‘younger’ and ‘older’ ages with opposite effects, but on balance increased lifespan inequality during the period (Figure 2). For males, the reduction in lifespan inequality was larger than for females, mainly driven by a reduction in infant mortality and early-life external mortality (Supplementary Figure 1).

Between 1995 and 2014, Danish female and male life expectancy increased (from 77.8 to 82.7 and 72.7 to 78.6, respectively) due to almost all causes, particularly cardiovascular conditions which were concentrated over adult ages. As for lifespan inequality, for both sexes all ages and all causes up to around the life expectancy reduced inequality, while a reduction in cardiovascular mortality at ages higher than life expectancy increased inequality.

[Figure 2 about here]

*Decomposition of current differences in life expectancy and lifespan inequality between Denmark and Sweden*



Currently (2014)<sup>2</sup>, life expectancy is higher in Sweden than in Denmark for both sexes due to almost all causes at all ages, with the major exception of external mortality being higher in Sweden than in Denmark at all ages, in particular ages 15-35 (Figure 3). Two major classes of mortality where Denmark is doing worse than Sweden could be identified. First, infant mortality is higher in Denmark than in Sweden (by a factor two). Second, mid- and old-age cancer mortality is higher in Denmark than in Sweden. Other recent years showed the same pattern.

For lifespan inequality, the same holds: infant mortality and mid-life cancer mortality increase Denmark's disadvantage relative to Sweden, somewhat offset by lower external mortality between ages 15 and 35 (Figure 3). However, Denmark's life expectancy disadvantage relative to Sweden is mainly due to mid- and high-age cancer mortality, while Denmark's lifespan inequality disadvantage is mainly due to higher infant mortality (Figure 3).

[Figure 3 about here]

### ***Potential gains in Danish life expectancy if lifespan inequality were reduced towards Swedish levels***

Reducing mortality from cancers below age 85 would decrease the gap in lifespan inequality by 31% and 22% for females and males, respectively (Table 1). This translates into gains in life expectancy of 0.57 years for females and 0.66 years for males, respectively 44% and 37% of the overall life expectancy gap. Reducing infant mortality (from all causes) to

---

<sup>2</sup> Results for any year from 1960-2014 and for Norway vs Sweden available in [online app](#).

Swedish levels would reduce lifespan inequality by 46% for females and 49% for males. This would be translated into gains in life expectancy of .14 years for females and .16 years for males, respectively 10% and 9% of the total gap.

Achieving Swedish levels in cardiovascular conditions would decrease the gap in lifespan inequality by almost 10% in both sexes and increase life expectancy by about 3 months. Conversely, if Sweden were to achieve the level of Danish external mortality, it would benefit by two additional months of life expectancy for both sexes. Mortality above age 85 has negligible effect on the difference between Denmark and Sweden in lifespan inequality.

[Table 1 about here]

## **Discussion**

In this study, we found that the same causes and age groups that held back Danish life expectancy in 1975-1995, especially for females, also held back lifespan equality in the same period. This suggests important social development, but also a clear policy target. Although lifespan inequality has been reduced and life expectancy has increased since the late 1990s, Denmark still lags its Scandinavian counterparts, despite similarities in social and healthcare systems. The comparison with Sweden suggests that Denmark can now reduce inequality in lifespans and increase life expectancy through the same policy targets: cancer and infant mortality.

Reducing lifespan inequality cannot be the only policy goal, since this would neglect the interests of those who have already lived to higher ages: The effect of mortality reduction on lifespan inequality is large and negative at age zero, decreases with increasing age, and re-

verses at a unique threshold age, so that mortality reductions above this threshold age increase lifespan inequality.(5, 35) Therefore, the causes that extend lifespan and the causes that reduce inequality are not necessarily the same.(36) Smoking-related mortality is a clear example of this. In Denmark, life expectancy stagnated over the 1975-1995 period because mortality reduction from most causes of death were offset by mortality increase from smoking-related causes. These increases in smoking-related had a smaller net impact on lifespan inequality compared to life expectancy over the same period, since smoking-related mortality occurred below and above the threshold age. By the latest period 1995-2014, however, reduction in smoking-related mortality was comparatively more important for decreases in lifespan inequality (19.4%) than increases in life expectancy (11.2%). In general, the impact of smoking on lifespan inequality is dependent on both the age of smokers compared to non-smokers (the maturity of the smoking epidemic), as well as the actual impact of smoking on mortality at different ages.(17) Similar to what was found in a comparison of G7 countries ,(36) reductions in injuries and child mortality were relatively more important for lifespan inequality decrease than for life expectancy increase.

Causes of death that drive cross-sectional differences in lifespan inequality are not necessarily the same as the causes of death that drive contemporary gaps between countries. (37) The comparison with Sweden suggests that Denmark can simultaneous increase life expectancy and decrease lifespan inequality by targeting two main causes of death: cancer and infant mortality. Reducing lifespan inequality towards Sweden by these conditions would lead to an increase of 0.7 and 0.8 years in life expectancy for females and males in Denmark, respectively. To put this in perspective, in 2014 the infant mortality rate in Denmark is twice as

high as in Sweden, which is one of the lowest among developed countries.(12) Although mortality at very young ages may be affected by different registration practices in high income countries (e.g. non-viable live births registered as stillbirths),(38) the Nordic countries do not show evidence of such patterns.(39) Moreover, even after controlling for gestational age Sweden showed lower infant mortality rates than Denmark.(40) Thus, infant mortality is the largest single contributor to the gap with Sweden in terms of lifespan inequality. Preventive policies focusing on prenatal risk factors and improving maternal health before and during pregnancy,(41) as well as efforts to reduce the risk of sudden infant death syndrome (42) could help to reduce infant mortality towards Swedish levels.

Targeting cancer is another clear public health intervention to reduce lifespan inequality and increase life expectancy in Denmark. Our results show that improvements in cancer mortality have had an effect on both health indicators over the last 20 years. However, Denmark had the highest mortality rates from all neoplasms in the European region, and the female population exhibited the highest lung cancer mortality rates.(24) This is in line with our comparison with Sweden and with previous evidence highlighting the role of smoking behaviors on life expectancy trends.(14)

For Sweden, the decomposition results suggest that young-age external mortality can be further reduced. According to the WHO, males in Denmark have lower age-standardized external mortality rates (39 per 100,000) than Sweden and Norway (50.6 and 52 respectively) in 2014.(43) Our results further show that these differences are concentrated between ages 15 and 40. Moreover, since the late 1990s, Swedish and Norwegian males have experienced higher suicide rates between ages 15 and 24.(44)

Indeed, the mere observation that Sweden is doing better than Denmark for most causes of death does not mean that Denmark could easily do better. However, it does provide a starting point for public health intervention. For instance, previous evidence suggests that focusing on vulnerable and less socially advantaged subgroups may reduce suicide rates among the young.(44, 45)

For other countries that lag a comparable country, similar decompositions can be made. This may not result in a clear and consistent message: causes of death that hold back life expectancy may not be the same as the causes of death that hold back equality. Yet if it does, as in the case of Denmark when compared to Sweden, the benefits are substantial, because the policy goals can be so clearly stated. We therefore suggest that this method could be a valuable tool for epidemiologists and policy makers alike.

As any cause of death analysis, our study has the limitations that: 1) causes of death are treated as mutually exclusive, while they may not be (e.g., poor sight due to diabetes may lead to an accident); 2) medical doctors and even coroners have imperfect knowledge about causes of death; and 3) trends in awareness of certain diseases and changing insights in disease processes affect classification. Yet through using otherwise high-quality data and broad categories of causes of death, we believe we have achieved a useful, workable grouping of causes of death. In addition, we performed a sensitivity analysis to assure consistency of grouping across ICD versions and did not find significant variation when ICD revisions changed (Supplementary Figures 2-4).

Lifespan inequality is an important dimension of population health. By looking at this dimension we could disclose how lifespans differ within Denmark, Sweden and Norway.

Moreover, our decomposition by age- and cause-of-death allowed us to identify conditions and ages that contribute the most to lifespan inequality changes, and we were able to translate them into potential gains in life expectancy if efforts were concentrated in these ages and causes of death.

## **Conclusions**

Lifespan inequality together with life expectancy gives a broader perspective on the effect of mortality changes on population health. Our results show that life expectancy and lifespan inequality have been negatively correlated since at least 1960 in Denmark. Currently, Denmark lags Sweden both in terms of high life expectancy and low inequality due to two main causes: infant mortality and cancer. Denmark therefore has a clear and consistent public health policy target: reduce infant mortality and cancer mortality. Our approach demonstrates how reduction in lifespan inequality as a policy target can be translated into gains in life expectancy.

## **List of abbreviations**

CoV. Coefficient of variation.

ICD. International Classifications of Diseases

## **Declarations**

**Ethics approval and consent to participate:** We use secondary, grouped, publicly available data. Therefore, no ethics approval from our institutions was needed.

**Consent to Publish:** Not applicable.

**Availability of data and materials:** All data used in this study is publicly available through websites of Human Mortality Database and WHO mortality database (references 12 and 20, respectively). Code to replicate the results available at <https://github.com/jmaburto/Lifespan-inequality-Denmark/tree/master/R>

**Competing interests:** None declared.

**Funding** All authors thank their supporting institutions This work was supported by a starting grant from the European Research Council [grant No. 716323 to AvR].

**Authors contributions:** JMA and MW designed the study. JMA, MW, AvR interpreted results. All authors contributed to writing the manuscript.

**Acknowledgements:** Not applicable.

## References

1. Organization WH. World Health Statistics 2016: Monitoring Health for the SDGs Sustainable Development Goals: World Health Organization; 2016.
2. Tuljapurkar S. The final inequality. *Demography and the Economy*. 2011;209.
3. Marmot M. Inequalities in health. *New England Journal of Medicine*. 2001; 345(2):134-5.
4. Mackenbach JP, Kulhánová I, Artnik B, et al. Changes in mortality inequalities over two decades: register based study of European countries. *bmj*. 2016; 353:i1732.
5. Vaupel JW, Zhang Z, van Raalte AA. Life expectancy and disparity: an international comparison of life table data. *BMJ open*. 2011; 1(1):e000128.
6. Edwards RD, Tuljapurkar S. Inequality in life spans and a new perspective on mortality convergence across industrialized countries. *Population and Development Review*. 2005; 31(4):645-74.
7. Smits J, Monden C. Length of life inequality around the globe. *Social Science & Medicine*. 2009; 68(6):1114-23.
8. Brønnum-Hansen H. Socially disparate trends in lifespan variation: a trend study on income and mortality based on nationwide Danish register data. *BMJ open*. 2017; 7(5):e014489.
9. Firebaugh G, Acciai F, Noah AJ, Prather C, Nau C. Why lifespans are more variable among blacks than among whites in the United States. *Demography*. 2014; 51(6):2025-45.
10. van Raalte AA, Kunst AE, Deboosere P, et al. More variation in lifespan in lower educated groups: evidence from 10 European countries. *International Journal of Epidemiology*. 2011:dyr146.
11. Aburto JM, van Raalte A. Lifespan dispersion in times of life expectancy fluctuation: the case of Central and Eastern Europe. MPIDR working paper. 2017.
12. Human Mortality Database. University of California BU, and Max Planck Institute for Demographic Research (Germany). Human Mortality Database. 2017.

13. Lindahl-Jacobsen R, Oeppen J, Rizzi S, et al. Why did Danish women's life expectancy stagnate? The influence of interwar generations' smoking behaviour. *European Journal of Epidemiology*. 2016;1-5.
14. Lindahl-Jacobsen R, Rau R, Jeune B, et al. Rise, stagnation, and rise of Danish women's life expectancy. *Proceedings of the National Academy of Sciences*. 2016; 113(15):4015-20.
15. Ramström L, Wikmans T. Mortality attributable to tobacco among men in Sweden and other European countries: an analysis of data in a WHO report. *Tobacco induced diseases*. 2014; 12(1):14.
16. Drefahl S, Ahlbom A, Modig K. Losing Ground-Swedish Life Expectancy in a Comparative Perspective. *PloS one*. 2014; 9(2):e88357.
17. van Raalte AA, Myrskylä M, Martikainen P. The role of smoking on mortality compression: An analysis of Finnish occupational social classes, 1971-2010. *Demographic Research*. 2015; 32:589.
18. Janssen F, Rousson V, Paccaud F. The role of smoking in changes in the survival curve: an empirical study in 10 European countries. *Annals of epidemiology*. 2015; 25(4):243-9.
19. Magnusson J. The Scandinavian healthcare system. *Medical solutions*. 2009:63-8.
20. Organization WH. Health statistics and information systems. 2017.  
[http://www.who.int/healthinfo/mortality\\_data/en/](http://www.who.int/healthinfo/mortality_data/en/).
21. Rizzi S, Thinggaard M, Engholm G, et al. Comparison of non-parametric methods for ungrouping coarsely aggregated data. *BMC medical research methodology*. 2016; 16(1):59.
22. Rizzi S, Gampe J, Eilers PH. Efficient estimation of smooth distributions from coarsely grouped data. *American journal of epidemiology*. 2015; 182(2):138-47.
23. Organization WH. International statistical classification of diseases and related health problems: World Health Organization; 2004.
24. Hashim D, Boffetta P, La Vecchia C, et al. The global decrease in cancer mortality: trends and disparities. *Annals of Oncology*. 2016; 27(5):926-33.
25. Smallman-Raynor M, Phillips D. Late stages of epidemiological transition: health status in the developed world. *Health & place*. 1999; 5(3):209-22.
26. Organization WH. The world health report 2000: health systems: improving performance: World Health Organization; 2000.
27. Organization WH, Cancer IAfRo. Personal Habits and Indoor Combustions Volume 100e A Review Of Human Carcinogens. IARC Monographs on the Evaluation of Carcinogenic Risks to Humans. 2012; 100:1-441.
28. Rosenberg HM. Cause of death as a contemporary problem. *Journal of the history of medicine and allied sciences*. 1999; 54(2):133-53.
29. Janssen F, Kunst AE. ICD coding changes and discontinuities in trends in cause-specific mortality in six European countries, 1950-99. *Bulletin of the World Health Organization*. 2004; 82(12):904-13.
30. van Raalte AA, Caswell H. Perturbation analysis of indices of lifespan variability. *Demography*. 2013; 50(5):1615-40.
31. Wrycza TF, Missov TI, Baudisch A. Quantifying the shape of aging. *PloS one*. 2015; 10(3):e0119163.
32. Wilmoth JR, Horiuchi S. Rectangularization revisited: Variability of age at death within human populations. *Demography*. 1999; 36(4):475-95.
33. Colchero F, Rau R, Jones OR, et al. The emergence of longevous populations. *Proceedings of the National Academy of Sciences*. 2016. doi:10.1073/pnas.1612191113
34. Horiuchi S, Wilmoth JR, Pletcher SD. A decomposition method based on a model of continuous change. *Demography*. 2008; 45(4):785-801.
35. Gillespie DO, Trotter MV, Tuljapurkar SD. Divergence in age patterns of mortality change drives international divergence in lifespan inequality. *Demography*. 2014; 51(3):1003-17.
36. Seligman B, Greenberg G, Tuljapurkar S. Equity and length of lifespan are not the same. *Proceedings of the National Academy of Sciences*. 2016; 113(30):8420-3.



37. Shkolnikov VM, Andreev EM, Zhang Z, Oeppen J, Vaupel JW. Losses of expected lifetime in the United States and other developed countries: methods and empirical analyses. *Demography*. 2011; 48(1):211-39.
38. Blencowe H, Cousens S, Oestergaard MZ, et al. National, regional, and worldwide estimates of preterm birth rates in the year 2010 with time trends since 1990 for selected countries: a systematic analysis and implications. *The Lancet*. 2012; 379(9832):2162-72.
39. Deb-Rinker P, León JA, Gilbert NL, et al. Differences in perinatal and infant mortality in high-income countries: artifacts of birth registration or evidence of true differences? *BMC pediatrics*. 2015; 15(1):112.
40. MacDorman MF, Mathews T. Behind international rankings of infant mortality: how the United States compares with Europe: Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Health Statistics; 2009.
41. Zylbersztejn A, Gilbert R, Hardelid P, Hjern A. Why do more infants die in the UK than in Sweden? An intercountry comparison of birthweight-specific infant mortality. *The Lancet*. 2015; 386:S83.
42. Wennergren G, Nordstrand K, Alm B, et al. Updated Swedish advice on reducing the risk of sudden infant death syndrome. *Acta Paediatrica*. 2015; 104(5):444-8.
43. Organization WH. European Health Information Gateway.  
[https://gateway.euro.who.int/en/indicators/h2020\\_15-mortality-from-external-causes-males/](https://gateway.euro.who.int/en/indicators/h2020_15-mortality-from-external-causes-males/). 2017. Accessed 21/11 2017.
44. Titelman D, Oskarsson H, Wahlbeck K, et al. Suicide mortality trends in the Nordic countries 1980–2009. *Nordic journal of psychiatry*. 2013; 67(6):414-23.
45. Bergen H, Hawton K, Kapur N, et al. Shared characteristics of suicides and other unnatural deaths following non-fatal self-harm? A multicentre study of risk factors. *Psychological medicine*. 2012; 42(4):727-41.

## Figures and Tables

Figure 1. Life expectancy (panel A) and lifespan inequality (panel B) trends from 1960 to 2015 for Denmark, Sweden and Norway by sex. The shaded area refers to the period of life expectancy stagnation in Danish females 1975-1995.

Figure 2. Age and cause contributions to changes in life expectancy (panel A) and lifespan inequality (panel B) between 1960-1975, 1975-1995 and 1995-2014 for Danish females. Note: Age 0 is truncated in panel B since it accounts for the largest contribution.

Figure 3. Age and cause contributions to the gap in life expectancy (Panel A) and lifespan inequality (Panel B) with Sweden in 2014 by sex.

Table 1. Potential gains in life expectancy in Denmark if inequality is reduced (%) to Swedish levels in 2014 by cause of death.


Sex	Cause of death category and mortality above age 85	Reduce gap with Sweden in CoV (%)	Reduction in life expectancy (%)	Potential Gains in life expectancy (years)
Females	1 Smoking-related cancer	18%	25%	0.35
	2 Non-Smoking related cancer	13%	16%	0.22
	3 Cardiovascular	10%	15%	0.21
	4 Respiratory-Infectious	2%	2%	0.03
	5 Respiratory-Non-infectious	7%	17%	0.23
	6 External	-26% *	-11% **	-0.15
	7 Other	71%	40%	0.55
	Above age 85	5%	-3% **	-0.05
Males	1 Smoking-related cancer	15%	26%	0.47
	2 Non-Smoking related cancer	7%	10%	0.19
	3 Cardiovascular	10%	19%	0.33
	4 Respiratory-Infectious	1%	3%	0.05
	5 Respiratory-Non-infectious	5%	7%	0.12
	6 External	-26% *	-11% **	-0.19
	7 Other	92%	43%	0.77
	Above age 85	0.0	0.0	0.04

\* Increases the gap with Sweden. Represents potential gains for Sweden if they achieve the levels of Denmark.


\*\* Increases the gap with Sweden in life expectancy.

Note: the sum of percentages differ to 100% due to rounding.


## A Life expectancy at birth




## B Lifespan inequality (CoV)



A Decomposition of life expectancy




B Decomposition of lifespan inequality (CoV)




# A Decomposition of life expectancy

Sweden - Denmark, 2014.



# B Decomposition of lifespan inequality (CoV)

Denmark - Sweden, 2014.



**Supplemental material to: Potential gains in life expectancy by reducing inequality of lifespans in Denmark: An international comparison and cause of death analysis.**

**Authors:** José Manuel Aburto <sup>a\*</sup>, Maarten Wensink <sup>a\*</sup>, Alyson van Raalte <sup>b</sup> & Rune Lindahl-Jacobsen <sup>a</sup>

**Author affiliations:**

<sup>a</sup> Institute of Public Health- Unit of Biodemography, University of Southern Denmark, Odense 5000, Denmark.

<sup>b</sup> Max Planck Institute for Demographic Research, Rostock 18057, Germany

\* These authors contributed equally to the paper

**Corresponding authors:** JMA and MW:

Institute of Public Health- Unit of Biodemography, University of Southern Denmark

J.B. Winsløws Vej 9, DK-5000 Odense C

T: +45 65509416

[jmaburto@health.sdu.dk](mailto:jmaburto@health.sdu.dk)

**Table 1. ICD code for the cause of death classification.**

<b>Cause of Death</b>	<b>ICD-7</b>	<b>ICD-8</b>	<b>ICD-9 (Sweden, Norway)</b>	<b>ICD-10</b>
Cancer, smoking-related	A044-A050, A052, 157, 180-181	A045-A051, A055, 157, 188-189	B08, B090-B094, B096, B100-B101, 180, 188-189	C00-C21, C25, C30-C34, C53, C64-C68
Cancer, not smoking-related	A051, A053-A056, A058-A059, 155-156, 158-160, 164-165, 175-176, 178-179, 192-195, 198-199	A052-A054, A056-A057, A059-A60, 155-156, 158-160, 163, 171, 183-184, 186-187, 190-199	B095, B099, B109, B11, B13-B14, 179, 181-187	C22-C24, C26, C37-C39, C40-C41, C43-C52, C54-C58, C60-C63, C69-C97
Cardiovascular	A063, A070, A079-A086	A064, A080-A088	B181, B25-B30	E10-E14, I00-I99
Respiratory, infectious	A087-A092, A095	A089-A092, A095	B310-B312, B320-B322	J00-J06, J09-J18, J20-J22, J34.0, J36, J39.0, J39.1, J85, J86
Respiratory, non-infectious	A093, A094, A096, A097	A093, A094, A096	B313-B315, B319, B323-B327, B329	J30-J33, J34.1-J34.3, J34.8, J35, J37, J38, J39.2, J39.3, J39.8, J39.9, J40-J47, J60-J70, J80-J82, J840-J841, J848-J849, J90-J99
External	A138-A150	A138-A150	B47-B56	S00-T89, V01-Y84
Other	A001-A043, A060-A062, A064-A069, A071-A078, A098-A137	A001-A044, A061-A063, A065-A079, A097-A137	B01-B07, B184-B185, B15-B17, B180, B182-B183, B189, B19-B23, B33-B46	A00-B89, B99, D00-D48, D50-D89, E00-E07, E15-E16, E20-E35, E40-E46, E50-E68, E70-E90, F00-F99, G00-G99, H00-H59, H60-H95, K00-K93, L00-L99, M00-M99, N00-N99, O00-O99, P00-P96, Q00-Q99, R00-R99

### **1) Details on the classification**

Primary malignancies that are sensitive to smoking are found predominantly in the respiratory, digestive and genitourinary tracts, in line with the principle that where smoke or its products pass, the risk of cancer rises. Primary malignancies in the gastrointestinal tract from mouth to anus were classified as sensitive to smoking, with the exception of liver cancer, for which detail could not be reconstructed across ICD versions (see below). Cancer in the respiratory tract was also classified as sensitive to smoking. In addition, it has been proven that smoking causes cancer of the uterine cervix, the ovaries (mucinous carcinoma), the bladder, the kidney (pelvis and body) and the ureter. For mucinous carcinoma of the ovaries, detail could not be reconstructed across ICD versions (see below). Malignancies in the urinary tract were classified as being sensitive to smoking.

The resolution of the ICD classification has grown substantially over the years. As we analyzed deaths from 1960 through 2014, we used ICD-7 through ICD-10. The specifically identified categories “cancer sensitive to smoking” and “respiratory infectious”, are based on the smallest common denominator: only if a specific disease could be separately identified across ICD versions did we include it in these groups. For instance, myeloid leukemia is associated with smoking, but ICD-7 and -8 contain only a category ‘leukemia’, without subclassification. Hence, for reasons of consistency across classifications, myeloid leukemia is considered as not sensitive to smoking throughout. Also, ICD-7 and ICD-8 have an overall rest group for malignant neoplasms, while ICD-9 and ICD-10 have rest groups for each tract, if known. Because ICD-7 and ICD-8 do not have these detailed rest groups, rest groups were classified as not sensitive to smoking for all ICDs.




## 2) Brief description of the indicator

In lifetable notation, it is:


$$CV = \frac{\sqrt{\int_a^\omega (x - e_a)^2 f(x) dx}}{\int_a^\omega \ell(x) dx} = \frac{\sigma}{e_a}. \quad (1)$$

Where  $f(x)$ ,  $e_0$  and  $\omega$  denote the age at death density function, life expectancy at age  $a$ , and the open-aged interval (110+ in our case), respectively.


Figure 1.



Figures 2-4 Sensitivity analysis of cause of death differences in ICD coding



Sweden  
Years of change in ICDs: 1969, 1987 and 1997



# Norway

Years of change in ICDs: 1969, 1986 and 1996

