

The threshold age of the lifetable entropy

May 27, 2019

Abstract

BACKGROUND

Indicators of relative inequality of lifespans are important because they capture the dimensionless shape of aging. They are markers of inequality at the population level and express the uncertainty at the time of death at the individual level. In particular, the lifetable entropy \bar{H} represents the elasticity of life expectancy to a change in mortality and it has been used as an indicator of lifespan variation. However, it is unknown how this measure changes over time and whether a threshold age exists, as it does for other lifespan variation indicators.

RESULTS

The time derivative of \bar{H} can be decomposed into changes in life disparity e^\dagger and life expectancy at birth e_o . Likewise, changes over time in \bar{H} are a weighted average of age-specific rates of mortality improvements. These weights reflect the sensitivity of \bar{H} and show how mortality improvements can increase (or decrease) the relative inequality of lifespans. Further, we prove that in the assumption that mortality is reduced at all ages, \bar{H} , as well as e^\dagger , has a threshold age below which saving lives reduces entropy, whereas improvements above that age increase entropy.

CONTRIBUTION

We give a formal expression for changes over time of \bar{H} , and provide a formal proof of the existence of a unique threshold age that separates reductions and increases in lifespan inequality as a result age-specific mortality improvements.

1 Relationship

The lifetable entropy is a dimensionless indicator of the relative variation in the length of life compared to life expectancy at birth (Keyfitz 1968; Demetrius 1974; Keyfitz 1977; Demetrius 1978). It is usually defined as

$$\bar{H}(t) = -\frac{\int_0^\infty \ell(a, t) \ln \ell(a, t) da}{\int_0^\infty \ell(a, t) da} = \int_0^\infty c(a, t) H(a, t) da = \frac{e^\dagger(t)}{e_o(t)},$$

where $e^\dagger(t) = -\int_0^\infty \ell(a, t) \ln \ell(a, t) da$ is the life disparity or number of life-years lost as a result of death (Vaupel and Canudas-Romo 2003), $e_o(t) = \int_0^\infty \ell(a, t) da$ is the life expectancy at birth at time t , $\ell(a, t)$ is the lifetable survival function, $c(a, t) = \ell(a, t) / \int_0^\infty \ell(x, t) dx$ is the population structure, and $H(a, t) = \int_0^a \mu(x, t) dx$ is the cumulative hazard to age a , where $\mu(x, t)$ is the force of mortality (hazard rate or risk of death) at age x at time t . Note that $\bar{H}(t)$ can be interpreted as an *average value* of $H(a, t)$ in the population at time t .

Goldman and Lord (1986) and Vaupel (1986) proved that

$$e^\dagger(t) = \int_0^\infty d(a, t) e(a, t) da,$$

where $d(a, t)$ represents the distribution of deaths and $e(a, t) = \int_a^\infty \ell(x, t) dx / \ell(a, t)$ the remaining life expectancy at age a at time t . This provides an alternative expression for the lifetable entropy as

$$\bar{H}(t) = \frac{\int_0^\infty d(a, t) e(a, t) da}{\int_0^\infty \ell(a, t) da}.$$

Let $\dot{\bar{H}}$ denote the partial derivative of \bar{H} with respect to time.¹ We define $\rho(x) =$

¹ In the following, a dot over a function will denote its partial derivative with respect to time t , but

$-\dot{\mu}(x)/\mu(x)$ as the age-specific rates of mortality improvements. Then, the relative derivative of \bar{H} can be expressed as a weighted average of age-specific rates of mortality improvement,

$$(1) \quad \dot{\bar{H}}/\bar{H} = \int_0^\infty \rho(x) w(x) W(x) dx ,$$

with weights

$$w(x) = \mu(x) \ell(x) e(x) \quad \text{and} \quad W(x) = \frac{1}{e^\dagger} (H(x) + \bar{H}(x) - 1) - \frac{1}{e_o} .$$

Function $\bar{H}(x)$ is the lifetable entropy conditioned on surviving to age x , defined as

$$\bar{H}(x) = \frac{e^\dagger(x)}{e(x)} = \frac{\int_x^\infty d(a) e(a) da}{\int_x^\infty \ell(a) da} .$$

where $e^\dagger(x) = \int_x^\infty d(a) e(a) da / \ell(x)$ refers to life disparity above age x , and $e(x)$ is the remaining life expectancy at age x .

Note that the lifetable entropy \bar{H} is a measure of relative lifespan inequality. Thus, higher values represent more lifespan disparity, whereas lower values denote less variation of lifespans. If mortality improvements over time occur at all ages, there exists a unique threshold age a^H that separates *positive* from *negative* contributions to the lifetable entropy \bar{H} resulting from those mortality improvements. This threshold age a^H is reached when

$$(2) \quad H(a^H) + \bar{H}(a^H) = 1 + \bar{H} .$$

variable t will be omitted for simplicity.

2 Proof

Fernández and Beltrán-Sánchez (2015) showed that the relative derivative of \bar{H} can be expressed as

$$(3) \quad \dot{\bar{H}} / \bar{H} = \frac{\dot{e}^\dagger}{e^\dagger} - \frac{\dot{e}_o}{e_o},$$

This formula indicates that relative changes in \bar{H} over time are given by the difference between relative changes in e^\dagger (dispersion component) and relative changes in e_o (translation component). We will first provide expressions for \dot{e}_o and \dot{e}^\dagger to prove that (1) and (3) are equivalent. Next, we will prove the existence of threshold age for \bar{H} and its uniqueness.

2.1 Relative changes over time in \bar{H}

Vaupel and Canudas-Romo (2003) showed that changes over time in life expectancy at birth are a weighted average of the total rates of mortality improvements, given by

$$(4) \quad \dot{e}_o = \int_0^\infty \rho(x) w(x) dx,$$

where $\rho(x) = -\dot{\mu}(x) / \mu(x)$ are the age-specific rates of mortality improvement, and $w(x) = \mu(x) \ell(x) e(x) = d(x) e(x)$ is a measure of the importance of death at age x .

Since $d(x) = \mu(x) \ell(x)$ and $\ell(x) e(x) = \int_x^\infty \ell(a) da$, the partial derivative with respect

to time of $e^\dagger = \int_0^\infty d(x) e(x) dx$ can be expressed as

$$\begin{aligned} \dot{e}^\dagger &= \int_0^\infty \dot{\mu}(x) \ell(x) e(x) dx + \int_0^\infty \mu(x) \int_x^\infty \dot{\ell}(a) da dx \\ &= - \int_0^\infty \rho(x) w(x) dx + \int_0^\infty \dot{\ell}(a) \int_0^a \mu(x) dx da \\ &= - \int_0^\infty \rho(x) w(x) dx - \int_0^\infty \int_0^a \dot{\mu}(x) dx \ell(a) H(a) da , \end{aligned}$$

where $H(a)$ is the cumulative hazard to age a . By reversing the order of integration and doing some additional manipulations, we get

$$\begin{aligned} \dot{e}^\dagger &= - \int_0^\infty \rho(x) w(x) dx - \int_0^\infty \dot{\mu}(x) \int_x^\infty \ell(a) H(a) da dx \\ &= - \int_0^\infty \rho(x) w(x) dx + \int_0^\infty \rho(x) w(x) \frac{\int_x^\infty \ell(a) H(a) da}{\ell(x) e(x)} dx \\ (5) \quad &= \int_0^\infty \rho(x) w(x) \left(\frac{\int_x^\infty \ell(a) (H(a) - H(x) + H(x)) da}{\ell(x) e(x)} - 1 \right) dx \\ &= \int_0^\infty \rho(x) w(x) \left(H(x) \frac{\int_x^\infty \ell(a) da}{\ell(x) e(x)} + \frac{\int_x^\infty \ell(a) (H(a) - H(x)) da}{\ell(x) e(x)} - 1 \right) dx \\ &= \int_0^\infty \rho(x) w(x) \left(H(x) + \frac{\int_x^\infty \ell(a) (H(a) - H(x)) da}{\ell(x) e(x)} - 1 \right) dx . \end{aligned}$$

In Proposition 1 in the Appendix, we prove that

$$(6) \quad e^\dagger(x) = \frac{1}{\ell(x)} \int_x^\infty d(a) e(a) da = \frac{1}{\ell(x)} \int_x^\infty \ell(a) (H(a) - H(x)) da .$$

Replacing (6) into (5) yields

$$\begin{aligned}
(7) \quad \dot{e}^\dagger &= \int_0^\infty \rho(x) w(x) \left(H(x) + \frac{e^\dagger(x)}{e(x)} - 1 \right) dx \\
&= \int_0^\infty \rho(x) w(x) (H(x) + \bar{H}(x) - 1) dx .
\end{aligned}$$

Finally, replacing the expressions of \dot{e}_o and \dot{e}^\dagger from (4) and (7) into (3), we get

$$\begin{aligned}
\dot{\bar{H}} / \bar{H} &= \frac{1}{e^\dagger} \int_0^\infty \rho(x) w(x) (H(x) + \bar{H}(x) - 1) dx - \frac{1}{e_o} \int_0^\infty \rho(x) w(x) dx \\
&= \int_0^\infty \rho(x) w(x) \left(\frac{1}{e^\dagger} (H(x) + \bar{H}(x) - 1) - \frac{1}{e_o} \right) dx \\
&= \int_0^\infty \rho(x) w(x) W(x) dx ,
\end{aligned}$$

which proves (1) and shows that relative changes over time in the lifetable entropy are the average of the rates of mortality improvement weighted by the product $w(x) W(x)$. \square

2.2 The threshold age for \bar{H}

Using (1), changes over time in the lifetable entropy \bar{H} are given by

$$(8) \quad \dot{\bar{H}} = \bar{H} \int_0^\infty \rho(x) w(x) W(x) dx .$$

Whenever $\dot{\bar{H}} > 0$, lifespan inequality increases over time, whereas $\dot{\bar{H}} < 0$ implies that variation of lifespans decreases over time. Because $\ell(x)$ is a positive function bounded between 0 and 1, we have that $\bar{H} > 0$. Moreover, assuming age-specific death rates $\mu(x)$ improve over time at all ages, then $\dot{\mu}(x) < 0$ and $\rho(x) > 0$ at any age x . Therefore, (8) implies that

1. Those ages x in which $w(x)W(x) > 0$ will contribute *positively* to the lifetable entropy \bar{H} and increase lifespan variation;
2. Those ages x in which $w(x)W(x) < 0$ will contribute *negatively* to the lifetable entropy \bar{H} and favor lifespan equality;
3. Those ages x in which $w(x)W(x) = 0$ will have no effect on the variation over time of \bar{H} .

Our goal is to prove that if mortality improvements occur for all ages and $\rho(x) > 0$, there exists a unique threshold age a^H such that $w(a^H)W(a^H) = 0$. That threshold age will separate *positive* from *negative* contributions to \bar{H} resulting from mortality improvements.

The product $w(x)W(x)$ can be re-expressed as

$$\begin{aligned} w(x)W(x) &= \mu(x)\ell(x)e(x)\left(\frac{1}{e^\dagger}(H(x) + \bar{H}(x) - 1) - \frac{1}{e_o}\right) \\ &= \frac{\mu(x)\ell(x)e(x)}{e^\dagger}(H(x) + \bar{H}(x) - \bar{H} - 1) . \end{aligned}$$

Since $\mu(x)$, $\ell(x)$, $e(x)$ and e^\dagger are all positive functions, the threshold age of \bar{H} occurs whenever

$$(9) \quad g(x) := H(x) + \bar{H}(x) - \bar{H} - 1 = 0 .$$

When x is close to 0, $g(x)$ takes negative values since

$$g(0) = H(0) + \bar{H}(0) - \bar{H} - 1 = 0 + \bar{H} - \bar{H} - 1 = -1 < 0 .$$

Likewise, $g(x)$ takes positive values when x becomes arbitrary large. Note that \bar{H} does

not depend on age, and therefore

$$\lim_{x \rightarrow \infty} g(x) = \lim_{x \rightarrow \infty} (H(x) + \bar{H}(x)) - \bar{H} - 1 = \infty$$

because $\lim_{x \rightarrow \infty} H(x) = \infty$. By definition, $\bar{H}(x) \geq 0$ for all x , so regardless of the behavior of $\bar{H}(x)$ when x is arbitrarily large, the limit of $g(x)$ tends to infinity. Hence, given that $g(0) = -1$ and $\lim_{x \rightarrow \infty} g(x) = \infty$, in a continuous framework the intermediate value theorem guarantees the existence of at least one age a^H at which $g(a^H) = 0$.

Moreover, as shown in Proposition 2 in the Appendix, $g(x)$ is a strictly increasing function, and therefore a one-to-one function assuming continuity. As a result, there is a unique threshold age a^H that separates *positive* from *negative* contributions to the lifetable entropy \bar{H} , and that threshold age is reached when

$$w(x)W(x) = 0 \iff g(x) = 0 \iff H(x) + \bar{H}(x) = 1 + \bar{H},$$

which proves (2). □

3 Related results

Demographers have developed a battery of indicators to measure how lifespans vary in populations (Colchero et al. 2016; van Raalte and Caswell 2013). The most common indexes are the variance (Edwards and Tuljapurkar 2005; Tuljapurkar and Edwards 2011), standard deviation (van Raalte, Sasson, and Martikainen 2018) and coefficient of variation (Aburto et al. 2018) of the age at death distribution, the Gini coefficient (Shkolnikov, Andreev, and Begun 2003; Archer et al. 2018; Gigliarano, Basellini, and Bonetti 2017),

the Theil index (Smits and Monden 2009), and the years of life lost (Vaupel, Zhang, and van Raalte 2011; Aburto and van Raalte 2018), among others. However, only few studies have analytically derived formulas for *threshold age* below and above which mortality improvements respectively decrease and increase lifespan variation. Zhang and Vaupel (2009) showed that the threshold age (a^\dagger) for life disparity (e^\dagger) occurs when $H(x) + \bar{H}(x) = 1$. Similarly, Gillespie, Trotter, and Tuljapurkar (2014) determined a threshold age for the variance of the age at death distribution. Van Raalte and Caswell (2013) also showed that it is possible to determine the threshold age by performing an empirical sensitivity analysis of lifespan variation indicators.

In this article, we contribute to the lifespan variation literature by deriving the threshold age a^H for the life table entropy \bar{H} . This age separates negative from positive contributions of age-specific mortality improvements. We analytically proved its existence and—in the assumption that mortality improves over time for all ages—also its uniqueness. In Section 4 we empirically show that it differs from the threshold age of e^\dagger .

4 Applications

4.1 Numerical findings

Figure 1 depicts the threshold ages for the two related measures, e^\dagger and \bar{H} . Calculations were performed using data from the Human Mortality Database (2018) for females in the United States and Italy in 2005. The blue line represents $g(x)$ from Equation (9). The threshold age a^H occurs when $g(x)$ crosses zero. The red and grey line display the same functions that Zhang and Vaupel (2009) used to find the threshold age for e^\dagger . The

intersection of these two lines denotes the threshold age a^\dagger . Finally, the dashed black line depicts the life expectancy at birth. Vaupel, Zhang, and van Raalte (2011) noted that a^\dagger tends to fall just below life expectancy. The threshold age for the lifetable entropy a^H is greater than a^\dagger and is very close above life expectancy for these countries. Note the similarity between the formulas for a^\dagger , given by $H(a^\dagger) + \bar{H}(a^\dagger) = 1$, and a^H , given by $H(a^H) + \bar{H}(a^H) = 1 + \bar{H}$.




Figure 1: Threshold ages for life disparity (a^\dagger) and for the lifetable entropy (a^H). United States and Italy in 2005. Values in Panel A: $e_o = 80.13$, $a^\dagger = 78.51$, and $a^H = 80.86$. Values in Panel B: $e_o = 83.67$, $a^\dagger = 81.76$, and $a^H = 83.28$. Source: Human Mortality Database (2018)

Panels A and B in Figure 2 illustrate the evolution over time of the threshold ages for e^\dagger and \bar{H} in French and Swedish females, respectively. We chose these countries because they portray large series of reliable data available at the Human Mortality Database (2018).

Values for a^\dagger are close to life expectancy throughout the period. However, around 1950 there is a crossover between a^\dagger and e_o such that a^\dagger remained close to life expectancy but below it. This result shows that the threshold age a^\dagger being below life expectancy is a modern feature of ageing populations with high life expectancy. From the beginning of the period of observation to the 1950s, the threshold age for the lifetable entropy was above life expectancy for both countries. During some periods a^\dagger was roughly constant

whereas life expectancy trended upwards. After the 1950s, a^\dagger converged towards life expectancy. The code and data to reproduce these results are publicly available through this repository [link not given to avoid identification of authors].




Figure 2: Threshold ages for life disparity (a^\dagger) and for the lifetable entropy (a^H) compared to life expectancy at birth. French and Swedish females, 1800–2016. Source: Human Mortality Database (2018)

4.2 Decomposition of the relative derivative of \bar{H}

The relative derivative of \bar{H} defined in Equation (1) can be decomposed between components before and after the threshold age a^H as follows:

$$\begin{aligned}
 \dot{\bar{H}} / \bar{H} &= \int_0^\infty \rho(x) w(x) W(x) dx \\
 (10) \quad &= \int_0^{a^H} \rho(x) w(x) W(x) dx + \int_{a^H}^\infty \rho(x) w(x) W(x) dx \\
 &= \underbrace{\left\{ \frac{\dot{e}^\dagger[x|x < a^H]}{e^\dagger} - \frac{\dot{e}_o[x|x < a^H]}{e_o} \right\}}_{\text{Early life component}} + \underbrace{\left\{ \frac{\dot{e}^\dagger[x|x > a^H]}{e^\dagger} - \frac{\dot{e}_o[x|x > a^H]}{e_o} \right\}}_{\text{Late life component}}
 \end{aligned}$$

If mortality reductions occur at every age, the *early life component* in Equation (10) is always negative (contributing to reduce entropy) while the *late life component* is positive (contributing to increasing entropy). Thus, it is clear that a negative relationship between

life expectancy and entropy over time occurs if the early life component outpaces the late life component. This decomposition is based on the additive properties of the derivatives of life expectancy and e^\dagger as previously shown in Vaupel and Canudas-Romo (2003) and Fernández and Beltrán-Sánchez (2015).

5 Conclusion

Several authors have been interested in decomposing changes over time in life expectancy (Arriaga 1984; Vaupel 1986; Pollard 1988; Vaupel and Canudas-Romo 2003; Beltrán-Sánchez, Preston, and Canudas-Romo 2008; Beltrán-Sánchez and Soneji 2011). Recently, authors have also investigated how life disparity fluctuations over time can be decomposed (Wagner 2010; Zhang and Vaupel 2009; Shkolnikov et al. 2011; Aburto and van Raalte 2018; Aburto and Beltrán-Sánchez 2019). In this paper, we bring both perspectives together and shed light on the dynamics behind changes in the lifetable entropy.

Keyfitz (1977) first proposed \bar{H} as a lifetable function “that measures the change in life expectancy at birth consequent on a proportional change in age-specific rates” (p. 413). Since then, several authors have been interested in this measure and its use (Demetrius 1978, 1979; Mitra 1978; Goldman and Lord 1986; Vaupel 1986; Hakkert 1987; Hill 1993; Fernández and Beltrán-Sánchez 2015). The lifetable entropy has become an appropriate indicator of lifespan variation that permits comparison of the shape of ageing across different species and over time (Baudisch et al. 2013; Wrycza, Missov, and Baudisch 2015). In this article, we uncovered the mathematical regularities behind the changes over time in the lifetable entropy. In particular, this study contributes to the existing literature by showing that (1) the lifetable entropy can be decomposed as a weighted

average of rates of mortality improvements, and (2) there exists a unique threshold age that separates positive and negative contributions of reductions in mortality over time.

References

- Aburto, J.M. and Beltrán-Sánchez, H. (2019). Upsurge of homicides and its impact on life expectancy and life span inequality in Mexico, 2005–2015. *American journal of public health* 109(3): 483–489.
- Aburto, J.M. and van Raalte, A.A. (2018). Lifespan dispersion in times of life expectancy fluctuation: The case of Central and Eastern Europe. *Demography* 55(6): 2071–2096.
- Aburto, J.M., Wensink, M., van Raalte, A., and Lindahl-Jacobsen, R. (2018). Potential gains in life expectancy by reducing inequality of lifespans in Denmark: An international comparison and cause-of-death analysis. *BMC Public Health* 18(1): 831.
- Archer, C.R., Basellini, U., Hunt, J., Simpson, S.J., Lee, K.P., and Baudisch, A. (2018). Diet has independent effects on the pace and shape of aging in *Drosophila melanogaster*. *Biogerontology* 19(1): 1–12.
- Arriaga, E.E. (1984). Measuring and explaining the change in life expectancies. *Demography* 21(1): 83–96.
- Baudisch, A., Salguero-Gómez, R., Jones, O.R., Wrycza, T., Mbeau-Ache, C., Franco, M., and Colchero, F. (2013). The pace and shape of senescence in angiosperms. *Journal of Ecology* 101(3): 596–606.
- Beltrán-Sánchez, H., Preston, S.H., and Canudas-Romo, V. (2008). An integrated approach to cause-of-death analysis: Cause-deleted life tables and decompositions of life expectancy. *Demographic Research* 19: 1323–1350.
- Beltrán-Sánchez, H. and Soneji, S. (2011). A unifying framework for assessing changes in life expectancy associated with changes in mortality: The case of violent deaths. *Theoretical Population Biology* 80(1): 38–48.
- Colchero, F., Rau, R., Jones, O.R., Barthold, J.A., Conde, D.A., Lenart, A., Németh, L., Scheuerlein, A., Schoeley, J., Torres, C., ..., Alberts, S.C., and Vaupel, J.W. (2016). The emergence of longevous populations. *Proceedings of the National Academy of Sciences* 113(48): E7681–E7690.
- Demetrius, L. (1974). Demographic parameters and natural selection. *Proceedings of the National Academy of Sciences* 71(12): 4645–4647.
- Demetrius, L. (1978). Adaptive value, entropy and survivorship curves. *Nature* 275: 213–214.
- Demetrius, L. (1979). Relations between demographic parameters. *Demography* 16(2): 329–338.
- Edwards, R.D. and Tuljapurkar, S. (2005). Inequality in life spans and a new perspective on mortality convergence across industrialized countries. *Population and Development Review* 31(4): 645–674.

- Fernández, O.E. and Beltrán-Sánchez, H. (2015). The entropy of the life table: A reappraisal. *Theoretical Population Biology* 104: 26–45.
- Gigliarano, C., Basellini, U., and Bonetti, M. (2017). Longevity and concentration in survival times: The log-scale-location family of failure time models. *Lifetime Data Analysis* 23(2): 254–274.
- Gillespie, D.O., Trotter, M.V., and Tuljapurkar, S.D. (2014). Divergence in age patterns of mortality change drives international divergence in lifespan inequality. *Demography* 51(3): 1003–1017.
- Goldman, N. and Lord, G. (1986). A new look at entropy and the life table. *Demography* 23(2): 275–282.
- Hakkert, R. (1987). Life table transformations and inequality measures: Some noteworthy formal relationships. *Demography* 24(4): 615–622.
- Hill, G. (1993). The entropy of the survival curve: An alternative measure. *Canadian Studies in Population* 20(1): 43–57.
- Human Mortality Database (2018). University of California, Berkeley, and Max Planck Institute for Demographic Research, Rostock. URL <http://www.mortality.org>.
- Keyfitz, N. (1968). *Introduction to the mathematics of populations*. Addison-Wesley Series in Behavioral Science: Quantitative Methods.
- Keyfitz, N. (1977). What difference would it make if cancer were eradicated? An examination of the Taeuber paradox. *Demography* 14(4): 411–418.
- Mitra, S. (1978). A short note on the Taeuber paradox. *Demography* 15(4): 621–623.
- Pollard, J.H. (1988). On the decomposition of changes in expectation of life and differentials in life expectancy. *Demography* 25(2): 265–276.
- Shkolnikov, V.M., Andreev, E.M., Zhang, Z., Oeppen, J., and Vaupel, J.W. (2011). Losses of expected lifetime in the United States and other developed countries: Methods and empirical analyses. *Demography* 48(1): 211–239.
- Shkolnikov, V.M., Andreev, E.E., and Begun, A.Z. (2003). Gini coefficient as a life table function: Computation from discrete data, decomposition of differences and empirical examples. *Demographic Research* 8(11): 305–358.
- Smits, J. and Monden, C. (2009). Length of life inequality around the globe. *Social Science & Medicine* 68(6): 1114–1123.
- Tuljapurkar, S. and Edwards, R.D. (2011). Variance in death and its implications for modeling and forecasting mortality. *Demographic Research* 24(21): 497–526.
- van Raalte, A.A. and Caswell, H. (2013). Perturbation analysis of indices of lifespan variability. *Demography* 50(5): 1615–1640.

- van Raalte, A.A., Sasson, I., and Martikainen, P. (2018). The case for monitoring life-span inequality. *Science* .
- Vaupel, J.W. (1986). How change in age-specific mortality affects life expectancy. *Population Studies* 40(1): 147–157.
- Vaupel, J.W. and Canudas-Romo, V. (2003). Decomposing change in life expectancy: A bouquet of formulas in honor of Nathan Keyfitz’s 90th birthday. *Demography* 40(2): 201–216.
- Vaupel, J.W., Zhang, Z., and van Raalte, A.A. (2011). Life expectancy and disparity: An international comparison of life table data. *BMJ Open* bmjopen–2011–000128.
- Wagner, P. (2010). Sensitivity of life disparity with respect to changes in mortality rates. *Demographic Research* 23(3): 63–72.
- Wrycza, T.F., Missov, T.I., and Baudisch, A. (2015). Quantifying the shape of aging. *PLoS ONE* 10(3): e0119163.
- Zhang, Z. and Vaupel, J.W. (2009). The age separating early deaths from late deaths. *Demographic Research* 20(29): 721–730.

Appendix

Proposition 1. Let $e^\dagger(x) = \int_x^\infty d(a) e(a) da / \ell(x)$ be a measure of lifespan disparity above age x , where $d(a)$ accounts for the distribution of deaths, $e(a)$ the remaining life expectancy at age a , and $\ell(x)$ is the probability of surviving from birth to age x . Then,

$$(A1) \quad e^\dagger(x) = \frac{1}{\ell(x)} \int_x^\infty \ell(a) (H(a) - H(x)) da ,$$

where $H(x)$ is the cumulative hazard to age x .

Proof. Note that

$$\frac{1}{\ell(x)} \int_x^\infty \ell(a) (H(a) - H(x)) da = \frac{1}{\ell(x)} \int_x^\infty \ell(a) \int_x^a \mu(y) dy da ,$$

where function $\mu(y)$ is the force of mortality or hazard rate. By reversing the order of integration, and using that $e(y) = \int_y^\infty \ell(a) da / \ell(y)$ and $d(y) = \mu(y) \ell(y)$, we get

$$\begin{aligned} \frac{1}{\ell(x)} \int_x^\infty \ell(a) \int_x^a \mu(y) dy da &= \frac{1}{\ell(x)} \int_x^\infty \mu(y) \int_y^\infty \ell(a) da dy \\ &= \frac{1}{\ell(x)} \int_x^\infty \mu(y) \ell(y) e(y) dy \\ &= \frac{1}{\ell(x)} \int_x^\infty d(y) e(y) dy \\ &= e^\dagger(x) , \end{aligned}$$

which proves (A1). □

Proposition 2. Let $\ell(x)$ be the probability of surviving from birth to age x . Let \bar{H} be the lifeable entropy and $\bar{H}(x) = e^\dagger(x) / e(x)$ the lifeable entropy conditioned on reaching age x . Let $H(x)$ be the cumulative hazard to age x . Then, $g(x) = H(x) + \bar{H}(x) - 1 - \bar{H}$ is a strictly increasing function.

Proof. In order to demonstrate that $g(x)$ is a strictly increasing function it is sufficient to show that its first derivative is always positive. We must prove that

$$(A2) \quad \frac{\partial}{\partial x} g(x) = \frac{\partial}{\partial x} (H(x) + \bar{H}(x) - 1 - \bar{H}) = \frac{\partial}{\partial x} H(x) + \frac{\partial}{\partial x} \bar{H}(x) > 0$$

for all ages x .

By the fundamental theorem of calculus,

$$(A3) \quad \frac{\partial}{\partial x} H(x) = \frac{\partial}{\partial x} \int_0^x \mu(a) da = \mu(x) ,$$

whereas

$$\frac{\partial}{\partial x} \bar{H}(x) = \frac{\partial}{\partial x} \left(\frac{e^\dagger(x)}{e(x)} \right) = \frac{1}{e(x)^2} \left(e(x) \frac{\partial}{\partial x} e^\dagger(x) - e^\dagger(x) \frac{\partial}{\partial x} e(x) \right) .$$

First, note that

$$\begin{aligned} \frac{\partial}{\partial x} e^\dagger(x) &= \frac{\partial}{\partial x} \left(\frac{1}{\ell(x)} \int_x^\infty d(a) e(a) da \right) \\ &= \frac{1}{\ell(x)^2} \left(\ell(x) \frac{\partial}{\partial x} \left(\int_x^\infty d(a) e(a) da \right) - \int_x^\infty d(a) e(a) da \frac{\partial}{\partial x} \ell(x) \right) \\ (A4) \quad &= \frac{1}{\ell(x)^2} \left(\ell(x) (-d(x) e(x)) - \int_x^\infty d(a) e(a) da (-\mu(x) \ell(x)) \right) \\ &= -\frac{\mu(x) \ell(x) e(x)}{\ell(x)} + \mu(x) \frac{\int_x^\infty d(a) e(a) da}{\ell(x)} \\ &= \mu(x) (e^\dagger(x) - e(x)) . \end{aligned}$$

On the other hand,

$$\begin{aligned} \frac{\partial}{\partial x} e(x) &= \frac{\partial}{\partial x} \left(\frac{1}{\ell(x)} \int_x^\infty \ell(a) da \right) \\ (A5) \quad &= \frac{1}{\ell(x)^2} \left(\ell(x) \frac{\partial}{\partial x} \left(\int_x^\infty \ell(a) da \right) - \int_x^\infty \ell(a) da \frac{\partial}{\partial x} \ell(x) \right) \\ &= \frac{1}{\ell(x)^2} \left(\ell(x) (-\ell(x)) - \int_x^\infty \ell(a) da (-\mu(x) \ell(x)) \right) \\ &= e(x) \mu(x) - 1 . \end{aligned}$$

Therefore, using (A4) and (A5), we get

$$\begin{aligned} \frac{\partial}{\partial x} \bar{H}(x) &= \frac{1}{e(x)^2} \left(e(x) \mu(x) (e^\dagger(x) - e(x)) - e^\dagger(x) (e(x) \mu(x) - 1) \right) \\ (A6) \quad &= \frac{1}{e(x)^2} (e^\dagger(x) e(x) \mu(x) - e(x)^2 \mu(x) - e^\dagger(x) e(x) \mu(x) + e^\dagger(x)) \\ &= \frac{e^\dagger(x)}{e(x)^2} - \mu(x) . \end{aligned}$$

Finally, replacing (A3) and (A6) in (A2) yields

$$\frac{\partial}{\partial x} g(x) = \mu(x) + \frac{e^\dagger(x)}{e(x)^2} - \mu(x) = \frac{e^\dagger(x)}{e(x)^2} > 0 ,$$

which holds true for all ages since by definition $e^\dagger(x) > 0$ for all $x \geq 0$. Hence, $g(x)$ is a strictly increasing function. \square