

Randomly activated proximal methods for nonsmooth convex minimization

Patrick L. Combettes & Javier I. Madariaga

Department of Mathematics
North Carolina State University
Raleigh, NC 27695, USA

TAGMaC Fall 2024
November 02, 2024

NC STATE UNIVERSITY

Background and motivation

- Throughout, H, G_1, \dots, G_p are separable real Hilbert spaces.

Background and motivation

- Throughout, H, G_1, \dots, G_p are separable real Hilbert spaces.
- $\Gamma_0(H)$ denotes the class of lower semicontinuous convex functions $f: H \rightarrow]-\infty, +\infty]$ such that

$$\text{dom } f = \{x \in H \mid f(x) < +\infty\} \neq \emptyset.$$

Background and motivation

- Throughout, H, G_1, \dots, G_p are separable real Hilbert spaces.
- $\Gamma_0(H)$ denotes the class of lower semicontinuous convex functions $f: H \rightarrow]-\infty, +\infty]$ such that

$$\text{dom } f = \{x \in H \mid f(x) < +\infty\} \neq \emptyset.$$

- Let $f \in \Gamma_0(H)$. The subdifferential of f is the operator

$$\partial f: H \rightarrow 2^H: x \mapsto \{x^* \in H \mid (\forall z \in H) \langle z - x \mid x^* \rangle + f(x) \leq f(z)\}$$

and the proximity operator of f is

$$\text{prox}_f: H \rightarrow H: x \mapsto \underset{z \in H}{\operatorname{argmin}} \left(f(z) + \frac{1}{2} \|x - z\|^2 \right).$$

Background and motivation

- Let C be a nonempty closed convex subset of H . Then I_C denotes the indicator function of C and $\text{proj}_C = \text{prox}_{I_C}$ the projection operator onto C .

Background and motivation

- Let C be a nonempty closed convex subset of H . Then ι_C denotes the indicator function of C and $\text{proj}_C = \text{prox}_{\iota_C}$ the projection operator onto C .
- The underlying probability space is (Ω, \mathcal{F}, P) . \mathcal{B}_H denotes the Borel σ -algebra of H . An H -valued random variable is a measurable mapping $x: (\Omega, \mathcal{F}) \rightarrow (H, \mathcal{B}_H)$.

Background and motivation

- Let C be a nonempty closed convex subset of H . Then ι_C denotes the indicator function of C and $\text{proj}_C = \text{prox}_{\iota_C}$ the projection operator onto C .
- The underlying probability space is (Ω, \mathcal{F}, P) . \mathcal{B}_H denotes the Borel σ -algebra of H . An H -valued random variable is a measurable mapping $x: (\Omega, \mathcal{F}) \rightarrow (H, \mathcal{B}_H)$.
- We use sans-serif letters to denote deterministic variables and italicized serif letters to denote random variables.

Background and motivation

Problem 1

H is a separable real Hilbert space and $f \in \Gamma_0(H)$. For every $k \in \{1, \dots, p\}$, G_k is a separable real Hilbert space, $g_k \in \Gamma_0(G_k)$, and $0 \neq L_k: H \rightarrow G_k$ is linear and bounded. It is assumed that

$$\text{zer}\left(\partial f + \sum_{k=1}^p L_k^* \circ \partial g_k \circ L_k\right) \neq \emptyset.$$

The task is to

$$\underset{x \in H}{\text{minimize}} \quad f(x) + \sum_{k=1}^p g_k(L_k x)$$

and the set of solutions is denoted by Z .

Background and motivation

Problem 1

H is a separable real Hilbert space and $f \in \Gamma_0(H)$. For every $k \in \{1, \dots, p\}$, G_k is a separable real Hilbert space, $g_k \in \Gamma_0(G_k)$, and $0 \neq L_k: H \rightarrow G_k$ is linear and bounded. It is assumed that

$$\text{zer}\left(\partial f + \sum_{k=1}^p L_k^* \circ \partial g_k \circ L_k\right) \neq \emptyset.$$

The task is to

$$\underset{x \in H}{\text{minimize}} \quad f(x) + \sum_{k=1}^p g_k(L_k x)$$

and the set of solutions is denoted by Z .

It covers a wide range of minimization models in data analysis and it is an essential tool in signal processing, statistics, inverse problems, and machine learning.

Background and motivation

We aim at designing **proximal algorithms** which are **stochastic** in the sense that they activate a randomly selected block of functions at each iteration

Background and motivation

We aim at designing **proximal algorithms** which are **stochastic** in the sense that they activate a randomly selected block of functions at each iteration

Furthermore, they should satisfy:

R1: They **guarantee the convergence** of the sequence of iterates to a solution to Problem 1 without any additional assumptions.

Background and motivation

We aim at designing **proximal algorithms** which are **stochastic** in the sense that they activate a randomly selected block of functions at each iteration

Furthermore, they should satisfy:

- R1:** They **guarantee the convergence** of the sequence of iterates to a solution to Problem 1 without any additional assumptions.
- R2:** At each iteration, more than one randomly selected function (f, g_1, \dots, g_p) can be activated.

Background and motivation

We aim at designing **proximal algorithms** which are **stochastic** in the sense that they activate a randomly selected block of functions at each iteration

Furthermore, they should satisfy:

- R1:** They **guarantee the convergence** of the sequence of iterates to a solution to Problem 1 without any additional assumptions.
- R2:** At each iteration, more than one randomly selected function (f, g_1, \dots, g_p) can be activated.
- R3:** Knowledge of bounds on the norms of the linear operators $(L_k)_{1 \leq k \leq p}$ is not required.

Background and motivation

There does not seem to exist methods that satisfy simultaneously **R1–R3**.

Background and motivation

There does not seem to exist methods that satisfy simultaneously **R1–R3**.

Our main contribution is to proposed three algorithmic frameworks that comply with **R1–R3**.

Background and motivation

There does not seem to exist methods that satisfy simultaneously **R1–R3**.

Our main contribution is to proposed three algorithmic frameworks that comply with **R1–R3**.

The strategy consists in embedding Problem 1 into a **multivariate problem**.

Problem 2

- Let $(X_i)_{1 \leq i \leq m}$ and $(Y_j)_{1 \leq j \leq r}$ be families of separable real Hilbert spaces with direct Hilbert sums $\mathbf{X} = X_1 \oplus \cdots \oplus X_m$ and $\mathbf{Y} = Y_1 \oplus \cdots \oplus Y_r$.

Problem 2

- Let $(X_i)_{1 \leq i \leq m}$ and $(Y_j)_{1 \leq j \leq r}$ be families of separable real Hilbert spaces with direct Hilbert sums $X = X_1 \oplus \cdots \oplus X_m$ and $Y = Y_1 \oplus \cdots \oplus Y_r$.
- For every $i \in \{1, \dots, m\}$, let $f_i \in \Gamma_0(X_i)$ and, for every $j \in \{1, \dots, r\}$, let $h_j \in \Gamma_0(Y_j)$, and let $M_{ji}: X_i \rightarrow Y_j$ be linear and bounded.

Problem 2

- Let $(X_i)_{1 \leq i \leq m}$ and $(Y_j)_{1 \leq j \leq r}$ be families of separable real Hilbert spaces with direct Hilbert sums $\mathbf{X} = X_1 \oplus \cdots \oplus X_m$ and $\mathbf{Y} = Y_1 \oplus \cdots \oplus Y_r$.
- For every $i \in \{1, \dots, m\}$, let $f_i \in \Gamma_0(X_i)$ and, for every $j \in \{1, \dots, r\}$, let $h_j \in \Gamma_0(Y_j)$, and let $M_{ji}: X_i \rightarrow Y_j$ be linear and bounded.

The task is to

$$\underset{\mathbf{x} \in \mathbf{X}}{\text{minimize}} \quad \sum_{i=1}^m f_i(x_i) + \sum_{j=1}^r h_j\left(\sum_{i=1}^m M_{ji}x_i\right).$$

The set of solutions to Problem 2 is denoted by \mathbf{Z} .

General Framework

The set of solutions to Problem 2 is denoted by \mathbf{Z} . Further, the projection operator onto the subspace

$$\mathbf{V} = \left\{ (\mathbf{x}, \mathbf{y}) \in \mathbf{X} \oplus \mathbf{Y} \mid (\forall j \in \{1, \dots, r\}) y_j = \sum_{i=1}^m M_{ji} x_i \right\}$$

is decomposed as $\text{proj}_{\mathbf{V}}: (\mathbf{x}, \mathbf{y}) \mapsto (\mathbf{Q}_i(\mathbf{x}, \mathbf{y}))_{1 \leq i \leq m+r}$, where for every $i \in \{1, \dots, m\}$, $\mathbf{Q}_i: \mathbf{X} \oplus \mathbf{Y} \rightarrow \mathbf{X}_i$ and, for every $j \in \{1, \dots, r\}$, $\mathbf{Q}_{m+j}: \mathbf{X} \oplus \mathbf{Y} \rightarrow \mathbf{Y}_j$.

Theorem 3...

Douglas–Rachford method yields the following result.

Theorem 3...

Douglas–Rachford method yields the following result.

➤ Let $\gamma \in]0, +\infty[$,

Theorem 3...

Douglas–Rachford method yields the following result.

- Let $\gamma \in]0, +\infty[$,
- let $(\lambda_n)_{n \in \mathbb{N}}$ be a sequence in $]0, 2[$ such that $\inf_{n \in \mathbb{N}} \lambda_n > 0$ and $\sup_{n \in \mathbb{N}} \lambda_n < 2$,

Theorem 3...

Douglas–Rachford method yields the following result.

- Let $\gamma \in]0, +\infty[$,
- let $(\lambda_n)_{n \in \mathbb{N}}$ be a sequence in $]0, 2[$ such that $\inf_{n \in \mathbb{N}} \lambda_n > 0$ and $\sup_{n \in \mathbb{N}} \lambda_n < 2$,
- let \mathbf{x}_0 and \mathbf{z}_0 be \mathbf{X} -valued random variables,

Theorem 3...

Douglas–Rachford method yields the following result.

- Let $\gamma \in]0, +\infty[$,
- let $(\lambda_n)_{n \in \mathbb{N}}$ be a sequence in $]0, 2[$ such that $\inf_{n \in \mathbb{N}} \lambda_n > 0$ and $\sup_{n \in \mathbb{N}} \lambda_n < 2$,
- let \mathbf{x}_0 and \mathbf{z}_0 be \mathbf{X} -valued random variables,
- let \mathbf{y}_0 and \mathbf{w}_0 be \mathbf{Y} -valued random variables,

Theorem 3...

Douglas–Rachford method yields the following result.

- Let $\gamma \in]0, +\infty[$,
- let $(\lambda_n)_{n \in \mathbb{N}}$ be a sequence in $]0, 2[$ such that $\inf_{n \in \mathbb{N}} \lambda_n > 0$ and $\sup_{n \in \mathbb{N}} \lambda_n < 2$,
- let \mathbf{x}_0 and \mathbf{z}_0 be **X**-valued random variables,
- let \mathbf{y}_0 and \mathbf{w}_0 be **Y**-valued random variables,
- set $D = \{0, 1\}^{m+r} \setminus \{\mathbf{0}\}$, and let $(\varepsilon_n)_{n \in \mathbb{N}}$ be identically distributed D -valued random variables. (**)

...Theorem 3

Iterate

```
for n = 0, 1, ...  
  for i = 1, ..., m  
     $x_{i,n+1} = x_{i,n} + \varepsilon_{i,n}(\mathbf{Q}_i(\mathbf{z}_n, \mathbf{w}_n) - x_{i,n})$   
     $z_{i,n+1} = z_{i,n} + \varepsilon_{i,n}\lambda_n(\text{prox}_{\gamma f_i}(2x_{i,n+1} - z_{i,n}) - x_{i,n+1})$   
  for j = 1, ..., r  
     $y_{j,n+1} = y_{j,n} + \varepsilon_{m+j,n}(\mathbf{Q}_{m+j}(\mathbf{z}_n, \mathbf{w}_n) - y_{j,n})$   
     $w_{j,n+1} = w_{j,n} + \varepsilon_{m+j,n}\lambda_n(\text{prox}_{\gamma h_j}(2y_{j,n+1} - w_{j,n}) - y_{j,n+1}).$ 
```

...Theorem 3

Iterate

$$\begin{array}{l} \text{for } n = 0, 1, \dots \\ \quad \left[\begin{array}{l} \text{for } i = 1, \dots, m \\ \quad \left[\begin{array}{l} x_{i,n+1} = x_{i,n} + \varepsilon_{i,n} (\mathcal{Q}_i(\mathbf{z}_n, \mathbf{w}_n) - x_{i,n}) \\ z_{i,n+1} = z_{i,n} + \varepsilon_{i,n} \lambda_n (\text{prox}_{\gamma f_i}(2x_{i,n+1} - z_{i,n}) - x_{i,n+1}) \end{array} \right] \\ \text{for } j = 1, \dots, r \\ \quad \left[\begin{array}{l} y_{j,n+1} = y_{j,n} + \varepsilon_{m+j,n} (\mathcal{Q}_{m+j}(\mathbf{z}_n, \mathbf{w}_n) - y_{j,n}) \\ w_{j,n+1} = w_{j,n} + \varepsilon_{m+j,n} \lambda_n (\text{prox}_{\gamma h_j}(2y_{j,n+1} - w_{j,n}) - y_{j,n+1}). \end{array} \right] \end{array} \right. \end{array}$$

Then $(\mathbf{x}_n)_{n \in \mathbb{N}}$ converges weakly P-a.s. to a \mathbf{Z} -valued random variable.

How can we use this general framework to solve Problem 1?

Framework 1

How can we use this general framework to solve Problem 1?

$$\underset{\mathbf{x} \in \mathbf{X}}{\text{minimize}} \quad \sum_{i=1}^m f_i(x_i) + \sum_{j=1}^r h_j \left(\sum_{i=1}^m M_{ji} x_i \right). \quad (\text{GF})$$

$$\underset{x \in \mathbf{H}}{\text{minimize}} \quad f(x) + \sum_{k=1}^p g_k(L_k x). \quad (\text{P1})$$

Framework 1

How can we use this general framework to solve Problem 1?

$$\underset{\mathbf{x} \in \mathbf{X}}{\text{minimize}} \quad \sum_{i=1}^m \mathbf{f}_i(\mathbf{x}_i) + \sum_{j=1}^r \mathbf{h}_j \left(\sum_{i=1}^m \mathbf{M}_{ji} \mathbf{x}_i \right). \quad (\text{GF})$$

$$\underset{\mathbf{x} \in \mathbf{H}}{\text{minimize}} \quad \mathbf{f}(\mathbf{x}) + \sum_{k=1}^p \mathbf{g}_k(\mathbf{L}_k \mathbf{x}). \quad (\text{P1})$$

The same for $m = 1$, $r = p$, $\mathbf{X}_1 = \mathbf{H}$, $\mathbf{f}_1 = \mathbf{f}$, and $(\forall k \in \{1, \dots, p\})$
 $\mathbf{Y}_k = \mathbf{G}_k$, $\mathbf{M}_{k,1} = \mathbf{L}_k$, and $\mathbf{h}_k = \mathbf{g}_k$.

Proposition 4

Set $D = \{0, 1\}^{1+p} \setminus \{\mathbf{0}\}$ and consider Theorem 3 with

Proposition 4

Set $D = \{0, 1\}^{1+p} \setminus \{\mathbf{0}\}$ and consider Theorem 3 with

for $n = 0, 1, \dots$

$$\left[\begin{array}{l} \mathbf{q}_n = (\text{Id} + \sum_{k=1}^p \mathbf{L}_k^* \circ \mathbf{L}_k)^{-1} (z_n + \sum_{k=1}^p \mathbf{L}_k^* w_{k,n}) \\ \mathbf{Q}_1(\mathbf{z}_n, \mathbf{w}_n) = \mathbf{q}_n \end{array} \right.$$

Proposition 4

Set $D = \{0, 1\}^{1+p} \setminus \{\mathbf{0}\}$ and consider Theorem 3 with

$$\begin{array}{l} \text{for } n = 0, 1, \dots \\ \left[\begin{array}{l} \mathbf{q}_n = (\text{Id} + \sum_{k=1}^p \mathbf{L}_k^* \circ \mathbf{L}_k)^{-1} (\mathbf{z}_n + \sum_{k=1}^p \mathbf{L}_k^* \mathbf{w}_{k,n}) \\ \mathbf{Q}_1(\mathbf{z}_n, \mathbf{w}_n) = \mathbf{q}_n \\ \text{for } k = 1, \dots, p \\ \quad \left[\begin{array}{l} \mathbf{Q}_{1+k}(\mathbf{z}_n, \mathbf{w}_n) = \mathbf{L}_k \mathbf{q}_n. \end{array} \right. \end{array} \right. \end{array}$$

Then $(x_n)_{n \in \mathbb{N}}$ converges weakly P-a.s. to a Z -valued random variable.

Can we use (GF) to solve the original problem differently?

Framework 2

Can we use (GF) to solve the original problem differently? Set $\mathbf{G} = \mathbf{G}_1 \oplus \dots \oplus \mathbf{G}_p$ and

$$\mathbf{W} = \left\{ \mathbf{x} \in \mathbf{H} \oplus \mathbf{G} \mid (\forall k \in \{1, \dots, p\}) x_{k+1} = \mathbf{L}_k x_1 \right\}.$$

Framework 2

Can we use (GF) to solve the original problem differently? Set $\mathbf{G} = \mathbf{G}_1 \oplus \dots \oplus \mathbf{G}_p$ and

$$\mathbf{W} = \left\{ \mathbf{x} \in \mathbf{H} \oplus \mathbf{G} \mid (\forall k \in \{1, \dots, p\}) x_{k+1} = \mathbf{L}_k x_1 \right\}.$$

Problem 5

Set $\mathbf{f}_1 = \mathbf{f}$ and, for every $i \in \{2, \dots, p+1\}$, $\mathbf{f}_i = \mathbf{g}_{i-1}$. Denote by $\mathbf{x} = (x_1, \dots, x_{p+1})$ a generic element in $\mathbf{H} \oplus \mathbf{G}$. The task is to

$$\underset{\mathbf{x} \in \mathbf{H} \oplus \mathbf{G}}{\text{minimize}} \quad \sum_{i=1}^{p+1} \mathbf{f}_i(x_i)$$

Framework 2

Can we use (GF) to solve the original problem differently? Set $\mathbf{G} = \mathbf{G}_1 \oplus \cdots \oplus \mathbf{G}_p$ and

$$\mathbf{W} = \left\{ \mathbf{x} \in \mathbf{H} \oplus \mathbf{G} \mid (\forall k \in \{1, \dots, p\}) x_{k+1} = \mathbf{L}_k x_1 \right\}.$$

Problem 5

Set $\mathbf{f}_1 = \mathbf{f}$ and, for every $i \in \{2, \dots, p+1\}$, $\mathbf{f}_i = \mathbf{g}_{i-1}$. Denote by $\mathbf{x} = (x_1, \dots, x_{p+1})$ a generic element in $\mathbf{H} \oplus \mathbf{G}$. The task is to

$$\underset{\mathbf{x} \in \mathbf{H} \oplus \mathbf{G}}{\text{minimize}} \quad \sum_{i=1}^{p+1} \mathbf{f}_i(x_i) + \iota_{\mathbf{W}}(\mathbf{x}).$$

Proposition 6

Set $D = \{0, 1\}^{p+2} \setminus \{\mathbf{0}\}$ and consider Theorem 3 with

Proposition 6

Set $D = \{0, 1\}^{p+2} \setminus \{\mathbf{0}\}$ and consider Theorem 3 with

for $n = 0, 1, \dots$

 for $i = 1, \dots, p + 1$

$Q_i(\mathbf{z}_n, \mathbf{w}_n) = \frac{1}{2}(z_{i,n} + w_{i,n})$

Proposition 6

Set $D = \{0, 1\}^{p+2} \setminus \{\mathbf{0}\}$ and consider Theorem 3 with

for $n = 0, 1, \dots$

$$\left[\begin{array}{l} \text{for } i = 1, \dots, p+1 \\ \quad \mathbf{Q}_i(\mathbf{z}_n, \mathbf{w}_n) = \frac{1}{2}(\mathbf{z}_{i,n} + \mathbf{w}_{i,n}) \\ \mathbf{Q}_{p+2}(\mathbf{z}_n, \mathbf{w}_n) = \frac{1}{2}(\mathbf{z}_n + \mathbf{w}_n) \\ \mathbf{q}_n = (\text{Id} + \sum_{k=1}^p \mathbf{L}_k^* \circ \mathbf{L}_k)^{-1} (2\mathbf{y}_{1,n+1} - \mathbf{w}_{1,n} + \sum_{k=1}^p \mathbf{L}_k^* (2\mathbf{y}_{k+1,n+1} - \mathbf{w}_{k+1,n})) \\ \text{prox}_{\mathbf{L}\mathbf{w}}(2\mathbf{y}_{k,n+1} - \mathbf{w}_{k,n}) = (\mathbf{q}_n, \mathbf{L}_1 \mathbf{q}_n, \dots, \mathbf{L}_p \mathbf{q}_n) \end{array} \right.$$

Proposition 6

Set $D = \{0, 1\}^{p+2} \setminus \{\mathbf{0}\}$ and consider Theorem 3 with

for $n = 0, 1, \dots$

$$\left[\begin{array}{l} \text{for } i = 1, \dots, p+1 \\ \quad \mathbf{Q}_i(\mathbf{z}_n, \mathbf{w}_n) = \frac{1}{2}(\mathbf{z}_{i,n} + \mathbf{w}_{i,n}) \\ \mathbf{Q}_{p+2}(\mathbf{z}_n, \mathbf{w}_n) = \frac{1}{2}(\mathbf{z}_n + \mathbf{w}_n) \\ \mathbf{q}_n = (\text{Id} + \sum_{k=1}^p \mathbf{L}_k^* \circ \mathbf{L}_k)^{-1} (2\mathbf{y}_{1,n+1} - \mathbf{w}_{1,n} + \sum_{k=1}^p \mathbf{L}_k^* (2\mathbf{y}_{k+1,n+1} - \mathbf{w}_{k+1,n})) \\ \text{prox}_{\mathbf{w}}(2\mathbf{y}_{k,n+1} - \mathbf{w}_{k,n}) = (\mathbf{q}_n, \mathbf{L}_1 \mathbf{q}_n, \dots, \mathbf{L}_p \mathbf{q}_n) \end{array} \right.$$

Then $(x_{1,n})_{n \in \mathbb{N}}$ converges weakly P-a.s. to a \mathbb{Z} -valued random variable.

We extend the same idea...

Problem 7

Consider the problem

$$\underset{\mathbf{x} \in \mathbf{H} \oplus \mathbf{G}}{\text{minimize}} \quad \sum_{i=1}^{p+1} f_i(x_i)$$

Framework 3

We extend the same idea...

Problem 7

Consider the problem

$$\underset{\mathbf{x} \in \mathbb{H} \oplus \mathbb{G}}{\text{minimize}} \quad \sum_{i=1}^{p+1} f_i(x_i) + \sum_{k=1}^p \ell_{\{0\}} \left(\sum_{i=1}^{p+1} C_{ki} x_i \right),$$

where

$$C_{ki} = \begin{cases} L_k, & \text{if } i = 1; \\ -Id, & \text{if } i = k + 1; \\ 0, & \text{otherwise.} \end{cases}$$

Proposition 8

Set $D = \{0, 1\}^{2p+1} \setminus \{\mathbf{0}\}$ and consider Theorem 3 with

Proposition 8

Set $D = \{0, 1\}^{2p+1} \setminus \{\mathbf{0}\}$ and consider Theorem 3 with

for $n = 0, 1, \dots$

$$\mathbf{q}_n = (2\text{Id} + \sum_{k=1}^p \mathbf{L}_k^* \circ \mathbf{L}_k)^{-1} (2\mathbf{z}_{1,n} + \sum_{k=1}^p \mathbf{L}_k^* (\mathbf{z}_{k+1,n} + \mathbf{w}_{k,n}))$$

$$\mathbf{Q}_1(\mathbf{z}_n, \mathbf{w}_n) = \mathbf{q}_n$$

for $i = 1, \dots, p$

$$\mathbf{Q}_{1+i}(\mathbf{z}_n, \mathbf{w}_n) = \frac{1}{2} (\mathbf{L}_i \mathbf{q}_n + \mathbf{z}_{i+1,n} - \mathbf{w}_{i,n})$$

for $k = 1, \dots, p$

$$\mathbf{Q}_{1+p+k}(\mathbf{z}_n, \mathbf{w}_n) = \frac{1}{2} (\mathbf{L}_k \mathbf{q}_n - \mathbf{z}_{k+1,n} - \mathbf{w}_{k,n})$$

$$\text{prox}_{\iota_{\{0\}}}(2\mathbf{y}_{k,n+1} - \mathbf{w}_{k,n}) = \mathbf{0}.$$

Proposition 8

Set $D = \{0, 1\}^{2p+1} \setminus \{\mathbf{0}\}$ and consider Theorem 3 with

for $n = 0, 1, \dots$

$$\left[\begin{array}{l} \mathbf{q}_n = (2\text{Id} + \sum_{k=1}^p \mathbf{L}_k^* \circ \mathbf{L}_k)^{-1} (2\mathbf{z}_{1,n} + \sum_{k=1}^p \mathbf{L}_k^* (\mathbf{z}_{k+1,n} + \mathbf{w}_{k,n})) \\ \mathbf{Q}_1(\mathbf{z}_n, \mathbf{w}_n) = \mathbf{q}_n \\ \text{for } i = 1, \dots, p \\ \quad \mathbf{Q}_{1+i}(\mathbf{z}_n, \mathbf{w}_n) = \frac{1}{2} (\mathbf{L}_i \mathbf{q}_n + \mathbf{z}_{i+1,n} - \mathbf{w}_{i,n}) \\ \\ \text{for } k = 1, \dots, p \\ \quad \mathbf{Q}_{1+p+k}(\mathbf{z}_n, \mathbf{w}_n) = \frac{1}{2} (\mathbf{L}_k \mathbf{q}_n - \mathbf{z}_{k+1,n} - \mathbf{w}_{k,n}) \\ \quad \text{prox}_{\iota_{\{0\}}}(2\mathbf{y}_{k,n+1} - \mathbf{w}_{k,n}) = \mathbf{0}. \end{array} \right.$$

Then $(x_{1,n})_{n \in \mathbb{N}}$ converges weakly P-a.s. to a \mathbb{Z} -valued random variable.

Consider the **overlapping group lasso regression** described as follows:

Numerical experiment

Consider the **overlapping group lasso regression** described as follows: $H = \mathbb{R}^N$ and, for every $k \in \{1, \dots, q\}$, $\emptyset \neq I_k \subset \{1, \dots, N\}$ and

$$L_k: \mathbb{R}^N \rightarrow \mathbb{R}^{\text{card } I_k}: x = (\xi_j)_{1 \leq j \leq N} \mapsto (\xi_j)_{j \in I_k}.$$

Further, $\bigcup_{k=1}^q I_k = \{1, \dots, N\}$. The goal is to

$$\underset{x \in \mathbb{R}^N}{\text{minimize}} \quad \frac{\alpha}{2} \|Ax - b\|^2 + \frac{1}{q} \sum_{k=1}^q \|L_k x\|,$$

where $A \in \mathbb{R}^{M \times N}$, $b \in \mathbb{R}^M$, and $\alpha \in]0, +\infty[$.

Numerical experiment

- In the experiment $M = 1000$, $N = 3610$, $q = 40$, and $\lambda = 5/q^2$.

Numerical experiment

- In the experiment $M = 1000$, $N = 3610$, $q = 40$, and $\lambda = 5/q^2$.
- The entries of A are i.i.d. samples from a $\mathcal{N}(0, 1)$ distribution, and the entries of b are i.i.d. samples from a $\mathcal{N}(100, 100)$ distribution.

Numerical experiment

- In the experiment $M = 1000$, $N = 3610$, $q = 40$, and $\lambda = 5/q^2$.
- The entries of A are i.i.d. samples from a $\mathcal{N}(0, 1)$ distribution, and the entries of b are i.i.d. samples from a $\mathcal{N}(100, 100)$ distribution.
- We split the function $\|Ax - b\|^2$ into 30 blocks of 40 entries each, where the entries are selected in order without overlap.

Numerical experiment

- In the experiment $M = 1000$, $N = 3610$, $q = 40$, and $\lambda = 5/q^2$.
- The entries of A are i.i.d. samples from a $\mathcal{N}(0, 1)$ distribution, and the entries of b are i.i.d. samples from a $\mathcal{N}(100, 100)$ distribution.
- We split the function $\|Ax - b\|^2$ into 30 blocks of 40 entries each, where the entries are selected in order without overlap.
- Finally,

$$(\forall k \in \{1, \dots, p\}) \quad I_k = \{90k - 89, \dots, 90k + 10\}.$$

Numerical experiment




Figure 1: Normalized error $20 \log(\|x_{1,n} - x_\infty\| / \|x_{1,0} - x_\infty\|)$ (dB) versus execution time (s). **Green**: Framework 1. **Orange**: Framework 2. **Blue**: Framework 3.

The frameworks differ in terms of storage requirements, use of proximity operators, and use of linear operators.

The frameworks differ in terms of storage requirements, use of **proximity** operators, and use of **linear** operators.

- Framework 1: It stores $2p + 3$ vectors and, for each of the $p + 1$ random activation indices, there is one **proximity** evaluation.

The frameworks differ in terms of storage requirements, use of **proximity** operators, and use of **linear** operators.

- Framework 1: It stores $2p + 3$ vectors and, for each of the $p + 1$ random activation indices, there is one **proximity** evaluation.
- Framework 2: It stores $4p + 5$ vectors. The first $p + 1$ indices involve a **proximity** operator. The **linear** operators are used only if index $p + 2$ is activated.

The frameworks differ in terms of storage requirements, use of **proximity** operators, and use of **linear** operators.

- Framework 1: It stores $2p + 3$ vectors and, for each of the $p + 1$ random activation indices, there is one **proximity** evaluation.
- Framework 2: It stores $4p + 5$ vectors. The first $p + 1$ indices involve a **proximity** operator. The **linear** operators are used only if index $p + 2$ is activated.
- Framework 3: It stores $4p + 3$ vectors. Moreover, out of the $2p+1$ random activation indices, those in $\{1, \dots, p+1\}$ involve a **proximity** operator, while those in $\{p+2, \dots, p+r+1\}$ do not require it.

Although Framework 1 is the most efficient in terms of storage, it may not always be the fastest, especially when **proximity** operators are computationally expensive or when the **linear** operators are costly, which is the case in the experiment.

Randomly activated proximal methods for nonsmooth convex minimization

Patrick L. Combettes & Javier I. Madariaga

Department of Mathematics
North Carolina State University
Raleigh, NC 27695, USA

TAGMaC Fall 2024
November 02, 2024

NC STATE UNIVERSITY