

# Geometrical scheme for stochastic Fejér monotonicity

Javier I. Madariaga

Department of Mathematics  
North Carolina State University  
Raleigh, NC 27695, USA

**NLA Student Seminar**  
November 14, 2024

**NC STATE** UNIVERSITY

- Throughout,  $H$  is a separable real Hilbert space with scalar product  $\langle \cdot | \cdot \rangle$  and norm  $\|\cdot\|$ .

# Background

- Throughout,  $H$  is a separable real Hilbert space with scalar product  $\langle \cdot | \cdot \rangle$  and norm  $\|\cdot\|$ .
- The sets of strong and weak sequential cluster points of a sequence  $(x_n)_{n \in \mathbb{N}}$  in  $H$  are denoted by  $\mathfrak{S}(x_n)_{n \in \mathbb{N}}$  and  $\mathfrak{W}(x_n)_{n \in \mathbb{N}}$ , respectively.

# General problem

## Problem 1

Let  $Z$  be a nonempty closed convex subset of  $H$ .

# General problem

## Problem 1

Let  $Z$  be a nonempty closed convex subset of  $H$ . The task is to

$$\text{Find } \bar{x} \in H \text{ such that } \bar{x} \in Z. \quad (1)$$

# General problem

## Problem 1

Let  $Z$  be a nonempty closed convex subset of  $H$ . The task is to

$$\text{Find } \bar{x} \in H \text{ such that } \bar{x} \in Z. \quad (1)$$

Problem 1 covers many problems in analysis and optimization.

# Example I

For every  $n \in \mathbb{N}$ , let  $T_n: H \rightarrow H$  be a *firmly nonexpansive operator*, i.e.,

$$(\forall (x, y) \in H \times H) \quad \|T_n x - T_n y\|^2 \leq \langle T_n x - T_n y | x - y \rangle. \quad (1)$$

# Example I

For every  $n \in \mathbb{N}$ , let  $T_n: H \rightarrow H$  be a *firmly nonexpansive operator*, i.e.,

$$(\forall (x, y) \in H \times H) \quad \|T_n x - T_n y\|^2 \leq \langle T_n x - T_n y | x - y \rangle. \quad (1)$$

The common fixed point problem, i.e.,

$$\text{Find } \bar{x} \in H \text{ such that } (\forall n \in \mathbb{N}) T_n \bar{x} = \bar{x},$$

is an example of Problem 1.



## Example II

Let  $f: \mathbb{H} \rightarrow ]-\infty, +\infty]$  be a proper, lower semicontinuous convex function.

## Example II

Let  $f: H \rightarrow ]-\infty, +\infty]$  be a proper, lower semicontinuous convex function. The minimization problem

$$\underset{x \in H}{\text{minimize}} \quad f(x).$$

is an example of Problem 1.

# Fejér monotonicity

A sequence  $(x_n)_{n \in \mathbb{N}}$  of vectors in  $H$  is said to be **Fejér monotone** with respect to  $Z$  if

$$(\forall z \in Z)(\forall n \in \mathbb{N}) \quad \|x_{n+1} - z\| \leq \|x_n - z\|.$$

# Fejér monotonicity

A sequence  $(x_n)_{n \in \mathbb{N}}$  of vectors in  $H$  is said to be **Fejér monotone** with respect to  $Z$  if

$$(\forall z \in Z)(\forall n \in \mathbb{N}) \quad \|x_{n+1} - z\| \leq \|x_n - z\|.$$

Some consequences are:

# Fejér monotonicity

A sequence  $(x_n)_{n \in \mathbb{N}}$  of vectors in  $H$  is said to be **Fejér monotone** with respect to  $Z$  if

$$(\forall z \in Z)(\forall n \in \mathbb{N}) \quad \|x_{n+1} - z\| \leq \|x_n - z\|.$$

Some consequences are:

- 1  $(x_n)_{n \in \mathbb{N}}$  is bounded.

# Fejér monotonicity

A sequence  $(x_n)_{n \in \mathbb{N}}$  of vectors in  $H$  is said to be **Fejér monotone** with respect to  $Z$  if

$$(\forall z \in Z)(\forall n \in \mathbb{N}) \quad \|x_{n+1} - z\| \leq \|x_n - z\|.$$

Some consequences are:

- 1  $(x_n)_{n \in \mathbb{N}}$  is bounded.
- 2  $(\forall z \in Z) (\|x_n - z\|)_{n \in \mathbb{N}}$  converges.

# Fejér monotonicity

A sequence  $(x_n)_{n \in \mathbb{N}}$  of vectors in  $H$  is said to be **Fejér monotone** with respect to  $Z$  if

$$(\forall z \in Z)(\forall n \in \mathbb{N}) \quad \|x_{n+1} - z\| \leq \|x_n - z\|.$$

Some consequences are:

- 1  $(x_n)_{n \in \mathbb{N}}$  is bounded.
- 2  $(\forall z \in Z) (\|x_n - z\|)_{n \in \mathbb{N}}$  converges.
- 3 Suppose that  $\mathfrak{B}(x_n)_{n \in \mathbb{N}} \subset Z$ .

# Fejér monotonicity

A sequence  $(x_n)_{n \in \mathbb{N}}$  of vectors in  $H$  is said to be **Fejér monotone** with respect to  $Z$  if

$$(\forall z \in Z)(\forall n \in \mathbb{N}) \quad \|x_{n+1} - z\| \leq \|x_n - z\|.$$

Some consequences are:

- 1  $(x_n)_{n \in \mathbb{N}}$  is bounded.
- 2  $(\forall z \in Z) (\|x_n - z\|)_{n \in \mathbb{N}}$  converges.
- 3 Suppose that  $\mathfrak{B}(x_n)_{n \in \mathbb{N}} \subset Z$ . Then  $(x_n)_{n \in \mathbb{N}}$  converges weakly to some vector in  $Z$ .



# Fejérian scheme

Let  $x_0 \in H$ . Iterate


# Fejérian scheme

Let  $x_0 \in H$ . Iterate

for  $n = 0, 1, \dots$

$$\left[ \begin{array}{l} t_n^* \in H \text{ and } \eta_n \in \mathbb{R} \text{ satisfy} \\ \quad \alpha_n = \begin{cases} \frac{\langle x_n | t_n^* \rangle - \eta_n}{\|t_n^*\|^2}, & \text{if } t_n^* \neq 0 \text{ and } \langle x_n | t_n^* \rangle > \eta_n; \\ 0, & \text{otherwise.} \end{cases} \\ \quad (\forall z \in Z) \quad \langle z | \alpha_n t_n^* \rangle \leq \alpha_n \eta_n \\ x_{n+1} = x_n - \alpha_n t_n^*. \end{array} \right.$$

# Fejérian scheme



**Main issue:** How to construct/select  $t_n^*$  and  $\eta_n$  such that

$$\mathfrak{B}(x_n)_{n \in \mathbb{N}} \subset \mathbb{Z} ?$$


# Fejérian scheme



When transforming those algorithms into stochastic algorithms,  
**Fejér monotonicity is no longer true.**


# Stochastic algorithms

Let  $f_1: x \mapsto (x+1)^2$  and  $f_2: x \mapsto (x-1)^2$ . The task is to minimize  $f_1 + f_2$  using the stochastic gradient method.



# Stochastic algorithms

Let  $f_1: x \mapsto (x+1)^2$  and  $f_2: x \mapsto (x-1)^2$ . The task is to minimize  $f_1 + f_2$  using the stochastic gradient method.



No Fejér monotonicity (in general)!



- The underlying probability space is  $(\Omega, \mathcal{F}, P)$ . Denote as  $\mathcal{B}_H$  the Borel  $\sigma$ -algebra of  $H$ . An  $H$ -valued random variable is a measurable mapping  $x: (\Omega, \mathcal{F}) \rightarrow (H, \mathcal{B}_H)$ .

# Background

- The underlying probability space is  $(\Omega, \mathcal{F}, P)$ . Denote as  $\mathcal{B}_H$  the Borel  $\sigma$ -algebra of  $H$ . An  $H$ -valued random variable is a measurable mapping  $x: (\Omega, \mathcal{F}) \rightarrow (H, \mathcal{B}_H)$ .
- Given  $p \in [1, +\infty[$ ,  $L^p(\Omega, \mathcal{F}, P; H)$  denotes the space of  $H$ -valued random variables  $x: (\Omega, \mathcal{F}) \rightarrow (H, \mathcal{B}_H)$  such that

$$E\|x\|^p < +\infty.$$

# Background

- The underlying probability space is  $(\Omega, \mathcal{F}, P)$ . Denote as  $\mathcal{B}_H$  the Borel  $\sigma$ -algebra of  $H$ . An  $H$ -valued random variable is a measurable mapping  $x: (\Omega, \mathcal{F}) \rightarrow (H, \mathcal{B}_H)$ .
- Given  $p \in [1, +\infty[$ ,  $L^p(\Omega, \mathcal{F}, P; H)$  denotes the space of  $H$ -valued random variables  $x: (\Omega, \mathcal{F}) \rightarrow (H, \mathcal{B}_H)$  such that

$$E\|x\|^p < +\infty.$$

Further,

$$(\forall A \in \mathcal{B}_H) \quad L^p(\Omega, \mathcal{F}, P; A) = \{x \in L^p(\Omega, \mathcal{F}, P; H) \mid x \in A \text{ P-a.s.}\}$$

# Stochastic quasi-Fejér monotonicity

- In the late 60s, Ermol'ev introduce some notion of Random Fejérity.

# Stochastic quasi-Fejér monotonicity

- In the late 60s, Ermol'ev introduce some notion of Random Fejérity.
- This was highly ignored until 10 years ago.

# Stochastic quasi-Fejér monotonicity

- In the late 60s, Ermol'ev introduce some notion of Random Fejérity.
- This was highly ignored until 10 years ago.
- One of the biggest contributor is (Combettes and Pesquet 2015)

# Stochastic quasi-Fejér monotonicity

- In the late 60s, Ermol'ev introduce some notion of Random Fejérity.
- This was highly ignored until 10 years ago.
- One of the biggest contributor is (Combettes and Pesquet 2015)

A sequence  $(x_n)_{n \in \mathbb{N}}$  of  $H$ -valued random variables is said to be **stochastic quasi-Fejér monotone** with respect to  $Z$  if there exists a sequence of  $[0, +\infty[$ -valued random variables  $(\varepsilon_n)_{n \in \mathbb{N}}$  such that

$$(\forall z \in Z)(\forall n \in \mathbb{N}) \quad \begin{cases} x_n = \sigma(x_0, \dots, x_n) \\ E(\|x_{n+1} - z\| \mid \mathcal{X}_n) \leq \|x_n - z\| + 2\varepsilon_n. \end{cases}$$

No geometrical interpretation of this!



# Stochastic quasi-Fejér monotonicity

No geometrical interpretation of this!

We propose a geometrical framework...

# Stochastic quasi-Fejérian scheme

Let  $x_0 \in L^2(\Omega, \mathcal{F}, P; \mathbb{H})$ . Iterate

for  $n = 0, 1, \dots$

$$\left[ \begin{array}{l} \mathcal{X}_n = \sigma(x_0, \dots, x_n) \\ t_n^* \in L^2(\Omega, \mathcal{F}, P; \mathbb{H}), \eta_n \in L^1(\Omega, \mathcal{F}, P; \mathbb{R}), \text{ and } \varepsilon_n \in L^1(\Omega, \mathcal{F}, P; [0, +\infty[) \\ \text{satisfy} \\ \left\{ \begin{array}{l} 1_{[t_n^* \neq 0]} \eta_n / (\|t_n^*\| + 1_{[t_n^* = 0]}) \in L^2(\Omega, \mathcal{F}, P; \mathbb{R}) \\ \alpha_n = \frac{1_{[t_n^* \neq 0 \text{ and } \langle x_n | t_n^* \rangle > \eta_n]} (\langle x_n | t_n^* \rangle - \eta_n)}{\|t_n^*\|^2 + 1_{[t_n^* = 0]}} \\ (\forall Z \in \mathbb{Z}) \langle Z | E(\alpha_n t_n^* | \mathcal{X}_n) \rangle \leq E(\alpha_n \eta_n | \mathcal{X}_n) + \varepsilon_n \text{ P-a.s.} \end{array} \right. \\ x_{n+1} = x_n - \alpha_n t_n^*. \end{array} \right.$$

# Some results

- ① Let  $n \in \mathbb{N}$  and  $z \in \mathbb{Z}$ . Then  
$$E(\|x_{n+1} - z\|^2 \mid \mathcal{X}_n) \leq \|x_n - z\|^2 + 2\varepsilon_n \text{ P-a.s.}$$

# Some results

- 1 Let  $n \in \mathbb{N}$  and  $z \in \mathbb{Z}$ . Then  
$$E(\|x_{n+1} - z\|^2 \mid \mathcal{X}_n) \leq \|x_n - z\|^2 + 2\varepsilon_n \text{ P-a.s.}$$
- 2 Let  $n \in \mathbb{N}$  and  $z \in L^2(\Omega, \mathcal{X}_n, P; \mathbb{Z})$ . Then  
$$E(\|x_{n+1} - z\|^2 \mid \mathcal{X}_n) \leq \|x_n - z\|^2 + 2\varepsilon_n \text{ P-a.s.}$$

- 1 Let  $n \in \mathbb{N}$  and  $z \in \mathcal{Z}$ . Then
$$E(\|x_{n+1} - z\|^2 \mid \mathcal{X}_n) \leq \|x_n - z\|^2 + 2\varepsilon_n \text{ P-a.s.}$$
- 2 Let  $n \in \mathbb{N}$  and  $z \in L^2(\Omega, \mathcal{X}_n, P; \mathcal{Z})$ . Then
$$E(\|x_{n+1} - z\|^2 \mid \mathcal{X}_n) \leq \|x_n - z\|^2 + 2\varepsilon_n \text{ P-a.s.}$$
- 3 Let  $n \in \mathbb{N}$  and  $z \in L^2(\Omega, \mathcal{X}_n, P; \mathcal{Z})$ . Then
$$\|x_{n+1} - z\|_{L^2(\Omega, \mathcal{F}, P; \mathcal{H})}^2 \leq \|x_n - z\|_{L^2(\Omega, \mathcal{F}, P; \mathcal{H})}^2 + 2E\varepsilon_n.$$

Suppose that  $\sum_{n \in \mathbb{N}} \varepsilon_n < +\infty$  P-a.s. Then the following hold:

- 1 The sequence  $(\|x_n\|)_{n \in \mathbb{N}}$  is bounded P-a.s.

# Some results

Suppose that  $\sum_{n \in \mathbb{N}} \varepsilon_n < +\infty$  P-a.s. Then the following hold:

- 1 The sequence  $(\|x_n\|)_{n \in \mathbb{N}}$  is bounded P-a.s.
- 2 Let  $z \in L^2(\Omega, \mathcal{F}, P; \mathbb{Z})$ . Then  $(\|x_n - z\|)_{n \in \mathbb{N}}$  converges P-a.s.

# Some results

Suppose that  $\sum_{n \in \mathbb{N}} \varepsilon_n < +\infty$  P-a.s. Then the following hold:

- 1 The sequence  $(\|x_n\|)_{n \in \mathbb{N}}$  is bounded P-a.s.
- 2 Let  $z \in L^2(\Omega, \mathcal{F}, P; \mathbb{Z})$ . Then  $(\|x_n - z\|)_{n \in \mathbb{N}}$  converges P-a.s.
- 3 Suppose that  $\mathfrak{B}(x_n)_{n \in \mathbb{N}} \subset \mathbb{Z}$  P-a.s. Then  $(x_n)_{n \in \mathbb{N}}$  converges weakly P-a.s. to a  $\mathbb{Z}$ -valued random variable.



# Some results

Suppose that  $\sum_{n \in \mathbb{N}} E\varepsilon_n < +\infty$ . Then the following hold:

# Some results

Suppose that  $\sum_{n \in \mathbb{N}} E \varepsilon_n < +\infty$ . Then the following hold:

- 1 The sequence  $(\|x_n\|_{L^2(\Omega, \mathcal{F}, P; H)})_{n \in \mathbb{N}}$  is bounded.

# Some results

Suppose that  $\sum_{n \in \mathbb{N}} E \varepsilon_n < +\infty$ . Then the following hold:

- 1 The sequence  $(\|x_n\|_{L^2(\Omega, \mathcal{F}, P; \mathbb{H})})_{n \in \mathbb{N}}$  is bounded.
- 2 Let  $z \in L^2(\Omega, \mathcal{F}, P; \mathbb{Z})$ . Then  $(E\|x_n - z\|)_{n \in \mathbb{N}}$  converges.

# Some results

Suppose that  $\sum_{n \in \mathbb{N}} E \varepsilon_n < +\infty$ . Then the following hold:

- 1 The sequence  $(\|x_n\|_{L^2(\Omega, \mathcal{F}, P; H)})_{n \in \mathbb{N}}$  is bounded.
- 2 Let  $z \in L^2(\Omega, \mathcal{F}, P; Z)$ . Then  $(E\|x_n - z\|)_{n \in \mathbb{N}}$  converges.
- 3 Let  $x$  be an  $H$ -valued random variable and suppose that  $(x_n)_{n \in \mathbb{N}}$  converges weakly P-a.s. to  $x$ .

# Some results

Suppose that  $\sum_{n \in \mathbb{N}} E \varepsilon_n < +\infty$ . Then the following hold:

- 1 The sequence  $(\|x_n\|_{L^2(\Omega, \mathcal{F}, P; H)})_{n \in \mathbb{N}}$  is bounded.
- 2 Let  $z \in L^2(\Omega, \mathcal{F}, P; Z)$ . Then  $(E\|x_n - z\|)_{n \in \mathbb{N}}$  converges.
- 3 Let  $x$  be an  $H$ -valued random variable and suppose that  $(x_n)_{n \in \mathbb{N}}$  converges weakly  $P$ -a.s. to  $x$ . Then  $(x_n)_{n \in \mathbb{N}}$  converges weakly in  $L^2(\Omega, \mathcal{F}, P; H)$  to  $x$ .

# Some results

Suppose that  $\sum_{n \in \mathbb{N}} E \varepsilon_n < +\infty$ . Then the following hold:

- 1 The sequence  $(\|x_n\|_{L^2(\Omega, \mathcal{F}, P; H)})_{n \in \mathbb{N}}$  is bounded.
- 2 Let  $z \in L^2(\Omega, \mathcal{F}, P; Z)$ . Then  $(E\|x_n - z\|)_{n \in \mathbb{N}}$  converges.
- 3 Let  $x$  be an  $H$ -valued random variable and suppose that  $(x_n)_{n \in \mathbb{N}}$  converges weakly  $P$ -a.s. to  $x$ . Then  $(x_n)_{n \in \mathbb{N}}$  converges weakly in  $L^2(\Omega, \mathcal{F}, P; H)$  to  $x$ .
- 4 Let  $x$  be an  $H$ -valued random variable. Then  $(x_n)_{n \in \mathbb{N}}$  converges strongly  $P$ -a.s. to  $x$  if and only if  $(x_n)_{n \in \mathbb{N}}$  converges strongly in  $L^1(\Omega, \mathcal{F}, P; H)$  to  $x$ .

# Geometrical scheme for stochastic Fejér monotonicity

Javier I. Madariaga

Department of Mathematics  
North Carolina State University  
Raleigh, NC 27695, USA

**NLA Student Seminar**  
November 14, 2024

**NC STATE** UNIVERSITY