

Pneumonia*	
Presentation	Fever, cough, dyspnea, pleuritic pain, respiratory distress
Etiology	<ul style="list-style-type: none"> • Neonatal: GBS, E. coli, K. pneumoniae, HSV • Infants: viral, S. pneumoniae, C. trachomatis • Pre-school age: viral, S. pneumoniae, S. pyogenes, S. aureus, B. pertussis • School-aged: M. pneumoniae, C. pneumoniae, S. pneumoniae, S. aureus
Differential	Asthma, pleural effusion/empyema, FB aspiration
Workup	CXR, respiratory viral panel including flu, blood culture if inpatient, ESR/CRP, procalcitonin
When to Hospitalize	Moderate-severe respiratory distress, SpO2 <90%, infants <6 mos, concern for virulent pathogen (MRSA), unable to tolerate PO intake
Treatment	<ul style="list-style-type: none"> • Outpatient: amoxicillin • Inpatient: ampicillin • Alternatives: add azithromycin if concern for atypicals, vancomycin if concern for s. aureus • Duration: 10 days, 2-4 weeks if parapneumonic effusion

Pleural Effusions							
Presentation	<ul style="list-style-type: none"> • Pain w/ inspiration, hypoxemia, hypercarbia • Exam: decreased breath sounds, dullness to percussion 						
Differential	<table border="1"> <tr> <td>Transudative</td><td>Decreased plasma oncotic pressure (nephrotic syndrome, cirrhosis, hypoalbuminemia)</td></tr> <tr> <td>Exudative</td><td>Increased capillary permeability (parapneumonic effusions, TB, AI disease, malignancy)</td></tr> <tr> <td>Chylothorax</td><td>Secondary to lymphatic abnormalities</td></tr> </table>	Transudative	Decreased plasma oncotic pressure (nephrotic syndrome, cirrhosis, hypoalbuminemia)	Exudative	Increased capillary permeability (parapneumonic effusions, TB, AI disease, malignancy)	Chylothorax	Secondary to lymphatic abnormalities
Transudative	Decreased plasma oncotic pressure (nephrotic syndrome, cirrhosis, hypoalbuminemia)						
Exudative	Increased capillary permeability (parapneumonic effusions, TB, AI disease, malignancy)						
Chylothorax	Secondary to lymphatic abnormalities						
Workup	<ul style="list-style-type: none"> • Imaging: CXR, US, CT • Diagnostic thoracentesis (consider if >10 mm fluid from lung to chest wall, need for definitive diagnosis, respiratory compromise) <ul style="list-style-type: none"> ■ Light's Criteria: Exudative if 1+ of (1) Pleural fluid protein:serum protein ratio ≥ 0.5, (2) Pleural fluid LDH:Serum LDH ratio >0.6, (3) Pleural fluid LDH $>66\%$ ULN of normal serum LDH 						
Treatment	<ul style="list-style-type: none"> • Transudative: address underlying problem • Chylothorax: Drainage, restrict to medium chain TGs as main source of dietary fat • Paraneumonic effusions (pleural fluid + pneumonia, abscess or bronchiectasis) <ul style="list-style-type: none"> ■ Uncomplicated: Antibiotics ■ Complicated: Antibiotics + drainage +/- fibrinolytics +/- VATS • Consider chest tube if: persistent fever, toxic appearing, large effusion, complicated pleural effusion or empyema 						

Obstructive Sleep Apnea	
Presentation	<ul style="list-style-type: none"> • Snoring (>3 nights/wk), labored breathing, morning headaches, daytime sleepiness, learning difficulties • Exam: tonsillary hypertrophy, adenoidal faces, micrognathia, HTN, overweight
Differential	Central sleep apnea, narcolepsy
Workup	Polysomnography to assess severity via apnea-hypopnea index (AHI) → >5 AHI warrants treatment
Treatment	CPAP , adenotonsillectomy if adenotonsillar hypertrophy, topical intranasal steroids or montelukast