

The Double Pendulum: Creating a Better Baseball Bat

Jared Baur

(Dated: 1 March 2019)

abstract goes here

I. INTRODUCTION

The double pendulum is a classic example of chaotic motion¹. The trajectories of various trials will show that the motion of a double pendulum is highly unpredictable. The chaos in a classic double pendulum is described by Equation 1. The exponent λ is a positive constant, Δx is the trajectory of the bottom arm of the pendulum, and t is time. For small times t , the trajectories are relatively the same, however for increasing times, the trajectories exponentially increase in distance from trial to trial.

$$\Delta x(t) \sim \Delta x(t_0)e^{\lambda t} \quad (1)$$

The double pendulum is used in real life situations such as in double-trailer trucks (officially known as “long combination vehicles”) where two large-wheeled vehicles are combined at a hinge point. These vehicles are essentially semi-trucks with two trailers attached instead of one. The swinging motion in sports is also a common instance of a double



FIG. 1. Chaos in a double pendulum. This figure depicts the trajectory of the end of the bottom arm in the double pendulum. The light is traced along the route of the bottom arm. As seen, there is no evident pattern in the bottom arm’s trajectory.

pendulum. For this paper, the double pendulum will be compared to a baseball swing. In a baseball swing, a bat (striking implement) is used in conjunction with the arms to strike a baseball. In general, the arms are held up and back from the player’s head, with the bat pointed upwards. The arms initiate the swing, moving towards the center of the player’s body; the bat follows the arms, and then swings through as the transfer of energy from the arms to the bat occurs. This motion occurs mostly in the horizontal plane, but nonetheless acts as a double pendulum with the player’s arms and the bat as the upper and lower “arms” of the pendulum, respectively.

Since the swing of a baseball bat only occurs once per trial (does not swing back and forth and time t is relatively low), this motion is equivalent to the first half cycle of a double pendulum swing. Thus, the chaotic motion prevalent in classic double pendulums will not be applicable for the purpose of the baseball swing.

Although the focus of this paper will be on the baseball swing, the double pendulum motion can be applied to the motion of a swing of a baseball bat, golf club, tennis racquet, or any other “swinging” motion that occurs with a striking implement. The goal of this paper will be to quantitatively describe the double pendulum in the context of a baseball swing and determine whether all of the energy in the striking implement can be transferred to the ball on impact. From this, we will decide whether we can design a baseball bat or ball that would allow for this perfect transfer of energy to occur.

II. EQUATIONS OF MOTION

Describing the motion of a double pendulum has been argued as highly complicated². For this reason, Lagrangians have been used in past studies to represent this motion. However, when we look at the forces on each arm of the pendulum with respect to each arm’s center of mass, Newtonian equations of motion can be used to describe the motion.

Figure 2 depicts the components in a simple double pendulum. The x, y axes for position and forces are shown in the figure. The upper arm of the pendulum is the first arm attached to a fixed axis A . This arm has length L_1 , center of mass at point G_1 , and distance to its center of mass h_1 (measured from point A). This arm makes an angle θ with the vertical axis and has a gravitational force acting on its center of mass equal to M_1g , where M_1 is the mass of the arm and g is the acceleration due to gravity.

The second arm (lower arm) rotates on a “fixed” axis at point B . If we isolate the second arm from the system, it would rotate about point B as if it were fixed, however point B is rotating around point A since it is at the end of the first arm.




FIG. 2. The components in a double pendulum. Since the equations that describe a double pendulum can be rather confusing, we use the center of mass G_1 and G_2 to describe the forces acting on the pendula arms.

The second arm has a length L_2 and its center of mass at point G_2 , a distance of h_2 away from point B . The arm makes an angle of ϕ with the vertical axis and has a gravitational force of M_2g acting on its mass M_2 . Point C is the end of the second arm, and will act as a point of interest for our paper as we track the trajectory of this point. The velocity vector V is the direction of motion for the second arm's center of mass through the swinging of a double pendulum.

We start writing our equations of motion by describing the coordinates of our center of mass on the second arm. We want the coordinates of the center of mass since we will write our forces with respect to that (as opposed to point C). Point G_2 is described in Equation 2, where the first half of the equation is the x component and the second half is the y component. The y component is negative since the double pendulum's origin at $(x, y) = (0, 0)$ is assumed at point A .

$$G_2(x, y) = \left([L_1 \sin \theta + h_2 \sin \phi], [-L_1 \cos \theta - h_2 \cos \phi] \right) \quad (2)$$

to-do: describe velocities

$$V_x = \frac{dx}{dt} = -L_1 \omega_1 \cos \theta - h_2 \omega_2 \cos \phi \quad (3)$$

$$V_y = \frac{dy}{dt} = -L_1 \omega_1 \sin \theta - h_2 \omega_2 \sin \phi \quad (4)$$

to-do: describe how to get force

$$\begin{aligned} F_x &= M_2 \frac{dV_x}{dt} \\ &= -M_2 \left[L_1 \cos \theta \frac{d\omega_1}{dt} + L_1 \omega_1^2 \sin \theta \right. \\ &\quad \left. + h_2 \cos \phi \frac{d\omega_2}{dt} + h_2 \omega_2^2 \sin \phi \right] \end{aligned} \quad (5)$$

$$\begin{aligned} F_y - M_2 g &= M_2 \frac{dV_y}{dt} \\ &= -M_2 \left[L_1 \sin \theta \frac{d\omega_1}{dt} - L_1 \omega_1^2 \cos \theta \right. \\ &\quad \left. + h_2 \sin \phi \frac{d\omega_2}{dt} - h_2 \omega_2^2 \cos \phi \right] \end{aligned} \quad (6)$$

III. EXPERIMENTAL RESULTS

IV. TORQUE

V. ENERGY TRANSFER

VI. CONCLUSION

¹Shinbrot, "Chaos in a double pendulum," American Journal of Physics **60**, 491–498 (1992).

²Jorgensen, "On the dynamics of the swing of a golf club," American Journal of Physics, 644–651 (1970).

³Cross, "A double pendulum swing experiment," American Journal of Physics **73**, 330–339 (2005).