

## Sheet 11

Solutions to be handed in before class on Wednesday June 26

**Problem 51.** Consider  $\mathfrak{g} = \mathfrak{sl}_2(\mathbb{C})$  with the usual diagonal Cartan subalgebra  $\mathfrak{h}$  and associated root system  $R = \{\pm\alpha\}$ . Take  $R^+ = \{\alpha\}$  as a system of positive roots, and  $\rho = \alpha/2$ . Let  $z \in \mathbb{C}$ .

1. Describe all  $U(\mathfrak{g})$ -submodules of the Verma module  $M(z\rho)$ . (2 points)
2. Show that the simple quotient  $L(z\rho)$  of  $M(z\rho)$  has finite dimension (equal to  $z + 1$ ) if and only if  $z \in \mathbb{N}_0 = \{0, 1, 2, \dots\}$ . (2 points)

**Problem 52.** Consider the natural representation  $V = \mathbb{C}^n$  of  $\mathfrak{g} = \mathfrak{sl}_n(\mathbb{C})$ .

1. Show that the exterior power  $\bigwedge^d V$  is an irreducible representation of  $\mathfrak{g}$ , for all  $d = 0, \dots, \dim_{\mathbb{C}} V$ . (2 points)
2. Determine the highest weight of  $\bigwedge^d V$  with respect to the standard diagonal Cartan subalgebra  $\mathfrak{h}$  and choose a basis of  $R$  such that  $\mathfrak{b} = \mathfrak{h} \oplus \bigoplus_{\alpha \in R^+} \mathfrak{g}_{\alpha}$  is given by the upper triangular matrices in  $\mathfrak{g}$ . (2 points)

**Problem 53.** Let  $\mathfrak{b} \subset \mathfrak{g}$  be an inclusion of Lie algebras over a field  $k$ . Let  $M$  be a representation of  $\mathfrak{b}$  and  $E$  a representation of  $\mathfrak{g}$ . Show that there is a canonical isomorphism of  $\mathfrak{g}$ -representations

$$U(\mathfrak{g}) \otimes_{U(\mathfrak{b})} (E \otimes_k M) \xrightarrow{\sim} E \otimes_k (U(\mathfrak{g}) \otimes_{U(\mathfrak{b})} M). \quad (40)$$

(4 points)

**Hint** Here  $E$  on the left is considered as a representation of  $\mathfrak{b}$  by restriction. Either define an isomorphism explicitly or deduce it from the fact that the left adjoint of a composition is the composition of the left adjoints, provided they exist.

**Problem 54.** Let  $\mathfrak{g}$  be a complex semisimple Lie algebra with a Cartan subalgebra  $\mathfrak{h}$  and  $R^+ \subseteq R = R(\mathfrak{g}, \mathfrak{h})$  a system of positive roots. We say that a  $U(\mathfrak{g})$ -module  $M$  has a Verma flag if there is a sequence

$$0 = M_0 \subseteq M_1 \subseteq \dots \subseteq M_n = M \quad (41)$$

of submodules such that each quotient  $M_i/M_{i-1}$  is isomorphic to a Verma module. It can be shown that the tensor product  $E \otimes_{\mathbb{C}} \Delta(\lambda)$  of a Verma module  $\Delta(\lambda)$  with a finite-dimensional  $U(\mathfrak{g})$ -module  $E$  has a Verma flag.

Specialising  $\mathfrak{g}$  to  $\mathfrak{sl}_2(\mathbb{C})$  with the notation of problem 51, find Verma flags for  $V \otimes_{\mathbb{C}} \Delta(0)$  and  $V \otimes_{\mathbb{C}} \Delta(-\rho)$ , where  $\rho = \alpha/2$ . Are these tensor products direct sums of Verma modules? (2 points)

**Problem 55.** Consider  $\mathfrak{g} = \mathfrak{sl}_2(\mathbb{C})$  with the Cartan subalgebra and positive roots as in problem 51. Let  $\mathcal{C}$  be the smallest subcategory of the category of all  $U(\mathfrak{g})$ -modules which contains all Verma modules and is closed under tensor products with finite-dimensional modules. A short exact sequence in  $\mathcal{C}$  is by definition a short exact sequence of  $U(\mathfrak{g})$ -modules whose terms lie in  $\mathcal{C}$ .

Show that there is a non-split short exact sequence

$$0 \rightarrow M(0) \rightarrow P \rightarrow M(s_\alpha \cdot 0) \rightarrow 0 \quad (42)$$

in  $\mathcal{C}$ . (2 points)