

Ablation Study 1 -- Layer Weighting Formulation & Sensitivity

Method: Procrustes-DS (feedback-refined, `final.py`). **Ablated variable:** the L=12 layer-weight vector `w` that drives the feature blend. Every other component of the pipeline (scale-matched relative-Mahalanobis, mid-layer off-manifold residual, logit-energy evidence, asymmetric additive log-evidence fusion) is held **identical** to the main experiment, so any difference is attributable solely to the layer weighting.

Setup

- **ID dataset:** CIFAR-100 (balanced)
- **Backbone:** IN21K-ViT-B/16 + AdaptFormer PEFT (the exact X-Maha checkpoint reused by `final.py`)
- **CIFAR-100 top-1 accuracy:** 92.95%
- **OOD datasets (6/6):** Texture, SVHN, CIFAR-10, Tiny ImageNet, LSUN, Places365 (SCOOD, `id_name=cifar100`)

Variants compared

#	Scheme	Formula
1	Uniform	<code>w_l = 1/L</code>
2	Trace-only (X-Maha)	<code>w_l = trace-variance_l</code>
3	HSKP softmax	<code>w_l = softmax(HSKP_l / 0.5)</code>
4	Pure Fisher ratio	<code>w_l = (S_b/S_w)_l^2</code>
5	Fisher-discriminability (proposed)	<code>w_l = trace_l * (S_b/S_w)_l^gamma</code> , gamma in {0.5, 1.0, 2.0, 3.0}