

Accurate binding of calcium to phospholipid bilayers by effective inclusion of electronic polarization

Josef Melcr and Hector Martinez-Seara Monne

Institute of Organic Chemistry and Biochemistry, Academy of Sciences of the Czech Republic, Prague 6, Czech Republic

Pavel Jungwirth

*Institute of Organic Chemistry and Biochemistry, Academy of Sciences of the Czech Republic, Prague 6, Czech Republic and
Department of Physics, Tampere University of Technology, P.O. Box 692, FI-33101 Tampere, Finland*

O. H. Samuli Ollila*

*Institute of Organic Chemistry and Biochemistry, Academy of Sciences of the Czech Republic, Prague 6, Czech Republic and
Institute of Biotechnology, University of Helsinki*

(Dated: October 10, 2017)

1. Abstract directly from Joe's conference abstracts. To be rewritten. Classical molecular dynamics simulations give detailed information about membrane structure and dynamics. However, there is still a room for improvements in current force fields it is known from the literature, that the binding of ions, especially cations, to phospholipid membranes is overestimated in all classical models [1]. We suggest that the membrane-ion interactions can be corrected by including implicit electronic polarizability into the lipid models through the electronic continuum correction (ECC) [2], which was already applied to monovalent and divalent ions yielding models that feature correct ion pairing [3]. Using the electrometer concept [3, 4] and x-ray scattering form factors, our simulations point out that our hypothesis is correct and ECC is indeed a missing important contribution in current classical lipid models. Moreover, the solid physical principles behind ECC are found not to hamper other relevant properties of a phospholipid bilayer. The new lipid model, "ECC-lipids", shows accurate binding affinity to sodium and calcium cations and head group order parameter response to bound charge. We also provide for the first time a realistic stoichiometry of bound calcium cations to a POPC membrane, and their binding sites. This work will continue as an open collaboration project NMRLipids VI (<http://nmrlipids.blogspot.fi>).

I. INTRODUCTION

Cation interactions with cellular membranes play a key role in several biological processes, such as in signal propagation in neurons and vesicle fusion. Since direct measurements of ion-membrane interactions from biological systems are difficult, lipid bilayers are often used as model systems for cellular membranes. The detailed results from simple model systems can be then used to understand the role of ions in complex biological systems.

Zwitterionic phosphocholine (PC) lipid bilayers are commonly used model systems for cellular membranes. Interactions of biological cations, especially Na^+ and Ca^{2+} , with PC bilayers are widely studied in experiments [1–8] and classical MD simulations [9–13]. The details of ion binding are, however, not fully consistent in the literature. Non-invasive spectroscopic methods, like nuclear magnetic resonance (NMR), scattering and infrared spectroscopy, give accurate information about ion binding in PC lipid bilayers [1, 2, 6–8, 14–16]. Interpretation most of these experiments suggests that Na^+ ions exhibit negligible binding in PC lipid bilayers with submolar concentrations, while Ca^{2+} specifically binds to phosphate groups of two lipid molecules. Atomistic resolution molecular dynamics (MD) simulation models, however, predict significantly stronger binding for the cations than NMR experiments [17]. On the other hand, some experiments have also been interpreted to support the predictions from MD sim-

ulations [9, 18]. Interactions of Calcium ions with 3–4 lipids, including also interactions with carbonyl oxygens, have been reported from simulations [9, 10, 12, 13].

Recent work published by the NMRLipids project (nmrlipids.blogspot.fi) [17] made an attempt to resolve the apparent controversies. This study presented a direct comparison between ion binding affinity between simulations and experiments by using NMR data for lipid headgroup order parameters and the electrometer concept [19]. Using massive amounts of data collected by Open Collaboration method, it was concluded that the accuracy of the current state of the art lipid models for MD simulations is not sufficient for the detailed interpretation of the cation interactions with PC lipid bilayers [17].

In this work we show that the cation binding behavior in MD simulations of 1-Palmitoyl-2-oleoylphosphatidylcholine (POPC) bilayer can be significantly improved by implicitly including electronic polarizability in the polar region of lipid molecules. The electronic polarizability is included by using the electronic continuum correction (ECC) [20], which has been previously shown to improve the behaviour of MD simulations of ions in bulk water [21–23]. As a starting point we use the parameters from the Lipid14 model [24], which gave the best cation binding behaviour in the previous study [17]. The developed ECC-lipid parameters reproduce the experimentally measurable structural parameters of an ion-free POPC lipid bilayer with the accuracy comparable to the other state of the art lipid models, while surpassing them significantly for reproducing the membrane binding affinities and induced structural effects of Sodium and Calcium ions.

*samuli.ollila@helsinki.fi

II. METHODS

A. Electronic continuum correction for lipid bilayers

The lack of electronic polarizability in the standard MD simulation force fields has been considered a highly relevant issue since the early days of lipid bilayer simulations. In this work we circumvent the rather demanding explicit inclusion of electronic polarization effects [25] by implicitly including electronic polarizability in lipid bilayer simulations by using the electronic continuum correction (ECC) [20]. Technically, it is a similar approach to the phenomenological charge-scaling as applied in the early studies where a scaling factor one half was used [26, 27]. **2.We should also cite papers where empirical scaling was used ionic liquids - but there the factor is not 0.5.** However, the present concept of ECC is physically well justified and rigorously derived [20, 28, 29].

According to ECC, electronic polarizability can be implicitly included in classical MD simulations by placing all particles into a homogeneous dielectric continuum with a dielectric constant ϵ_{el} , which is the electronic part of the dielectric constant of the media [20]. Measurements of high frequency dielectric constant gives values of approximately $\epsilon_{el} \approx 2$ for almost any biomaterial [20?]. Such a dielectric continuum can be easily included in standard MD simulation by a formal transformation of partial charges

$$Q^{ECC} = f_q \cdot Q \quad (1)$$

with a constant scaling factor $f_q = \epsilon_{el}^{-1/2}$ effectively representing the newly introduced electronic continuum. Assuming globally a high frequency dielectric constant as measured in water (corresponding to the square of the refraction index), $\epsilon_{el} = 1.78$, results in a scaling factor of $f_q = 0.75$ [20?]. This scaling factor has been successfully used to improve the performance of force field for ions in solution [22? , 23] which then agree quantitatively with neutron scattering data [21–23].

While the scaling factor of $f_q = 0.75$ for ions in water improves their performance and is physically justifiable within the ECC theory [?], it is not clear whether the same factor should be used for partial charges in molecules, e.g., lipids in our case. Unlike the total charge of an atom or molecules, atomic partial charges within each molecule are not physical observables. There are several schemes for the assignment of partial charges for biomolecules. [30] Currently, the most commonly employed scheme is the restrained electrostatic potential (RESP) method [31, 32]. In practice, partial charges currently implemented in force fields may already include to some extent some of the solvent electronic polarizability effects, i.e., the RESP charges are often scaled to fit some experimental observables **3.This needs a citation.** Thus, we expect that the application of the ECC scaling factor, f_q , to the molecular partial charges included in the available force fields does not necessarily have to follow the relation $f_q = \epsilon_{el}^{-1/2}$, but instead it lies between 0.75 (no electronic polarizability in the partial charge calculation) and 1 (full electronic polarizability already included in the partial charge calculation).

In this work, we develop a phospholipid model for classical MD simulations that accurately describes the lipid head group response to varying concentrations of monovalent and divalent cations. This is a biologically highly relevant membrane feature, which is poorly reproduced by currently available models which can affect not only on membrane properties in the presence of ions but also modulates the interaction with charged moieties in the surface. Importantly, this response from simulations can be accurately compared against experimental NMR data [1, 2, 33], as discussed in Ref. 17 in section II B. To this end, we empirically explore the scaling factor parameter space, $f_q \in [0.75, 1.0]$ for the Lipid14 [24] force field. We selected this force field as a starting point because its response to bound ions was apparently the most realistic against NMR data in recent work by NMRlipids project (see Fig. 5 in Ref. 17). Also glycerol backbone and head group structures in Lipid14 model were relatively realistic when compared with other state of the art lipid models [34]. The ECC correction was applied to Lipid14 parameters by scaling partial charges of the head group, glycerol backbone and carbonyl regions, which are the most polar parts in lipids and are expected to have the largest contribution to the cation binding. However, we do not modify the hydrocarbon chain parameters, because they do not come in contact with salt ions and are already highly optimized and give generally a good description for hydrophobic part of lipid bilayers in most lipids models, including Lipid14 [35]. This is in contrasts with the behavior in glycerol backbone and head group order parameters which call for improvements in all available lipid models [34].

Exploring different scaling factor values, we found out that ion binding and related head group order parameter responses become weakened in general. The optimal behaviour of ion binding was observed with the scaling factor of $f_q = 0.8$.

4.JOE: following discussion shall be modified in the enlightenment of our recent ECC-discussions. Interestingly, this scaling factor is in line with the estimate given by “implicitly polarized charges” (IPolQ) [36] combined with RESP calculations in vacuum and implicit solvent reported in [37]. IPolQ charges are obtained as the average of partial charges given by RESP calculation [31] in vacuum and in a solvent. Applying the scaling factor of 0.75 to IPolQ charges calculated from the data in Ref. [37], gives similar partial charges to ones obtained by scaling Lipid14 charges with a factor 0.8.

While, the charge scaling improved the behaviour of lipid-ion interactions, it reduced the area per molecule of the lipid bilayer without ions below experimental values. Simulations with Lipid14 parameters having partial charges of head group, glycerol backbone and carbonyls scaled with 0.8 gave the area per molecule value of $\approx 60 \text{ \AA}^2$, which is smaller than the experimental value 64.3 \AA^2 (**5.missing REF for APL experiment**) and the original Lipid14 value $(65.6 \pm 0.5) \text{ \AA}^2$ [24]. The decrease of the area per lipid was found to arise from a lower hydration of the lipid head group region, which can be explained by the increased solvation free energy due to the lower polarity of molecules with scaled charges. The hydration can be increased back by reducing the effective radius of atoms by changing the σ parameters in the Lennard-Jones potential for the selected atoms similarly as done for free ions

in solution [21–23]. 6. We should discuss how this can potentially affect the intermolecular interaction when mixing scaled and non scaled molecules. JOE: I think that we rather increasingly see that there's nothing like "fully non-scaled" with the exception of ions with integer charges. So the discussion shall be rather more about the interaction of our "scaled" (I'd still rather call it ECC-corrected or whatever) and "semi-scaled" models. This decreases the solvation free energy by allowing water molecules to approach closer to lipid atoms and have stronger electrostatic interactions with them. After reducing the σ parameters by a factor of $f_\sigma = 0.89$ for the same atoms for which charges were scaled, the area per molecule value was back in agreement with the experimental value (see Table I).

B. Electrometer concept

Ion binding between experiments and simulations was compared by using lipid head group order parameters and the "electrometer concept" [17, 19]. The concept is based on the experimental observation that the C-H bond order parameters of α and β carbons in PC lipid head group (see Fig. 1) are proportional to the amount of unit charge bound per lipid, X^\pm [19]. Change in the order parameters measured with varying aqueous ion concentration can be then related to the amount of bound ions.

The change of the head group order parameters is empirically quantified as [19, 39]

$$\Delta S_{\text{CH}}^i = S_{\text{CH}}^i(X^\pm) - S_{\text{CH}}^i(0) \approx \frac{4m_i}{3\chi} X^\pm, \quad (2)$$

where $S_{\text{CH}}^i(0)$ denote the order parameter in the absence of bound charge, i refers to either α or β carbon, m_i is an empirical constant depending on the valency and position of the bound charge, and the experimental value [40, 41], $\chi \approx 167 \text{ kHz}$, is used for the quadrupole coupling constant. Atomic absorption spectra and ^2H NMR data gave $m_\alpha = -20.5 \text{ kHz}$ and $m_\beta = -10.0 \text{ kHz}$ for Ca^{2+} binding to POPC bilayer (in the presence of 100 mM NaCl) [2, 17, 35]. The slopes are negative, because recent analysis concluded that the order parameters decrease with bound positive charge and increase with bound negative charge when the signs are taken in account [17, 35]. This is rationalized as a change of lipid head group dipole tilt toward water phase with bound positive charge and *vice versa* with negative charge [19].

The concept can be used to compare the ion binding affinity in lipid bilayers between MD simulations and NMR experiments, because the order parameters can be accurately determined from both techniques [35]. The order parameters for all C-H bonds in lipid molecules, including α and β segments in head group, can be accurately measured using ^2H NMR or ^{13}C NMR techniques. From MD simulations the order parameters can be calculated using the definition

$$S_{\text{CH}} = \frac{3}{2} \langle \cos^2 \theta - 1 \rangle, \quad (3)$$

where θ is the angle between the bond and membrane normal and the average is taken over all sampled configurations [35].

The measured order parameter change depends on the response of the headgroup on the bound charge, i.e. m_i in Eq. 2, and the ion binding affinity. Thus, the former property has to be well quantified before using the electrometer concept to analyze binding affinities. This is done experimentally for a wide range of systems [19, 42]. To calibrate the head group order parameter response also in simulations, we use experimental data for dihexadecyldimethylammonium ($2\text{C}_{16}^+\text{N}2\text{C}_1\text{Br}^-$) in a POPC bilayer [33]. Dihexadecyldimethylammonium is a cation surfactant having two acyl chains and bearing a unit charge in the hydrophilic end. Thus, it is expected to locate in the bilayer similarly to the phospholipids and the molar ratio then gives directly the amount of bound unit charge per lipid X^\pm in these systems [43].

C. Salt concentrations and binding affinity

Experimental studies providing the lipid head group order parameters report salt concentrations in two different ways. For DPPC, salt concentrations before solvating the lipids are used [1]. Alternatively, for POPC, the salt concentration of supernatant after solvation is measured using atomic absorption [2]. 7. JOE: I'm afraid that the above description might sound to the reader like for some reasons one method is used with DPPC, but a different one with POPC. I think simply writing one way is...another way is ... (or similarly) will be less confusing. In this work, we will use the latter definition. For this we will calculate the salt concentration in the aqueous bulk region, i.e., the farthest point from both lipid leaflets in the water phase. Note that in the previous study, Ref. 17, the concentration was calculated as in the DPPC experiments. Despite of the measurable differences between these two concentrations for CaCl_2 systems, the qualitative conclusions are affected neither in this nor in our previous work [17].

To quantify the ion binding affinity to a membrane, we calculated the relative surface excess of ions with respect to water Γ_i^w [44]. This quantity can be used to compare binding affinity between different models, because it does not depend on the definition of the interfacial region or the position of the Gibbs dividing plane between two bulk regions. 8. JOE: following is very unclear to me. I think we can write it in a more concise way. Therefore, we consider the hydrophobic interior of a membrane having zero concentrations of ions and water as a bulk region, which is separated by a monolayer interface from a another region with bulk concentration of ions in water. Since the boundary of the latter region is set to the edge of simulation box, all molecules in simulation are within the defined interfacial region. This setup provides a simplified relation for Γ_i^w in lipid bilayers simulations

$$\Gamma_i^w = \frac{1}{2A_b} \left(n_i - n_w \frac{C_i}{C_w} \right), \quad (4)$$

where n_w and n_i are the total number of waters and ions in the system; C_w and C_i are their respective bulk concentrations in the aqueous phase; and A_b is the size of the box in the membrane plane. The total area of the interface is twice the

area of the membrane, $2A_b$, because bilayers have an interface at both leaflets.

D. Validation of lipid bilayer structure against experiments

Lipid bilayer structure without ions was validated against NMR and x-ray scattering experiments by calculating order parameters for C-H bonds and form factors from our simulations. The former validates the structures sampled by individual lipid molecules in simulations with atomic resolution, while the latter validates the dimensions of the lipid bilayer (thickness and area per molecule) [35].

The order parameters were calculated from simulations for all C-H bonds in lipid molecules by using Eq. 3. Form factors were calculated from equation

$$F(q) = \int_{-D/2}^{D/2} \left(\sum_{\alpha} f_{\alpha}(q_z) n_{\alpha}(z) - \rho_s \right) \exp(izq_z) dz, \quad (5)$$

where $f_{\alpha}(q_z)$ is the density of atomic scattering length, ρ_s is the density of solvent scattering length in the bulk region, $n_{\alpha}(z)$ is the number density of atom α and z is the distance from the membrane centre along its normal spanning until the water bulk region, D .

E. Simulation details

1. Simulations of POPC bilayers in aqueous ions


The simulated systems consisted of POPC bilayer in pure water or in varying salt concentrations. In particular, the periodic orthorhombic simulation box contained 128 POPC molecules and approximately 50 water molecules per each lipid. As a default, water molecules were described by the OPC3 force field [45] which is currently the most accurate three site rigid water model. In order to test transferability of our newly developed ECC-lipids model, we also performed several additional simulations with OPC [46], SPC/E [47], TIP3p-FB and TIP4p-FB [48], and TIP4p/2005 [49] water models presented in Supporting Information (SI). We used the ECC-ions model for Sodium, Calcium and Chloride ions [21, 23]. Simulations with Lipid14 use ion models by Dang [50–52], and by Åqvist [53]. MD simulations were performed using the GROMACS [54] simulation package (version 5.1.4). The simulation settings used in this work are summarized in Table III. Simulation trajectories and parameters are available at [?] [9.To be uploaded to Zenodo.](#)

2. Simulations of POPC bilayers with cationic surfactants

An automated topology builder [55] was first used to create the structure of dihexadecyldimethylammonium

TABLE I: Area per lipid (APL) from different models of POPC with no ions

model	APL (Å ²)	Temperature [K]
Lipid14	65.1 ± 0.6	300
Lipid14 [24]	65.6 ± 0.5	303
ECC-lipids	62.2 ± 0.6	300
experiment [59] 10.REF	64.3	303



which is one of the cationic surfactants used to experimentally quantify the electrometer concept [33]. The AmberTools program [56] was then used to generate the Amber-type force field parameters. The parameters were converted to the Gromacs format by using the acpype tool [57]. The partial charges were then manually modified to approximately correspond to their equivalent segments in Lipid14 [24]. The surfactants were randomly placed among the lipids to form bilayer structures with mole fractions of 10%, 20%, 30%, 42%, or 50% of surfactant in the POPC bilayer. All systems contained 50 POPC molecules per leaflet, 6340 TIP3P water molecules and 6, 14, 21, 35, or 50 surfactants per leaflet. The systems were simulated for 200 ns using Lipid14 model for POPC where reasonable lipid neighbor exchange occurs. First 20 ns were omitted from the analysis.

The same systems were also simulated with the ECC-lipid model for POPC using the same setup. In these simulations the ECC correction was also applied to the cationic surfactant by scaling all charges with the same factor as for ECC-lipids, i.e., $f_q = 0.8$, and by using the atom types with reduced σ parameters from ECC-lipids.

III. RESULTS AND DISCUSSION

A. POPC membrane structure and dynamics

The x-ray scattering form factor (Fig. 1) and the area per lipid (Table I) of POPC bilayer simulated with the ECC-lipid model are in good agreement with the experimental results, as well as with the Lipid14 model. Thus, we conclude that the lipid bilayer dimensions are well captured by the developed ECC-lipid model. Areas per lipid of ECC-lipids with various other water models are documented in SI, table II.

As in the original Lipid14 model [24], the acyl chain order parameters of the ECC-lipid model agree well with the experimental values in Fig. 1. Notably, the experimentally measured forking and small order parameter values of C_2 segment in sn -2 chain are relatively well reproduced by the both models. This has been suggested to indicate that the carbonyl region of sn -2 chain is directed towards the water phase, in contrast to the carbonyl in sn -1 chain, which would orient more along the bilayer plane [60–62]. While this may be an important feature




FIG. 1: Top: Chemical structure of POPC with definitions of different order parameters calculated. Middle: Order parameters of head group, glycerol backbone and sn-1 and sn-2 tails from simulations with Lipid14 [24] and ECC-lipids models compared with experimental order parameters from [58]. Error bars are approximately the size of the points for head group order parameters. Point size for the order parameters of tails are decreased by a factor of 3 to improve the clarity of the plot. Bottom: X-ray scattering form factors from experiments [?] and simulations using Lipid14 [24] and ECC-lipids models.

for the ion binding details, it is not necessarily reproduced by the available lipid models [35]

The headgroup order parameters of α and β carbons are slightly larger in the ECC-lipid model than in Lipid14, which is apparently related to the smaller P-N vector angle with respect to the membrane normal in Fig. 2. With the current data we cannot, however, conclude which one of the models give the more realistic headgroup conformations. The ECC-lipid model gives the β carbon order parameter value closer to experiments, while value for α carbon is better in Lipid14. Despite of some deviations from the experimental order parameter values in Fig. 1, the accuracy of the both models in the glycerol backbone region is comparable to the other state of art lipid models available in literature [34].

11.Dynamics check is missing: MSD (Hector/Joe)




FIG. 2: Headgroup order parameter changes and P-N vector orientation as a function of cationic surfactant (dihexadecyldimethylammonium bromide, $2C_{16}^{+}N2C_1Br^{-}$) in PC bilayer from simulations and experiments [33].

B. Response of POPC head groups to bound charge

Before proceeding to the ion binding affinity, we quantify the response of headgroup order parameters to the amount of bound charge by using mixtures of monovalent cationic surfactants and POPC [33]. The amount of bound charge per PC in these systems is given by the molar fraction of cationic surfactants, because essentially all surfactants locate in the lipid bilayers. Experimental data for these systems can be used to validate the sensitivity of lipid headgroup order parameters to the amount of bound charge in simulations.

The headgroup order parameter changes with an increasing amount of the cationic surfactant dihexadecyldimethylammonium bromide is compared between experiments [33] and simulations in Fig. 2. The observed order parameter decrease in simulations and experiments can be approximated to be linear at least for mole fractions below $\sim 30\%$, as expected from Eq. 2. The slope is, however, too steep in the Lipid14 model indicating that the head group order parameters are too sensitive to a bound charge. The ECC-lipids model gives a slope in very good agreement with experiments for the α segment, while the slope is slightly underestimated for the β segment.

12.SAMULI: We could calculate the slopes from simulations, but I am not sure if we would actually learn anything useful from this.

The headgroup P-N vector angle with respect to the membrane normal is also shown as a function of cationic surfactant mole fraction in Fig. 2. The headgroup orients more towards the water phase with an increasing amount of bound cations, as previously reported in Ref. 19. The effect is more pronounced in Lipid14 than in ECC-lipids model, which is in line with the order parameter results and the reduced charge-dipole interactions in the latter model. The response of α -order parameter to bound positive charge in ECC-lipid model is in good agreement with experiments. The model can be thus used to study changes of lipid P-N vector in varying conditions.

C. Cation binding affinities to POPC

The binding affinities of aqueous cations to lipid bilayers can be measured using the headgroup order parameters, because they decrease proportionally to the bound positive charge [17, 19]. The headgroup order parameter responses to increasing aqueous CaCl_2 concentrations from experiments (DPPC [1] and POPC [2]) and different simulation models for POPC are shown in Fig. 3. Responses to NaCl concentrations are in SI, Fig. S9.

Negligible changes of the headgroup order parameters are measured with submolar concentrations of NaCl due to the very low affinity of Na^+ in PC bilayers [1]. While Na^+ binding and the related headgroup order parameter changes were overestimated in almost all the available simulation models, the low affinity and negligible order parameter changes were reproduced by Lipid14 model when simulated with Åqvist ions [17]. However, the same combination of force field parameters overestimated the headgroup order parameter response to CaCl_2 concentration, which was the case also in all other models tested in Ref. 17. Using the ion model by Dang et al. [50–52] or ECC-ions [21, 23?] with more realistic bulk behaviour did not improve the results for the CaCl_2 interactions with a POPC bilayer described by the Lipid14 model, as seen in Figs. 3 and S8 (in SI), respectively. 13.Add OP-response of Lipid14+ECC-ions plot in SI. The results support the conclusion of the previous work [17] that improvements also in lipid models are needed to correctly describe divalent cation binding to PC bilayers.

Significant improvement can be achieved using the ECC approach also for lipids. The headgroup order parameters change as a function of CaCl_2 concentration for the ECC-lipid model with ECC-ions exhibit a good agreement with experiments in Fig. 3 and S9. As discussed in previous section, the model gives also a good agreement with experiments for the headgroup response to bound charge.

The binding affinities are quantified using the water density profiles along the membrane normal shown in Fig. 4. The density profiles show a larger Ca^{2+} density peak in lipid headgroup region for the Lipid14 model with Dang and ECC-ions than for the ECC-lipid model. The relative surface excess calculated from Eq. 4 is $\Gamma_i^w = 0.07 \pm 0.01 \text{ nm}^{-2}$




FIG. 3: Changes of head group order parameters of POPC bilayer as a function of CaCl_2 concentrations are shown from simulations with different force fields together with experimental data (DPPC [1] and POPC [2]). Ion concentrations in bulk water are shown in x-axis. Values from simulations are calculated from the of cation number density C_{np} from the region at the simulation box edge with the constant ion concentration as $[\text{ion}] = C_{np}/0.602$. Simulation data with Lipid14 and Åqvist ion parameters is taken directly from Ref. [17].

for the ECC-lipid model, which is significantly smaller than $\Gamma_i^w = 0.13 \pm 0.01 \text{ nm}^{-2}$ for Lipid14 with Åqvist and $\Gamma_i^w = 0.3 \pm 0.03 \text{ nm}^{-2}$ with Dang ions.

14.Below analysis is done in a stupid way to get some idea. I would fine it useful to do this analysis by using the density profiles, but it is not necessary. The reason why I want to do something like this, is that it would be useful to have some easier way to estimate the correct binding affinity than the electrometer concept, which is

quite tedious to apply in practise (simulations with different concentrations, cationic surfactant check, etc.). I would like to be able to estimate much faster from a simulation if the binding affinity is reasonable or not. This may not be the correct way for that anyway. Rough estimates for the free energy difference between bound and unbound cation are given by

$$\Delta G = -k_b T \log\left(\frac{p_i}{p_o}\right), \quad (6)$$

where p_o and p_i are estimated from the Ca^{2+} densities in bulk water and in the maximum density in bilayer, respectively. The density profiles in Fig. 4 give $\sim 0.8 k_b T$ for the free energy difference between bound and unbound Ca^{2+} ions in ECC-lipid model and $\sim 1.4 k_b T$ in Lipid14 with Åqvist.

15.SAMULI: Maybe we should discuss the repeat distances and area per molecules measured at [7, 8, 63]

Since the lipid headgroup order parameter responds to the amount of bound charge and to the aqueous ion concentrations are both in good agreement with experiments in the ECC-lipid model with ECC-ion parameters, we consider the Na^+ and Ca^{2+} binding affinities to be realistically described in this model. In contrast, the Ca^{2+} binding affinity is overestimated by Lipid14 model when simulated with all the tested ion models. Similar conclusions were previously made based only on the headgroup order parameter data with aqueous cations [17]. However, the discrepancies with experiments in previous work could partly arise also from the inaccurately described sensitivity of the headgroup to bound charge. Here we quantify this effect and conclude that the improvement due to ECC is to a large extent, but not entirely, caused by a more realistic headgroup sensitivity to bound charge. This indicates that the issue should be carefully considered also when the electrometer concept is used to compare ion binding between experiments and simulations with other models.

D. Binding stoichiometry

This section is rough and will likely require editing. Data – up to noted exceptions – shall be all there, however. **18.SAMULI:** I think the we should make clear that the current binding constants in, e.g., Marsh's handbook are based on the headgroup order parameter changes interpreted with ternary complex binding model. I.e. the experimental raw data is exactly the same as we have in this work. Our simulations enable much more versatile interpretation of the binding phenomena. I think that this is one the main points of this work, so this comment applies to the previous section, introduction, abstract and conclusions as well.

Binding stoichiometry of Ca^{2+} and POPC was thoroughly studied in the experimental work [2], in which the head group order parameter changes to cation binding are determined. Several binding models were proposed and tested of which only one, ternary complex binding model, provided a good fit of the experimental observations.

Simulations allow us to directly evaluate the stoichiometry by calculating relative propensities of various $\text{Ca}^{2+}:n \times \text{POPC}$ clusters by evaluating contacts between cations and lipids with a cut off radius 0.3 nm. In Figure 5 we see that ternary complex is indeed the most probable binding mode of calcium at




FIG. 4: Number density of Ca^{2+} and Cl^- as a function of membrane normal axis for different force fields. Data for Lipid14 with Åqvist ions are taken directly from Ref. 17. Densities of Cl^- and water are divided with 2 and 200, respectively, to visualize them with the same scale as Ca^{2+} . The molar concentration of the ions in water is 350 mM in all systems presented here.

16.PAVEL: draw phosphate position with its variance, add water density (scaled) and include the number of T-surface access.

17.JOE: Change the figure so that it contains a membrane background

285 mM concentration. Apart from this complex, we also find complexes with 1 and 3 lipids occurring with only a slightly lower but similar probability. The fractions of $\text{Ca}^{2+}:n \times \text{POPC}$ complexes at 285 mM concentration are then in order: 42% for two lipids, 30% for one lipid, and 28% for three lipids.

Several binding models were proposed and tested [2] of which only one, ternary complex binding model, provided a good fit of the experimental observations. In such a model, it is assumed that Ca^{2+} cations bind to a POPC membrane with a stoichiometry 2 POPC:1 Ca^{2+} . In a later work [64], a Langmuir adsorption model (i.e. stoichiometry 1 POPC:1 Ca^{2+}) was found to provide as good fit as ternary complex model, when only low concentrations of CaCl_2 are considered. Ternary complex model also provides a good fit to our simulations with ECC-lipids (see Fig. 7 in SI and its




FIG. 5: Relative probabilities of existence of Na^+ or Ca^{2+} complexes with a certain number of POPC lipids. Na^+ complexes were evaluated from the simulation with 1 M concentration; and Ca^{2+} complexes were evaluated from the simulation with 287 mM concentration.

caption for details). The symmetry of the distribution of complexes from simulation – i.e. almost equal probabilities of complexes with 1 or 3 lipids that behave in the total average picture as complexes with 2 lipids – provides clues why ternary complex binding model fits both simulation and experimental results relatively well, although it is apparently incorrect.

In addition, we estimated relative binding affinities of several moieties in POPC towards Ca^{2+} . The dominant contribution to the binding of Ca^{2+} to POPC membranes comes from the phosphate group. Such a finding is supported by the calculated number of contacts between Ca^{2+} and POPC oxygen atoms. The average number of contacts between Ca^{2+} and any oxygen atom in POPC is 59.8, whereas if only phosphate oxygen atoms are considered the average number of contacts decreases only by a tiny amount to 57.2. We corroborate this finding with the probability isodensity contours in Fig. 6, which are easier to interpret.

The residence time of Ca^{2+} bound to a POPC membrane is experimentally estimated to be lower than $10 \mu\text{s}$ [2]. From recent theoretical work with long enough simulations this time can be roughly estimated in the order of $1\text{--}10 \mu\text{s}$ [13]. This is in contrast to our model, ECC-lipids, which gives a mean residence time in the order of 10 ns, **19.evaluate this number, mean residence time, accurately based on the contacts data.** i.e. at least two orders of magnitude lower than previous estimates. Such a finding changes the point of view of calcium binding from very tight long-term stable binding with rare exchanges to a relatively frequent exchange of cations in equilibrium between membrane and solvent.

20.Finalize stoichiometry analysis for Na^+ , Ca^{2+} , their interaction energies with the lipid membrane, etc, and finalize the discussion after these results.

IV. CONCLUSIONS

By using the electrometer concept we show that the binding of Na^+ and Ca^{2+} ions to a POPC lipid bilayer can be accurately described with the classical MD simulation force field, where the electronic polarization is implicitly included




FIG. 6: Contours of probability isodensities of Ca^{2+} with respect to various moieties fixed in space (highlighted with transparent spheres): phosphate moiety, side chain 1 carbonyl group and side chain 2 carbonyl group. Shown contours suggest that the dominant contribution to Ca^{2+} binding comes from the phosphate oxygens, whereas the interactions with any of the two carbonyl groups are considerably milder.

21.JOE: I'll update this figure with some ensemble of configuration to support binding preference of Ca^{2+}

using the electronic continuum correction (ECC) [20]. The proposed ECC-lipids model is a significant improvement over other available lipid models, which all overestimate specific cation binding affinities [17]. While the structural details of a POPC lipid bilayer simulated with the ECC-lipids model agree with experiments with the comparable accuracy to the other state of the art lipid models, it also reproduces the experimental lipid head group order parameter responses to the cationic surfactant, NaCl and CaCl_2 concentrations.

The good agreement with experiments enables the atomic resolution interpretation of NMR experiments by using MD simulations. In agreement with previous interpretations of experimental data [6, 14–16], the Ca^{2+} ions mainly interact with phosphate oxygens. However, the stoichiometry of the binding is significantly more complicated than in the previous interpretation of the NMR data based on the ternary complex model, where one calcium binds to two POPC molecules [2]. The complexes of one calcium bound to two lipids are the most probable also in the ECC-lipids model, but the complexes of one or three lipids per one calcium were observed to be almost equally likely. While the success of the ternary complex model is understandable based on the simulation results, a simple binding model cannot detect the complex binding observed in the simulation.

The improved cation binding behaviour to POPC bilayer pave the way for simulations of complex biochemical systems

with correctly described electrostatic interactions in the vicinity of cellular membranes. The ECC-lipids model is build by scaling the partial charges and the LJ-radius of the head-group, glycerol backbone and carbonyl atoms of the Lipid14 POPC model [24]. While the Lipid14 model is compatible with the AMBER force field family, the compatibility of the ECC-lipids model may be compromised due to the changes in intermolecular interactions of the scaled atoms. On the other hand, a fully consistent ECC-force field should include the correction also in other than lipid molecules, including water. The work toward this direction and the extension to other lipid molecules and force fields is left for the future studies.

This work can be reached as a repository containing all data at [zenodo.org:\dots\dots\dots](https://zenodo.org/records/10000000000000000000) and as project NMRlipids VI in nmrlipids.blogspot.fi.

Acknowledgments

P.J. acknowledges support from the Czech Science Foundation (grant no. 16-01074S) and 600 from the Academy of Finland via the FiDiPro award.




FIG. 7: Ternary complex binding model of Ca^{2+} to a POPC membrane that assumes the stoichiometry of 2 POPC:1 Ca^{2+} (details in reference 2) provides a good fit to experimental measurements [2] and it also provides a good fit to our simulation data. Note that the units in the reference 2 are different from the units presented here, and, hence, the observed slope of the linear relationship is different.

SUPPLEMENTARY INFORMATION

It was found in the original work [2] that a ternary complex binding model (i.e. 2 POPC:1 Ca^{2+}) provides the best fit to experimental measurements of all considered models in that study. In such a model, there is a linear relationship between quantities C_b , mole fraction of bound Ca^{2+} per POPC, and $\sqrt{C_b/C_I}$, where C_I is the concentration of free cations at the plane of ion binding [2]. The concentration C_b was obtained from an extrapolation of linear relation between deuterium NMR measurements and atomic absorption spectroscopy for low concentrations of CaCl_2 . Such an extrapolation is valid as long as the mode of Ca^{2+} binding remains constant throughout the extrapolation range. The concentration C_I is determined by using the surface potential by using the Boltzmann equation. However, Boltzmann theory yields inaccurate results for divalent cations like Ca^{2+} [67]. An atomistic simulation, on the other hand, provides these quantities directly without severe assumptions. **22. Did you really calculate the C_I from simulations without severe assumptions? Note that this concentration at the plane of binding, which do not equal the concentration of free cations.** Hence we hypothesise that the discrepancy between the results in the experiment [2] and our simulations likely lays in the fact that the assumptions and relations used for determining concentrations C_b and C_I in the experiment [2] gradually do not hold for higher concentrations of Ca^{2+} .

-
- [1] H. Akutsu and J. Seelig, *Biochemistry* **20**, 7366 (1981).
 - [2] C. Altenbach and J. Seelig, *Biochemistry* **23**, 3913 (1984).
 - [3] J. Seelig, *Cell Biol. Int. Rep.* **14**, 353 (1990), URL [http://dx.doi.org/10.1016/0309-1651\(90\)91204-H](http://dx.doi.org/10.1016/0309-1651(90)91204-H).
 - [4] G. Cevc, *Biochim. Biophys. Acta - Rev. Biomemb.* **1031**, 311 (1990).
 - [5] J.-F. Tocanne and J. Teissie, *Biochim. Biophys. Acta - Reviews on Biomembranes* **1031**, 111 (1990).
 - [6] H. Binder and O. Zschörnig, *Chem. Phys. Lipids* **115**, 39 (2002).
 - [7] G. Pabst, A. Hodzic, J. Strancar, S. Danner, M. Rappolt, and P. Laggner, *Biophys. J.* **93**, 2688 (2007).
 - [8] D. Uhrkov, N. Kuerka, J. Teixeira, V. Gordeliy, and P. Balgav, *Chemistry and Physics of Lipids* **155**, 80 (2008).
 - [9] R. A. Böckmann, A. Hac, T. Heimburg, and H. Grubmüller, *Biophys. J.* **85**, 1647 (2003).
 - [10] R. A. Böckmann and H. Grubmüller, *Ang. Chem. Int. Ed.* **43**, 1021 (2004).
 - [11] M. L. Berkowitz and R. Vacha, *Acc. Chem. Res.* **45**, 74 (2012).
 - [12] A. Melcrov, S. Pokorna, S. Pullanchery, M. Kohagen, P. Jurkiewicz, M. Hof, P. Jungwirth, P. S. Cremer, and L. Cwiklik, *Sci. Reports* **6**, 38035 (2016).




FIG. 8: Changes of head group order parameters of POPC bilayer as a function of CaCl_2 concentrations are shown from simulations with different force fields and water models together with experimental data (DPPC [1] and POPC [2]). Ion concentrations in bulk water are shown in x-axis. Values from simulations are calculated from the of cation number density C_{np} from the region at the simulation box edge with the constant ion concentration as $[\text{ion}] = C_{np}/0.602$.




FIG. 9: Changes of head group order parameters of POPC bilayer as a function of NaCl concentrations are shown from simulations with different force fields together with experimental data [1]. Ion concentrations in bulk water are shown in x-axis. Values from simulations are calculated from the of cation number density C_{np} from the region at the simulation box edge with the constant ion concentration as $[\text{ion}] = C_{np}/0.602$. Simulation data with Lipid14 and Åqvist ion parameters is taken directly from Ref. [17].

TABLE II: Area per lipid (APL) from different models of POPC with no ions

model	APL (\AA^2)	Temperature [K]
Lipid14	65.1 ± 0.6	300
Lipid14 [24]	65.6 ± 0.5	303
ECC-lipids		
OPC3	62.2 ± 0.6	300
OPC3	64.2 ± 0.6	313
SPC/E	65.1 ± 0.6	313
OPC	64.4 ± 0.6	313
TIP4p/2005	66.8 ± 0.6	313
experiment	62.7	293
experiment [59] ^{23,REF}	64.3	303
experiment	67.3	323
experiment	68.1	333

TABLE III: Simulation parameters

simulation property	parameter
time-step	2 fs
equilibration time	100 ns
simulation time	200 ns
temperature	313 K
thermostat	v-rescale [68]
barostat	Parrinello-Rahman, semi-isotropic [69]
long-range electrostatics	PME [70]
cut-off scheme	Verlet [71]
Coulomb and VdW cut-off	1.0 nm
constraints	LINCS, only hydrogen atoms [72]
constraints for water	SETTLE [73]

- [13] M. Javanainen, A. Melcrova, A. Magarkar, P. Jurkiewicz, M. Hof, P. Jungwirth, and H. Martinez-Seara, *Chem. Commun.* **53**, 5380 (2017), URL <http://dx.doi.org/10.1039/C7CC02208E>.
- [14] H. Hauser, M. C. Phillips, B. Levine, and R. Williams, *Nature* **261**, 390 (1976).
- [15] H. Hauser, W. Guyer, B. Levine, P. Skrabal, and R. Williams, *Biochim. Biophys. Acta - Biomembranes* **508**, 450 (1978), ISSN 0005-2736, URL <http://www.sciencedirect.com/science/article/pii/0005273678900913>.
- [16] L. Herbette, C. Napolitano, and R. McDaniel, *Biophys. J.* **46**, 677 (1984).
- [17] A. Catte, M. Giryach, M. Javanainen, C. Loison, J. Melcr, M. S. Miettinen, L. Monticelli, J. Maatta, V. S. Oganessian, O. H. S. Ollila, et al., *Phys. Chem. Chem. Phys.* **18** (2016).
- [18] R. Vacha, S. W. I. Siu, M. Petrov, R. A. Böckmann, J. Barucha-Kraszewska, P. Jurkiewicz, M. Hof, M. L. Berkowitz, and P. Jungwirth, *J. Phys. Chem. A* **113**, 7235 (2009).
- [19] J. Seelig, P. M. MacDonald, and P. G. Scherer, *Biochemistry* **26**, 7535 (1987).
- [20] I. Leontyev and A. Stuchebrukhov, *Phys. Chem. Chem. Phys.* **13**, 2613 (2011).
- [21] E. Pluhaová, H. E. Fischer, P. E. Mason, and P. Jungwirth, *Molecular Physics* **112**, 1230 (2014), ISSN 0026-8976, URL <http://www.tandfonline.com/doi/abs/10.1080/00268976.2013.875231>.
- [22] M. Kohagen, P. E. Mason, and P. Jungwirth, *J. Phys. Chem. B* **118**, 7902 (2014).
- [23] M. Kohagen, P. E. Mason, and P. Jungwirth, *J. Phys. Chem. B* **120**, 1454 (2016).
- [24] C. J. Dickson, B. D. Madej, A. Skjevik, R. M. Betz, K. Teigen, I. R. Gould, and R. C. Walker, *J. Chem. Theory Comput.* **10**, 865 (2014).
- [25] J. Chowdhary, E. Harder, P. E. M. Lopes, L. Huang, A. D. MacKerell, and B. Roux, *J. Phys. Chem. B* **117**, 9142 (2013).
- [26] B. Jonsson, O. Edholm, and O. Teleman, *J. Chem. Phys.* **85**, 2259 (1986).
- [27] E. Egberts, S.-J. Marrink, and H. J. C. Berendsen, *European Biophysics Journal* **22**, 423 (1994).
- [28] I. V. Leontyev and A. A. Stuchebrukhov, *The Journal of chemical physics* **130**, 085102 (2009), ISSN 1089-7690, URL <http://scitation.aip.org/content/aip/journal/jcp/130/8/10.1063/1.3060164>.
- [29] I. V. Leontyev and A. A. Stuchebrukhov, *Journal of Chemical Theory and Computation* **6**, 1498 (2010), ISSN 1549-9618, URL <http://dx.doi.org/10.1021/ct9005807>.
- [30] H. Hu, Z. Lu, and and Weitao Yang*, *Journal of Chemical Theory and Computation* **3**, 1004 (2007), ISSN 1549-9618, URL <http://dx.doi.org/10.1021/ct600295n>.
- [31] C. C. I. Bayly, P. Cieplak, W. D. Cornell, and P. a. Kollman, *The Journal of Physical ...* **97**, 10269 (1993), ISSN 0022-3654, 93/2091- 10269\$04.00/0, URL <http://pubs.acs.org/doi/abs/10.1021/j100142a004>.
- [32] U. C. Singh and P. A. Kollman, *Journal of Computational Chemistry* **5**, 129 (1984), ISSN 1096987X.
- [33] P. G. Scherer and J. Seelig, *Biochemistry* **28**, 7720 (1989).
- [34] A. Botan, F. Favela-Rosales, P. F. J. Fuchs, M. Javanainen, M. Kanduć, W. Kulig, A. Lamberg, C. Loison, A. Lyubartsev, M. S. Miettinen, et al., *J. Phys. Chem. B* **119**, 15075 (2015).
- [35] O. S. Ollila and G. Pabst, *Atomistic resolution structure and dynamics of lipid bilayers in simulations and experiments* (2016), in Press, URL <http://dx.doi.org/10.1016/j.bbamem.2016.01.019>.
- [36] D. S. Cerutti, J. E. Rice, W. C. Swope, and D. A. Case, *The Journal of Physical Chemistry B* **117**, 2328 (2013), PMID: 23379664, <http://dx.doi.org/10.1021/jp311851r>, URL <http://dx.doi.org/10.1021/jp311851r>.
- [37] A. Maciejewski, M. Pasenkiewicz-Gierula, O. Cramariuc, I. Vattulainen, and T. Rog, *J. Phys. Chem. B* **118**, 4571 (2014).
- [38] (????).
- [39] T. M. Ferreira, R. Sood, R. Bärenwald, G. Carlström, D. Topgaard, K. Saalwächter, P. K. J. Kinnunen, and O. H. S. Ollila, *Langmuir* **32**, 6524 (2016).
- [40] A. Seelig and J. Seelig, *Biochemistry* **16**, 45 (1977).
- [41] J. H. Davis, *Biochim. Biophys. Acta - Reviews on Biomembranes* **737**, 117 (1983).
- [42] G. Beschiaschvili and J. Seelig, *Biochim. Biophys. Acta - Biomembranes* **1061**, 78 (1991).
- [43] P. Scherer and J. Seelig, *The EMBO journal* **6** (1987).
- [44] D. K. Chattoraj and K. S. Birdi, *Adsorption at the Liquid Interface from the Multicomponent Solution* (Springer US, Boston, MA, 1984), pp. 83–131, ISBN 978-1-4615-8333-2, URL https://doi.org/10.1007/978-1-4615-8333-2_4.
- [45] S. Izadi and A. V. Onufriev, *Journal of Chemical Physics* **145**, 074501 (2016), ISSN 00219606, URL <http://aip.scitation.org/doi/10.1063/1.4960175>.

- [46] S. Izadi, R. Anandakrishnan, and A. V. Onufriev, The Journal of Physical Chemistry Letters **5**, 3863 (2014), ISSN 1948-7185, 1408.1679, URL <http://pubs.acs.org/doi/10.1021/jz501780a>.
- [47] H. J. C. Berendsen, J. R. Grigera, and T. P. Straatsma, Journal of Physical Chemistry **91**, 6269 (1987), ISSN 0022-3654, URL <http://links.isiglobalnet2.com/gateway/Gateway.cgi?GWVersion=2\&SrcAuth=mekentosj\&SrcApp=Papers\&DestLinkType=FullRecord\&DestApp=WOS\&KeyUT=A1987K994100038\&5Cnpapers2://publication/uuid/17978EF7-93C9-4CB5-89B3-086E5D2B9169\&5Cnhttp://pubs.acs.org/doi/pdf/10.1021/jz500737m>.
- [48] L. P. Wang, T. J. Martinez, and V. S. Pande, Journal of Physical Chemistry Letters **5**, 1885 (2014), ISSN 19487185, URL <http://pubs.acs.org/doi/abs/10.1021/jz500737m>.
- [49] J. L. Abascal and C. Vega, The Journal of chemical physics **123**, 234505 (2005), ISSN 00219606, URL <http://aip.scitation.org/doi/10.1063/1.2121687>.
- [50] D. E. Smith and L. X. Dang, J. Chem. Phys **100** (1994).
- [51] T.-M. Chang and L. X. Dang, J. Phys. Chem. B **103**, 4714 (1999), ISSN 1520-6106, URL <http://dx.doi.org/10.1021/jp982079o>.
- [52] L. X. Dang, G. K. Schenter, V.-A. Glezakou, and J. L. Fulton, J. Phys. Chem. B **110**, 23644 (2006), ISSN 1520-6106, URL <http://dx.doi.org/10.1021/jp064661f>.
- [53] J. Aqvist, The Journal of Physical Chemistry **94**, 8021 (1990), URL <http://dx.doi.org/10.1021/j100384a009>.
- [54] M. J. Abraham, T. Murtola, R. Schulz, S. Páll, J. C. Smith, B. Hess, and E. Lindahl, SoftwareX **1-2**, 19 (2015), ISSN 23527110, URL <http://www.sciencedirect.com/science/article/pii/S2352711015000059>.
- [55] A. K. Malde, L. Zuo, M. Breeze, M. Stroet, D. Poger, P. C. Nair, C. Oostenbrink, and A. E. Mark, Journal of Chemical Theory and Computation **7**, 4026 (2011).
- [56] D. Case, D. Cerutti, T. Cheatham, III, T. Darden, R. Duke, T. Giese, H. Gohlke, A. Goetz, D. Greene, et al., *AMBER 2017* (2017), university of California, San Francisco.
- [57] A. W. SOUSA DA SILVA and W. F. VRANKEN, *ACPYPE - AnteChamber PYthon Parser interface*. (2017), manuscript submitted.
- [58] T. M. Ferreira, F. Coreta-Gomes, O. H. S. Ollila, M. J. Moreno, W. L. C. Vaz, and D. Topgaard, Phys. Chem. Chem. Phys. **15**, 1976 (2013).
- [59] J. P. M. Jämbbeck and A. P. Lyubartsev, J. Phys. Chem. B **116**, 3164 (2012).
- [60] (???)
- [61] (???)
- [62] (???)
- [63] H. I. Petrache, S. Tristram-Nagle, D. Harries, N. Kucerka, J. F. Nagle, and V. A. Parsegian, J. Lipid Res. **47**, 302 (2006).
- [64] P. M. Macdonald and J. Seelig, Biochemistry **26**, 1231 (1987).
- [65] L.-P. Wang, K. A. McKiernan, J. Gomes, K. A. Beauchamp, T. Head-Gordon, J. E. Rice, W. C. Swope, T. J. Martínez, and V. S. Pande, The Journal of Physical Chemistry B **121**, 4023 (2017), ISSN 1520-6106, URL <http://pubs.acs.org/doi/abs/10.1021/acs.jpcb.7b02320>.
- [66] W. L. Jorgensen, J. Chandrasekhar, J. D. Madura, R. W. Impey, and M. L. Klein, J. Chem. Phys **79** (1983).
- [67] D. Andelman, in *Handbook of biological physics* (Elsevier Science, 1995), vol. 1, chap. 12, pp. 603–642, URL <http://hwiki.liebel-lab.org/wiki/images/9/90/AndelmannReview.pdf>.
- [68] G. Bussi, D. Donadio, and M. Parrinello, J. Chem. Phys **126** (2007).
- [69] M. Parrinello and A. Rahman, J. Appl. Phys. **52**, 7182 (1981).
- [70] T. Darden, D. York, and L. Pedersen, J. Chem. Phys **98** (1993).
- [71] S. Páll and B. Hess, Computer Physics Communications **184**, 2641 (2013), ISSN 0010-4655, URL <http://www.sciencedirect.com/science/article/pii/S0010465513001975>.
- [72] B. Hess, H. Bekker, H. J. C. Berendsen, and J. G. E. M. Fraaije, J. Comput. Chem. **18**, 1463 (1997).
- [73] S. Miyamoto and P. A. Kollman, J. Comput. Chem **13**, 952 (1992).

ToDo

- | | P. |
|--|----|
| 1. Abstract directly from Joe's conference abstracts. To be rewritten. | 1 |
| 2. We should also cite papers where empirical scaling was used ionic liquids - but there the factor is not 0.5. . | 2 |
| 3. This needs a citation | 2 |
| 4. JOE: following discussion shall be modified in the enlightenment of our recent ECC-discussions. | 2 |
| 5. missing REF for APL experiment | 2 |
| 6. We should discuss how this can potentially affect the intermolecular interaction when mixing scaled and non scaled molecules. JOE: I think that we rather increasingly see that there's nothing like "fully non-scaled" with the exception of ions with integer charges. So the discussion shall be rather more about the interaction of our "scaled" (I'd still rather call it ECC-corrected or whatever) and "semi-scaled" models. | 3 |
| 7. JOE: I'm afraid that the above description might sound to the reader like for some reasons one method is used with DPPC, but a different one with POPC. I think simply writing one way is... another way is... (or similarly) will be less confusing. | 3 |
| 8. JOE: following is very unclear to me. I think we can write it in a more concise way. | 3 |
| 9. To be uploaded to Zenodo | 4 |
| 10. put original references, not Slipids param. paper. . | 4 |
| 11. Dynamics check is missing: MSD (Hector/Joe) . . | 5 |
| 12. SAMULI: We could calculate the slopes from simulations, but I am not sure if we would actually learn anything useful from this. | 6 |
| 13. Add OP-response of Lipid14+ECC-ions plot in SI . | 6 |
| 14. Below analysis is done in a stupid way to get some idea. I would find it useful to do this analysis by using the density profiles, but it is not necessary. The reason why I want to do something like this, is that it would be useful to have some easier way to estimate the correct binding affinity than the electrometer concept, which is quite tedious to apply in practise (simulations with different concentrations, cationic surfactant check, etc.). I would like to be able to estimate much faster from a simulation if the binding affinity is reasonable or not. This may not be the correct way for that anyway. . . . | 7 |

15. SAMULI: Maybe we should discuss the repeat distances and area per molecules measured at [7, 8, 63] . . .	7	17. JOE: Change the figure so that it contains a membrane background	7
18. SAMULI: I think the we should make clear that the current binding constants in, e.g., Marsh's handbook are based on the headgroup order parameter changes interpreted with ternary complex binding model. I.e. the experimental raw data is exactly the same as we have in this work. Our simulations enable much more versatile interpretation of the binding phenomena. I think that this is one the main points of this work, so this comment applies to the previous section, introduction, abstract and conclusions as well.	7	19. evaluate this number, mean residence time, accurately based on the contacts data.	8
16. PAVEL: draw phosphate position with its variance, add water density (scaled) and include the number of Γ -surface access.	7	20. Finalize stoichiometry analysis for Na^+ , Ca^{2+} , their interaction energies with the lipid membrane, etc, and finalize the discussion after these results.	8
		21. JOE: I'll update this figure with some ensemble of configuration to support binding preference of Ca^{2+} . .	8
		22. Did you really calculate the C_I from simulations without severe assumptions? Note that this concentration at the plane of binding, which do not equal the concentration of free cations.	9
		23. put original references, not Slipids param. paper. .	11