

Propositional logic

Solution 1

(a) $false(as(true \Rightarrow false) \Leftrightarrow false)$

(b) $true(as(false \Rightarrow false) \Leftrightarrow true)$

(c) $true(as(false \Rightarrow true) \Leftrightarrow true)$

(d)

$$true(as(false \Rightarrow false) \Leftrightarrow true)$$

(Assuming that pigs can't fly . . .)

Solution 2

(a)

p	q	$p \wedge q$	$(p \wedge q) \Rightarrow p$
t	t	t	t
t	f	f	t
f	t	f	t
f	f	f	t

(b)

p	q	$p \wedge q$	$\neg p$	$\neg p \Rightarrow (p \wedge q)$	$(\neg p \Rightarrow (p \wedge q)) \Leftrightarrow p$
t	t	t	f	t	t
t	f	f	f	t	t
f	t	f	t	f	t
f	f	f	t	f	t

(c)

p	q	$p \Rightarrow q$	$p \wedge (p \Rightarrow q)$	$(p \wedge (p \Rightarrow q)) \Rightarrow q$
t	t	t	t	t
t	f	f	f	t
f	t	t	f	t
f	f	t	f	t

Solution 3

(a)

$$\begin{aligned}
 p \Rightarrow \neg p & \\
 & \Leftrightarrow \neg p \vee \neg p && [\Rightarrow] \\
 & \Leftrightarrow \neg p && [\text{idempotence}]
 \end{aligned}$$

(b)

$$\begin{aligned}\neg p \Rightarrow p & \\ \Leftrightarrow \neg \neg p \vee p & \quad [\Rightarrow] \\ \Leftrightarrow p \vee p & \quad [\neg \neg] \\ \Leftrightarrow p & \quad [\text{idempotence}]\end{aligned}$$

(c)

$$\begin{aligned}p \Rightarrow (q \Rightarrow r) & \\ \Leftrightarrow \neg p \vee (q \Rightarrow r) & \quad [\Rightarrow] \\ \Leftrightarrow \neg p \vee \neg q \vee r & \quad [\Rightarrow] \\ \Leftrightarrow \neg p \vee \neg q \vee r & \quad [\text{associativity}] \\ \Leftrightarrow \neg (p \wedge q) \vee r & \quad [\text{De Morgan}] \\ \Leftrightarrow p \wedge q \Rightarrow r & \quad [\Rightarrow]\end{aligned}$$

(d)

$$\begin{aligned}q \Rightarrow (p \Rightarrow r) & \\ \Leftrightarrow \neg q \vee (p \Rightarrow r) & \quad [\Rightarrow] \\ \Leftrightarrow \neg q \vee \neg p \vee r & \quad [\Rightarrow] \\ \Leftrightarrow \neg p \vee \neg q \vee r & \quad [\text{associativity} \wedge \text{commutativity}] \\ \Leftrightarrow \neg p \vee (q \Rightarrow r) & \quad [\Rightarrow] \\ \Leftrightarrow p \Rightarrow (q \Rightarrow r) & \quad [\Rightarrow]\end{aligned}$$

(e)

$$\begin{aligned}p \wedge q \Leftrightarrow p & \\ \Leftrightarrow (p \wedge q \Rightarrow p) \wedge (p \Rightarrow p \wedge q) & \quad [\Leftrightarrow] \\ \Leftrightarrow (\neg (p \wedge q) \vee p) \wedge (\neg p \vee p \wedge q) & \quad [\Rightarrow] \\ \Leftrightarrow (\neg p \vee \neg q \vee p) \wedge (\neg p \vee p \wedge q) & \quad [\text{De Morgan}] \\ \Leftrightarrow (\neg q \vee \neg p \vee p) \wedge (\neg p \vee p \wedge q) & \quad [\text{associativity} \wedge \text{comm.}] \\ \Leftrightarrow (\neg q \vee \text{true}) \wedge (\neg p \vee p \wedge q) & \quad [\text{excluded middle}] \\ \Leftrightarrow \text{true} \wedge (\neg p \vee p \wedge q) & \quad [\vee \wedge \text{true}] \\ \Leftrightarrow \neg p \vee p \wedge q & \quad [\wedge \wedge \text{true}] \\ \Leftrightarrow (\neg p \vee p) \wedge (\neg p \vee q) & \quad [\text{distribution}] \\ \Leftrightarrow \text{true} \wedge (\neg p \vee q) & \quad [\text{excluded middle}] \\ \Leftrightarrow \neg p \vee q & \quad [\wedge \wedge \text{true}] \\ \Leftrightarrow p \Rightarrow q & \quad [\Rightarrow]\end{aligned}$$

(f)

$$\begin{aligned}
p \vee q &\Leftrightarrow p \\
&\Leftrightarrow (p \vee q \Rightarrow p) \wedge (p \Rightarrow p \vee q) && [\Leftrightarrow] \\
&\Leftrightarrow (\neg (p \vee q) \vee p) \wedge (\neg p \vee p \vee q) && [\Rightarrow] \\
&\Leftrightarrow (\neg p \wedge \neg q \vee p) \wedge (\neg p \vee p \vee q) && [\text{De Morgan}] \\
&\Leftrightarrow (\neg p \vee p) \wedge (\neg q \vee p) \wedge (\neg p \vee p \vee q) && [\text{distribution}] \\
&\Leftrightarrow \text{true} \wedge (\neg q \vee p) \wedge (\neg p \vee p \vee q) && [\text{excluded middle}] \\
&\Leftrightarrow (\neg q \vee p) \wedge (\neg p \vee p \vee q) && [\wedge \wedge \text{true}] \\
&\Leftrightarrow (\neg q \vee p) \wedge (\neg p \vee p \vee q) && [\text{associativity}] \\
&\Leftrightarrow (\neg q \vee p) \wedge (\text{true} \vee q) && [\text{excluded middle}] \\
&\Leftrightarrow (\neg q \vee p) \wedge \text{true} && [\vee \wedge \text{true}] \\
&\Leftrightarrow \neg q \vee p && [\wedge \wedge \text{true}] \\
&\Leftrightarrow q \Rightarrow p && [\Rightarrow]
\end{aligned}$$

Solution 4

- (a) $(p \text{ or } q) \Leftrightarrow ((\text{not } p \text{ or not } q) \text{ and } q)$ is not a tautology. You might illustrate this via a truth table or via a chain of equivalences, showing that the proposition is not equivalent to true. Alternatively, you might try and find a combination of values for which the proposition is false. (In this case, the proposition is false when p and q are both true.)
- (b) $(p \text{ or } q) \Leftrightarrow ((\text{not } p \text{ and not } q) \text{ or } q)$ is not a tautology. In this case, the proposition is false when p is true and q is false.

Solution 5

- (a) $\exists d: \text{Dog} \bullet \text{gentle}(d) \wedge \text{well_trained}(d)$
- (b) $\forall d: \text{Dog} \bullet \text{neat}(d) \wedge \text{well_trained}(d) \Rightarrow \text{attractive}(d)$
- (c) (Requires nested quantifier in implication - parser limitation)

Solution 6

- (a) This is a true *proposition* : *whatever the value of x, the expression* $x^2 - x + 1$ *denotes a natural number. If we choose y to be this natural number, we will find that p is true.*
- (b) This is a false proposition. We cannot choose a large enough value for y such that p will hold for any value of x.
- (c) This is a false proposition. It is an implication whose antecedent part is true and whose consequent part is false.
- (d) This is a true proposition. It is an implication whose antecedent part is false and whose consequent part is true.

Solution 7

- (a) We must define a predicate p that is false for at least one value of x , and is true for at least one other value. A suitable solution would be $p \Leftrightarrow x > 1$.
- (b) With the above choice of p , we require only that q is sometimes false when p is true (for else the universal quantification would hold). A suitable solution would be $q \Leftrightarrow x > 3$.

Solution 8

- (a) $\forall x: N \bullet x \geq z$

Equality**Solution 9**

- (d)

$$\begin{aligned}
& \exists x: N \bullet x = 1 \wedge x > y \vee x = 2 \wedge x > z \\
& \Leftrightarrow \exists x: N \bullet x = 1 \wedge x > y \vee \exists x: N \bullet x = 2 \wedge x > z \\
& \Leftrightarrow 1 \in N \wedge 1 > y \vee \exists x: N \bullet x = 2 \wedge x > z \\
& \Leftrightarrow 1 \in N \wedge 1 > y \vee 2 \in N \wedge 2 > z \\
& \Leftrightarrow 1 > y \vee 2 > z
\end{aligned}$$

Solution 10

As discussed, the quantifier \exists_1 can help give rise to a 'test' or 'precondition' to ensure that an application of μ will work.

So, as a simple example, as the proposition

$$\exists_1 n: N \bullet \forall m: N \bullet n \leq m$$

is equivalent to true, we can be certain that the statement

$$\mu n: N \bullet \forall m: N \bullet n \leq m$$

will return a result (which happens to be 0).

Solution 11

- (a) $(\mu a: N \mid a = a) = 0$ is a provable statement, since 0 is the only natural number with the specified property.

- (b) $(\mu b: N \bullet b = b) = 1$ is not provable. The specified property is true of both 0 and 1, and thus the value of the mu-expression is undefined.
- (c) $(\mu c: N \bullet c > c) = (\mu c: N \bullet c > c)$ is a provable statement. Neither expression is properly defined, but we may conclude that they are equal; there is little else that we can prove about them.
- (d) $(\mu d: N \bullet d = d) = 1$ is not a provable statement. We cannot confirm that 1 is the only natural number with the specified property; we do not know what value is taken by undefined operations.

Solution 12

(Requires mu-operator with expression part - not yet implemented)

- (a) $\mu m: Mountain \mid \forall n: Mountain \bullet height(n) \leq height(m) \bullet height(m)$
 (b) $\mu c: Chapter \mid \exists_1 d: Chapter \bullet length(d) > length(c) \bullet length(c)$
 (c) Assuming the existence of a suitable function, max: $(\mu n: N \bullet n = \max(\{m: N \mid 8 * m < 100.8 * m\}) \cdot 100 - n)$

Deductive proofs

Solution 13

$$\frac{\frac{\frac{\vdash p \wedge (p \Rightarrow q) \neg^{[1]}}{p \wedge (p \Rightarrow q) \Rightarrow p \wedge q} [\Rightarrow\text{-intro}^{[1]}] \quad \frac{\frac{\frac{p \wedge (p \Rightarrow q)}{p} [\wedge\text{-elim}^{[1]}] \quad \frac{\frac{p \wedge (p \Rightarrow q)}{p \Rightarrow q} [\wedge\text{-elim}^{[1]}] \quad \frac{p \wedge (p \Rightarrow q)}{p} [\wedge\text{-elim}^{[1]}]}{q} [\wedge\text{intro}]}}{} {}$$

Solution 14

In one direction:

[illegible]

and the other:

$$\frac{\frac{\frac{\frac{\Gamma p \wedge q^{\neg[2]} \quad \Gamma p^{\neg[2]}}{p \wedge q \Rightarrow p} [\Rightarrow\text{-intro}^{[2]}] \quad \frac{\frac{\Gamma p^{\neg[3]} \quad \Gamma p \wedge q^{\neg[1]}}{p \Rightarrow p \wedge q} [\Rightarrow\text{-intro}^{[3]}]}{p \wedge q \Leftrightarrow p} [\Leftrightarrow\text{intro}]}{\Gamma p \Rightarrow q^{\neg[1]} \quad p \wedge q \Leftrightarrow p} [\Rightarrow\text{-intro}^{[1]}]$$

We can then combine these two proofs *with \Leftrightarrow intro*.

Solution 15

[illegible]

Solution 16

In one direction:

$$\begin{array}{c}
\frac{\frac{\frac{}{\vdash p \neg[1]} \quad \frac{}{\neg r}}{p \wedge r} [\wedge \text{ intro}] \quad \frac{}{p \wedge q \vee p \wedge r} [\vee \text{ intro}]}{\frac{}{\vdash p \wedge (q \vee r) \neg[1]} \quad \frac{}{p \wedge q \vee p \wedge r} [\vee \text{ elim}^{[2]}]}{\frac{}{p \wedge (q \vee r) \Rightarrow p \wedge q \vee p \wedge r} [\Rightarrow\text{-intro}^{[1]}]} \\
\\
\frac{\frac{\frac{}{\vdash p \neg[1]} \quad \frac{}{\neg q} [\text{case assumption}]}{p \wedge q} [\wedge \text{ intro}] \quad \frac{}{p \wedge q \vee p \wedge r} [\vee \text{ intro}]}{\frac{}{\vdash q \vee r \neg[1]}}
\end{array}$$

In the other:

[illegible]

Solution 17

In one direction:

$$\frac{\frac{\lceil p \vee q \wedge r \neg^{[3]} \rceil \quad \overline{(p \vee q) \wedge (p \vee r)}}{p \vee q \wedge r \Rightarrow (p \vee q) \wedge (p \vee r)} [\vee \text{ elim } \wedge \wedge \text{ intro}]}{p \vee q \wedge r \Rightarrow (p \vee q) \wedge (p \vee r)} [\Rightarrow\text{-intro}^{[3]}]$$

and the other:

$$\frac{\frac{\lceil (p \vee q) \wedge (p \vee r) \neg^{[1]} \rceil \quad \lceil p \vee q \wedge r \neg^{[2]} \rceil}{(p \vee q) \wedge (p \vee r) \Rightarrow p \vee q \wedge r} [\Rightarrow\text{-intro}^{[1]}]}{p \vee q \wedge r \Rightarrow (p \vee q) \wedge (p \vee r)}$$

Solution 18

In one direction:

$$\frac{\frac{\lceil p \Rightarrow q \neg^{[1]} \rceil \quad \neg p \vee q}{(p \Rightarrow q) \Rightarrow \neg p \vee q} [\Rightarrow\text{-intro}^{[1]}]}{p \Rightarrow q \Rightarrow \neg p \vee q}$$

and the other:

$$\frac{\frac{\lceil \neg p \vee q \neg^{[3]} \rceil \quad \frac{\frac{\lceil p \neg^{[4]} \rceil \quad \lceil q \neg^{[3]} \rceil}{p \Rightarrow q} [\Rightarrow\text{-intro}^{[4]}]}{\neg p \vee q \Rightarrow (p \Rightarrow q)} [\Rightarrow\text{-intro}^{[3]}]}{\neg p \vee q \Rightarrow (p \Rightarrow q)}$$

Sets and types

Solution 19

- (a) 1 in $\{4, 3, 2, 1\}$ is true.
- (b) $\{1\}$ in $\{1, 2, 3, 4\}$ is undefined.
- (c) $\{1\}$ in $\{\{1\}, \{2\}, \{3\}, \{4\}\}$ is true.
- (d) The empty set in $\{1, 2, 3, 4\}$ is undefined.

Solution 20

(a)

$$\{1\} \times \{2, 3\}$$

is the set $\{(1, 2), (1, 3)\}$

(b) The empty set cross $\{2, 3\}$ is the empty set

(c)

$$\mathbb{P} \emptyset \times \{1\}$$

is the set $\{(\emptyset, 1)\}$

(d) $\{(1, 2)\}$ cross $\{3, 4\}$ is the set $\{((1, 2), 3), ((1, 2), 4)\}$

Solution 21

There are various ways of describing these sets via set comprehensions. Examples are given below.

(a) $\{z: Z \mid 0 \leq z \wedge z \leq 100\}$

(b) $\{z: Z \mid z = 10\}$

(c) $\{z: Z \mid z \bmod 2 = 0 \vee z \bmod 3 = 0 \vee z \bmod 5 = 0\}$

Solution 22

(a) $\{n: N \mid n \leq 4 \bullet n^2\}$

(b) $\{n: N \mid n \leq 4 \bullet (n, n^2)\}$

(c) $n: P0, 1$

(d) $\{n: \mathbb{P}\{0, 1\} \mid true \bullet (n, \#n)\}$

Solution 23

(a)

$$\begin{aligned} x &\in a \cap a \\ &\Leftrightarrow x \in a \wedge x \in a \\ &\Leftrightarrow x \in a \end{aligned}$$

(b)

$$\begin{aligned} x &\in a \cup a \\ &\Leftrightarrow x \in a \vee x \in a \\ &\Leftrightarrow x \in a \end{aligned}$$

Solution 24

(a)

The set of all pairs of integers is Z cross Z . To give it a name, we could write:

(b)

StrictlyPositivePairs == { m, n : Z | $m > 0 \wedge n > 0 \bullet (m, n)$ }

(c)

[*Person*]

(d)

$$\text{Couples} == \{ s : \mathbb{P} \text{ Person} \mid \#s = 2 \}$$

(Requires generic set notation and Cartesian product)

(Requires generic parameters and relation type notation)

Solution 27

(a)

$$\{\emptyset, \{(0,0)\}, \{(0,1)\}, \{(1,0)\}, \{(1,1)\}, \{(1,0), (1,1)\}, \{(0,0), (0,1)\}, \{(0,1), (1,1)\}, \{(0,1), (1,0)\}, \{(0,0), (1,1)\}, \{(0,0)$$

(b) $\{\emptyset, \{(0, 0)\}, \{(0, 1)\}, \{(0, 0), (0, 1)\}\}$

(c) $\{\emptyset\}$

(d) $\{\emptyset\}$

Solution 28

- (a) $\text{dom } R = \{0, 1, 2\}$
- (b) $\text{ran } R = \{1, 2, 3\}$
- (c) $\{1, 2\} \triangleleft R = \{1 \mapsto 2, 1 \mapsto 3, 2 \mapsto 3\}$

Solution 29

- (a) $\{2 \mapsto 4, 3 \mapsto 3, 3 \mapsto 4, 4 \mapsto 2\}$
- (b) $\{1 \mapsto 3, 2 \mapsto 2, 2 \mapsto 3, 3 \mapsto 1\}$
- (c) $\{1 \mapsto 1, 2 \mapsto 2, 2 \mapsto 3, 3 \mapsto 2, 3 \mapsto 3, 4 \mapsto 4\}$
- (d) $\{1 \mapsto 4, 2 \mapsto 2, 2 \mapsto 3, 3 \mapsto 2, 3 \mapsto 3, 4 \mapsto 1\}$

Solution 30

$$\mid \text{ childOf} : \text{Person} \leftrightarrow \text{Person}$$

(a)

$$\text{parentOf} == \text{childOf}^{-1}$$

This is a good example of how there are many different ways of writing the same thing. An alternative abbreviation is:

$$\text{parentOf} == \{ x, y : \text{Person} \mid x \mapsto y \in \text{childOf} \bullet y \mapsto x \}$$

Or, via an axiomatic definition:

$$\frac{\mid \text{ parentOf} : \text{Person} \leftrightarrow \text{Person}}{\mid \text{ parentOf} = \text{childOf}^{-1}}$$

(b)

$$\text{siblingOf} == (\text{childOf} \circ \text{parentOf}) \setminus \text{id}$$

(c)

$$\text{cousinOf} == \text{childOf} \circ \text{siblingOf} \circ \text{parentOf}$$

(d)

$$\text{ancestorOf} == \text{parentOf}^+$$

Solution 31

(Requires compound identifiers with operators - \mathbb{R}^+ , \mathbb{R}^*)

(a)

$$\mathbb{R} == \{ a, b : \mathbb{N} \mid b = a \vee b = a \}$$

(b)

$$S == \{ a, b : N \mid b = a \vee b = a \}$$

$$(c) R+ == \{ a, b : N \mid b > a \}$$

$$(d) R^* == \{ a, b : N \mid b \geq a \}$$

Solution 32

(a)

$$x \mapsto y \in A \triangleleft B \triangleleft R$$

$$\Leftrightarrow x \in A \wedge x \mapsto y \in (B \triangleleft R)$$

$$\Leftrightarrow x \in A \wedge x \in B \wedge x \mapsto y \in R$$

$$\Leftrightarrow x \in A \cap B \wedge x \mapsto y \in R$$

$$\Leftrightarrow x \mapsto y \in A \cap B \triangleleft R$$

(b)

$$x \mapsto y \in R \cup S \triangleright C$$

$$\Leftrightarrow x \mapsto y \in R \cup S \wedge y \in C$$

$$\Leftrightarrow (x \mapsto y \in R \vee x \mapsto y \in S) \wedge y \in C$$

$$\Leftrightarrow x \mapsto y \in R \wedge y \in C \vee x \mapsto y \in S \wedge y \in C$$

$$\Leftrightarrow x \mapsto y \in R \triangleright C \vee x \mapsto y \in S \triangleright C$$

$$\Leftrightarrow x \mapsto y \in (R \triangleright C) \cup (S \triangleright C)$$

Functions

Solution 33

The set of 9 functions:

$$\{\emptyset, \{(0,0)\}, \{(0,1)\}, \{(1,1)\}, \{(1,0)\}, \{(0,0), (1,1)\}, \{(0,1), (1,1)\}, \{(1,0), (0,0)\}, \{(0,1), (1,0)\}\}$$

(a)

The set of total functions:

$$\{\{(0,0), (1,1)\}, \{(0,1), (1,1)\}, \{(1,0), (0,0)\}, \{(0,1), (1,0)\}\}$$

(b)

The set of functions which are neither injective nor surjective:

$$\{\{(0, 1), (1, 1)\}, \{(0, 0), (1, 0)\}\}$$

(c)

The set of functions which are injective but not surjective:

$$\{\emptyset, \{(0, 0)\}, \{(0, 1)\}, \{(1, 0)\}, \{(1, 1)\}\}$$

(d) There are no functions (of this type) which are surjective but not injective.

(e)

The set of bijective functions:

$$\{\{(0, 0), (1, 1)\}, \{(0, 1), (1, 0)\}\}$$

Solution 34

(a) $\{1 \mapsto a, 2 \mapsto b, 3 \mapsto c, 4 \mapsto b\}$

(b) $\{1 \mapsto c, 2 \mapsto b, 3 \mapsto c, 4 \mapsto d\}$

(c) $\{1 \mapsto c, 2 \mapsto b, 3 \mapsto c, 4 \mapsto b\}$

(d) $\{1 \mapsto c, 2 \mapsto b, 3 \mapsto c, 4 \mapsto b\}$

Solution 35

(Requires power set notation \mathbb{P} and relational image)

(a)

$$\frac{\text{children} : \text{Person} \rightarrow \mathbb{P} \text{Person}}{\text{children} = \{p : \text{Person} \bullet p \mapsto \text{parentOf}(\llbracket \{p\} \rrbracket)\}}$$

(b)

$$\frac{\text{number_of_grandchildren} : \text{Person} \rightarrow N}{\text{number_of_grandchildren} = \{p : \text{Person} \bullet p \mapsto \# \text{parentOf} \circ \text{parentOf}(\llbracket \{p\} \rrbracket)\}}$$

Solution 36

(Note: This solution demonstrates relation types in quantifier domains)

$$\frac{\text{number_of_drivers} : \text{Drivers} \leftrightarrow \text{Cars} \rightarrow (\text{Cars} \rightarrow N)}{\text{number_of_drivers} = \lambda r : \text{Drivers} \leftrightarrow \text{Cars} \bullet \{c : \text{ran } r \bullet c \mapsto \#\{d : \text{Drivers} \mid d \mapsto c \in r\}\}}$$

Sequences

Solution 37

- (a) $\langle a \rangle$
- (b) $\{1 \mapsto a, 2 \mapsto b, 2 \mapsto a, 3 \mapsto c, 3 \mapsto b, 4 \mapsto d\}$
- (c) $\{2 \mapsto b, 3 \mapsto c, 4 \mapsto d\}$
- (d) $\{1, 2, 3, 4\}$
- (e) $\{a, b\}$
- (f) $\{a \mapsto 1, b \mapsto 2, c \mapsto 3, d \mapsto 4\}$
- (g) $\langle a, b \rangle$
- (h) $\{3 \mapsto b\}$
- (i) $\{a\}$
- (j) c

Solution 38

- (a)

$$\frac{f : Place \rightarrow \mathbb{P} Place}{\forall p : Place \bullet f(p) = \{q : Place \mid p \mapsto q \in \text{ran } \text{trains}\}}$$

- (b) $\{p : Place \mid \exists_1 x : \text{dom } \text{trains} \bullet \text{trains}(x).2 = p\}$
- (c) $\mu p : Place \bullet \forall q : Place \bullet p \neq q \wedge \#\{x : \text{dom } \text{trains} \mid \text{trains}(x).2 = p\} > \#\{x : \text{dom } \text{trains} \mid \text{trains}(x).2 = q\}$

Solution 39

- (a)

$$\text{large_coins} : Collection \rightarrow N$$

$$\forall c : Collection \bullet \text{large_coins}(c) = c(\text{large})$$

(Blocked by : *underscoreinidentifierforfuzzcompatibility*)

- (b)

$$\text{add_coin} : Collection * Coin \rightarrow Collection$$

$$\forall c : Collection \bullet \forall d : Coin \bullet \text{add_coin}(c, d) = c \cup \llbracket d \rrbracket$$

(Blocked by : *underscoreinidentifierandbagunion*)

Modelling

Solutions 40-52 are work in progress - many require features not yet implemented

Solution 40

(Work in progress - requires semicolon-separated bindings in set comprehensions)

(a)

$hd : seq(Title * Length * Viewed)$

$cumulative_total(hd) \leq 12000$

$\forall p: \text{ran } hd \bullet p.2 \leq 360$

Note that $cumulative_total$ is defined in part (d).

(b) $\{p: \text{ran } hd \mid p.2 > 120 \bullet p.1\}$

(c)

These can be defined recursively:

$$\left| \begin{array}{l} viewed : seq \ Programme \rightarrow seq \ Programme \\ viewed(\langle \rangle) = \langle \rangle \wedge \forall x: Programme \bullet \forall s: seq \ Programme \bullet viewed(\langle x \rangle \frown s) = (\text{if } x.3 = \text{yes then } \langle x \rangle \frown viewed(s)) \end{array} \right|$$

or otherwise (omitted - requires semicolon-separated bindings in set comprehension)

(d)

$$\left| \begin{array}{l} cumulative_total : seq \ Title * Length * Viewed \rightarrow N \\ cumulative_total(\langle \rangle) = 0 \wedge \forall x: Title * Length * Viewed \bullet \forall s: seq \ Title * Length * Viewed \bullet cumulative_total(\langle x \rangle \frown s) = 1 + cumulative_total(s) \end{array} \right|$$

(e)

$(\mu p : \text{ran } hd \mid \forall q: \text{ran } hd \bullet p \neq q \wedge p.2 > q.2 \mid p.1)$

(This, of course, assumes that there is a unique element with this property.)

(f)

(f) Omitted - requires semicolon-separated bindings in nested set comprehension

(g)

axdef

$g : \text{seq}(\text{Title} * \text{Length} * \text{Viewed}) \rightarrow \text{seq}(\text{Title} * \text{Length} * \text{Viewed})$

where

$\forall s : \text{seq } \text{Title} * \text{Length} * \text{Viewed} \bullet g(s) = s \triangleright \{x : \text{ran } s \mid x \neq \text{longest_viewed}(s)\}$

end

Where longest_viewed is defined as

axdef

$\text{longest_viewed} : \text{seq}(\text{Title} * \text{Length} * \text{Viewed})^+ \rightarrow \text{Title} * \text{Length} * \text{Viewed}$

where

$\forall s : \text{seq } \text{Title} * \text{Length} * \text{Viewed} \bullet \text{longest_viewed}(s) = (\mu p : \text{ran } s \bullet p.3 = \text{yes and}$
 $\forall q : \text{ran } s \bullet p \neq q \wedge q.3 = \text{yes} \wedge p.2 > q.2)$

end

This, of course, assumes that there is at least one viewed programme (and one of a unique maximum length).

(h)

$s : \text{seq } \text{Title} * \text{Length} * \text{Viewed} \rightarrow \text{seq } \text{Title} * \text{Length} * \text{Viewed}$	
$\forall x : \text{seq } \text{Title} * \text{Length} * \text{Viewed} \bullet \text{items}(s(x)) = \text{items}(x) \wedge \forall i, j : \text{dom } s(x) \bullet i < j \Rightarrow s(x)(i).2 \geq s(x)(j).2$	

Solution 41

(a)

axdef

$records : Year \mapsto \text{Table}$

where

$\text{dom}(records) = 1993..current$

$\forall y : \text{dom } records \bullet \#records(y) \leq 50$

$\forall y : \text{dom}(records) \mid \forall e : \text{ran } records(y) \bullet year(e.1) = y$

$\forall r : \text{ran}(records) \mid \forall i1, i2 : \text{dom } r \bullet i1 \neq i2 \wedge r(i1).1 = r(i2).1 \Rightarrow r(i1).3 \neq r(i2).3$

end

(b)

(i)

$\{e : \text{Entry} \mid \exists r : \text{ran } records \bullet e \in \text{ran } r \wedge e.3 = 479\}$

ii

$\{e : \text{Entry} \mid \exists r : \text{ran } records \bullet e \in \text{ran } r \wedge e.6 > e.5\}$

iii

$\{e : \text{Entry} \mid \exists r : \text{ran } records \bullet e \in \text{ran } r \wedge e.7 \geq 70\}$

iv

$\{c : \text{Course} \mid \forall r : \text{ran } records \bullet \forall e : \text{ran } r \bullet e.2 = c \Rightarrow e.7 \geq 70\}$

v

$\{y : \text{Year} \mid y \in \text{dom } records \bullet y \mapsto \{l : \text{Lecturer} \mid \#\{c : \text{ran } records(y) \mid c.4 = l\} > 6\}\}$

(c)

axdef

where

$\forall x: Entry \bullet \forall s: seq\ Entry \bullet 479_courses(\langle \rangle) = \langle \rangle$ and $479_courses(\langle x \rangle^s) = if x.3 = 479 then \langle x \rangle^{\cdot 479_courses(s)} else 479_courses(s)$

end

(Blocked by : underscore in identifier – use camelCase for fuzz compatibility)

(d)

$\overline{\quad} \quad \forall x: Entry \bullet \forall s: seq\ Entry \bullet total(\langle \rangle) = 0 \wedge total(\langle x \rangle \frown s) = x.5 + total(s)$

Solution 42

$[Person]$

axdef

$State : P(seq(iseq(Person)))$

where

$\forall s: State \mid \forall i, j: \text{dom } s \bullet i \neq j \wedge \text{ran } s(i) \cap \text{ran } s(j) = \{\}$

end

(b)

axdef

$add : N * Person * State \mapsto State$

where

$\forall n: N \bullet \forall p: Person \bullet \forall s: State \bullet n \in \text{dom } s \wedge p \notin \bigcup \text{ran } s \mid$

add(n, p, s) = s ++ n \mapsto s(n) $\langle p \rangle$

end

(Blocked by: \mapsto operator not implemented)

Solution 43

(a)

(i) $\forall i : \text{dombookings} \mid \forall x, y : \text{bookings}(i) \bullet x \neq y \wedge x.2 \dots x.3 \cap y.2 \dots y.3 = \{\}$

(ii) $\forall i : \text{dombookings} \mid \forall x : \text{bookings}(i) \mid \{x.2, x.3\} \text{ subseq } 1.. \text{max}(i.1)$

(iii) $\forall i : \text{dombookings} \mid \forall b : \text{bookings}(i) \bullet b.2 \leq b.3$

(iv) This is enforced by the constraint for part (i).

(b)

(i) $\{i : \text{dom bookings} \mid i.1 = \text{Banbury} \bullet i.2\}$

(ii) $\{i : \text{dom bookings} \mid i.1 = \text{Banbury} \wedge \exists b : \text{bookings}(i) \bullet 50 \in b.2 \dots b.3\}$

(iii) $r : \text{Room}; s : N \mid \exists i : \text{dom bookings} \bullet i.1 = r \wedge i.2 = s. (r, s)$

(iv) $r : \text{Room} \mid \exists i : \text{dom bookings} \bullet i.1 = r \wedge \# \text{bookings}(i) \geq 10$

Free types and induction

$[N]$

$Tree ::= stalk \mid leaf \langle\langle N \rangle\rangle \mid branch \langle\langle Tree \times Tree \rangle\rangle$

Solution 44

The two cases of the proof are established by equational *reasoning* : *thefirstby*

$$\text{reverse } (\langle \rangle^t) = \text{reverset}[cat.1a] = (\text{reverset})\langle \rangle \text{ [cat.1b]}$$

where cat.1a is $\langle \rangle s = \text{sandcat.1biss} \langle \rangle = s$

and the second by

$$\text{reverse } ((\langle x \rangle^u)^t) = \text{reverse}(\langle x \rangle^{\langle u^t \rangle})[cat.2]$$

$$= \text{reverse } (u^t) \langle x \rangle \text{ [reverse.2]}$$

$$= (\text{reverse } t^r \text{everseu}) \langle x \rangle \text{ [anti-distributive]}$$

$$= \text{reverse } t^r (\text{reverseu} \langle x \rangle) \text{ [cat.2]}$$

$$= \text{reverse } t^r \text{everse}(\langle x \rangle^u) \text{ [reverse.2]}$$

Solution 45

The base case:

$$\text{reverse } (\text{reverse } \langle \rangle) = \text{reverse } \langle \rangle \text{ [reverse.1]} = \langle \rangle \text{ [reverse.1]}$$

The inductive step:

$$\begin{aligned}
& \text{reverse } (\text{reverse } (\langle x \rangle^t)) \\
&= \text{reverse } ((\text{reverse } t) \langle x \rangle) [\text{reverse.2}] \\
&= \text{reverse } (\langle x \rangle)^{\text{reverse}(\text{reverse } t)} [\text{anti-distributive}] \\
&= \text{reverse } (\langle x \rangle \langle \rangle)^{\text{reverse}(\text{reverse } t)} [\text{cat.1}] \\
&= ((\text{reverse } \langle \rangle) \langle x \rangle)^{\text{reverse}(\text{reverse } t)} [\text{reverse.2}] \\
&= (\langle \rangle \langle x \rangle)^{\text{reverse}(\text{reverse } t)} [\text{reverse.1}] \\
&= \langle x \rangle^{\text{reverse}(\text{reverse } t)} [\text{cat.1}] \\
&= \langle x \rangle^t [\text{reverse}(\text{reverse } t) = t]
\end{aligned}$$

Solution 46

(a)

$$\text{count} : \text{Tree} \rightarrow N$$

$$\text{count stalk} = 0$$

$$\forall n : N \bullet \text{count}(\text{leaf}(n)) = 1$$

$$\forall t1, t2 : \text{Tree} \bullet \text{count}(\text{branch}(t1, t2)) = \text{count}(t1) + \text{count}(t2)$$

(Blocked by : *recursive freetypes and pattern matching*)

(b)

$$\text{flatten} : \text{Tree} \rightarrow \text{seq } N$$

$$\text{flatten stalk} = \langle \rangle$$

$$\forall n: N \bullet \text{flatten}(\text{leaf}(n)) = \langle n \rangle$$

$$\forall t1, t2: \text{Tree} \bullet \text{flatten}(\text{branch}(t1, t2)) = \text{flatten}(t1^{\text{flatten}})(t2)$$

(Blocked by : *recursive freetypes and pattern matching*)

Solution 47

First, exhibit the induction principle for the free type:

$$\mathbb{P} \text{ stalk and } (\forall n: N \bullet \mathbb{P} \text{ leaf}(n)) \text{ and } (\forall t1, t2: \text{Tree} \bullet \mathbb{P} t1 \wedge \mathbb{P} t2 \Rightarrow \mathbb{P} \text{ branch}(t1, t2))$$

implies $\forall t: \text{Tree} \bullet \mathbb{P} t$

This gives three cases for the proof:

$$(\text{flatten stalk}) = \langle \rangle [\text{flatten}] = 0 [] = \text{count stalk} [\text{count}]$$

(Remaining cases omitted - require equational reasoning with recursive functions)

Supplementary material : assignment practice

Solution 48

$$[\text{SongId}, \text{UserId}, \text{PlaylistId}, \text{Playlist}]$$

$$\frac{\text{songs} : \mathbb{F} \text{ SongId} \text{ users} : \mathbb{F} \text{ UserId} \text{ playlists} : \text{PlaylistId} \rightarrow \text{Playlist} \text{ playlistOwner} : \text{PlaylistId} \rightarrow \text{UserId} \text{ playlistOwner} : \text{PlaylistId} \rightarrow \text{UserId}}{\forall i: \text{dom playlists} \bullet \text{ran playlists}(i)(\text{subseq})(\text{songs}) \text{ dom playlistOwner}(\text{subseq})(\text{dom playlists}) \text{ ran playlistOwner}(i)(\text{subseq})(\text{dom playlists})}$$

Solution 49

$$\frac{\text{hated} : \text{UserId} \rightarrow \mathbb{F} \text{ SongId} \text{ loved} : \text{UserId} \rightarrow \mathbb{F} \text{ SongId}}{\text{dom hated}(\text{subseq})(\text{users}) \forall i: \text{dom hated} \bullet \text{hated}(i)(\text{subseq})(\text{songs}) \text{ dom loved}(\text{subseq})(\text{users}) \forall i: \text{dom loved} \bullet \text{loved}(i)(\text{subseq})(\text{songs})}$$

Solution 50

(a)

$$A == \text{users} \setminus \bigcup \text{ran } \text{playlistSubscribers}$$

(b)

$$B == \{ p : \text{dom } \text{playlistSubscribers} \mid \# \text{playlistSubscribers}(p) \geq 100 \}$$

(c)

$$C == \mu u : \text{dom } \text{loved} \bullet \forall v : \text{dom } \text{loved} \bullet u \neq v \wedge \# \text{loved}(u) > \# \text{loved}(v)$$

(d)

$$D == \mu s : \text{songs} \bullet \forall t : \text{songs} \bullet s \neq t \wedge \# \{ u : \text{UserId} \mid s \in \text{loved}(u) \} > \# \{ u : \text{UserId} \mid t \in \text{loved}(u) \}$$

Solution 51

(a)

Let's first define two helper functions:

$$\text{loveHateScore} : \text{SongId}+ \rightarrow N$$

$$\forall i : \text{songs} \mid \{ u : \text{UserId} \mid i \in \text{loved}(u) \} \geq \{ u : \text{UserId} \mid i \in \text{hated}(u) \} \Rightarrow$$

$$\text{loveHateScore}(i) = \{ u : \text{UserId} \mid i \in \text{loved}(u) \} - \{ u : \text{UserId} \mid i \in \text{hated}(u) \}$$

and

$$\forall i : \text{songs} \mid \{ u : \text{UserId} \mid i \in \text{loved}(u) \} < \{ u : \text{UserId} \mid i \in \text{hated}(u) \} \Rightarrow$$

$$\text{loveHateScore}(i) = 0$$

$$\left| \begin{array}{l} \text{playlistCount} : \text{SongId} \leftrightarrow N \\ \hline \forall i : \text{songs} \bullet \text{playlistCount}(i) = \# \{ p : \text{dom } \text{playlist} \mid i \in \text{ran } \text{playlist}(p) \} \end{array} \right|$$

We then have:

$$\frac{\text{length} : \text{SongId} \rightarrow N \text{ popularity} : \text{SongId} \rightarrow N}{\text{dom length}(\text{subseq})(\text{songs}) \text{ dom popularity}(\text{subseq})(\text{songs}) \forall i : \text{songs} \bullet \text{popularity}(i) = \text{loveHateScore}(i) + 1}$$

(b)

mostPopular : *SongId*

$(\exists_1 i : \text{songs} \mid \forall j : \text{songs} \bullet i \neq j \wedge \text{popularity}(i) > \text{popularity}(j)) \Rightarrow$

$\text{mostPopular} = (\mu i : \text{songs} \mid \forall j : \text{songs} \bullet i \neq j \wedge \text{popularity}(i) > \text{popularity}(j))$

and

$\neg \exists_1 i : \text{songs} \bullet \forall j : \text{songs} \bullet i \neq j \wedge \text{popularity}(i) > \text{popularity}(j) \Rightarrow \text{mostPopular} = \text{nullSong}$

(c) $\text{playlistsContainingMostPopularSong} == \{i : \text{dom playlists} \mid \text{mostPopular} \in \text{ran playlists}(i)\}$

Solution 52

(a)

premiumPlays : *seq(Play)* → *seq(Play)*

premiumPlays(⟨⟩) = ⟨⟩

$\forall x : \text{Play}; s : \text{seq}(\text{Play}) \mid$

$\text{premiumPlays}(\langle x \rangle^s) = \langle x \rangle^{\text{premiumPlays}(s) \text{ if } \text{userStatus}(x.2) = \text{premium}}$

$\text{premiumPlays}(s) \text{ if } \text{userStatus}(x.2) = \text{standard}$

(Note: Uses camelCase for fuzz compatibility)

(b)

standardPlays : *seq(Play)* → *seq(Play)*

standardPlays(⟨⟩) = ⟨⟩

$\forall x : Play; s : seq(Play) \mid$

$standardPlays(\langle x \rangle^s) = \langle x \rangle^s standardPlays(s) if userStatus(x.2) = standard$

$standardPlays(s) if userStatus(x.2) = premium$

(Note: Uses camelCase for fuzz compatibility)

(c)

$cumulativeLength : seq(Play) \rightarrow N$

$cumulativeLength(\langle \rangle) = 0$

$\forall x : Play; s : seq(Play) \mid$

$cumulativeLength(\langle x \rangle^s) = length(x.1) + cumulativeLength(s)$

(Note: Uses camelCase for fuzz compatibility)