

Lesson 12: Independence and Conditioning


Meike Niederhausen and Nicky Wakim

2025-11-03

Learning Objectives

1. Identify the formula for joint distributions for independent RVs and conditional distributions (PMFs/PDFs)
2. Find conditional pmf from a joint pmf and check if two RVs are independent.
3. Construct a joint distribution for two independent continuous RVs from their marginal distributions.
4. Calculate conditional probabilities and distributions for continuous RVs.

Where are we?



Learning Objectives

1. Identify the formula for joint distributions for independent RVs and conditional distributions (PMFs/PDFs)
2. Find conditional pmf from a joint pmf and check if two RVs are independent.
3. Construct a joint distribution for two independent continuous RVs from their marginal distributions.
4. Calculate conditional probabilities and distributions for continuous RVs.

How do we represent conditional pmfs/pdfs?

For events:

$$P(B|A) = \frac{P(A \cap B)}{P(A)}$$

$$P(A|B) = \frac{P(A \cap B)}{P(B)}$$

For discrete RVs:

$$p_{X|Y}(x|y) = P(X = x|Y = y) = \frac{p_{X,Y}(x,y)}{p_Y(y)}$$

$$p_{Y|X}(y|x) = P(Y = y|X = x) = \frac{p_{X,Y}(x,y)}{p_X(x)}$$

if denominator is greater than 0 ($p_Y(y) > 0$ or
 $p_X(x) > 0$)

For continuous RVs:

pdf
≠ prob

$$f_{X|Y}(x|y) = \frac{f_{X,Y}(x,y)}{f_Y(y)}$$

$$f_{Y|X}(y|x) = \frac{f_{X,Y}(x,y)}{f_X(x)}$$

if denominator is greater than 0 ($f_Y(y) > 0$ or
 $f_X(x) > 0$)

How do we represent independent RVs in a joint pmf/pdf?

What do we know about independence for events?

For events: If $A \perp B$

marginal
pmf of
 X

$$P(A \cap B) = P(A)P(B)$$

$$P(A|B) = P(A)$$

$$P(A|B) = \frac{P(A \cap B)}{P(B)} = \frac{P(A)P(B)}{P(B)} = P(A)$$

For discrete RVs: If $X \perp Y$

Joint CDF

$$p_{X,Y}(x,y) = p_X(x)p_Y(y)$$

marginal
pmf of
 y

$$\rightarrow F_{X,Y}(x,y) = \underline{F_X(x)} \underline{F_Y(y)}$$

$$p_{X|Y}(x|y) = p_X(x)$$

$$p_{Y|X}(y|x) = p_Y(y)$$

For continuous RVs: If $X \perp Y$

marginal pdf of X

$$f_{X,Y}(x,y) = f_X(x)f_Y(y)$$

marginal pdf of y

$$F_{X,Y}(x,y) = F_X(x)F_Y(y)$$

$$f_{X|Y}(x|y) = f_X(x)$$

$$f_{Y|X}(y|x) = f_Y(y)$$

Remember: our probability rules must hold for these!

For discrete RVs

For a valid joint pmf, we need:

- $0 \leq p_{X,Y}(x,y) \leq 1$ for all x, y
- $\sum_{\{all\} x} \sum_{\{all\} y} p_{X,Y}(x,y) = 1$

For a valid conditional pmf, we need:

- $0 \geq p_{X|Y}(x|y) \leq 1$ for all x, y
- $\sum_{\{all\} x} p_{X|Y}(x|y) = 1$

For continuous RVs

For a valid joint pdf, we need:

- $f_{X,Y}(x,y) \geq 0$ for all x, y
- $\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f_{X,Y}(x,y) dx dy = 1$

For a valid conditional pdf, we need:

- $f_{X|Y}(x|y) \geq 0$ for all x and y
- $\int_{-\infty}^{\infty} f_{X|Y}(x|y) dx = 1$

$$\int_{-\infty}^{\infty} f_{Y|X}(y|x) dy = 1$$

Extra notes

- If X_1, X_2, \dots, X_n are independent

$$P(X_1, X_2, X_3) = P(X_1 = x_1)P(X_2 = x_2) \cdot P(X_3 = x_3)$$

$$p_{X_1, X_2, \dots, X_n}(x_1, x_2, \dots, x_n) = P(X_1 = x_1, X_2 = x_2, \dots, X_n = x_n) = \prod_{i=1}^n p_{X_i}(x_i)$$

$$f_{X_1, X_2, \dots, X_n}(x_1, x_2, \dots, x_n) = \prod_{i=1}^n f_{X_i}(x_i)$$

$$F_{X_1, X_2, \dots, X_n}(x_1, x_2, \dots, x_n) = P(X_1 \leq x_1, X_2 \leq x_2, \dots, X_n \leq x_n) = \prod_{i=1}^n P(X_i \leq x_i) = \prod_{i=1}^n F_{X_i}(x_i)$$

- Don't forget, you can manipulate the conditional density to get the joint:

same goes for discrete RVS.

$$f_{X,Y}(x,y) = \frac{f_{X|Y}(x|y)f_Y(y)}{f_Y(y)f_{X|Y}(x|y)}$$

product of
marginal
CDF's

Learning Objectives

1. Identify the formula for joint distributions for independent RVs and conditional distributions (PMFs/PDFs)
discrete
2. Find conditional pmf from a joint pmf and check if two RVs are independent.
3. Construct a joint distribution for two independent continuous RVs from their marginal distributions.
4. Calculate conditional probabilities and distributions for continuous RVs.

Last class: joint distribution for two discrete random variables (1/2)

Example 1

Let X and Y be two random draws from a box containing balls labelled 1, 2, and 3 without replacement.

7. Find $p_{X|Y}(x|y)$.

joint pmf:

	1	2	3	$P_X(x)$
1	0	$\frac{1}{6}$	$\frac{1}{6}$	$\frac{1}{3}$
2	$\frac{1}{6}$	0	$\frac{1}{6}$	$\frac{1}{3}$
3	$\frac{1}{6}$	$\frac{1}{6}$	0	$\frac{1}{3}$
$P_Y(y)$	$\frac{1}{3}$	$\frac{1}{3}$	$\frac{1}{3}$	

$$P_{X|Y}(x|y) = \frac{P_{X,Y}(x,y)}{P_Y(y)}$$

ex1) $P_{X|Y}(X=1 | Y=1) = \frac{P(X=1, Y=1)}{P(Y=1)}$

$$= \frac{0}{\frac{1}{3}} = 0$$

ex2) $P_{X|Y}(X=2 | Y=3) = \frac{P(X=2, Y=3)}{P(Y=3)} = \frac{\frac{1}{6}}{\frac{1}{3}} = \frac{1}{2}$

$P_{X|Y}(x|y)$ = $\begin{cases} \frac{1}{2} & x \neq y \\ 0 & x = y \end{cases}$

Last class: joint distribution for two discrete random variables (2/2)

Example 1

Let X and Y be two random draws from a box containing balls labelled 1, 2, and 3 without replacement.

8. Are X and Y independent? Why or why not?

Remark:

- To show that X and Y are *not* independent, we just need to find one counter example
- However, to show that they *are* independent, we need to verify this for all possible pairs of x and y

		1	2	3	$P_X(x)$
x	1	0	$\frac{1}{6}$	$\frac{1}{6}$	$\frac{1}{3}$
	2	$\frac{1}{6}$	0	$\frac{1}{6}$	$\frac{1}{3}$
	3	$\frac{1}{6}$	$\frac{1}{6}$	0	$\frac{1}{3}$
$P_{Y X}(y x)$		$\frac{1}{3}$	$\frac{1}{3}$	$\frac{1}{3}$	

if $X \perp Y$:

$$P(X=x, Y=y) =$$

$$\underline{P(X=x)} \underline{P(Y=y)}$$

$$\underline{x=1, y=1} \cdot$$

$$P(X=x, Y=y) = \underline{0}$$

$$P(X=1, Y=1) = 0$$

$$P(\underline{X=1}) = \underline{\frac{1}{3}}$$

$$\neq \frac{1}{3} \cdot \frac{1}{3} = \frac{1}{9}$$

$$P(\underline{Y=1}) = \underline{\frac{1}{3}}$$

$\Rightarrow X$ & Y are
not independent

Learning Objectives

1. Identify the formula for joint distributions for independent RVs and conditional distributions (PMFs/PDFs)
2. Find conditional pmf from a joint pmf and check if two RVs are independent.
3. Construct a joint distribution for two independent continuous RVs from their marginal distributions.
4. Calculate conditional probabilities and distributions for continuous RVs.

Constructing a joint pdf from two independent, continuous RVs

Example 1.1

Let X and Y be independent r.v.'s with $f_X(x) = \frac{1}{2}$, for $0 \leq x \leq 2$ and $f_Y(y) = 3y^2$, for $0 \leq y \leq 1$.

1. Find $\underline{f_{X,Y}(x,y)}$

$$f_{X,Y}(x,y) = f_X(x)f_Y(y)$$

$$= \frac{1}{2} \cdot 3y^2 = \frac{3}{2}y^2$$

$$f_{X,Y}(x,y) = \frac{3}{2}y^2 \quad \text{for}$$

$$\begin{cases} 0 \leq x \leq 2, \\ 0 \leq y \leq 1 \end{cases}$$

↳ when $X \perp Y$

ex) domain w/ x dep on y

$$0 \leq x \leq y/2$$

immediately know $X \not\perp Y$

Probability from joint pdf from two independent, continuous RVs

Example 1.2

Let X and Y be independent rv's with $f_X(x) = \frac{1}{2}$, for $0 \leq x \leq 2$ and $f_Y(y) = 3y^2$, for $0 \leq y \leq 1$.

2. Find

$$\mathbb{P}(0 \leq X \leq 1, 0 \leq Y \leq \frac{1}{2})$$

$$*F_{X,Y}(x,y) = f_X(x)F_Y(y)$$

$$\mathbb{P}(0 \leq X \leq 1, 0 \leq Y \leq \frac{1}{2})$$

$$= \frac{1}{16}$$

$$f_{X,Y}(x,y) = \frac{3}{2}y^2$$

$$\underline{\text{OPT 1}}: \quad P(0 \leq X \leq 1, 0 \leq Y \leq \frac{1}{2}) = \int_0^1 \int_0^{1/2} \frac{3}{2}y^2 dy dx$$

$$\underline{\text{OPT 2}}: \quad P(0 \leq X \leq 1, 0 \leq Y \leq \frac{1}{2}) =$$

$$\begin{aligned} & F(X=1, Y=\frac{1}{2}) - F(X=0, Y=0) \\ &= F(X=1, Y=\frac{1}{2}) \end{aligned}$$

$$\begin{aligned} F &= 0 & P(X \leq 0, Y \leq 0) \\ &= 0 \end{aligned}$$


$$\left\{ \begin{aligned} &= F_X(X=1)F_Y(Y=\frac{1}{2}) \\ &= \left[\int_0^1 \frac{1}{2} dx \right] \left[\int_0^{1/2} \frac{3y^2}{f_Y(y)} dy \right] \\ &= \left(\frac{1}{2}x \Big|_{x=0}^1 \right) \left[y^3 \Big|_{y=0}^{1/2} \right] = \frac{(\frac{1}{2}-0)((\frac{1}{2})^3-0)}{16} \end{aligned} \right.$$

Showing independence from joint pdf

Example 2.1

Let $f_{X,Y}(x,y) = 18x^2y^5$, for $0 \leq x \leq 1, 0 \leq y \leq 1$.

1. Are X and Y independent?



marg of x :
integrate over y

$$f_{X,Y}(x,y) = f_X(x)f_Y(y)$$

$$\begin{aligned} f_X(x) &= \int_{-\infty}^{\infty} f_{X,Y}(x,y) dy = \int_{y=0}^{y=1} 18x^2y^5 dy \\ &= 3x^2y^6 \Big|_{y=0}^{y=1} = 3x^2(1)^6 - 3x(0)^6 \end{aligned}$$

$$\begin{aligned} f_X(x) &= 3x^2 \text{ for } 0 \leq x \leq 1 \\ f_Y(y) &= \int_{x=0}^{x=1} 18x^2y^5 dx = 6x^3y^5 \Big|_{x=0}^{x=1} \\ &= 6y^5 \text{ for } 0 \leq y \leq 1 \end{aligned}$$

$$f_X(x)f_Y(y) = [3x^2][6y^5] = 18x^2y^5$$

$$\begin{aligned} f_{X,Y}(x,y) &= 18x^2y^5 = f_X(x)f_Y(y) \\ \text{for } 0 \leq x \leq 1, 0 \leq y \leq 1 &\Rightarrow X \perp Y \end{aligned}$$

- ① find marginals
- ② mult them together
- ③ look if equal to joint

Finding CDF from two independent RVs

Example 2.2

Let $f_{X,Y}(x,y) = 18x^2y^5$, for $0 \leq x \leq 1, 0 \leq y \leq 1$.

2. Find $F_{X,Y}(x,y)$.


STEPS

- ① given joint pdf
- ② get marg pdf
- ③ int marg pdfs to get CDFs
- ④ mult marg CDFs

$$\textcircled{3} \quad F_X(x) = P(X \leq x) = \int_0^x f_X(s) ds = \int_0^x 3s^2 ds \\ = s^3 \Big|_{s=0}^{s=x} = x^3 \text{ for } 0 \leq x \leq 1$$

$$F_Y(y) = P(Y \leq y) = \int_0^y f_Y(t) dt = \int_0^y 6t^5 dt \\ = \dots = y^6 \text{ for } 0 \leq y \leq 1$$

$$\textcircled{4} \quad F_{X,Y}(x,y) = F_X(x)F_Y(y)$$



$$F_{X,Y}(x,y) = \begin{cases} 0 & x < 0 \text{ OR } y < 0 \\ x^3 y^6 & 0 \leq x \leq 1 \text{ & } 0 \leq y \leq 1 \\ x^3 & 0 \leq x \leq 1 \text{ & } y > 1 \\ y^6 & 0 \leq y \leq 1 \text{ & } x > 1 \\ 1 & x > 1 \text{ & } y > 1 \end{cases}$$

Showing independence from joint pdf

Do this problem at home for extra practice. The solution is available in Meike's video!

Example 3

Let $f_{X,Y}(x,y) = 2e^{-(x+y)}$, for $0 \leq x \leq y$. Are X and Y independent?

Final statement on independence

1. If $f_{X,Y}(x,y) = \underline{g(x)h(y)}$, where $g(x)$ and $h(y)$ are pdf's, then X and Y are independent.

- The domain of the joint pdf needs to be independent as well!!

transformations also independent

2. If $F_{X,Y}(x,y) = \underline{G(x)H(y)}$, where $G(x)$ and $H(y)$ are cdf's, then \underline{X} and \underline{Y} are independent.

- The domain of the joint CDF needs to be independent as well!!

- Make sure that:

- X domain does NOT depend on Y
 - Y domain does NOT depend on X

Learning Objectives

1. Identify the formula for joint distributions for independent RVs and conditional distributions (PMFs/PDFs)
2. Find conditional pmf from a joint pmf and check if two RVs are independent.
3. Construct a joint distribution for two independent continuous RVs from their marginal distributions.
4. Calculate conditional probabilities and distributions for continuous RVs.

Example starting from a joint pdf: first try!

$$P(A|B) = \frac{P(A \cap B)}{P(B)}$$

Example 1.1

Let $f_{X,Y}(x,y) = 5e^{-x-3y}$, for $0 < y < \frac{x}{2}$.

1. Find

$$\mathbb{P}(2 < X < 10 | Y = 4)$$

WAY NOT TO DO THIS:

$$\mathbb{P}(2 < X < 10 | Y = 4) = \frac{\mathbb{P}(2 < X < 10, Y = 4)}{\mathbb{P}(Y = 4)}$$

$$\mathbb{P}(Y = 4) = 0$$



$\mathbb{P}(Y = 4) = 0$
for a cont RV

RIGHT WAY: first find $f_Y(y)$
then $f_{X|Y}(x|y) \rightarrow \frac{f_{X,Y}(x,y)}{f_Y(y)}$
then we can consider
intervals & probability

Example starting from a joint pdf: second try! (1/2)

$$e^{x+y} = e^x e^y$$

Example 1.1

Let $f_{X,Y}(x,y) = 5e^{-x-3y}$, for $0 < y < \frac{x}{2}$.

1. Find

$$\mathbb{P}(2 < X < 10 | Y = 4)$$

$$f_{X|Y}(x|y=4) = \frac{e^{-x} e^8}{\text{red plane}} \quad 4 < \frac{x}{2} \quad x > 8$$

to get green plane: $\mathbb{P}(2 < X < 10 | Y = 4)$

when $y=4$
 x must be
 > 8

$$= \int_8^{10} e^8 e^{-x} dx = \dots = 1 - e^{-2}$$

$$f_{X|Y}(x|y) = \frac{f_{X,Y}(x,y)}{f_Y(y)}$$

$$= \frac{5e^{-x-3y}}{5e^{-5y}} \quad \begin{cases} 0 < y < \frac{x}{2} \\ y > 0 \end{cases}$$

$$= e^{-x} e^{-3y} - (-5y)$$

$$f_{X|Y}(x|y) = e^{-x} e^{2y} \quad \text{for } 0 < y < \frac{x}{2}$$

$$f_Y(y) = \int_{2y}^{\infty} 5e^{-x-3y} dx$$

bounds of x :

$$y < \frac{x}{2}$$


$$2y < x < \infty$$

$$= \int_{2y}^{\infty} 5e^{-x} e^{-3y} dx$$

$$= 5e^{-3y} \int_{2y}^{\infty} e^{-x} dx$$

$$= \dots = 5e^{-5y}$$

$$f_{X|Y}(x|y) \quad y > 0$$



Example starting from a joint pdf: second try! (2/2)

$$\begin{aligned} \text{why not } & F_{X,Y}(x=10, y=4) - F_{X,Y}(x=2, y=4) \\ = & \int_8^{10} \int_4^4 f_{X,Y}(x,y) dy dx \\ & \hookrightarrow \text{ends up @ 0 again} \end{aligned}$$

when $P(\underline{\quad} \mid X=x)$, must find conditional
pdf ($f_{Y|X}(y|x)$)
first!

Example starting from a joint pdf

Do this problem at home for extra practice. The solution is available in Meike's video!

Example 1.2

Let $f_{X,Y}(x,y) = 5e^{-x-3y}$, for $0 < y < \frac{x}{2}$.

2. Find $\mathbb{P}(X > 20 | Y = 5)$

Finding probability with conditional domain and pdf

Do this problem at home for extra practice. The solution is available in Meike's video!

Example 2

Randomly choose a point X from the interval $[0, 1]$, and given $X = x$, randomly choose a point Y from $[0, x]$. Find $\mathbb{P}(0 < Y < \frac{1}{4})$.