

Is $n=3$ enough? How to approach sample size and power calculations

Jessica Minnier, PhD

Knight Cancer Institute Biostatistics Shared Resource

Oregon Health & Science University

AACR, March 30, 2019

 bit.ly/aacr-power

 [datapointier](#)



KNIGHT
CANCER
Institute

Outline

- Why calculate sample size?
- What are the important components?
- Examples
- Software

Goals of sample size and power calculations

- Design a study that will have enough information about underlying population to reject a hypothesis with high confidence.
- Calculate the number of sampling units (e.g. people, animals) you need to estimate statistics with a certain level of precision.

Design your study

Step 1: State your research hypothesis

- Define your:
 - Population
 - Outcome variables/measurements
 - Predictor variables (i.e. treatment, age, genetic mutation)
- Be specific!
- Example: Among *women* (population you sample from), the *BRCA1* mutation (predictor) is associated with an increased risk of developing *breast cancer* (outcome).
- **Question:** How many women do we need to sample/study to determine that BRCA1 is associated with breast cancer?

Your hypothesis and design inform your analysis method.

Step 2: Choose your analysis and test(s)

- You can't calculate sample size without knowing which test and model you will use.
- How will you measure your outcome? Continuous? Categorical? Binary (yes/no)?
 - choose outcomes with high sensitivity and low measurement error
- How many groups/experimental conditions/predictors?
 - the more you have the more samples you will need
- What test? t-test? Linear regression? Random effects model? Chi-square test?

Calculate sample size based on analysis method you will use.

Calculate power and sample size

Need to know (/tell your statistician!):

- Overall design (outcome, endpoint, hypothesis)
- Size/magnitude of effect of interest
 - What do you *hope* to detect
- Variability of measurements
 - Precision of your measurement, biological variability within population
- Level of type I error (false positive rate, significance level, α)
- Level of power (true positive rate)
- Other design details (number of groups, clustering, repeated measures)

Components of Sample Size

Need to know 3 of the 4 to determine the 4th:

Measure	Definition
Effect Size	Magnitude of difference or association; i.e. (difference in means)/(population standard deviation) = $\frac{\mu_1 - \mu_0}{\sigma} = \frac{\delta}{\sigma}$
Sample Size	N
Type I Error / Significance level	α = probability of rejecting null hypothesis when it is true
Power	$1 - \beta$ = 1 - Type II error = probability of rejecting null hypothesis when it is false

Components of Sample Size

Need to know 3 of the 4 to determine the 4th:

Do We Know?	Measure	Definition
??	Effect Size	Magnitude of difference or association; i.e. (difference in means)/(population standard deviation) = $\frac{\mu_1 - \mu_0}{\sigma} = \frac{\delta}{\sigma}$
??	Sample Size	N
0.05, 0.01	Type I Error / Significance level	α = probability of rejecting null hypothesis when it is true
0.9, 0.8	Power	$1 - \beta$ = 1 - Type II error = probability of rejecting null hypothesis when it is false

What is effect size?

- Summarizes the outcome of interest
- Magnitude of difference or association
- Specification depends on study design and statistical model/test

Examples:

- Difference in treatment and control mean outcomes, relative to variance (standard deviation)
- Correlation coefficient of two biomarkers
- Risk ratio of breast cancer comparing BRCA carriers to non-carriers
- Magnitude of regression coefficient

Effect size must be

- pre-specified
- based on what is meaningful biologically or clinically (not statistical significance)
- based on pilot data or literature review if available

Simple example: T-test

Outcome = Continuous measurement, normal distribution

Predictor = Treatment yes/no (treatment vs control group)

Test: two sample T-test, equal variance

Effect Size: difference in means divided by standard deviation of population


$$\frac{\mu_{trt} - \mu_{ctrl}}{\sigma}$$

Null Hypothesis: Difference in means = 0

Alternative Hypothesis: Difference in means \neq 0

"Given a desired effect size, what sample size gives us enough information to reject the null hypothesis with power 90%, type I error 5?"


Underlying data distributions



$n=25$, effect size = 1


Power = 0.93

(Significance based on two sample t-test for difference in means)



$n=25$, effect size = 0.33

Increase standard deviation from 1 to 3, divides effect size by 3
Power = 0.21



To detect an effect size of 0.33 with power = 0.9 and type I error = 0.05, what sample size would we need? n=194 in each group!

```
power.t.test(delta = 0.33, sd = 1, sig.level = 0.05, power = 0.9)
```

Two-sample t test power calculation


```
      n = 193.9392
delta = 0.33
  sd = 1
sig.level = 0.05
  power = 0.9
alternative = two.sided
```

NOTE: n is number in *each* group

$n=10$, effect size = 1


Decrease sample size

Power = 0.56



$n=3$, effect size = 1

Decrease sample size even more
Power = 0.16



$n = 3$, power = 0.9, effect size = ?

In the R output below, the effect size is $\text{delta}/\text{sd} = 3.59/1 = 3.59$.


```
power.t.test(n=3, sd=1, sig.level=0.05, power=0.9)
```

Two-sample t test power calculation

```
      n = 3
  delta = 3.589209
     sd = 1
sig.level = 0.05
   power = 0.9
alternative = two.sided
```

NOTE: n is number in *each* group

Underlying data distributions



Other reasons to calculate sample size

Precision of statistics

- Sample sizes can also be calculated for a specific maximum width in confidence interval around an estimate
- i.e. we will estimate the proportion with a 95% confidence interval of width 0.1 such as $[0.2, 0.3]$

Prediction models

- Large sample sizes are needed for complex prediction models.
- Stability of prediction model accuracy measures depends on sample size.

Important to remember:

Sample size estimates are **ESTIMATES**.

- based on assumptions that could be incorrect
- based on pilot data that could be a poor sample or too small
- the more you don't know, the more conservative you should be (inflate your n)
- good to provide multiple estimates for a variety of scenarios/effects

Free online software

G*power

(examples of how to use it: <http://www.ats.ucla.edu/stat/gpower/>)

Shiny Dashboard for Sample Size and Power Calculations




Others

- [TrialDesign.org](#)
- [GLIMMPSE](#)
- [CRAB Stat tools](#)
- [The Shiny CRT Calculator: Power and Sample size for Cluster Randomised Trials](#)
- [Cal Poly Stats Dept Apps](#)
- Statistical software such as R, SAS, STATA

Take home message:

Do your research before you do your research!

Thank you!

Contact me:  minnier-[at]-ohsu.edu,  datapointier,  jminnier

Slides available: bit.ly/aacr-power

Slide code available at: github.com/jminnier/talks-etc

References

- Some of this talk adapted from: [David Yanez's Sample Size](#) talk at [OCTRI Research Forum \(OHSU\)](#)
- [Statistical Rules of Thumb, Chapter 2](#)