

# Epstein-Barr Virus (EBV) Reactivation and ME/CFS

## A Comprehensive Research Report

Compiled: May 27, 2026 | For informational purposes only. This is not medical advice.

### SECTION 1: EBV BASICS

#### What Is EBV (HHV-4)?

Epstein-Barr virus (EBV), formally designated Human Herpesvirus 4 (HHV-4), is a member of the Gammaherpesviridae subfamily within the Herpesviridae family. Identified in 1964 by Michael Epstein and Yvonne Barr, EBV is among the most pervasive human viruses ever identified: **approximately 90-95% of the global adult population carries the virus**. Transmission occurs primarily through oral secretions (saliva) — giving rise to its colloquial label as the "kissing disease" — though it can also spread via blood transfusion, organ transplantation, and sexual contact.

The virus preferentially infects **B lymphocytes** (via the CD21/CR2 receptor) and oropharyngeal epithelial cells, and to a lesser extent T cells and NK cells.

#### Primary Infection: Infectious Mononucleosis

When EBV is first acquired — typically in childhood in lower-income settings, but often in adolescence or young adulthood in higher-income settings — it can cause **infectious mononucleosis (IM)**, commonly called "mono." Primary infection in young children is usually asymptomatic or presents as mild illness.

**Classic IM symptoms include:** - Extreme fatigue and malaise - Fever (often 38-40°C) - Severe sore throat and pharyngitis (often with exudate) - Markedly swollen lymph nodes, especially cervical - Splenomegaly (~50% of cases); splenic rupture is a rare but serious complication - Hepatomegaly and transient liver function abnormalities - Skin rash (particularly if ampicillin/amoxicillin is given — a classic diagnostic clue)

**Duration:** Acute IM symptoms typically resolve within 2-4 weeks, but profound fatigue can persist for weeks to several months. Most otherwise healthy individuals recover fully within 2-3 months, though a minority go on to develop prolonged post-infectious fatigue or ME/CFS.

#### Latency: A Lifelong Relationship

Following primary infection, EBV establishes **lifelong latency** in a small pool of resting memory B cells. The virus is never truly cleared — it persists indefinitely as a circular episome in the nucleus, tethered to host chromatin through EBNA-1 protein, replicating passively alongside the host cell. Under normal immune surveillance, cytotoxic T cells (CD8+) and NK cells keep latently infected B cells tightly in check.

## Reactivation: From Latency to Lytic Cycle

Reactivation is the switch from latency back to the **lytic (productive) cycle**, where the virus actively replicates and produces new viral particles. The key molecular trigger is expression of the viral **BZLF1 (ZEBRA/Zta) protein**, which drives lytic gene expression when EBV-infected memory B cells undergo terminal differentiation into plasma cells — a normal immunological event that EBV exploits.

**Triggers of EBV reactivation include:**

- **Psychological stress** — A meta-analysis of 20 immune outcomes identified EBV reactivation as the strongest, most reliable immunological correlate of psychological stress
- **Physical illness** — Concurrent infections (notably SARS-CoV-2, as documented extensively in Long COVID research)
- **Immune suppression** — Transplant recipients, HIV/AIDS patients, patients on biologics or chemotherapy
- **Pregnancy** — Cell-mediated immunity is physiologically modulated during pregnancy
- **Sleep deprivation and overexertion**
- **Nutritional deficiencies** — Zinc, selenium, vitamins D, B12

In immunocompetent individuals, reactivation is typically **asymptomatic or subclinical** — the immune system rapidly suppresses the lytic cycle before significant viral amplification occurs.

## How Reactivation Is Diagnosed: The Antibody Panel

EBV serology is **pattern-based** — no single antibody test is sufficient; the full panel must be interpreted together.

| Antibody        | Biological Significance                    | Timing                                           |
|-----------------|--------------------------------------------|--------------------------------------------------|
| <b>VCA IgM</b>  | Acute primary infection                    | Appears early; undetectable by 4–8 weeks         |
| <b>VCA IgG</b>  | Prior or ongoing EBV infection             | Appears in acute phase; <b>persists for life</b> |
| <b>EBNA IgG</b> | Confirms past (not recent acute) infection | Develops 6–12 weeks after primary infection      |
| <b>EA-D IgG</b> | Active viral replication                   | Appears during acute infection and reactivation  |

**Interpretation Patterns:**

- **Acute primary infection:** VCA IgM+, VCA IgG+, EBNA–, EA-D+/-
- **Past infection (no active replication):** VCA IgM–, VCA IgG+, EBNA+, EA-D–
- **Reactivation:** VCA IgM–, VCA IgG+ (often elevated), EBNA+, **EA-D+** ← key distinguishing marker

**Additional diagnostics:**

- **EBV DNA PCR in blood:** Detects actual viral DNA load; most useful in immunocompromised patients
- **EBV DNA in CSF:** Used when CNS involvement is suspected

## SECTION 2: EBV REACTIVATION SYMPTOMS

### Symptom Profile

When EBV reactivation becomes symptomatic, it can closely resemble primary mononucleosis but is generally **less severe** in immunocompetent individuals:

**Most common:**

- Profound, often disproportionate fatigue and malaise
- Low-grade fever or temperature dysregulation
- Sore throat and pharyngeal irritation
- Swollen lymph nodes (cervical, axillary, inguinal)
- Muscle aches and myalgias
- Headaches
- Cognitive impairment ("brain fog") — difficulty concentrating, word retrieval issues
- Sleep disturbances — unrefreshing sleep despite excessive duration

**Less common but reported:**

- Mild splenomegaly
- Liver function abnormalities
- Skin rashes
- Joint pains
- Abdominal discomfort
- Depression and anxiety

### Duration and Influencing Factors

- **Acute symptomatic phase:** 2–4 weeks (typical)

- **Residual fatigue:** Several additional weeks, commonly up to 2–3 months
- **Prolonged cases:** A minority experience symptoms lasting 6 months or more

**Factors influencing duration and severity:** - Age at primary infection (adolescent/adult onset → more severe initial illness) - Immune competence (medications, concurrent illness, malnutrition, stress) - Genetic factors (specific HLA alleles affect EBV-specific T cell responses) - Sex (women appear more susceptible to persistent post-viral fatigue syndromes) - Baseline sleep quality and nutritional status - Psychological factors (depression and high perceived stress → greater reactivation and slower immune recovery)

### Primary Infection vs. Reactivation: Key Differences

| Feature          | Primary Infection (Mono)                       | Reactivation                                   |
|------------------|------------------------------------------------|------------------------------------------------|
| Typical severity | Often severe, especially in adolescents/adults | Usually mild to subclinical in immunocompetent |
| Lymphadenopathy  | Prominent, often generalized                   | Mild or absent                                 |
| Splenomegaly     | Common (~50%)                                  | Unusual                                        |
| Fever            | Typically high-grade                           | Often low-grade or absent                      |
| VCA IgM          | Positive                                       | Usually negative                               |
| EBNA IgG         | Absent (develops over weeks)                   | Already positive                               |
| EA-D IgG         | May be positive                                | Typically positive during active reactivation  |
| Duration         | 2–4 weeks acute; months for full recovery      | Highly variable; days to months                |

## SECTION 3: EBV AND ME/CFS — THE CONNECTION

### Historical Background

The story of EBV and ME/CFS stretches back to the **1980s**. The term "chronic Epstein-Barr virus (CEBV) syndrome" was used in the mid-1980s following case series suggesting that individuals with unexplained chronic fatigue had elevated EBV antibody titers. Between 1985 and 1988, **five studies** reported increased EBV antibodies in ME/CFS patients.

However, a definitive controlled study failed to show EBV antibody profiles were specific to CFS versus other fatigued or healthy populations, and no study could link antibody titers to symptom severity — a key requirement for establishing causation. The CDC formally concluded in the late 1980s that "CFS" should not be equated with CEBV syndrome.

Interest declined but never disappeared. Improved immunological tools and a deeper understanding of post-viral syndromes led to a **significant resurgence** in EBV-ME/CFS research since ~2010, accelerating dramatically post-COVID.

### Prevalence of EBV Evidence in ME/CFS

Multiple lines of evidence using different methodologies converge on a significant EBV-associated subset of ME/CFS patients:

- Approximately **30% of ME/CFS cases associated with a recognized infectious onset** are attributed to EBV-triggered infectious mononucleosis
- **9–13% of individuals who develop IM** meet ME/CFS criteria at 6 months post-infection (Hickie et al., White et al., Katz et al.)

- A study assessing latent/persistent EBV found it in **40% of ME/CFS patients vs. 0% of controls**
- An EBV-induced gene subtype (EBI2 upregulation) may account for **38-55% of ME/CFS cases**
- **Up to 75% of ME/CFS patients have reduced levels of B cells** specifically tasked with keeping EBV in check
- Shikova et al. (2020, Journal of Medical Virology): **EBV DNA found in 24.1% of ME/CFS plasma samples** vs. 4% in healthy controls (p = 0.0027)

These figures indicate EBV is a **significant but not universal** factor — pointing toward an EBV-associated subtype rather than EBV being the sole cause.

## Proposed Biological Mechanisms

### 1. Abortive Lytic Reactivation and dUTPase-Mediated Immune Dysregulation

Rather than full reactivation, key research by **Maria Ariza and colleagues at Ohio State University** proposes that EBV is most dangerous during **partial/abortive lytic reactivation** — expressing early lytic proteins like **dUTPase** without producing complete virions.

EBV dUTPase functions as a **pathogen-associated molecular pattern (PAMP)** protein, activating NF-κB through TLR2/MyD88-dependent signaling and inducing neuroinflammatory mediators. A landmark 2022 JCI Insight study found significant elevations in **activin A and IL-21** in ME/CFS cohorts, correlating with seropositivity to EBV and HHV-6 dUTPase antibodies — pointing toward abnormal germinal center and extrafollicular B cell activity as a mechanistic pathway.

### 2. Immune Dysregulation and T/NK Cell Exhaustion

Chronic EBV infection drives persistent antigenic stimulation, leading to upregulation of immune checkpoint molecules — **PD-1, LAG-3, TIM-3, CTLA-4** — hallmarks of **immune exhaustion**. Key abnormalities documented in ME/CFS include:

- Reduced CD8+ T cell counts in early-stage disease
- Impaired NK cell cytotoxicity — particularly reduced **perforin secretion**
- Expanded populations of exhausted CD28– CD8+ T cells and CD56dim NK cells
- Deficient EBV-specific cytotoxic T lymphocyte (CTL) responses
- A 2023 JCI Insight study documented overlapping immune exhaustion patterns in ME/CFS and Long COVID

This framework explains why EBV may persist in a smoldering, partially active state — not fully reactivating (which robust immunity would suppress), but not truly latent either.

### 3. Mitochondrial Dysfunction and Energy Metabolism

EBV infection has been shown to disrupt **mitochondrial quality control** by: - Reducing autophagy (cellular housekeeping removing damaged organelles) - Creating a feedback loop: impaired autophagy → dysfunctional mitochondria → impaired cellular repair

Dysfunctional mitochondria cannot maintain adequate ATP production during physiological stress, producing the characteristic clinical picture of **post-exertional malaise (PEM)** — the hallmark of ME/CFS. A 2024 Dutch study examining skeletal muscle biopsies found severe muscle tissue damage, macrophage infiltration, a sharp drop in amino acid levels, and **amyloid-containing deposits** in ME/CFS patients after exercise.

### 4. Neuroinflammation

PET scanning studies have found evidence of **microglial activation** in ME/CFS patients, particularly in the frontal cortex, insula, thalamus, and limbic areas. EBV contributes through: - Direct neurotropism in some cases - Peripheral cytokine production crossing the blood-brain barrier - EBV dUTPase specifically inducing neuroinflammatory mediators

## 5. Autoimmunity via Molecular Mimicry

EBV EBNA-1 protein contains sequences that cross-react with human self-antigens, forming a basis for **molecular mimicry** — where EBV-directed antibodies inadvertently target host tissues. In ME/CFS, autoantibodies targeting beta-2 adrenergic receptors and muscarinic receptors have been documented.

## 6. HPA Axis and Autonomic Nervous System Dysregulation

EBV-driven immune dysregulation affects **hypothalamic-pituitary-adrenal (HPA) axis** function and autonomic regulation. ME/CFS patients frequently have attenuated cortisol responses, orthostatic intolerance, POTS, and autonomic dysfunction — partially explained by neuroinflammation and central sensitization following viral infection.

### Key Research Studies

| Study                                        | Key Finding                                                                                                                     |
|----------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------|
| <b>Hickie et al., 2006</b> (Lancet)          | Prospective: 11% of acute IM patients met ME/CFS criteria at 6 months                                                           |
| <b>Katz et al., 2009</b>                     | 13% of college students met ME/CFS criteria 6 months after IM; 4% at 24 months                                                  |
| <b>Lerner et al., 2010</b>                   | 36-month valacyclovir follow-up: near-normal return to life in EBV-subset CFS (p=0.0001)                                        |
| <b>Montoya et al., 2013</b> (J Med Virology) | RCT: valganciclovir showed significant improvements in mental fatigue, FSS score, cognitive function in EBV/HHV-6 ME/CFS subset |
| <b>Shikova et al., 2020</b> (J Med Virology) | EBV DNA in 24.1% ME/CFS plasma vs. 4% controls (p=0.0027)                                                                       |
| <b>Ariza et al., 2022</b> (JCI Insight)      | EBV/HHV-6 dUTPases drive abnormal germinal center activity; elevated activin A and IL-21 in ME/CFS                              |

### Cause, Trigger, or Perpetuating Factor?

The most accurate current formulation: **EBV frequently triggers ME/CFS (in the right host context), and may perpetuate it in a biologically distinct subset, while not being the universal or sole cause.**

- **EBV as sole cause:** Largely rejected — EBV is nearly universal (95% seroprevalence), yet only a small fraction of IM patients develop ME/CFS
- **EBV as a trigger:** Well-supported — EBV-associated IM is a **proven risk factor** for ME/CFS onset
- **EBV as a perpetuating factor:** Increasingly supported by newer research — chronic partial EBV activity may sustain ongoing immune activation, neuroinflammation, and metabolic disruption

### The Role of Other Herpesviruses: HHV-6, CMV, and Others

**HHV-6 (Human Herpesvirus 6):** - A systematic review and meta-analysis found HHV-6 infection significantly associated with elevated CFS risk - HHV-6 integrates into the human genome chromosomally in ~1% of the population (ciHHV-6), creating unique susceptibility - In the Ariza dUTPase research, HHV-6A dUTPase was found to drive the same aberrant immune activation as EBV dUTPase, suggesting **synergistic pathogenic mechanisms** - Patients with dual EBV + HHV-6 seropositivity showed the greatest benefit from antiviral treatment in the Montoya studies

**The "Multiple-Hit" Model:** One or more herpesviruses (most commonly EBV, often with HHV-6) may act in combination to trigger and perpetuate ME/CFS, particularly in individuals with underlying genetic vulnerabilities in immune control. A 2012 study found **active HHV-6, HHV-7, and Parvovirus B19**

**coinfection in 64.8% of 108 ME/CFS patients**, correlating with more frequent symptoms and elevated proinflammatory cytokines.

## SECTION 4: LONG COVID CONNECTION

### EBV Reactivation as a Feature of Long COVID

SARS-CoV-2 infection is a potent trigger for EBV reactivation, via: - Profound suppression of T cell and NK cell activity during acute COVID-19 - COVID-19-associated lymphopenia reducing B cell surveillance - Direct CD4+ T cell dysregulation impairing EBV-specific surveillance

#### Key Research Findings:

**Gold et al. (2021, Pathogens):** In a cohort of 185 patients with prior SARS-CoV-2 infection, 30.3% reported Long COVID symptoms. Among Long COVID patients, **66.7% were positive for EBV reactivation**, compared to only **10% of COVID-19 patients who recovered fully** — a statistically significant difference.

**Zubchenko et al.:** EBV reactivation (by PCR) was found in **42.6% of Long COVID patients**.

**Stein et al. (294-patient cohort):** Long COVID symptoms including fatigue and neurocognitive dysfunction were significantly associated with **serological evidence of recent EBV reactivation** but not ongoing EBV viremia — suggesting that the immune response to the reactivation event, rather than active viral replication per se, may be the pathological driver.

### EBV's Place in the Broader Long COVID Framework

EBV reactivation is not a prerequisite for Long COVID but is a **significant contributor** to the post-COVID fatiguing illness syndrome. Other contributing mechanisms include: - SARS-CoV-2 viral persistence (residual antigen in tissues) - Autoantibody formation - Microbiome disruption (gut dysbiosis) - Endothelial dysfunction and microvascular damage - Mast cell activation syndrome

The symptom overlap between Long COVID, post-COVID ME/CFS, and classic ME/CFS is extensive — Anthony Fauci and other researchers have noted that Long COVID may be producing ME/CFS at a large scale.

## SECTION 5: MANAGEMENT AND TREATMENT

There is no FDA-approved cure for either EBV reactivation in the context of ME/CFS, or for ME/CFS itself. Management is **multimodal, individualized, and primarily symptom-focused**.

### Antiviral Medications

**Valacyclovir (Valtrex):** - Inhibits viral DNA polymerase during the lytic phase - **Lerner et al. (36-month RCT):** EBV-subset CFS patients returned to near-normal life ( $p=0.0001$ ) — among the most cited positive antiviral results in ME/CFS - **Straus et al. (1988):** Earlier trial with acyclovir in unselected CFS cohort showed no benefit — underscoring the importance of patient selection - **Key caveat:** Minimal activity against **latent EBV**; most likely to help patients with demonstrable lytic reactivation

**Valganciclovir (Valcyte):** - Broader herpesvirus coverage, including HHV-6 and CMV in addition to EBV - **Montoya et al. RCT (2013):** 30 CFS patients with elevated EBV + HHV-6 titers; 6 months of valganciclovir showed significant improvements in mental fatigue ( $p=0.039$ ), Fatigue Severity Scale ( $p=0.006$ ), and cognitive function ( $p=0.025$ ) - **Montoya observational cohort (106 patients):** 75% responded to long-term valacyclovir/valganciclovir treatment - **Important limitations:** Serious adverse effects including bone

marrow suppression (neutropenia, thrombocytopenia); requires regular CBC monitoring; high cost; small trial sizes

**Overall evidence quality:** Promising but inconclusive. Antivirals appear most likely to benefit the **EBV/HHV-6-seropositive subset** with demonstrable markers of active viral replication. Larger, well-powered RCTs with rigorous patient stratification are urgently needed.

---

## Immune Support and Immunomodulatory Approaches

**Low Dose Naltrexone (LDN):** At doses of 0.1–7 mg, LDN acts primarily as a **microglial inhibitor and immune modulator**: - Binds TLR4 and reduces central and peripheral neuroinflammation - Stimulates endorphin production via transient opioid receptor blockade - A 2025 study confirmed LDN restored TRPM3 channel function in NK cells from Long COVID patients (which are dysfunctional in ME/CFS)

Evidence: - Retrospective analysis of 218 ME/CFS patients treated with LDN: **73.9% positive response rate**, with improved physical and cognitive performance - At least three ongoing clinical trials of LDN in Long COVID / post-COVID ME/CFS

**BC007:** A novel drug designed to **neutralize pathogenic autoantibodies** against G protein-coupled receptors (beta-adrenergic and muscarinic receptors). A small trial found statistically significant improvements in fatigue following a single infusion. Larger RCTs advancing — one of the most promising new immunological approaches.

**Rituximab:** Early Norwegian open-label trial (Fluge & Mella, 2011) reported dramatic improvements in ME/CFS. However, the large **RituxME RCT (2019) showed no significant benefit over placebo**, suggesting ME/CFS is not primarily a straightforward B cell-driven autoimmune condition in the broad unselected cohort.

**Rintatolimod (Ampligen):** Experimental double-stranded RNA immune modulator (TLR3 agonist) stimulating innate antiviral immunity. Approved for ME/CFS only in Argentina. Has shown some efficacy in trials but failed to gain FDA approval. Under investigation for Long COVID.

---

## ME/CFS-Specific Management: Pacing (Most Critical Intervention)

**Pacing and Energy Envelope Management** is the **cornerstone of evidence-based ME/CFS management** and the most universally recommended intervention.

The core principle: maintain activity levels within the "energy envelope" — never pushing to the point of triggering post-exertional malaise (PEM).

**Heart Rate Monitoring:** Because ME/CFS patients switch from aerobic to anaerobic metabolism at abnormally low heart rates, heart rate monitoring provides an objective real-time tool for avoiding the anaerobic threshold. This threshold can be determined via a 2-day cardiopulmonary exercise test (CPET; Workwell Foundation). A common safe ceiling estimate:  **$(220 - \text{Age}) \times 0.55$** .

Exceeding the anaerobic threshold in ME/CFS does not build fitness — it triggers pathological PEM, potentially causing cumulative biological damage and permanent functional decline.

---

## Graded Exercise Therapy (GET) — What NOT to Do

Graded Exercise Therapy has been conclusively shown to be **harmful** for ME/CFS patients. Following extensive review:

- The **CDC removed GET recommendations** for ME/CFS in 2017
- **NICE updated its guidelines (October 2021)**, explicitly stating GET should not be used and is potentially harmful
- The PACE trial's widespread methodological flaws (post-hoc outcome measure changes, weakened recovery criteria, unblinded outcomes) have been extensively documented



## Symptom Management Strategies

**Orthostatic Intolerance / POTS:** - Increased fluid and salt intake - Compression garments - Pharmacological options: fludrocortisone, midodrine, beta-blockers, ivabradine

**Sleep:** - Sleep hygiene measures; polysomnography to exclude sleep apnea - Pharmacological support if needed: low-dose tricyclics, melatonin, trazodone

## Lifestyle Interventions

**Stress reduction:** Given that psychological stress is one of the most potent EBV reactivation triggers, **stress reduction is not optional but therapeutic**. Mind-body approaches adapted to avoid exertion-triggering PEM: gentle mindfulness meditation, diaphragmatic breathing, cognitive reframing.

**Nutrition and Supplementation** (evidence is limited but reasonable): - **CoQ10:** Evidence for mitochondrial support; 200 mg/day may reduce fatigue; some studies show benefit in NADH-CoQ10 combinations - **Magnesium:** Well-documented deficiency in ME/CFS; cofactor in all steps of ATP synthesis; supplementation has shown benefit - **Vitamin B12:** Anecdotal clinical reports of benefit even in the absence of frank deficiency; may relate to neurological support - **NADH:** Some studies show modest fatigue reduction - Anti-inflammatory dietary patterns (Mediterranean diet) are sensible but not yet validated in ME/CFS RCTs

## SECTION 6: PROGNOSIS AND RECOVERY

### Overall Prognosis

Prognosis for established ME/CFS is **generally poor but variable**: - Full recovery rate: estimated at only ~5%, **and probably no more than 10%** of adults - Proportion who improve (but do not fully recover): approximately **39.5%** across follow-up studies - A substantial proportion have stable chronic illness with significant functional impairment - A small subset experience progressive deterioration

**Children have dramatically better outcomes:** A 13-year follow-up study of 35 children showed **80% reported full recovery** — likely reflecting greater neurological plasticity and shorter illness duration before appropriate management.

### Post-Infectious Fatigue Syndrome vs. Full ME/CFS

|                           | Post-Infectious Fatigue Syndrome (PIFS) | ME/CFS                                                  |
|---------------------------|-----------------------------------------|---------------------------------------------------------|
| Duration                  | Weeks to several months                 | 6+ months; self-perpetuating                            |
| Prevalence post-infection | 10-30% of significant infections        | ~11% following IM (Hickie et al.)                       |
| Course                    | Typically self-limiting                 | Distinct biological state; not simply extended recovery |
| Prognosis                 | Generally good                          | Poor for full recovery in adults                        |

The **Hickie et al. 2006 prospective study** (tracking IM, Ross River virus, and Coxiella burnetii infections) found approximately **11% in each group** met ME/CFS criteria at 6 months — regardless of the specific infecting organism. This suggests that the **host's biological response to infection**, rather than the pathogen itself, is the primary determinant of ME/CFS development.



## Predictors of Better vs. Worse Outcomes

**Better outcomes:** - Younger age at onset (children >> adults) - Shorter duration before diagnosis and appropriate management - Milder initial illness severity - No significant pre-existing psychiatric comorbidity - Access to knowledgeable medical care (specifically: avoiding GET)

**Worse outcomes:** - Longer diagnostic delay (average 5+ years for ME/CFS diagnosis) - Continued exposure to the GET push-crash cycle (may cause permanent functional decline) - Older age - Comorbidities: POTS, mast cell activation syndrome, fibromyalgia, sleep apnea - Ongoing uncontrolled stress or immune triggers (including active EBV/HHV-6 reactivation)

## "Management" vs. "Cure"

The current reality: **ME/CFS has no established cure.** The treatment paradigm is:

- **Active management:** Pacing, symptom treatment, avoiding PEM, addressing comorbidities
- **Evidence-supported interventions for subsets:** Antivirals (in EBV/HHV-6-seropositive subsets with active replication), LDN, treatment of POTS and sleep disorders
- **Emerging treatments:** BC007, rintatolimod, immune modulators — in clinical trials
- **Goal:** Achieving the maximum sustainable functional baseline while avoiding the push-crash cycle that drives deterioration

## SECTION 7: CURRENT RESEARCH LANDSCAPE

### Major Research Centers

**United States:** - **Bateman Horne Center (Salt Lake City, UT):** Clinical care, research, and education in ME/CFS; active collaborations with Stanford, Columbia, Cornell, Harvard, CDC, and NIH - **Stanford ME/CFS Collaborative Research Center:** Immunological, genomic, and microbiome biomarkers; primarily funded by Open Medicine Foundation - **Cornell ME/CFS Research Center (Maureen Hanson, PhD):** NIH-funded through 2028; microbiome, gut permeability, metabolomics, immunology - **Columbia University ME/CFS Research Center (Ian Lipkin, MD):** NIH-funded through 2028; virology, immunology, and biomarkers - **Ohio State University (Maria Ariza group):** Herpesvirus dUTPases and immune dysregulation

**International:** - **Open Medicine Foundation (OMF):** Funds international research at Stanford, Harvard, University of Montreal, Australia, and Sweden; BioQuest project (10,000 protein/metabolite profiling of 1,200 samples) underway - **Norwegian ME Biobank and Research Center:** Autoantibodies, rituximab, and biomarkers - **ME Research UK:** UK-based research funding

### NIH Funding and the RECOVER Initiative

**NIH ME/CFS Funding:** - NIH funding for ME/CFS: approximately **\$13 million/year in 2022-2024**, declining to **~\$7.4-9.4 million** by 2025 - For comparison: NIH spends >\$700 million/year on HIV/AIDS and ~\$250 million on multiple sclerosis

**NIH RECOVER Initiative:** - \$1.15 billion invested for Long COVID research, including clinical trials - ME/CFS advocates have called on Congress and NIH to explicitly include ME/CFS within RECOVER's scope, given the massive post-COVID ME/CFS burden - RECOVER studies of antivirals (including Paxlovid for Long COVID) have direct relevance to ME/CFS post-viral mechanisms

## Diagnostic Breakthroughs

**EpiSwitch Blood-Based Diagnostic Test (2025, Journal of Translational Medicine):** Using 3D genomic regulatory profiling: - **96% overall accuracy** distinguishing ME/CFS patients from healthy controls - Sensitivity 92%, specificity 98% on independent validation cohort - Pathways identified: interleukins, TNF $\alpha$ , neuroinflammatory pathways, TLR signaling, JAK/STAT - Potentially the **first clinical-grade diagnostic test** for ME/CFS — ending decades without an objective biomarker

## Most Promising Research Directions

### 1. EBV Vaccine Development

**Moderna mRNA-1189:** Phase I/II trial launched December 2021. Encodes five EBV glycoproteins (gp350, gB, gH, gL, gp42). Interim data: well tolerated, elicited stronger immune response than natural infection. Advancing toward **pivotal trials**.

**mRNA-1195:** Second-generation EBV vaccine candidate in Phase I trials (NCT05831111; estimated completion October 2026), potentially targeting MS prevention as well.

The theoretical implications are profound: if an EBV vaccine prevents primary EBV infection, it could prevent not just IM but potentially **multiple sclerosis**, some lymphomas, and ME/CFS cases triggered by EBV.

**Bjornevik et al. (2022, Science):** Landmark study of 10 million US military personnel found prior EBV infection increased MS risk by **32-fold** without similar risk from other viruses. 801 of 801 MS cases were EBV-seropositive at MS onset. This massively reinvigorated the EBV-MS-ME/CFS nexus.

### 2. Targeted Antivirals and Immune Modulators

- Larger, virologically stratified RCTs of valganciclovir and valacyclovir
- New EBV-specific antivirals targeting novel viral enzymes
- BC007 (autoantibody neutralization) advancing toward larger RCTs
- Rintatolimod being explored in Long COVID/ME/CFS overlap populations

### 3. Precision Medicine / Immunological Phenotyping

- Stratifying ME/CFS patients by virological profile (EBV subset, HHV-6 subset), autoantibody profile, immune exhaustion markers, and metabolic phenotype
- T cell exhaustion reversal (checkpoint inhibitor-inspired approaches)
- JAK/STAT pathway modulation

### 4. Biomarker Development

- OMF's BioQuest project: 10,000 proteins and metabolites in 1,200 samples to define the ME/CFS "molecular signature"
- EpiSwitch blood test representing a potential near-term diagnostic tool
- 2-day CPET remains the gold standard for documenting the characteristic energy metabolism defect

## Key Research Milestones Timeline

| Year      | Milestone                                                                               |
|-----------|-----------------------------------------------------------------------------------------|
| 1985-1988 | 5 studies link elevated EBV antibodies to CFS; "CEBV syndrome" hypothesis proposed      |
| 1988-1994 | Controlled studies fail to confirm specificity; broader CFS case definitions adopted    |
| 2006      | Hickie et al. prospective post-infectious ME/CFS study; 11% IM → ME/CFS rate            |
| 2013      | Montoya valganciclovir RCT — first positive RCT of antiviral in EBV/HHV-6 ME/CFS subset |

| Year      | Milestone                                                                                          |
|-----------|----------------------------------------------------------------------------------------------------|
| 2019      | EBV dUTPase shown to induce neuroinflammatory mediators (Ariza)                                    |
| 2019      | NICE guideline review finds insufficient RCT evidence for GET; observational data for LDN positive |
| 2020      | Shikova: EBV DNA in 24% ME/CFS plasma vs. 4% controls                                              |
| 2021      | Gold et al.: 66.7% Long COVID patients EBV reactivation positive                                   |
| 2021      | NICE guidelines explicitly remove GET from ME/CFS recommendations                                  |
| 2022      | Bjornevik et al. (Science): EBV → 32-fold MS risk increase in 10-million-person military cohort    |
| 2022      | Ariza (JCI Insight): EBV/HHV-6 dUTPases drive abnormal germinal center activity in ME/CFS          |
| 2022      | Moderna mRNA-1189 EBV vaccine Phase I/II trial initiates                                           |
| 2023      | Immune exhaustion patterns documented in ME/CFS and Long COVID (JCI Insight)                       |
| 2024      | Dutch skeletal muscle biopsy study documents amyloid deposits and muscle damage in ME/CFS          |
| 2025      | EpiSwitch: 96%-accurate blood-based ME/CFS diagnostic biomarker test published                     |
| 2025      | LDN shown to restore TRPM3 function in NK cells from Long COVID patients                           |
| 2025-2026 | mRNA-1195 second-generation EBV vaccine Phase I underway                                           |
| 2026      | 7-year prospective follow-up data for post-IM ME/CFS published (Frontiers in Medicine)             |

## CONCLUDING SYNTHESIS

EBV is woven deeply into the story of ME/CFS — as a frequent trigger, likely perpetuating factor in a biologically defined subset, and a window into the broader immunological and neurological dysfunction that characterizes the disease.

The 1980s hypothesis that EBV "causes" CFS was too simple and was appropriately challenged; but the pendulum is now swinging back toward a more sophisticated model in which EBV — particularly through mechanisms of immune exhaustion, abortive lytic reactivation, dUTPase-mediated immune dysregulation, mitochondrial dysfunction, and neuroinflammation — plays a central and mechanistically specific role in a substantial proportion of ME/CFS cases.

The Long COVID pandemic has accelerated this research trajectory enormously. EBV reactivation in 42-67% of Long COVID patients, overlapping almost perfectly with the ME/CFS-like symptom cluster, provides a natural experiment confirming the biological plausibility of the viral trigger → immune dysregulation → post-viral syndrome pathway.

**The most urgent research needs are:** 1. Larger, virologically stratified antiviral RCTs in ME/CFS 2. Advancement of EBV vaccines (mRNA-1189) to prevent the initial trigger 3. Development of validated biomarkers for patient stratification 4. Substantially increased research funding commensurate with the scale of ME/CFS and Long COVID disability

## KEY SOURCES

- Hickie et al. (2006). "Post-infective and chronic fatigue syndromes precipitated by viral and non-viral pathogens." *Lancet*.
- Shikova et al. (2020). "Cytomegalovirus, Epstein-Barr virus, and human herpesvirus-6 infections in patients with myalgic encephalomyelitis/chronic fatigue syndrome." *Journal of Medical Virology*. PMC7687071

- Montoya et al. (2013). "Randomized clinical trial to evaluate the efficacy and safety of valganciclovir in a subset of patients with chronic fatigue syndrome." *Journal of Medical Virology*.
- Ariza et al. (2022). "EBV and HHV-6 dUTPases drive aberrant germinal center activity in ME/CFS." *JCI Insight*. PMC9220958
- Gold et al. (2021). "Patient-reported long-COVID symptoms and EBV reactivation." *Pathogens*. PMC8233978
- Bjornevik et al. (2022). "Longitudinal analysis reveals high prevalence of Epstein-Barr virus associated with multiple sclerosis." *Science*.
- Immune Exhaustion in ME/CFS and Long COVID (2023). *JCI Insight*. DOI:10.1172/jci.insight.183810
- EpiSwitch Diagnostic Biomarker for ME/CFS (2025). *Journal of Translational Medicine*.
- 7-Year Follow-Up Post-IM ME/CFS (2026). *Frontiers in Medicine*. DOI:10.3389/fmed.2026.1676628
- CDC EBV Laboratory Testing: [cdc.gov/epstein-barr/php/laboratories/index.html](https://cdc.gov/epstein-barr/php/laboratories/index.html)
- Workwell Foundation Pacing: [workwellfoundation.org](https://workwellfoundation.org)
- Bateman Horne Center LDN Resource: [batemanhornecenter.org](https://batemanhornecenter.org)
- NICE ME/CFS Guideline (2021): [nice.org.uk/guidance/ng206](https://nice.org.uk/guidance/ng206)
- Open Medicine Foundation: [omf.ngo](https://omf.ngo)
- Moderna mRNA-1189 EBV vaccine: [ClinicalTrials.gov NCT04645147](https://ClinicalTrials.gov/NCT04645147)
- mRNA-1195 Phase I Trial: [ClinicalTrials.gov NCT05831111](https://ClinicalTrials.gov/NCT05831111)
- NIH ME/CFS Funding data: [mecfsscience.org/nih-funding-for-me-cfs-in-2024/](https://mecfsscience.org/nih-funding-for-me-cfs-in-2024/)