

Model Fit

Jacob M. Montgomery

Quantitative Political Methodology

Model Fit

Poster questions?

Road map

Where we have been:

- ▶ Single-variable regression
- ▶ Multivariate regression
- ▶ Regression and causal inference

Road map

Where we have been:

- ▶ Single-variable regression
- ▶ Multivariate regression
- ▶ Regression and causal inference

Today:

- ▶ Review of correlation (r)
- ▶ RMSE and Model fit (r^2)
- ▶ F-tests
- ▶ Multivariate model fit
- ▶ Time for posters

Review of Correlation

Pearson's r

Review of Correlation

Pearson's r (Standardized slope)

Review of Correlation

Pearson's r (Standardized slope)



$$S_Y = \sqrt{\frac{\sum(Y_i - \bar{Y})^2}{n - 1}}$$

Review of Correlation

Pearson's r (Standardized slope)



$$S_Y = \sqrt{\frac{\sum(Y_i - \bar{Y})^2}{n - 1}}$$



$$S_X = \sqrt{\frac{\sum(X_i - \bar{X})^2}{n - 1}}$$

Review of Correlation

Pearson's r (Standardized slope)



$$S_Y = \sqrt{\frac{\sum (Y_i - \bar{Y})^2}{n - 1}}$$



$$S_X = \sqrt{\frac{\sum (X_i - \bar{X})^2}{n - 1}}$$



$$r = \left(\frac{S_X}{S_Y} \right) \hat{\beta}$$

Reminder: These are the main parameters

$$Y = \alpha + X\beta + \epsilon$$

$$\epsilon \sim N(0, \sigma^2)$$

Reminder: These are the main parameters

$$Y = \alpha + X\beta + \epsilon$$

$$\epsilon \sim N(0, \sigma^2)$$

$$\hat{\beta} = \frac{\sum_{i=1}^n ((X_i - \bar{X})(Y_i - \bar{Y}))}{\sum_{i=1}^n (X_i - \bar{X})^2}$$

Reminder: These are the main parameters

$$Y = \alpha + X\beta + \epsilon$$

$$\epsilon \sim N(0, \sigma^2)$$

$$\hat{\beta} = \frac{\sum_{i=1}^n ((X_i - \bar{X})(Y_i - \bar{Y}))}{\sum_{i=1}^n (X_i - \bar{X})^2}$$

$$\hat{\alpha} = \bar{Y} - \hat{\beta}\bar{X}$$

How good is our model?: Thinking about variance

- Unconditional variance: Estimate of total variance in the population

$$S^2 = \hat{\sigma}_Y^2 = \frac{\sum (Y_i - \bar{Y})^2}{n - 1} \Rightarrow S = \hat{\sigma}_Y = \sqrt{\frac{\sum (Y_i - \bar{Y})^2}{n - 1}}$$

How good is our model?: Thinking about variance

- Unconditional variance: Estimate of total variance in the population

$$S^2 = \hat{\sigma}_Y^2 = \frac{\sum (Y_i - \bar{Y})^2}{n - 1} \Rightarrow S = \hat{\sigma}_Y = \sqrt{\frac{\sum (Y_i - \bar{Y})^2}{n - 1}}$$

- Sum of Squared Error: A measure of “spread” around the line

$$SSE = \sum_{i=1}^n (Y_i - \hat{Y}_i)^2 = \sum_{i=1}^n (Y_i - \hat{\alpha} - \hat{\beta}X_i)^2$$

- Conditional Variance: Estimate of variance around line in population


$$\hat{\sigma}^2 = \frac{SSE}{n-2} = \frac{\sum(Y_i - \hat{Y}_i)^2}{n-2} \Rightarrow \hat{\sigma} = \sqrt{\frac{SSE}{n-2}} = \sqrt{\frac{\sum(Y_i - \hat{Y}_i)^2}{n-2}}$$

- Conditional Variance: Estimate of variance around line in population


$$\hat{\sigma}^2 = \frac{SSE}{n-2} = \frac{\sum(Y_i - \hat{Y}_i)^2}{n-2} \Rightarrow \hat{\sigma} = \sqrt{\frac{SSE}{n-2}} = \sqrt{\frac{\sum(Y_i - \hat{Y}_i)^2}{n-2}}$$

- $\hat{\sigma}^2$ is sometimes called “Mean squared error” (MSE) and $\hat{\sigma}$ is “Root mean squared error” (RMSE) or “Residual standard error” (in R) or “Standard error of the estimate” (in SPSS).


Histogram of Y



Regression of X and Y



Histogram of Y when X=2



A really, really good line will have small conditional variance.

Evaluating model fit: Hold onto some basic ideas

- ▶ $\sum(Y_i - \bar{Y})^2$

Evaluating model fit: Hold onto some basic ideas

- ▶ $\sum(Y_i - \bar{Y})^2 = \text{Total Sum of Squares}$
- ▶ Unconditional variance: $S^2 = \frac{\sum(Y_i - \bar{Y})^2}{n-1}$

Evaluating model fit: Hold onto some basic ideas

- ▶ $\sum(Y_i - \bar{Y})^2 = \text{Total Sum of Squares}$
- ▶ Unconditional variance: $S^2 = \frac{\sum(Y_i - \bar{Y})^2}{n-1}$
- ▶ $\sum(Y_i - \hat{Y}_i)^2$

Evaluating model fit: Hold onto some basic ideas

- ▶ $\sum(Y_i - \bar{Y})^2 = \text{Total Sum of Squares}$
- ▶ Unconditional variance: $S^2 = \frac{\sum(Y_i - \bar{Y})^2}{n-1}$
- ▶ $\sum(Y_i - \hat{Y}_i)^2 = \text{Sum of Squared Error}$


Evaluating model fit: Hold onto some basic ideas

- ▶ $\sum(Y_i - \bar{Y})^2 = \text{Total Sum of Squares}$
- ▶ Unconditional variance: $S^2 = \frac{\sum(Y_i - \bar{Y})^2}{n-1}$
- ▶ $\sum(Y_i - \hat{Y}_i)^2 = \text{Sum of Squared Error}$
- ▶ Conditional variance: $\hat{\sigma}^2 = \frac{SSE}{n-2} = \frac{\sum(Y_i - \hat{Y}_i)^2}{n-2}$

Evaluating model fit: Hold onto some basic ideas

- ▶ $\sum(Y_i - \bar{Y})^2 = \text{Total Sum of Squares}$
- ▶ Unconditional variance: $S^2 = \frac{\sum(Y_i - \bar{Y})^2}{n-1}$
- ▶ $\sum(Y_i - \hat{Y}_i)^2 = \text{Sum of Squared Error}$
- ▶ Conditional variance: $\hat{\sigma}^2 = \frac{SSE}{n-2} = \frac{\sum(Y_i - \hat{Y}_i)^2}{n-2}$
- ▶ We are going to say that IF we have a really good model, $\hat{\sigma}^2$ should be “a lot” smaller than S^2 .

Let's go back: Regression between GDP growth and election outcomes



We draw the line that reduces SSE


- Residuals:

$$e_i = (Y_i - \hat{Y}_i) = (y_i - \hat{\alpha} - \hat{\beta}X_i)$$

We draw the line that reduces SSE

► Residuals:

$$e_i = (Y_i - \hat{Y}_i) = (y_i - \hat{\alpha} - \hat{\beta}X_i)$$



So ... how good is your model? r^2

Define some preliminary terms:

- ▶ $TSS = \sum (Y_i - \bar{Y})^2$
- ▶ $SSE = \sum (Y_i - \hat{Y}_i)^2$

So ... how good is your model? r^2

Define some preliminary terms:

- ▶ $TSS = \sum (Y_i - \bar{Y})^2$
- ▶ $SSE = \sum (Y_i - \hat{Y}_i)^2$

Let's define r^2

$$r^2 = \frac{\text{Explained Variance}}{\text{Total Variance}} = \frac{\text{Total Variance} - \text{Unexplained Variance}}{\text{Total Variance}}$$

So ... how good is your model? r^2

Define some preliminary terms:

- ▶ $TSS = \sum (Y_i - \bar{Y})^2$
- ▶ $SSE = \sum (Y_i - \hat{Y}_i)^2$

Let's define r^2

$$r^2 = \frac{\text{Explained Variance}}{\text{Total Variance}} = \frac{\text{Total Variance} - \text{Unexplained Variance}}{\text{Total Variance}}$$

$$r^2 = \frac{TSS - SSE}{TSS}$$

Some notes on r^2

$$r^2 = \frac{TSS - SSE}{TSS}$$

- ▶ SSE=0 (Perfect fit) $r^2 = 1$

Some notes on r^2

$$r^2 = \frac{TSS - SSE}{TSS}$$

- ▶ $SSE=0$ (Perfect fit) $r^2 = 1$
- ▶ $SSE = TSS$ (No fit) $r^2 = 0$

Some notes on r^2

$$r^2 = \frac{TSS - SSE}{TSS}$$

- ▶ SSE=0 (Perfect fit) $r^2 = 1$
- ▶ SSE = TSS (No fit) $r^2 = 0$
- ▶ Does not depend on units of measurement

Some notes on r^2

$$r^2 = \frac{TSS - SSE}{TSS}$$

- ▶ $SSE=0$ (Perfect fit) $r^2 = 1$
- ▶ $SSE = TSS$ (No fit) $r^2 = 0$
- ▶ Does not depend on units of measurement
- ▶ Sometimes called the “coefficient of determination”

Some notes on r^2

$$r^2 = \frac{TSS - SSE}{TSS}$$

- ▶ $SSE=0$ (Perfect fit) $r^2 = 1$
- ▶ $SSE = TSS$ (No fit) $r^2 = 0$
- ▶ Does not depend on units of measurement
- ▶ Sometimes called the “coefficient of determination”
- ▶ It does not penalize for “model complexity.” Often “adjusted R-squared” is used.

Some notes on r^2

$$r^2 = \frac{TSS - SSE}{TSS}$$

- ▶ $SSE=0$ (Perfect fit) $r^2 = 1$
- ▶ $SSE = TSS$ (No fit) $r^2 = 0$
- ▶ Does not depend on units of measurement
- ▶ Sometimes called the “coefficient of determination”
- ▶ It does not penalize for “model complexity.” Often “adjusted R-squared” is used.
- ▶ In R-output this is labeled “Multiple R-squared”

Some notes on r^2

$$r^2 = \frac{TSS - SSE}{TSS}$$

- ▶ SSE=0 (Perfect fit) $r^2 = 1$
- ▶ SSE = TSS (No fit) $r^2 = 0$
- ▶ Does not depend on units of measurement
- ▶ Sometimes called the “coefficient of determination”
- ▶ It does not penalize for “model complexity.” Often “adjusted R-squared” is used.
- ▶ In R-output this is labeled “Multiple R-squared”
- ▶ Why do we use it?
 - ▶ Gives us an overall impression for how well our model is doing.
 - ▶ We can *informally* compare models.

R output

Call:

```
lm(formula = vote ~ q2gdp, data = Abram)
```

Residuals:

Min	1Q	Median	3Q	Max
-6.002	-3.409	0.084	2.078	8.496

Coefficients:

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	49.2560	1.4411	34.179	1.21e-15 ***
q2gdp	0.7549	0.2578	2.928	0.0104 *

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 4.481 on 15 degrees of freedom

Multiple R-squared: 0.3637, Adjusted R-squared: 0.3213

F-statistic: 8.573 on 1 and 15 DF, p-value: 0.01039

Primer on the F-statistic for regression

Before we said that r^2 is intuitively $r^2 = \frac{\text{Explained Variance}}{\text{Total Variance}}$.

Primer on the F-statistic for regression

Before we said that r^2 is intuitively $r^2 = \frac{\text{Explained Variance}}{\text{Total Variance}}$. It makes sense then that $(1 - r^2)$ is the percent of variance we haven't explained.

Primer on the F-statistic for regression

Before we said that r^2 is intuitively $r^2 = \frac{\text{Explained Variance}}{\text{Total Variance}}$. It makes sense then that $(1 - r^2)$ is the percent of variance we haven't explained. It turns out that:

- F-statistic for regression

$$F = \frac{r^2/p}{(1 - r^2)/[n - (p + 1)]}$$

Primer on the F-statistic for regression


Before we said that r^2 is intuitively $r^2 = \frac{\text{Explained Variance}}{\text{Total Variance}}$. It makes sense then that $(1 - r^2)$ is the percent of variance we haven't explained. It turns out that:

- ▶ F-statistic for regression


$$F = \frac{r^2/p}{(1 - r^2)/[n - (p + 1)]}$$

- ▶ Here p is the number of covariates (gdp, incumbent, etc.), and n is the number of observations. This will be distributed according to the F-distribution with $df_1 = p$, and $df_2 = n - (p + 1)$.

Example F-Distributions



And now you understand (almost) everything on a regression table.



Interpreting the F-test

- ▶ Is our model any good?

Interpreting the F-test

- ▶ Is our model any good?
- ▶ This is a *formalized* way of asking whether our model is any good.

Interpreting the F-test

- ▶ Is our model any good?
- ▶ This is a *formalized* way of asking whether our model is any good.
- ▶ We compare the amount of variance explained by the regression to the amount unexplained.

Interpreting the F-test

- ▶ Is our model any good?
- ▶ This is a *formalized* way of asking whether our model is any good.
- ▶ We compare the amount of variance explained by the regression to the amount unexplained.
- ▶ This is essentially a comparison of the following two models:
 - ▶ $H_0 : Y_i = \alpha + \epsilon_i$
 - ▶ $H_a : Y_i = \alpha + X_i\beta + \epsilon_i$

Interpreting the F-test

- ▶ Is our model any good?
- ▶ This is a *formalized* way of asking whether our model is any good.
- ▶ We compare the amount of variance explained by the regression to the amount unexplained.
- ▶ This is essentially a comparison of the following two models:
 - ▶ $H_0 : Y_i = \alpha + \epsilon_i$
 - ▶ $H_a : Y_i = \alpha + X_i\beta + \epsilon_i$
- ▶ This is more useful in multivariate regression:
 - ▶ $H_0 : Y_i = \alpha + \epsilon_i$
 - ▶ $H_a : Y_i = \alpha + X_{i1}\beta_1 + X_{i2}\beta_2 + X_{i3}\beta_3 + \dots + \epsilon_i$

R output

Call:

```
lm(formula = vote ~ q2gdp, data = Abram)
```

Residuals:

Min	1Q	Median	3Q	Max
-6.002	-3.409	0.084	2.078	8.496

Coefficients:

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	49.2560	1.4411	34.179	1.21e-15 ***
q2gdp	0.7549	0.2578	2.928	0.0104 *

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 4.481 on 15 degrees of freedom

Multiple R-squared: 0.3637, Adjusted R-squared: 0.3213

F-statistic: 8.573 on 1 and 15 DF, p-value: 0.01039