

# Effects of grass–endophyte symbiosis and herbivory on population demography across climatic and geographic gradients

Jacob K. Moutouama 0000-0003-1599-1671

<sup>\*1</sup>, Julia Martin<sup>1</sup>, Ulisses Rizo<sup>1</sup>, Malcolm Sherwood<sup>1</sup>, Emily Chong<sup>1</sup>,  
Dajanae Pearson<sup>1</sup>, Alexandra Jimenez Martín<sup>1</sup>, Josh Fowler<sup>2</sup>, Ali Campbell<sup>1</sup>,  
Chris Oxley<sup>1</sup>, Karl Schrader<sup>1</sup>, and Tom E.X. Miller 0000-0003-3208-6067<sup>1</sup>

<sup>1</sup>Program in Ecology and Evolutionary

Biology, Department of BioSciences, Rice University, Houston, TX USA

<sup>2</sup>University of Miami, Department of Biology, Miami, FL USA

**Running header:** Grass-endophyte-herbivory effects across gradients

**Keywords:** demography, range limits

**Submitted to:** *Ecology letters* (Letter)

**Data accessibility statement:** If the paper is accepted, all data and computer scripts supporting the results will be archived in a Zenodo package, with the DOI included at the end of the article. During peer review, our code (Stan and R) is available at <https://github.com/jmoutouama/ELVI-endophyte-density>.

**Conflict of interest statement:** None.

**Authorship statement:**

**Abstract:**

**Main Text:**

**Figures:** 6

**Tables:** X

**References:** X

---

<sup>\*</sup>Corresponding author: [jmoutouama@gmail.com](mailto:jmoutouama@gmail.com)

## Abstract

Interactions between plants and fungi are ubiquitous in nature and have significant effects on plant fitness. However, the extent to which variation in fungal frequency affects plant distribution remains understudied. Using three cool-season grass species, we combined field experiments on plant populations with variation in leaf endophytic fungi and Bayesian statistics to test whether plant-fungi interactions constrain host demography and, consequently, geographic range limits. We found that all vital rates decreased toward the species' range edges, indicating reduced demographic performance in these regions. This decline was more pronounced in individuals hosting endophytic symbionts, suggesting that plant-fungi interactions that are beneficial under benign conditions may become parasitic under stressful conditions. Our results highlight that range expansion may be hindered by the presence of a mutualist.

## Introduction

Plant-microbe symbioses are widespread and ecologically important. These interactions are famously context-dependent, where the direction and strength of the interaction outcome depends on the environment in which it occurs (Bronstein, 1994; Fowler et al., 2023; Hoeksema and Bruna, 2015). Plant-microbe interactions that are beneficial under stressful conditions may become parasitic under benign conditions (Giauque et al., 2019). Under biotic stress (e.g., herbivory), endophyte symbiosis can benefit host plants by facilitating the production of secondary compounds that deter feeding or exert direct toxicity, thereby reducing insect growth, survival, and oviposition (Atala et al., 2022; Bastias et al., 2017; Vega, 2008). Similarly under abiotic stress (e.g., drought), symbionts can increase their host tolerance (Clay and Schardl, 2002). However, in many plant-microbe interactions, host protection is not guaranteed solely by the presence of a symbionts; rather, the density of the symbiont can determine the effectiveness of this protection (Laughton et al., 2014). Higher endophyte densities may lead to increased resource exploitation by the symbiont, potentially imposing costs on the host, such as reduced growth or reproduction (Faeth, 2009).

Context dependence raises the hypothesis that plant-microbe interactions are likely to vary across environmental gradients, from range core to range edge, with significant implications for host range expansion. If microbial symbiosis provides greater benefits under environmental stress, then symbionts could enhance the suitability of range-edge environments, potentially extending host range limits (Allsup et al., 2023; Rudgers et al., 2020). For instance, fungal endophytes improve the survival of *Bromus laevipes* populations in dry conditions, increasing their drought resistance at the range edge and thereby extending the species' geographic range (Afkhami et al., 2014; David et al., 2019). Even if the symbiont does not directly improve host survival, it may still enhance host population growth over time by increasing relative growth and reproduction, potentially offsetting the negative effects of lower survival rates (Yule et al., 2013). Conversely, if microbial symbiosis is costly for the host at range edge, symbionts could constrain host range (Bennett and Groten, 2022; Benning and Moeller, 2021a,b).

Despite growing interest in the ecological and evolutionary roles of symbiosis, most studies focus on the host plant's perspective and rarely manipulate symbiont prevalence to evaluate how environmental variation shapes the symbiont's effects on host demography across geographic ranges. Yet, symbionts promote their own fitness by influencing host life history traits and enhancing resistance to stress (Giauque et al., 2019; Kazenel et al., 2015; Saikkonen et al., 1998). Theory predicts that exclusive vertical transmission fosters mutualism and leads to high symbiont prevalence in host populations (Fine, 1975). Field studies suggest that, under stressful conditions, beneficial symbionts may increase in prevalence and even

reach fixation if their fitness advantages outweigh reproductive costs (Donald et al., 2021). However, populations with lower prevalence may experience higher symbiont loss among offspring (Afkhami and Rudgers, 2008). Thus, overlooking symbiont dynamics could result in inaccurate predictions of host population responses to global change.

Additionally, few studies address the synergistic effects of abiotic and biotic stressors on host-symbiont dynamics across species' ranges. For example, herbivory not only affects the fitness benefits of symbiosis but can also increase endophyte prevalence through enhanced vertical transmission (Agrawal et al., 1999; Gundel et al., 2020). However, these findings are primarily based on greenhouse studies, which limits our understanding of how herbivory shapes natural population dynamics, especially under variable field conditions. If the ecological context in which herbivory occurs alters plant fitness, then interactions between herbivory and abiotic stress (e.g., drought) may substantially influence population dynamics and species range limitation.

Working across a precipitation gradient in the south-central US, we investigated how the demographic effects of endophyte symbiosis vary from core to edge of the host range. To answer that question, we studied the symbiotic association between cool-season grass species (*Agrostis hyemalis*, *Elymus virginicus* and *Poa autumnalis*) and their vertically transmitted fungal symbiont *Epichloë* spp. Our experiment was design to test the following hypotheses:

1. Because endophytes often produce defensive compounds (e.g., alkaloids), plants with endophytes (endophyte-infected plants) will experience less herbivore damage and maintain higher vital rate compared to plants without endophytes (endophyte-free plants) when grazed.
2. Because endophytes can improve drought tolerance (e.g., by enhancing water use efficiency or root growth), plants with endophytes will outperform plants without endophytes more in dry conditions than in wet conditions.
3. The combined stresses of herbivory and low precipitation will create conditions where the mutualistic benefits of endophytes are most pronounced, giving plants with endophytes a strong performance advantage over plants without endophytes.
4. The combined stresses of herbivory and low precipitation will amplify the physiological demands on the host plant, such that the additional resource costs of maintaining endophyte associations may reduce overall plant performance resulting in endophyte-infected plants performing worse than uninfected plants under extreme stress conditions.

## Materials and methods

### Study system


Our study was conducted across seven sites along a geographic and climatic gradient spanning Louisiana and Texas in the United States (Fig. 1; Fig.S-1). At each site, we established eight plots (1.5 m × 1.5 m), with each plot containing 18 individuals of a single species. We used three cool-season perennial grasses native to woodland and prairie habitats of eastern North America (Shaw, 2011): *Agrostis hyemalis*, *Elymus virginicus*, and *Poa autumnalis* (subfamily Pooideae). These species are capable of both self-pollination and outcrossing (Church, 1958). They typically host seed-transmitted fungal endophytes. Seeds were collected in summer–fall 2006 from naturally symbiotic populations (source populations) of the three focal host species. These seeds were either heat-treated to eliminate symbionts or left untreated, thereby generating symbiont-free ( $E^-$ ) and symbiotic ( $E^+$ ) plants from the same genetic lineages as the source populations.

Across all sites, each plot was randomly assigned one of four initial endophyte frequency treatments (80%, 60%, 40%, or 20%) and one of two herbivory treatments (herbivore exclusion or herbivore access). We ensured that all plots received comparable quantities of source plant material. This experimental design was informed by natural patterns of endophyte prevalence observed in Texas, where the proportion of symbiotic individuals has been reported as 86.55% in *A. hyemalis* (ranging from 77.16% to 93.5%) (Donald et al., 2021), 53% in *E. virginicus* (ranging from 10% to 100%) (Sneck et al., 2017), and 96% in *P. autumnalis* (Rudgers et al., 2009). Additionally, fungal genotyping has confirmed the presence of biosynthetic pathways for secondary metabolites such as peramine, loline, and ergot alkaloids, which may enhance host resistance to drought and herbivory (Beaudry, 1951). Full details of the study design are provided in the supplementary materials.

### Demographic data collection

We collected demographic data including survival, growth, and reproduction in May and June of 2023 and 2024, which coincided with the flowering season of *Agrostis hyemalis*, *Elymus virginicus*, and *Poa autumnalis*. For each individual, plant survival was recorded as a binary variable (alive or dead), and plant size was measured as the number of living tillers, indicated by the presence of green coloration. We recorded the number of inflorescences per plant and counted the number of spikelets on up to three inflorescences from three reproducing plants. Spikelet counts were limited to three reproducing tillers per plot due to the time-consuming

111 nature of this measurement. We used the number of spikelets from these three tillers to  
 112 estimate the average number of spikelets per plant.



**Figure 1:** Distribution of common garden sites along the longitudinal aridity gradient in the central and southern Great Plains for: A) *Agrostis hyemalis* (AGHY), B) *Elymus virginicus* (ELVI), and C) *Poa autumnalis* (POAU). Red dots indicate the locations of source populations, while grey dots represent GBIF occurrence records for each species within the study area. D) Relationship between the distance from the range center and longitude for each species.

## 113 Models building and models selection

114 To assess how stress associated with aridity affects plant–fungal interactions, we developed  
 115 four candidate models for each vital rate (survival, growth, and fertility). Each vital rate was  
 116 modeled using a grand mean intercept ( $\beta_0$ ), slopes for variation in each covariate ( $\beta_1$ ,  $\beta_2$  and  
 117  $\beta_3$ ), and interaction terms between covariates ( $\beta_4$ ,  $\beta_5$  and  $\beta_6$ ). Each model included normally

distributed random effects to account for site-to-site variation ( $\phi \sim N(0, \sigma_{site})$ ), plot-to-plot variation ( $\rho \sim N(0, \sigma_{plot})$ ), and source-to-source variation associated with the provenance of transplants used in the common garden ( $\omega \sim N(0, \sigma_{source})$ ) (Eq. 1).

$$\begin{aligned}
\text{Model 1: } \mu &= \beta_0^{sp} + \beta_P^{sp} P + \beta_{endo}^{sp} \text{endo} + \beta_{herb}^{sp} \text{herb} \\
&\quad + \beta_{endo \times P}^{sp} \text{endo} \cdot P + \beta_{herb \times endo}^{sp} \text{herb} \cdot \text{endo} + \beta_{herb \times P}^{sp} \text{herb} \cdot P \\
&\quad + \beta_{P^2}^{sp} P^2 + \beta_{endo \times P^2}^{sp} \text{endo} \cdot P^2 + \phi^{sp} + \omega^{sp} + \rho^{sp} \\
\text{Model 2: } \mu &= \beta_0^{sp} + \beta_{GD}^{sp} GD + \beta_{endo}^{sp} \text{endo} + \beta_{herb}^{sp} \text{herb} \\
&\quad + \beta_{endo \times GD}^{sp} \text{endo} \cdot GD + \beta_{herb \times endo}^{sp} \text{herb} \cdot \text{endo} + \beta_{herb \times GD}^{sp} \text{herb} \cdot GD \\
&\quad + \phi^{sp} + \omega^{sp} + \rho^{sp} \\
\text{Model 3: } \mu &= \beta_0^{sp} + \beta_{MD}^{sp} MD + \beta_{endo}^{sp} \text{endo} + \beta_{herb}^{sp} \text{herb} \\
&\quad + \beta_{endo \times MD}^{sp} \text{endo} \cdot MD + \beta_{herb \times endo}^{sp} \text{herb} \cdot \text{endo} + \beta_{herb \times MD}^{sp} \text{herb} \cdot MD \\
&\quad + \phi^{sp} + \omega^{sp} + \rho^{sp}
\end{aligned} \tag{1}$$


where  $sp \in \{1, 2, 3\}$  represents the species index,  $P$  denotes Precipitation,  $GD$  refers to distance from geographic center, and  $MD$  stands for mahalanobis distance,  $endo$  indicates Endophyte status (absence or present), and  $herb$  represents Herbivory (fenced and unfenced). The terms  $\phi^{(sp)}$ ,  $\omega^{(sp)}$ , and  $\rho^{(sp)}$  are plot-level, population-level, and site-level random effects, respectively, for species  $sp$ .

Survival was modeled using a Bernoulli distribution, growth with a Gaussian distribution, flowering with a negative binomial distribution, and fertility (number of spikelets) also with a negative binomial distribution. To assess the goodness-of-fit of the models, we performed posterior predictive checks (Berkhof et al., 2000; Gelman et al., 2000). All models effectively captured key aspects of the data, including means, standard deviations, and quantiles (Fig. ??, Fig. ??, Fig. ??, Fig. ??). To determine the best model for each vital rate, we compared the four models using leave-one-out cross-validation (LOOCV) (Vehtari et al., 2017). LOOCV combines validation and training by using one observation for testing and the remaining  $n - 1$  observations for training. This process is repeated  $n$  times once for each observation resulting in  $n$  fitted models (Silva and Zanella, 2024). The test error estimate from LOOCV is obtained by averaging the prediction errors across all  $n$  models (Eq. 2).

$$CV_n = \frac{1}{n} \sum_{i=1}^n (y_i - \hat{y}_i)^2 \tag{2}$$

139 All models were implemented in R (R Core Team, 2023) using Stan via the RStan  
140 interface (Stan Development Team, 2024).





**Figure 2:** XXX




Figure 3: XXX




Figure 4: XXX

## <sup>142</sup> **Acknowledgements**

<sup>143</sup> This research was supported by National Science Foundation Division of Environmental  
<sup>144</sup> Biology awards.

## References

- Afkhami, M. E., McIntyre, P. J., and Strauss, S. Y. (2014). Mutualist-mediated effects on species' range limits across large geographic scales. *Ecology letters*, 17(10):1265–1273.
- Afkhami, M. E. and Rudgers, J. A. (2008). Symbiosis lost: imperfect vertical transmission of fungal endophytes in grasses. *The American Naturalist*, 172(3):405–416.
- Agrawal, A. A., Laforsch, C., and Tollrian, R. (1999). Transgenerational induction of defences in animals and plants. *Nature*, 401(6748):60–63.
- Allsup, C. M., George, I., and Lankau, R. A. (2023). Shifting microbial communities can enhance tree tolerance to changing climates. *Science*, 380(6647):835–840.
- Atala, C., Acuña-Rodríguez, I. S., Torres-Díaz, C., and Molina-Montenegro, M. A. (2022). Fungal endophytes improve the performance of host plants but do not eliminate the growth/defence trade-off. *New Phytologist*, 235(2).
- Bastias, D. A., Martínez-Ghersa, M. A., Ballaré, C. L., and Gundel, P. E. (2017). Epichloë fungal endophytes and plant defenses: not just alkaloids. *Trends in Plant Science*, 22(11):939–948.
- Beaudry, J.-R. (1951). Seed development following the mating *elymus virginicus* l. x *agropyron repens* (l.) beauv. *Genetics*, 36(2):109.
- Bennett, A. E. and Groten, K. (2022). The costs and benefits of plant–arbuscular mycorrhizal fungal interactions. *Annual Review of Plant Biology*, 73(1):649–672.
- Benning, J. W. and Moeller, D. A. (2021a). Microbes, mutualism, and range margins: testing the fitness consequences of soil microbial communities across and beyond a native plant's range. *New Phytologist*, 229(5):2886–2900.
- Benning, J. W. and Moeller, D. A. (2021b). Plant–soil interactions limit lifetime fitness outside a native plant's geographic range margin. *Ecology*, 102(3):e03254.
- Berkhof, J., Van Mechelen, I., and Hoijsink, H. (2000). Posterior predictive checks: Principles and discussion. *Computational Statistics*, 15:337–354.
- Bronstein, J. L. (1994). Conditional outcomes in mutualistic interactions. *Trends in ecology & evolution*, 9(6):214–217.
- Chung, Y. A., Collins, S. L., and Rudgers, J. A. (2019). Connecting plant–soil feedbacks to long-term stability in a desert grassland. *Ecology*, 100(8):e02756.

- Church, G. L. (1958). Artificial hybrids of *elymus virginicus* with *e. canadensis*, *interruptus*, *riparius*, and *wiegandii*. *American Journal of Botany*, pages 410–417.
- Clay, K. and Schardl, C. (2002). Evolutionary origins and ecological consequences of endophyte symbiosis with grasses. *the american naturalist*, 160(S4):S99–S127.
- David, A. S., Quintana-Ascencio, P. F., Menges, E. S., Thapa-Magar, K. B., Afkhami, M. E., and Searcy, C. A. (2019). Soil microbiomes underlie population persistence of an endangered plant species. *The American Naturalist*, 194(4):488–494.
- Donald, M. L., Bohner, T. F., Kolis, K. M., Shadow, R. A., Rudgers, J. A., and Miller, T. E. (2021). Context-dependent variability in the population prevalence and individual fitness effects of plant–fungal symbiosis. *Journal of Ecology*, 109(2):847–859.
- Faeth, S. H. (2009). Asexual fungal symbionts alter reproductive allocation and herbivory over time in their native perennial grass hosts. *The American Naturalist*, 173(5):554–565.
- Fine, P. E. (1975). Vectors and vertical transmission: an epidemiologic perspective.
- Fowler, J. C., Donald, M. L., Bronstein, J. L., and Miller, T. E. (2023). The geographic footprint of mutualism: How mutualists influence species’ range limits. *Ecological Monographs*, 93(1):e1558.
- Gelman, A., Goegebeur, Y., Tuerlinckx, F., and Van Mechelen, I. (2000). Diagnostic checks for discrete data regression models using posterior predictive simulations. *Journal of the Royal Statistical Society: Series C (Applied Statistics)*, 49(2):247–268.
- Giauque, H., Connor, E. W., and Hawkes, C. V. (2019). Endophyte traits relevant to stress tolerance, resource use and habitat of origin predict effects on host plants. *New Phytologist*, 221(4):2239–2249.
- Gundel, P. E., Sun, P., Charlton, N. D., Young, C. A., Miller, T. E., and Rudgers, J. A. (2020). Simulated folivory increases vertical transmission of fungal endophytes that deter herbivores and alter tolerance to herbivory in *poa autumnalis*. *Annals of Botany*, 125(6):981–991.
- Hoeksema, J. D. and Bruna, E. M. (2015). Context-dependent outcomes of mutualistic interactions. *Mutualism*, 10:181–202.
- Kazenel, M. R., Debban, C. L., Ranelli, L., Hendricks, W. Q., Chung, Y. A., Pendergast IV, T. H., Charlton, N. D., Young, C. A., and Rudgers, J. A. (2015). A mutualistic endophyte alters the niche dimensions of its host plant. *AoB plants*, 7:plv005.

- Laughton, A. M., Fan, M. H., and Gerardo, N. M. (2014). The combined effects of bacterial symbionts and aging on life history traits in the pea aphid, *acyrthosiphon pisum*. *Applied and environmental microbiology*, 80(2):470–477.
- Medvigy, D. and Beaulieu, C. (2012). Trends in daily solar radiation and precipitation coefficients of variation since 1984. *Journal of Climate*, 25(4):1330–1339.
- Meshram, S. G., Singh, V. P., and Meshram, C. (2017). Long-term trend and variability of precipitation in chhattisgarh state, india. *Theoretical and Applied Climatology*, 129:729–744.
- Peacher, M. D. and Meiners, S. J. (2020). Inoculum handling alters the strength and direction of plant–microbe interactions.
- R Core Team (2023). *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing, Vienna, Austria.
- Rudgers, J. A., Afkhami, M. E., Bell-Dereske, L., Chung, Y. A., Crawford, K. M., Kivlin, S. N., Mann, M. A., and Nuñez, M. A. (2020). Climate disruption of plant-microbe interactions. *Annual review of ecology, evolution, and systematics*, 51:561–586.
- Rudgers, J. A., Afkhami, M. E., Rúa, M. A., Davitt, A. J., Hammer, S., and Huguet, V. M. (2009). A fungus among us: broad patterns of endophyte distribution in the grasses. *Ecology*, 90(6):1531–1539.
- Rudgers, J. A. and Swafford, A. L. (2009). Benefits of a fungal endophyte in *elymus virginicus* decline under drought stress. *Basic and Applied Ecology*, 10(1):43–51.
- Saikkonen, K., Faeth, S. H., Helander, M., and Sullivan, T. (1998). Fungal endophytes: a continuum of interactions with host plants. *Annual review of Ecology and Systematics*, 29(1):319–343.
- Shaw, R. B. (2011). *Guide to Texas grasses*. Texas A&M University Press.
- Silva, L. A. and Zanella, G. (2024). Robust leave-one-out cross-validation for high-dimensional bayesian models. *Journal of the American Statistical Association*, 119(547):2369–2381.
- Sneck, M. E., Rudgers, J. A., Young, C. A., and Miller, T. E. (2017). Variation in the prevalence and transmission of heritable symbionts across host populations in heterogeneous environments. *Microbial Ecology*, 74:640–653.
- Stan Development Team (2024). RStan: the R interface to Stan. R package version 2.32.6.

- 232 Vega, F. E. (2008). Insect pathology and fungal endophytes. *Journal of invertebrate pathology*,  
233 98(3):277–279.
- 234 Vehtari, A., Gelman, A., and Gabry, J. (2017). Practical bayesian model evaluation using  
235 leave-one-out cross-validation and waic. *Statistics and computing*, 27:1413–1432.
- 236 Yule, K. M., Miller, T. E., and Rudgers, J. A. (2013). Costs, benefits, and loss of vertically  
237 transmitted symbionts affect host population dynamics. *Oikos*, 122(10):1512–1520.



# Supporting Information

## S.1 Supporting methods

### Study design

**Experimental Design** To understand the demographic effects of endophyte symbiosis from core to edge of the host range, we established common gardens at 7 sites across the geographic range of *Elymus virginicus* (Fig. S-1). Experimental sites spanned an aridity gradient (temperature gradient). Common gardens were established in 8 plots per site. Plots were 1.5m \* 1.5m and the area was tilled of existing vegetation to control for native plant competition. Plots were also selected in shaded areas under tree canopy or near shrubs to mimic the natural environmental of the species. In each plot, we planted 15 individuals of *E. virginicus* approximately 15 cm deep in an evenly spaced 4\*4 grid pattern, with positions randomly assigned. For each plot, we randomly assigned a starting endophyte frequency (80%, 60%, 40%, 20%)<sup>1</sup> and herbivory treatment (herbivores exclusion and herbivores accessibility). We ensured that all plots had comparable quantities of source populations. After establishing the plots, we watered the plants and recorded initial tiller counts, flowering status and plot position, endophyte status, source population of each individual plant. For herbivory exclusion plots, we enclosed them with 1.2m tall mesh fencing to prevent browsing by vertebrate herbivores and sprayed the plots with insecticide. For herbivores accessibility plots (control treatment), we half enclosed the plots with the mesh netting.

**Source populations and Identification of individual endophyte status** Plants used in the common garden experiment were derived from natural populations throughout the native range in the south-central US. At each of these natural populations we collected seeds. Some of the seeds of *E. virginicus* were heat treated to produce endophyte negative plants ( $E^-$ ). To do so, we placed these seeds in a drying oven set at 60°C for approximately five days (120 hours). While this method eliminates the endophytes from all individuals, it does not affect seed viability. All seeds (both heat-treated and non-heat-treated) were planted in the Rice University greenhouse. Seedlings were regularly fertilized every two weeks. The seedlings were then vegetatively propagated to produce enough individuals for your experiment ( $N = 840$ ). Before planting in the field, we confirmed the endophyte status ( $E^+$  or  $E^-$ ) of all seedlings using either microscopy or an immunoblot assay. This was necessary due to the varying success of the heat treatment and differences in the prevalence of endophytes between

---

<sup>1</sup>Do we need a schematic of one replicate of the experimental design?

the natural populations. Leaf tissues were stained with aniline blue lactic acid and viewed under a compound microscope at 200x-400x to identify fungal hyphae. The immunoblot assay (Phytoscreen field tiller endophyte detection kit, Agrinostics Ltd. Co.) uses monoclonal antibodies that target proteins of *Epichloë* spp. and chromagen to visually indicate presence or absence. Both methods yield similar detection rates.

## S.2 Supporting Tables


**Table S-1:** Populations and Coordinates

Population	Location	Coordinates	NE+ / NE-
Sam Houston State University Field Station	Huntsville, TX	30°44'30.5"N, 95°28'28.2"W	NE+ = 39, NE- = 50
Texas Tech University	Junction, TX	30°28'18.2"N, 99°47'01.7"W	NE+ = 49, NE- = 83
Palmetto State Park	Palmetto, TX	29.591623, -97.584781	NE+ = 250, NE- = 192
Jean Lafitte National Historical Park and Preserve	Jean Lafitte, LA	29.785105, -90.114933	NE+ = 83, NE- = 94


**Table S-2:** Model comparison by vital rate using ELPD (Expected Log Pointwise Density) differences. The values in the **elpd\_diff** column represent the difference in the model's ELPD relative to the best model for each vital rate, with positive values indicating a worse model fit. The **se\_diff** column shows the standard error of the ELPD difference. Best models for each vital rate are bolded. Model selection is based on the ELPD difference and associated standard error, with smaller ELPD differences and lower standard errors indicating better model fit.

Vital rate	Model	elpd_diff	se_diff
Survival	<b>model1</b>	<b>0.0</b>	<b>0.0</b>
	model2	-1.0	1.5
	model3	-1.0	1.5
Growth	<b>model3</b>	<b>0.0</b>	<b>0.0</b>
	model2	-1.3	2.0
	model1	-2.1	2.2
Flowering	<b>model2</b>	<b>0.0</b>	<b>0.0</b>
	model3	0.0	3.2
	model1	-1.3	3.9


## S.3 Supporting Figures




**Figure S-1:** (A-C) Relationship between distance from geographic center and longitude for the focal species (*Agrostis hyemalis*, *Elymus virginicus*, and *Poa autumnalis*). A generalized additive model (GAM) with a smooth term was used to capture the non-linear trends in these relationships for each species. The solid lines represent the fitted GAM with shaded regions showing the 95% confidence interval. D) displays the relationship between the distance from geographic center and the distance from niche centroid for all species also fitted with a GAM on a log scale. The p-value for the smooth term in the GAM model is  $p=0.0166^{**}$ , indicating a significant relationship between log-transformed geographic distance and Mahalanobis distance.



**Figure S-2:** Coefficient estimates from Bayesian models linking survival and geographic distance. Points represent posterior means and lines indicate 95% credible intervals. Species abbreviations: *Agrostis hyemalis* (AGHY), *Elymus virginicus* (ELVI), and *Poa autumnalis* (PAOU). Horizontal dashed lines denote zero effect.



**Figure S-3:** Coefficient estimates from Bayesian models linking growth and geographic distance. Points represent posterior means and lines indicate 95% credible intervals. Species abbreviations: *Agrostis hyemalis* (AGHY), *Elymus virginicus* (ELVI), and *Poa autumnalis* (PAOU). Horizontal dashed lines denote zero effect.



**Figure S-4:** Coefficient estimates from Bayesian models linking flowering and geographic distance. Points represent posterior means and lines indicate 95% credible intervals. Species abbreviations: *Agrostis hyemalis* (AGHY), *Elymus virginicus* (ELVI), and *Poa autumnalis* (PAOU). Horizontal dashed lines denote zero effect.